

Crops and Soils Research Paper

Cite this article: Cataldo E, Fucile M, Manzi D, Peruzzi E, Mattii GB (2023). Effects of Zeowine and compost on leaf functionality and berry composition in Sangiovese grapevines. *The Journal of Agricultural Science* 1–16. <https://doi.org/10.1017/S002185962300028X>

Received: 24 January 2023
Revised: 12 April 2023
Accepted: 24 April 2023

Keywords:

Clinoptilolite; global warming; leaf temperature; low water availability; photosynthesis; *Vitis vinifera* L; water potential

Corresponding author:

E. Cataldo; E-mail: eleonora.cataldo@unifi.it

Effects of Zeowine and compost on leaf functionality and berry composition in Sangiovese grapevines

E. Cataldo¹ , M. Fucile¹, D. Manzi², E. Peruzzi² and G.B. Mattii¹

¹DAGRI, Department of Agriculture, Food, Environment and Forestry, University of Florence, 50019 Sesto Fiorentino (FI), Italy and ²CNR IRET, Via Moruzzi, 1, 56124 (PI), Italy

Abstract

Meteorological extremes such as heatwaves and water limitations during the ripening season could negatively impact vine ecophysiology and berry metabolism resulting in lower yield per vine. This project aimed to compare two different soil managements during two growing-production seasons (2021 and 2022) with respect to control without any treatment (control). The two managements were: Zeowine (30 t/ha; a soil conditioner made with clinoptilolite and compost proceeding of industrial wine-waste) and compost (20 t/ha). The trial was organized at Col d'Orcia Estate (Montalcino, Tuscan wine region, Italy). The purpose was twofold: (1) to evaluate the effects of Zeowine treatments on leaf gas exchanges, midday stem water potential, chlorophyll fluorescence and leaf temperature (ecophysiology); and (2) to determine any repercussions on the quality of the grapes (technological and phenolics analyses). The parameters plant yield, yeast assimilable nitrogen, fractionation of anthocyanins (cyanidin, delphinidin, malvidin, peonidin and petunidin), caffeic acid, coumaric acid, gallic acid, ferulic acid, kaempferol and quercetin were also analysed. Zeowine showed higher photosynthesis, less negative midday water potential and lower leaf temperature. Essentially, no significant difference was found between the compost and the control. Furthermore, Zeowine grapevines showed higher anthocyanin accumulation and less quercetin content. In general, compost applied together with zeolite could alleviate the adverse effects of water stress and improve plant growth, yield and quality. The control management strategy proved to be the least beneficial for the well-being of the plant and the final quality of the product, confirming the need for amendments in critical years.

Introduction

According to the Intergovernmental Panel on Climate Change, during the 20th century the Earth's surface warmed $\sim 0.75^\circ\text{C}$ (Raymond *et al.*, 2020). The increase of CO_2 as anthropic-origin carbon dioxide emissions is considered the primary starting point of the detected warming (Vaz *et al.*, 2022). Furthermore, the atmospheric CO_2 concentration is expected to continue to increase, resulting in higher Earth surface temperatures over the 21st century (Gutiérrez-Gamboa *et al.*, 2021). The persistent climate shift is having an overwhelming impact on global viticulture. In Europe, from 1950 to 2010, the temperatures during the growing season have increased by 1.7°C (Erlat and Türkeş, 2012). Moreover, by 2050, it is predicted that the temperature range for numerous wine regions will increase by 0.45°C per decade to a total of 2.05°C (Jones *et al.*, 2005).

Warming trends necessarily alter atmospheric composition, the balance of organic matter and the soil water balance (Smith *et al.*, 2008). The quantity of soil organic matter is affected by the inputs of waste production and added soluble organic material with decomposition and leaching as output. Climate alteration influences the input and output too through effects on net primary production by changing decomposition and leaching rates (Tóth *et al.*, 2007).

Ecological drought is defined as 'an episodic deficit in water availability that drives ecosystems beyond thresholds of vulnerability, impacts ecosystem services, and triggers feedback in natural and/or human systems' (Crausbay *et al.*, 2017). This deficit is driven by climate variability processes such as the occurrence of periods with below average precipitation amounts or with increased atmospheric evaporative demand (Christensen *et al.*, 2004; Kumar, 2012). Drought can trigger an alteration in hydrological processes (e.g. percolation, soil infiltration, rainfall interception and runoff) and influence the availability of surface and subsurface water resources (Brown *et al.*, 2005; Haj-Amor and Bouri, 2020). On one hand, climate change is expected to increase the frequency, intensity and duration of heat waves, and on the other hand, it is expected to change the normal distribution of rainfall throughout the year which will aggravate heat-related abiotic stress and drought (Gasparrini *et al.*, 2017; Heo *et al.*, 2019).

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Heat waves were defined as 'periods of at least three consecutive days when the maximum and the minimum temperature were simultaneously greater than their respective 95th percentile in Mediterranean environments: 30.0° and 17.3°C, respectively' (Rey *et al.*, 2007).

Environmental stress factors (high temperature and drought) cause changes in plant metabolism and induce the generation of reactive oxygen species (ROS) (Das and Roychoudhury, 2014). Cellular redox homeostasis is interrupted when ROS generation surpasses cellular scavenging capability (Reddy *et al.*, 2004), resulting in transitory excess of ROS, referred to as oxidative stress (Couée *et al.*, 2006). Under stress conditions, ROS generation leads to increased photorespiration, mitochondrial electron transport activity and fatty acid oxidation (Quan *et al.*, 2008; Sharma *et al.*, 2012). In fact, the closure of stomata because of water stress (diminishing CO₂ concentration in leaf mesophyll tissue) results in an accumulation of nicotinamide adenine dinucleotide phosphate (NADPH) and a decrease in NADP content. Oxygen operates as an alternate acceptor of electrons, forming the superoxide radical (O₂⁻) and H₂O₂, a reduction product of O₂⁻ and the hydroxyl radical (OH), which is produced by the Haber–Weiss reaction (Cadenas, 1989). Active oxygen species can provoke lipid peroxidation and consequentially membrane damage, enzyme inactivation, protein degradation, pigment bleaching and disruption of DNA strands (Imlay and Linn, 1988; Sairam and Saxena, 2000). In field conditions, water deficit is often associated with high irradiance, and it was suggested that high irradiance stress can cause additional damage to the plant, limiting crop productivity (Behera *et al.*, 2002). The relationship between soil water availability, grapevine water stress and stem water potential (or stomatal conductance) was widely described in the literature (Williams and Araujo, 2002; Williams and Baeza, 2007; Suter *et al.*, 2019). In brief, plant water transport follows four steps: (i) soil to root; (ii) root to shoot xylem; (iii) shoot to leaf through the petiole; (iv) leaf to the atmosphere through stomata. General plant water status depends on water potential in soil layers close to the root system, evaporative demand and canopy dimension. Internal vine water deficits fix xylem sap flow to leaf transpiration in relation to soil water availability (Chone *et al.*, 2001).

Drought and high temperatures are perceived as crucial challenges for viticulture, threatening the ascertained connection between the local microclimate, local wine grape cultivars and the representative wine styles within terroirs (Irimia *et al.*, 2018). The impact of this change on the grapevine is important; in fact, research over the last decades has demonstrated modification of vine phenology (Sánchez-Gómez *et al.*, 2020), an enhancement in must alcoholic potential with a decrease in total acidity (Jones and Davis *et al.*, 2000), less predictable dimension and quality of grape yields (Van Leeuwen and Destrac-Irvine, 2017), unbalanced ripening of berries with associated colour and aroma profile alterations (Van Leeuwen and Darriet, 2016; Lu *et al.*, 2022; Torres *et al.*, 2022) and modification of characteristic wine sensory templates (Jones and Alves, 2012; Delrot *et al.*, 2020).

This change could lead to alterations in both the variety of grapes cultivated and the location of suitable viticultural areas, necessitating cooler and higher altitude sites (Moriondo *et al.*, 2013). Alternative solutions to this invasive displacement of wine-growing areas should take the form of sustainable vineyard management strategies also applied systemically throughout a company.

Natural zeolites are mentioned as 'the magic rock' by mineralogists (Eroglu *et al.*, 2017) due to their numerous uses such as

dietary integration in animal feeds (Behin *et al.*, 2019), soil improvers in agronomy (Hamid *et al.*, 2019), insecticide or pesticides used for plant safeguarding (Abdelgaleil *et al.*, 2022; Singh *et al.*, 2022), components in radioactive waste site remediation and decontamination (Gupta *et al.*, 2021) and additives to water and soil for wastewater treatment (Bayuo *et al.*, 2022). Zeolites support agricultural productivity and directly influence the quality of food products (Elemike *et al.*, 2019).

The properties that are structure-related include the high potency of hydration (Depmeier, 2009), extensive porosity (Boettinger and Ming, 2002), low density when dehydrated (Sacerdoti, 2007), crystal structure stability when dehydrated (Di Iorio *et al.*, 2019), cation exchange capacity (Baek *et al.*, 2018), homogenous molecular-sized channels (Li and Pidko, 2019), electrical conductivity (Waqas *et al.*, 2019), adsorption of gases and vapours (Bellat *et al.*, 2019) and catalytic properties (Feliczak-Guzik, 2018).

This experiment was designed to improve the vegetative and productive performance of the plant by adding zeolite to the composted soil. The product called Zeowine was created to interact with the nutritional and water efficiency of the plant. For these reasons, this project aimed to verify if the treatment with zeolite was able to positively influence the ecophysiology of the vine by promoting a greater tolerance to drought and significant light radiation. To achieve these objectives, a comparison was drawn between vines treated with zeolite and compost (Zeowine), vines treated with compost only and untreated vines (control) on Sangiovese cv. in an open field (*Vitis vinifera* L.). This experimentation is linked to the European project ZeoWine (co-financed by the European Commission under the LIFE Program 2014–2020 – Environment and Resources Efficiency). It was specifically chosen to evaluate the effects of the product in two different wine-growing areas in order to exclude the 'terroir effect' (Montalcino and San Miniato). The final results of the vineyard located in San Miniato were published separately to give prominence to both viticultural areas in equal measure (Cataldo *et al.*, 2023). The adopted procedures and tools being, the same throughout the Life Zeowine project, were the same for the two different trials. In addition, the effect of control (control, no treatment applied) treatment alone was also evaluated in the trial in Montalcino (not considered in the other vineyard). Finally, rootstock, clone, vineyard and the management of the vineyard itself are different: organic in Montalcino and biodynamic in the San Miniato Estate.

Materials and methods

Location, experimental project and composting process

The trial was organized at Col d'Orcia Estate (Lat 43°06' N – Long 11°55' E) (CdO), Italy. The estate is located on the southern slope of the Montalcino territory and is an integral part of the Orcia Valley. The Val d'Orcia is a unique territory that was declared part of the Patrimony of Humanity in the year 2004 (UNESCO).

The site benefits from an extremely favourable south-facing position with a protective barrier provided by Mount Amiata (1.750 m) against meteorological events such as floods or hail, and a mild climate influenced by the Tyrrhenian coast in the west, where the sea is some 35 km away. The climate is typically Mediterranean with limited rainfalls concentrated in the months of March, April, November and December.

The experiment was executed on 16-year-old organic vines (*V. vinifera* L., 1753) in two plant cultivation vintages (i.e. 2021

and 2022). The plants taken into consideration are red Sangiovese cultivar (clonal selection SG-CDO-6), on 161-49 C rootstock (*Vitis berlandieri* × *Vitis riparia*). The non-irrigated vineyard is located on a south-facing, moderate longitudinal slope at 6%, with North-South oriented rows and vines trained to upward vertical shoot positioning pruned to spur cordon at a spacing of 2.3 m (inter-row) × 1.0 m (intra-row) for a density of 4350 vines/ha. Each vine had a bud-load of about eight nodes. Standard Tuscan regional protocol for organic viticulture was implemented in all the trial years. The canopy was mechanically trimmed once shoots outgrew the top foliage wire.

From the analysis of the company's soil, a clayey-calcareous soil with the presence of a rocky skeleton emerges (clay 40.5%; sand 27.6%; silt 31.9%; active limestone 173 g/kg; pH 8.3; CSC 27.3 meq/100 g; organic matter 1.7%).

Using a randomized block design with ten replications per treatment (each replication was made by two contiguous inter-rows; on the middle row the measurements were taken from one selected grapevine), the comparison between the control, compost and Zeowine was set up. Ten experimental vines per treatment were then randomly identified and assumed as sub-replicates for the entire trial duration. On test vines, healthy and mature leaves inserted at median shoot level (3–4th node) were chosen for measurements.

Zeowine is a product made by combining the properties of zeolite (clinoptilolite) with the stable organic substance of a compost obtained on a company scale from the reuse of processing waste from grapes, pomace and stalks with the following characteristics: 8.26 pH, 45.9 C mol c/kg CSC, 25.68 C% TOC, 17.35 C/N, 73 mg/kg N-NO₃, 611 mg/kg N-NH₄, 317 mg K/kg available K and 328 mg P/kg available P (Doni *et al.*, 2021).

CdO provided material from grape skins, stalks and vineyard pruning waste, which were shredded to 4–5 cm and processed for their composting. The compost had the following characteristics: 7.37 pH, 36.4 C mol c/kg CSC, 27.01 C% TOC, 21.1 C/N, 196 mg/kg N-NO₃ and 469 mg/kg N-NH₄. The optimal dimensions and typology of the zeolite (Zeocel Italia, PI, Italy) to be used for the production of Zeowine has been selected (85% clinoptilolite) with a granulometry of 0.2–2.5 mm identified in order to ensure better aeration of the heaps during composting. The application of treatments was executed on 1.6 ha of vineyard in production (February 2021 and 2022) with a manure spreader: Zeowine 30 t/ha and compost 20 t/ha (Doni *et al.*, 2021).

The agro-meteorological system Pre-meteo (Mybatec S.R.L., NO, Italy), situated near the vineyard (Montalcino, Italy), gathered the main parameters such as rainfall (mm) and air temperatures (°C).

Leaf gas exchange, chlorophyll fluorescence, water potential (stem) and leaf temperature

Ecophysiological surveys (between 11:10 a.m. and 13:10 p.m.) were conducted on ten replicates per treatment (on tagged vines) every week, from May to harvest: in 2021, 31 May, 15–30 June, 15–30 July, 17–30 August and 9–13 September; in 2022, 31 May, 17 June, 4–25 July, 3–17–26 August and 10–19 September. Data were collected for following parameters: °C (leaf temperature), PN (photosynthesis), gs (stomatal conductance) and E (transpiration), using the Ciras 3PP Systems gas analyser, USA (–400 ppm CO₂, surrounding temperature IR Thermometry, RGBW control 38%, 37%, 25%, 0% and 1300 μmol/m²/s photon flux) (Salvi *et al.*, 2020). Extrinsic water use efficiency (eWUE) was

estimated from the photosynthesis/transpiration ratio (Poni *et al.*, 2014). On the same leaves between 13:15 and 14:15 p.m., stem mid-day water potential (Ψ_{stem}) was evaluated using a Scholander pressure chamber (600-type, PMS Instrument Co, Albany, OR, USA) (Chone *et al.*, 2001). The method consists of increasing the pressure around a leaf petiole until xylem sap appears at the cut end of the petiole, which extends outside the Scholander chamber and is exposed to atmospheric pressure (Boyer, 1967). The surveys were conducted on the tagged vines every week at the beginning of the summer period, from June to harvest: in 2021, 30 June, 17–30 July, 17–30 August and 9–13 September; in 2022, 4–25 July, 3–17–26 August and 10–19 September.

On the same days, chlorophyll fluorescence was assessed with a Handy-PEA fluorometer (Handy-PEA®, UK) on leaves adapted to the dark for ~30 min (Christen *et al.*, 2007).

Technological parameters of berries

In each treatment (Zeowine, compost and control), 100 berries per replication were casually sampled to analyse technological maturity. The sample of 100 berries (ten berries for each tagged vine) was collected from the tagged vines. The berries were sampled from different areas of the bunch: central, upper, lower and lateral parts. Firstly, the berries of each treatment were individually weighed with the Kern PCD model (a precision-digital scale). The sample was crushed to analyse sugar content (° Brix), total acidity (g/l tartaric acid) and pH of the must. The following tools and products were employed for technological analysis: a portable-optical refractometer (RHA-503), a pH meter (HHTEC), bromothymol blue, glass burettes and sodium hydroxide solution (NaOH-0.1 M). In each treatment (Zeowine, compost and control), a further 100 berries per replication were casually sampled to analyse phenolic maturity. Total and extractable anthocyanins were estimated by Glories' method (Kontoudakis *et al.*, 2010). The determination of nine major anthocyanins (cyanidin-3-glucoside, delphinidin-3-glucoside, malvidin-3-acetylglucoside, malvidin-3-cumarylglucoside, malvidin-3-glucoside, peonidin-3-acetylglucoside, peonidin-3-cumarylglucoside, peonidin-3-glucoside and petunidin-3-glucoside) in musts was performed according to OIV-MA-AS315-11: R2007 1 Method OIV-MA-AS315-11 Type II method HPLC-Determination, by an external laboratory (ISVEA), under the analysis conditions proposed by Resolution Oeno 22/2003 and revised in Oeno 12/2007 (OIV, 2021). In addition, with HPLC-HRMS (high-performance liquid chromatography-high resolution mass spectrometry) (Sun *et al.*, 2018) coumaric acid, gallic acid, caffeic acid, ferulic acid, kaempferol-3-O-glucoside, quercetin-3-O-glucoside, quercetin-3-O-rutinoside, quercetin-3-O-galactoside and quercetin-3-O-glucuronide were evaluated. Berry samples were preserved at –80°C until the moment of analysis. The determination of yeast-assimilable nitrogen (as the sum of amino and ammoniacal nitrogen) in musts was performed with an enzymatic-colorimetric kit (Steroglass, Pg, Italy) (Suriñach Ros, 2017).

Finally, on tagged vines, the number of clusters per vine, the weight of bunch per vine and total yield/vine were determined at harvest with a digital scale (VAR model, Italy).

Statistical analysis

Data and graphs were processed with R and RStudio (R Development Core Team) (4.0.3. version) (Tidyverse packages; Lee *et al.*, 2020) with one-way analysis of variance (ANOVA)

($P < 0.05$). Means comparison was performed by the Tukey HSD test (Abdi and Williams, 2010) ($P \leq 0.05$).

Results

Meteorological parameters

The 2021–2022 climate situation of the experimental area is reported in Fig. 1. Daily minimum, average and maximum air temperatures were recorded in both seasons 2021–2022 (from April to September). The 2022 season was more arid and less rainy from April to August than in 2021. The rains in 2022 were mainly concentrated in the month of September. The rainfall summation was: 193.20 mm in April 2021, 116.1 mm in May 2021, 114.0 mm in June 2021, 105.6 mm in July 2021, 131.1 mm in August 2021 and 51.8 mm in September 2021; 54.6 mm in April 2022, 32.3 mm in May 2022, 7.80 mm in June 2022, 46.3 mm in July 2022, 103.7 mm in August 2022 and 112.4 mm in September 2022. In 2022, rainfall was concentrated in the final phase, late August and September. The monthly averages of max temperatures from April to September were: 14.5, 20.6,

28.3, 29.8, 29.4 and 25.4°C (2021); 16.4, 27.6, 30.4, 32.8, 30.1 and 23.8°C (2022). The days where temperatures exceeded 35°C were the following: in 2021, on August 11 at 35.5°C, on August 12 at 35.8°C, on August 13 at 36.2°C, on August 14 at 35.7°C, on August 15 at 37.2°C; in 2022, on July 3 at 37.4°C, on July 4 at 35.5°C, on July 17 at 36.2°C, on July 20 at 36.7°C, on July 23 at 35.3°C, on August 1 at 35.1°C and on August 4 at 35.5°C.

Leaf gas exchange, chlorophyll fluorescence, stem water potential and leaf temperature

The grapevine ecophysiological parameters according to three different land conductions (Zeowine, compost and control) are indicated in Tables 1 and 2, and Figs 2 and 3.

Significant differences in leaf temperatures, eWUE and transpiration were found, particularly during the hottest periods. Essentially no differences were seen between compost and control treatment. During the hottest periods in the 2021 season, the leaf temperature underwent a decrease of 2.66% in Zeowine compared to compost and a decrease of 4.19% compared to control (June 15); a 3.64% decrease in Zeowine *v.* compost and a 4.59% decrease




Figure 1. Meteorological parameters of the experiment location (Col d'Orcia, Montalcino, Italy). Monthly averages of mean, maximum and minimum air temperature (°C) and monthly total precipitation (mm) were measured from April to September (2021–2022 seasons).

Table 1. Ecophysiological parameters – 2021 season

| 2021 | Transpiration (mmol/m ² s) | | | Water use efficiency (μmol/mmol) | | | Leaf temperature (°C) | | | Fluorescence of chlorophyll (Fv/Fm) | | |
|--------------|---------------------------------------|---------------|---------------|----------------------------------|----------------|----------------|-----------------------|----------------|-----------------|-------------------------------------|---------------|---------------|
| | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control |
| 31 May | 1.33 ± 0.28 a | 1.26 ± 0.17 a | 1.24 ± 0.43 a | 7.85 ± 0.98 a | 6.43 ± 1.24 b | 7.42 ± 2.46 ab | 23.51 ± 0.97 a | 24.23 ± 0.73 a | 23.92 ± 0.54 a | 0.82 ± 0.04 a | 0.82 ± 0.01 a | 0.82 ± 0.03 a |
| 15 June | 4.81 ± 1.54 a | 4.83 ± 0.77 a | 4.93 ± 1.26 a | 3.20 ± 1.14 a | 2.12 ± 0.97 b | 1.96 ± 0.73 b | 33.14 ± 1.37 b | 34.02 ± 0.64 a | 34.53 ± 0.87 a | 0.82 ± 0.02 a | 0.80 ± 0.02 a | 0.80 ± 0.02 a |
| 30 June | 5.98 ± 0.98 c | 7.34 ± 0.74 b | 7.91 ± 0.83 a | 1.85 ± 0.44 a | 0.91 ± 0.30 b | 0.79 ± 0.34 b | 31.83 ± 1.31 b | 32.99 ± 0.52 a | 33.29 ± 0.53 a | 0.76 ± 0.03 a | 0.74 ± 0.04 a | 0.74 ± 0.03 a |
| 15 July | 4.26 ± 0.43 b | 4.91 ± 0.39 a | 5.06 ± 1.08 a | 2.80 ± 0.58 a | 2.13 ± 0.48 b | 1.93 ± 0.52 b | 26.50 ± 1.29 b | 28.88 ± 0.90 a | 28.79 ± 1.49 a | 0.71 ± 0.07 a | 0.70 ± 0.07 a | 0.68 ± 0.15 a |
| 30 July | 3.19 ± 0.68 c | 4.98 ± 0.76 a | 4.13 ± 0.71 b | 3.23 ± 0.60 a | 1.64 ± 0.30 b | 2.00 ± 0.53 b | 33.49 ± 0.51 b | 34.05 ± 0.23 a | 34.16 ± 0.56 a | 0.80 ± 0.02 a | 0.77 ± 0.02 b | 0.75 ± 0.03 b |
| 17 August | 2.40 ± 0.56 b | 2.68 ± 0.53 b | 3.43 ± 0.55 a | 4.15 ± 0.67 a | 2.71 ± 0.39 b | 1.97 ± 0.31 c | 32.27 ± 0.86 c | 33.08 ± 0.69 b | 33.88 ± 0.68 a | 0.75 ± 0.03 a | 0.67 ± 0.13 b | 0.68 ± 0.09 b |
| 30 August | 1.90 ± 0.13 a | 1.90 ± 0.26 a | 1.91 ± 0.31 a | 4.39 ± 0.79 a | 3.72 ± 1.08 ab | 3.57 ± 0.76 b | 24.22 ± 0.88 c | 25.19 ± 0.92 b | 25.82 ± 0.64 a | 0.79 ± 0.02 a | 0.70 ± 0.05 b | 0.70 ± 0.06 b |
| 9 September | 2.38 ± 0.34 a | 2.16 ± 0.29 a | 2.08 ± 0.27 a | 2.91 ± 1.01 a | 2.11 ± 1.00 ab | 2.10 ± 1.36 b | 25.34 ± 1.40 b | 26.13 ± 2.56 a | 26.59 ± 1.34 a | 0.79 ± 0.02 a | 0.77 ± 0.03 a | 0.78 ± 0.03 a |
| 13 September | 2.06 ± 0.48 a | 1.77 ± 0.52 a | 1.65 ± 0.56 a | 3.45 ± 0.96 a | 2.84 ± 1.05 b | 2.87 ± 0.77 ab | 24.65 ± 0.34 b | 25.63 ± 1.33 a | 25.22 ± 1.12 ab | 0.73 ± 0.02 a | 0.69 ± 0.05 a | 0.71 ± 0.06 a |

Transpiration (E), extrinsic water use efficiency (eWUE), leaf temperature (°C) and fluorescence of chlorophyll of *Vitis vinifera* with three different soil managements treated. Measurements were conducted from May 2021 to September 2021. Data (mean ± s.e., n = 10) were subjected to one-way ANOVA. Different letters indicate significant differences among Zeowine, compost and control (LSD test, P ≤ 0.05).

Table 2. Ecophysiological parameters – 2022 season

| 2022 | Transpiration (mmol/m ² s) | | | Water use efficiency (μmol/mmol) | | | Leaf temperature (°C) | | | Fluorescence of chlorophyll (Fv/Fm) | | |
|--------------|---------------------------------------|---------------|---------------|----------------------------------|----------------|---------------|-----------------------|---------------|---------------|-------------------------------------|---------------|---------------|
| | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control |
| 31 May | 1.88 ± 0.23 a | 2.01 ± 0.29 a | 2.00 ± 0.32 a | 5.71 ± 1.22 a | 4.95 ± 1.26 a | 5.13 ± 1.77 a | 23.3 ± 0.98 a | 23.6 ± 0.63 a | 23.9 ± 0.23 a | 0.83 ± 0.02 a | 0.82 ± 0.03 a | 0.82 ± 0.03 a |
| 17 June | 3.61 ± 1.21 a | 4.03 ± 1.05 a | 4.16 ± 1.10 a | 3.18 ± 0.32 a | 2.85 ± 0.68 a | 2.36 ± 0.83 a | 29.2 ± 0.89 b | 33.6 ± 0.68 a | 34.0 ± 0.67 a | 0.80 ± 0.03 a | 0.79 ± 0.02 a | 0.78 ± 0.02 a |
| 4 July | 6.28 ± 1.13 b | 8.33 ± 0.90 a | 7.99 ± 1.22 a | 1.45 ± 0.89 a | 0.78 ± 0.63 b | 0.65 ± 0.74 b | 34.0 ± 1.21 b | 38.3 ± 0.43 a | 38.6 ± 0.89 a | 0.77 ± 0.03 a | 0.74 ± 0.03 b | 0.73 ± 0.03 b |
| 25 July | 7.08 ± 0.88 b | 9.23 ± 0.45 a | 9.12 ± 0.54 a | 1.54 ± 0.32 a | 0.78 ± 0.87 b | 0.78 ± 0.56 b | 34.0 ± 1.14 b | 37.4 ± 1.05 a | 37.0 ± 1.25 a | 0.71 ± 0.07 a | 0.68 ± 0.02 b | 0.66 ± 0.04 b |
| 3 August | 4.14 ± 0.66 b | 6.42 ± 0.58 a | 6.96 ± 0.91 a | 2.30 ± 1.21 a | 1.26 ± 0.99 b | 1.07 ± 0.87 b | 33.3 ± 0.62 b | 35.1 ± 0.77 a | 35.2 ± 0.50 a | 0.82 ± 0.01 a | 0.78 ± 0.02 b | 0.76 ± 0.01 b |
| 17 August | 4.04 ± 0.36 a | 4.36 ± 0.67 a | 4.77 ± 0.54 a | 2.33 ± 1.03 a | 1.44 ± 1.08 b | 1.15 ± 0.75 b | 32.1 ± 0.23 a | 33.1 ± 0.44 a | 33.0 ± 0.38 a | 0.81 ± 0.02 a | 0.77 ± 0.04 b | 0.78 ± 0.02 b |
| 26 August | 3.25 ± 0.17 a | 3.12 ± 0.24 a | 3.53 ± 0.22 a | 3.07 ± 1.06 a | 2.20 ± 1.05 ab | 1.37 ± 0.71 b | 25.1 ± 0.95 b | 27.2 ± 0.75 a | 26.9 ± 0.32 a | 0.79 ± 0.02 a | 0.77 ± 0.05 a | 0.78 ± 0.06 a |
| 10 September | 2.27 ± 0.24 a | 2.13 ± 0.27 a | 2.32 ± 0.23 a | 3.67 ± 1.08 a | 3.12 ± 1.02 a | 1.88 ± 1.10 b | 23.2 ± 1.11 b | 25.5 ± 0.54 a | 25.3 ± 1.04 a | 0.80 ± 0.02 a | 0.79 ± 0.01 a | 0.79 ± 0.03 a |
| 19 September | 1.99 ± 0.28 a | 1.87 ± 0.33 a | 2.03 ± 0.44 a | 3.59 ± 1.13 a | 3.74 ± 1.08 a | 3.39 ± 1.00 a | 22.3 ± 0.64 a | 22.8 ± 0.54 a | 22.2 ± 0.67 a | 0.79 ± 0.02 a | 0.77 ± 0.03 a | 0.77 ± 0.04 a |

Transpiration (E), extrinsic water use efficiency (eWUE), leaf temperature (°C) and fluorescence of chlorophyll of *Vitis vinifera* with three different soil managements treated. Measurements were conducted from May 2022 to September 2022. Data (mean ± s.e., n = 10) were subjected to one-way ANOVA. Different letters indicate significant differences among Zeowine, compost and control (LSD test, P ≤ 0.05).

v. control was also demonstrated on June 30 (Table 1). During the hottest periods of the 2022 season, the leaf temperature underwent a decrease of 12.6% in Zeowine compared to compost and a decrease of 13.4% compared to control (July 4); a 10.1% decrease in Zeowine *v.* compost and an 8.79% decrease *v.* control was also demonstrated on July 25 (Table 2).

Significant differences in net photosynthesis and stomatal conductance during seasons were found (Fig. 2). Generally, no differences were ever found between the compost and the control treatments. The Zeowine treatment showed better values than the other two treatments, especially during the hottest periods. In general, the trends of photosynthesis and conductance

reflect the climatic situation showing reductions in the driest periods.

During the moments of decline in water potential, the compost and the control recorded a decrease of 9.49 and 13.3%, respectively, compared to Zeowine (17 August 2021) and on 25 July 2022, there were decreases of 10.4 and 10.7%, respectively (Fig. 3).

Grape composition and production

The treatment had a positive effect on the readily assimilable nitrogen content (Fig. 4). During the harvests (13th and 19th September), Zeowine recorded significantly higher values than




Figure 2. Ecophysiological parameters – 2021/2022 seasons. Net photosynthesis (PN) and stomatal conductance (gs) of *Vitis vinifera* with three different soil managements treated. Measurements were conducted from May to September (2021 and 2022). Data (mean \pm s.e., $n = 10$) were subjected to one-way ANOVA. The bars represent the standard deviation. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).




Figure 3. Ecophysiological parameters – 2021/2022 seasons. Stem water potential (Ψ_{stem}) of *Vitis vinifera* with three different soil managements treated. Measurements were conducted from June to September (2021 and 2022). Data (mean \pm s.e., $n = 10$) were subjected to one-way ANOVA. The bars represent the standard deviation. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).




Figure 4. Yeast assimilable nitrogen (YAN). YAN of *Vitis vinifera* treated with Zeowine, compost and control during two seasons (2021–2022). Measurements were conducted four times: full veraison (17 August 2021 and 17 August 2022), mid maturation (30 August 2021 and 26 August 2022), full maturation (9 September 2021 and 10 September 2022), and harvest (13 September 2021 and 19 September 2022). Data (mean \pm s.e., $n = 10$) were subjected to one-way ANOVA. The bars represent the standard deviation. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).




Figure 5. Technological maturity. Sugar content ($^{\circ}$ Brix), total acidity (TA), pH and berry weight of *Vitis vinifera* treated with Zeowine, compost and control during two seasons (2021–2022). Measurements were conducted four times: full veraison (17 August 2021 and 17 August 2022), mid maturation (30 August 2021 and 26 August 2022), full maturation (9 September 2021 and 10 September 2022) and harvest (13 September 2021 and 19 September 2022). Data (mean \pm s.e., $n = 10$) were subjected to one-way ANOVA. The bars represent the standard deviation. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).

the other treatments. No difference in nitrogen content was noted in relation to vintage.

The zeolitic amendment proved to be effective in improving the technological ripening of the grapes. Significantly higher values were found in the weight of the berry and in the sugar content compared to the other two treatments. No difference was found in both years at the time of harvest in the pH of the must (Fig. 5).

During the anthocyanin accumulation period (from veraison to harvest), the zeolitic amendment showed higher values of total and extractable anthocyanins. In particular, in the harvests (13 and 19 September), the following increases of extractable anthocyanins in Zeowine against compost and control were

recorded: +8.26, +8.52%, and +32.4, +25.7%. Considering the ripening season during 2021, higher values of anthocyanins were recorded in all treatments compared to 2022 (Fig. 6).

No significant differences were found in the percentage of anthocyanins among treatments. caffeic, ferulic and coumaric acids were found in trace amounts or not detected. Zeowine grapes harvested in 2021 revealed significantly lower values of quercetin-3-O-glucoside, quercetin-3-O-galactoside and quercetin-3-O-glucuronide compared to compost and control. Furthermore, the same plants in 2022 detected significantly lower values of quercetin-3-O-glucoside, quercetin-3-O-glucuronide and quercetin-3-O-rutinoside compared to compost and control (Tables 2–4).




Figure 5. Continued.

No differences were recorded in the number of bunches; however, both the production and the weight of the bunch showed higher values in the Zeowine treatment (Fig. 7).

Discussion

In our study Ψ stem, PN and gs were significantly strengthened and improved by zeolite addition in ZEOWINE grapevines with respect to compost and control. The Zeowine treatment positively influenced water stress. In fact, by valorising the soil chemical-physical features, such as potential infiltration, hydraulic conductivity and water-holding skill (Xiubin and Zhanbin, 2001), several authors reported that water stress in plants can be alleviated using

zeolite reinforced with substances of natural origin such as in *Aloe vera* L. (Hazrati *et al.*, 2017), *Oryza sativa* L. (Zheng *et al.*, 2018), *Trigonella foenum-graecum* L. (Baghbani-Arani *et al.*, 2017) and *Malva sylvestris* L. (Ahmadi *et al.*, 2015). These data confirmed what was observed in the parallel experimentation (San Miniato vineyard) (Cataldo *et al.*, 2023); the reproducibility of the treatment effect (Zeowine) in a different terroir is considered fundamental data in the monitoring of a product (experimental treatments may interact with different environmental conditions) (Richter *et al.*, 2009).

It is likely that in the compost and control grapevines, PN, and consequently eWUE, were restricted almost exclusively by impaired photochemistry (i.e. a decline in Fv/Fm ratio) due to




Figure 6. Total and extractable anthocyanins of *Vitis vinifera* treated with Zeowine, compost and control during two seasons (2021–2022). Measurements were conducted four times: full veraison (17 August 2021 and 17 August 2022), mid maturation (30 August 2021 and 26 August 2022), full maturation (9 September 2021 and 10 September 2022) and harvest (13 September 2021 and 19 September 2022). Data (mean \pm s.e., $n = 10$) were subjected to one-way ANOVA. The bars represent the standard deviation. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).

the hottest period during the growing season (Flexas *et al.*, 2004). The need for ‘heat escape’ through transpiration cooling dictated by high-temperature stress (Araújo *et al.*, 2019; Sadok *et al.*, 2021) and the need for water conservation under water-limited

conditions (Ye *et al.*, 2020) was counterbalanced in the compost and control treatments by a tendential increase in transpiration and a reduction in stomatal conductance. On the contrary, the plants treated with Zeowine faced heat stress and water stress in

Table 3. Phenolic maturity

| | 17 August 2021 | | | 30 August 2021 | | | 9 September 2021 | | | 13 September 2021 | | | m.u. |
|-----------------------------|----------------|-----------------|----------------|----------------|----------------|----------------|------------------|----------------|----------------|-------------------|----------------|-----------------|-------|
| | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | |
| Cyanidin-3-glucoside | 28.3 ± 3.15 a | 27.7 ± 2.22 a | 24.6 ± 3.11 a | 18.9 ± 1.32 b | 23.8 ± 2.47 a | 24.4 ± 3.28 a | 22.6 ± 3.25 a | 21.8 ± 2.41 a | 20.9 ± 2.82 a | 17.9 ± 2.10 b | 21.5 ± 2.33 b | 29.6 ± 1.18 a | % |
| Delphinidin-3-glucoside | 12.6 ± 1.24 a | 13.6 ± 1.18 a | 15.0 ± 2.05 a | 17.4 ± 1.10 a | 14.2 ± 2.31 a | 16.6 ± 1.78 a | 13.6 ± 2.62 a | 13.2 ± 3.09 a | 13.1 ± 2.48 a | 14.7 ± 2.56 a | 14.4 ± 2.54 a | 13.0 ± 2.37 a | % |
| Malvidin-3-acetylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | % |
| Malvidin-3-cumarylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 10.60 ± 2.74 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | % |
| Malvidin-3-glucoside | 26.5 ± 3.18 a | 26.7 ± 4.14 a | 28.8 ± 3.50 a | 33.0 ± 3.27 a | 29.6 ± 2.27 a | 27.2 ± 2.41 a | 30.1 ± 1.05 a | 31.6 ± 3.88 a | 32.4 ± 3.26 a | 34.1 ± 3.27 a | 30.8 ± 3.27 a | 24.1 ± 2.89 b | % |
| Peonidin-3-acetylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 0.50 ± 0.11 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | % |
| Peonidin-3-cumarylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | % |
| Peonidin-3-glucoside | 18.9 ± 1.45 a | 17.3 ± 1.77 a | 15.0 ± 1.68 a | 12.0 ± 2.92 a | 16.4 ± 2.84 a | 14.8 ± 2.55 a | 18.5 ± 2.03 a | 18.3 ± 2.12 a | 18.1 ± 2.36 a | 16.6 ± 2.81 a | 17.0 ± 2.27 a | 19.6 ± 2.10 a | % |
| Petunidin-3-glucoside | 13.7 ± 2.36 a | 14.8 ± 3.46 a | 16.5 ± 2.12 a | 18.7 ± 2.23 a | 16.0 ± 2.74 a | 16.9 ± 3.03 a | 15.2 ± 2.59 a | 15.1 ± 2.28 a | 15.5 ± 2.15 a | 16.7 ± 2.30 a | 16.3 ± 2.51 a | 13.7 ± 2.42 a | % |
| Caffeic acid | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | mg/kg |
| Coumaric acid | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | mg/kg |
| Ferulic acid | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | n.d. ± 0.00 a | mg/kg |
| Gallic acid | 3.70 ± 0.88 a | 2.73 ± 0.98 a | 2.76 ± 0.83 a | 3.07 ± 0.63 a | 1.75 ± 0.99 a | 1.60 ± 0.87 a | 1.26 ± 0.28 a | 1.54 ± 1.01 a | 1.88 ± 0.87 a | 1.57 ± 1.13 a | 1.72 ± 0.89 a | 2.38 ± 1.05 a | mg/kg |
| Quercetin-3-O-glucoside | 58.0 ± 4.55 a | 53.3 ± 4.21 a | 41.9 ± 4.90 b | 48.8 ± 4.85 b | 48.1 ± 5.12 b | 67.3 ± 6.21 a | 31.1 ± 3.71 c | 48.7 ± 6.18 b | 74.5 ± 9.12 a | 35.3 ± 3.20 b | 95.6 ± 19.54 a | 100.2 ± 22.88 a | mg/kg |
| Quercetin-3-O-galactoside | 12.0 ± 2.51 a | 10.8 ± 3.11 a | 7.31 ± 2.08 a | 8.91 ± 2.43 a | 8.62 ± 2.58 a | 13.1 ± 3.63 a | 4.29 ± 3.10 b | 10.2 ± 3.03 ab | 15.2 ± 3.62 a | 6.43 ± 1.23 b | 10.5 ± 2.87 b | 24.2 ± 4.65 a | mg/kg |
| Quercetin-3-O-glucuronide | 70.4 ± 8.35 c | 106.9 ± 17.11 a | 87.0 ± 16.42 b | 85.7 ± 9.55 a | 73.2 ± 8.32 b | 73.7 ± 8.38 b | 30.9 ± 6.12 c | 51.6 ± 6.14 b | 69.0 ± 7.12 a | 43.1 ± 5.60 c | 79.5 ± 8.32 b | 88.7 ± 7.47 a | mg/kg |
| Quercetin-3-O-rutinoside | 1.44 ± 1.07 a | 3.72 ± 1.55 a | 2.42 ± 1.23 a | 2.50 ± 1.67 a | 1.69 ± 1.32 a | 1.04 ± 1.17 a | 0.32 ± 0.25 a | 0.63 ± 0.16 a | 1.89 ± 0.34 a | 1.31 ± 0.22 a | 0.69 ± 0.27 a | 1.45 ± 0.59 a | mg/kg |
| Kaempferol-3-O-glucoside | 7.04 ± 1.87 a | 6.09 ± 1.57 a | 4.61 ± 1.68 a | 5.71 ± 1.29 a | 5.04 ± 1.12 a | 7.65 ± 1.99 a | 1.49 ± 1.01 b | 5.53 ± 1.75 ab | 9.31 ± 2.02 a | 14.1 ± 2.54 a | 5.66 ± 3.78 b | 3.25 ± 1.15 b | mg/kg |

Fractionation of anthocyanins (cyanidin-3-glucoside, delphinidin-3-glucoside, malvidin-3-acetylglucoside, malvidin-3-cumarylglucoside, malvidin-3-glucoside, peonidin-3-acetylglucoside, peonidin-3-cumarylglucoside, peonidin-3-glucoside and petunidin-3-glucoside) and coumaric acid, gallic acid, caffeic acid, ferulic acid, kaempferol-3-O-glucoside, quercetin-3-O-glucoside, quercetin-3-O-rutinoside, quercetin-3-O-galactoside and quercetin-3-O-glucuronide of *Vitis vinifera* treated with Zeowine, compost and control during the 2021 season. Measurements were conducted four times: full veraison (17 August 2021), mid maturation (30 August 2021), full maturation (9 September 2021) and harvest (13 September 2021). Data (mean ± s.e., $n = 5$) were subjected to one-way ANOVA. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).

Table 4. Phenolic maturity

| | 18 August 2022 | | | 28 August 2022 | | | 10 September 2022 | | | 15 September 2022 | | | m.u. |
|-----------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-------------------|-----------------|-----------------|-------------------|-----------------|-----------------|-------|
| | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | Zeowine | Compost | Control | |
| Cyanidin-3-glucoside | 18.8 ± 4.87 a | 23.4 ± 4.18 a | 14.8 ± 4.21 b | 19.4 ± 5.23 a | 19.5 ± 5.81 a | 16.3 ± 4.07 a | 22.0 ± 4.23 a | 17.3 ± 2.23 ab | 14.4 ± 3.28 b | 17.4 ± 2.18 a | 18.6 ± 3.63 a | 19.7 ± 3.43 a | % |
| Delphinidin-3-glucoside | 17.1 ± 3.76 a | 17.3 ± 3.82 a | 17.1 ± 3.54 a | 17.6 ± 2.87 a | 17.8 ± 2.45 a | 17.7 ± 2.83 a | 14.7 ± 3.56 a | 15.0 ± 2.45 a | 18.2 ± 3.12 a | 13.3 ± 3.74 a | 16.4 ± 2.21 a | 15.4 ± 2.66 a | % |
| Malvidin-3-acetylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 0.50 ± 0.00 a | 0.40 ± 0.00 a | 0.50 ± 0.00 a | <0.10 ± 0.00 a | 0.50 ± 0.09 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 0.40 ± 0.02 a | % |
| Malvidin-3-cumarylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 0.50 ± 0.02 a | 0.50 ± 0.02 a | 0.50 ± 0.03 a | 0.40 ± 0.04 a | 0.60 ± 0.03 a | 0.40 ± 0.02 a | 0.70 ± 0.04 a | 0.40 ± 0.03 a | 0.50 ± 0.01 a | % |
| Malvidin-3-glucoside | 33.6 ± 6.28 ab | 29.0 ± 5.34 b | 38.2 ± 6.14 a | 32.1 ± 4.23 a | 31.8 ± 5.98 a | 35.1 ± 5.21 a | 31.2 ± 4.87 a | 36.4 ± 4.21 a | 31.6 ± 5.32 a | 37.2 ± 3.37 a | 33.3 ± 4.23 a | 32.5 ± 4.67 a | % |
| Peonidin-3-acetylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | % |
| Peonidin-3-cumarylglucoside | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.10 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | <0.10 ± 0.00 a | 0.40 ± 0.05 a | <0.10 ± 0.00 a | 0.40 ± 0.02 a | % |
| Peonidin-3-glucoside | 12.3 ± 1.78 a | 12.2 ± 1.98 a | 9.90 ± 1.81 a | 11.4 ± 1.33 a | 11.5 ± 1.54 a | 10.7 ± 2.01 a | 15.3 ± 2.24 a | 12.7 ± 2.44 a | 11.2 ± 2.17 a | 14.9 ± 2.87 a | 13.0 ± 2.16 a | 13.7 ± 2.52 a | % |
| Petunidin-3-glucoside | 18.2 ± 2.81 a | 17.6 ± 2.57 a | 20.0 ± 3.04 a | 18.6 ± 4.10 a | 18.6 ± 4.71 a | 19.1 ± 3.07 a | 16.5 ± 2.86 a | 17.5 ± 2.24 a | 19.2 ± 2.67 a | 16.0 ± 2.76 a | 18.3 ± 2.71 a | 17.3 ± 2.87 a | % |
| Caffeic acid | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | n.d. ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | mg/kg |
| Coumaric acid | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | mg/kg |
| Ferulic acid | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | mg/kg |
| Gallic acid | 9.54 ± 2.27 ab | 7.53 ± 1.14 b | 15.2 ± 1.74 a | 18.0 ± 2.20 a | 8.9 ± 1.02 b | 15.2 ± 3.44 ab | 18.1 ± 2.63 a | 21.3 ± 3.55 a | 19.6 ± 2.19 a | 16.5 ± 1.87 b | 21.2 ± 3.87 a | 11.3 ± 2.37 b | mg/kg |
| Quercetin-3-O-glucoside | 89.8 ± 11.37 b | 126.2 ± 27.86 a | 72.4 ± 19.28 c | 105.1 ± 35.17 b | 129.8 ± 20.78 a | 130.9 ± 23.68 a | 157.9 ± 18.78 b | 170.3 ± 21.75 a | 179.0 ± 27.81 a | 118.1 ± 20.57 c | 164.1 ± 21.78 b | 175.8 ± 24.27 a | mg/kg |
| Quercetin-3-O-galactoside | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | <0.05 ± 0.00 a | mg/kg |
| Quercetin-3-O-glucuronide | 178.3 ± 27.24 b | 231.4 ± 20.37 a | 134.2 ± 14.66 c | 183.5 ± 16.42 b | 172.4 ± 17.32 c | 215.6 ± 28.74 a | 163.5 ± 17.47 b | 167.4 ± 16.54 b | 211.6 ± 26.30 a | 100.2 ± 14.57 c | 165.9 ± 18.13 a | 151.8 ± 17.58 b | mg/kg |
| Quercetin-3-O-rutinoside | 11.2 ± 1.53 a | 7.63 ± 1.87 b | 6.90 ± 1.93 b | 9.32 ± 1.73 b | 8.96 ± 1.76 b | 12.3 ± 1.88 a | 2.29 ± 0.34 b | 9.02 ± 1.23 a | 10.5 ± 2.67 a | 2.18 ± 0.23 b | 2.76 ± 1.01 b | 7.59 ± 2.15 a | mg/kg |
| Kaempferol-3-O-glucoside | 26.1 ± 4.42 ab | 32.1 ± 3.25 a | 20.8 ± 2.17 b | 34.2 ± 4.80 a | 28.3 ± 2.55 b | 30.8 ± 4.04 ab | 35.4 ± 2.28 b | 33.3 ± 3.21 b | 46.8 ± 4.27 a | 21.6 ± 3.11 b | 34.4 ± 3.76 a | 30.7 ± 3.27 a | mg/kg |

Fractionation of anthocyanins (cyanidin-3-glucoside, delphinidin-3-glucoside, malvidin-3-acetylglucoside, malvidin-3-cumarylglucoside, malvidin-3-glucoside, peonidin-3-acetylglucoside, peonidin-3-cumarylglucoside, peonidin-3-glucoside and petunidin-3-glucoside) and coumaric acid, gallic acid, caffeic acid, ferulic acid, kaempferol-3-O-glucoside, quercetin-3-O-glucoside, quercetin-3-O-rutinoside, quercetin-3-O-galactoside and quercetin-3-O-glucuronide of *Vitis vinifera* treated with Zeowine, compost and control during the 2022 season. Measurements were conducted four times: full veraison (17 August 2022), mid maturation (26 August 2022), full maturation (10 September 2022) and harvest (19 September 2022). Data (mean ± s.e., n = 5) were subjected to one-way ANOVA. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).




Figure 7. Production. Cluster weight, yield per vine and number of cluster per vine of *Vitis vinifera* treated with Zeowine, compost and control during the 2021–2022 seasons. Measurements were conducted at harvest (13 September 2021 and 19 September 2022). Data (mean \pm s.e., $n=5$) were subjected to one-way ANOVA. Different letters indicate significant differences among Zeowine, compost and control (LSD test, $P \leq 0.05$).

a more efficient way than the others, showing a lower leaf temperature and less negative water potential. In addition, we hypothesize that the zeolite addition to the soil made the roots

more humid in order to protect the leaves from photo-oxidative impairment (Wu *et al.*, 2019).

As reported in many studies, technological maturity was influenced by water and temperature stress (Zufferey *et al.*, 2017; Salvi *et al.*, 2020; Cataldo *et al.*, 2023). The compost and control treatments reported a more protracted ripening than Zeowine; in fact, a lower sugar content and a lower berry weight were found. According to Wang *et al.* (2003) water deficiency hampers sugar unloading in the berries. Conversely, Zeowine grapevines demonstrated a greater berry weight and higher production with increased cluster weight, confirming the positive impact of these particular aluminosilicates of alkaline and alkaline earth element on yield (Hazrati *et al.*, 2017). Absorption and controlled release of moisture by zeolite improves plant growth and yield under drought conditions (AL-Busaidi *et al.*, 2011). It has also been noted that zeolites can be effectively used in agriculture to encourage water infiltration and retention in the soil due to their porous and capillary skills that act as a slow-release source for water as well as macro or microelements (Rastogi *et al.*, 2019). Moreover, it was found that zeolite is capable of slowing urea release and, therefore, has the potential to be developed as controlling agent for the release of nitrogen from urea (Hidayat *et al.*, 2018). This considered, in addition to improving the administration of water resources, it was suggested that Zeowine increased the substrate cation exchange capacity which made nutrients available in the compost gradually and avoided losses due to leaching, affecting the final production (Confalonieri *et al.*, 2021; Zijun *et al.*, 2021). The pH parameter, however, was not influenced by treatment.

Regarding the anthocyanins, a general reduction in the content of totals and extractables was observed in all treatments in the 2022 season; this could be attributed to the scorching temperatures recorded in the moment of greatest synthesis (beginning of veraison – 1 August 2022, 35.1°C was enrolled) (Pirie and Mullins, 1977).

In extractable anthocyanins at the time of the two harvests, the following increases were recorded for treatment with Zeowine compared to compost and control, respectively: +7.68 and +7.87% on 13 September 2021; +24.5 and +20.5% on 13 September 2022. The increase in anthocyanins in the treatment with zeolites could in both years be attributed to the beneficial effect of this amendment on ripening. In fact, the treatment allowed a balanced and protected maturation from severe water deficit and intense thermal stress avoiding irreparable damage to the photosystem II (Shamili *et al.*, 2020) for correct storage of soluble solids. Furthermore, by improving the performance of the canopy, it improved thermal conditions by avoiding excessive increases in leaf temperature (probably verifiable also at cluster level).

Of the flavonols measured, as 3,5,7-trihydroxylated derivatives (hydroxylated in C3, C5 and C7), the glycosides from the following aglycons were identified: quercetin (3',4'-diOH) and kaempferol (4'-OH) (Castillo-Muñoz *et al.*, 2007), whereas the corresponding 3-O-galactoside derivatives were found to be minor compounds (Garrido and Borges, 2013). The highest concentrations of quercetin in grapes were found in the compost and control treatments. Quercetin-3-O-glucoside and kaempferol-3-O-glucoside have been found to be UV-B stress indicators in *V. vinifera* leaves in Pinot Noir and Riesling cultivars (Schoedl *et al.*, 2013; Boudérias *et al.*, 2020).

Therefore, it is believed that further research should be conducted, particularly to investigate the activity of anthocyanins

accumulated in the berry and the activity of phenylalanine as a precursor of various phenolic compounds.

Conclusions

In order to decrease the environmental impact of agriculture, the efficiency of fertilizer use and water use must be maximized. The use of zeolite is key to improving plant water holding capacity which minimizes the requirement for vine irrigation, as water is well retained within zeolite's structure. This experiment supplies new evidence that zeolite applications could impact both physiological profiles and berry skin metabolism (sugar and size) of vines, providing an improved ability to counteract low water availability during the season. These findings support the need for consequent further investigations to be carried out. Primarily, research activities should consider the sugars accumulated in the vacuole (activity of sucrose-metabolizing enzymes, sucrose transporters and monosaccharide transporters).

Acknowledgements. The authors acknowledge Col d'Orcia (CdO) winery. The experiment took place within the context of the LIFE EU project LIFE ZEOWINE LIFE17 ENV/IT/000427 'ZEOLite and WINery waste as an innovative product for wine production'.

Author contributions. The design was elaborated and planned by E. C., D. M., E. P. and G. B. M. In addition, E. C. and M. F. carried out plant and berries measurements and analyses. The analyses on the anthocyanin fractionation were made by an external analysis laboratory (ISVEA). E. P. and D. M. performed the analysis on the characterization of the product. E. C. carried out data and statistical analyses. Finally, E. C. wrote the draft text, which was supervised by G. B. M. All authors read and accepted this manuscript.

Financial support. This research was funded by the LIFE EU project LIFE ZEOWINE LIFE17 ENV/IT/000427 'ZEOLite and WINery waste as an innovative product for wine production'. The Life ZeoWine Project is co-funded with the contribution of the European Commission, under the LIFE Programme – Environment & Resources efficiency – Grant Agreement n. LIFE17 ENV/IT/000427.

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Abdelgaleil SA, Gad HA, Atta AA and Al-Anany MS (2022) Control of *Sitophilus granarius* and *Sitophilus oryzae* on stored wheat using low-rate combinations of natural zeolite with three insecticides. *Journal of Stored Products Research* **97**, 101975.
- Abdi H and Williams LJ (2010) Tukey's honestly significant difference (HSD) test. *Encyclopedia of Research Design* **3**, 1–5.
- Ahmadi Azar F, Hasanloo T and Feizi V (2015) Water stress and mineral zeolite application on growth and some physiological characteristics of Mallow (*Malva sylvestris*). *Journal of Plant Research (Iranian Journal of Biology)* **28**, 459–474.
- AL-Busaidi A, Yamamoto T, Tanigawa T and Rahman HA (2011) Use of zeolite to alleviate water stress on subsurface drip irrigated barley under hot environments. *Irrigation and Drainage* **60**, 473–480.
- Araújo M, Ferreira de Oliveira JMP, Santos C, Moutinho-Pereira J, Correia C and Dias MC (2019) Responses of olive plants exposed to different irrigation treatments in combination with heat shock: physiological and molecular mechanisms during exposure and recovery. *Planta* **249**, 1583–1598.
- Baek W, Ha S, Hong S, Kim S and Kim Y (2018) Cation exchange of cesium and cation selectivity of natural zeolites: chabazite, stilbite, and heulandite. *Microporous and Mesoporous Materials* **264**, 159–166.
- Baghbani-Arani A, Modarres-Sanavy SAM, Mashhadi-Akbar-Boojar M and Mokhtassi-Bidgoli A (2017) Towards improving the agronomic performance, chlorophyll fluorescence parameters and pigments in fenu-greek using zeolite and vermicompost under deficit water stress. *Industrial Crops and Products* **109**, 346–357.
- Bayuo J, Rwiza M and Mtei KA (2022) Comprehensive review on the decontamination of lead (II) from water and wastewater by low-cost biosorbents. *RSC Advances* **12**, 11233–11254.
- Behera RK, Mishra PC and Choudhury NK (2002) High irradiance and water stress induce alterations in pigment composition and chloroplast activities of primary wheat leaves. *Journal of Plant Physiology* **159**, 967–973.
- Behin J, Ghadamnan E and Kazemian H (2019) Recent advances in the science and technology of natural zeolites in Iran. *Clay Minerals* **54**, 131–144.
- Bellat JP, Weber G, Bezverkhy I and Lamonier JF (2019) Selective adsorption of formaldehyde and water vapors in NaY and NaX zeolites. *Microporous and Mesoporous Materials* **288**, 109563.
- Boettinger JL and Ming DW (2002) Zeolites. *Soil Mineralogy with Environmental Applications* **7**, 585–610.
- Bouderias S, Teszlák P, Jakab G and Kőrösi L (2020) Age- and season-dependent pattern of flavonol glycosides in Caber-net Sauvignon grapevine leaves. *Scientific Reports* **10**, 1–9.
- Boyer J (1967) Leaf water potentials measured with a pressure chamber. *Plant Physiology* **42**, 133–137.
- Brown AE, Zhang L, McMahon TA, Western AW and Vertessy RAA (2005) Review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation. *Journal of Hydrology* **310**, 28–61.
- Cadenas E (1989) Biochemistry of oxygen toxicity. *Annual Review of Biochemistry* **58**, 79–110.
- Castillo-Muñoz N, Gómez-Alonso S, García-Romero E and Hermosín-Gutiérrez I (2007) Flavonol profiles of *Vitis vinifera* red grapes and their single-cultivar wines. *Journal of Agricultural and Food Chemistry* **55**, 992–1002.
- Cataldo E, Fucile M, Manzi D, Masini CM, Doni S and Mattii GB (2023) Sustainable soil management: effects of clinoptilolite and organic compost soil application on eco-physiology, quercetin, and hydroxylated, methoxylated anthocyanins on *Vitis vinifera*. *Plants*, **12**, 708. <https://doi.org/10.3390/plants12040708>.
- Chone X, Van Leeuwen C, Dubourdieu D and Gaudillère JP (2001) Stem water potential is a sensitive indicator of grapevine water status. *Annals of Botany* **87**, 477–483.
- Christen D, Schönmann S, Jermini M, Strasser RJ and Défago G (2007) Characterization and early detection of grapevine (*Vitis vinifera*) stress responses to esca disease by *in situ* chlorophyll fluorescence and comparison with drought stress. *Environmental and Experimental Botany* **60**, 504–514.
- Christensen NS, Wood AW, Voisin N, Lettenmaier DP and Palmer RN (2004) The effects of climate change on the hydrology and water resources of the Colorado River basin. *Climatic Change* **62**, 337–363.
- Confalonieri G, Vezzalini G, Quattrini F, Quartieri S, Dejoie C and Arletti R (2021) Ce-exchange capacity of zeolite L in different cationic forms: a structural investigation. *Journal of Applied Crystallography* **54**, 1766–1774.
- Couée J, Sulmon C, Gouesbet G and El Amrani A (2006) Involvement of soluble sugars in reactive oxygen species balance and responses to oxidative stress in plants. *Journal of Experimental Botany* **57**, 449–459.
- Crausbay SD, Ramirez AR, Carter SL, Cross MS, Hall KR, Bathke DJ, Betancourt JL, Colt S, Cravens AE, Dalton MS, Dunham JB, Hay LE, Hayes MJ, McEvoy J, McNutt CA, Moritz MA, Nislow KH, Raheem N and Sanford T (2017) Defining ecological drought for the twenty-first century. *Bulletin of the American Meteorological Society* **98**, 2543–2550.
- Das K and Roychoudhury A (2014) Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in Environmental Science* **2**, 53.
- Delrot S, Grimplet J, Carbonell-Bejerano P, Schwandner A, Bert PF, Bavaresco L, Dalla Costa L, Di Gasparo G, Duchene E, Hausmann L, Malnoy M, Morgante M, Ollat N, Pecile M and Vezzulli S (2020) Genetic and genomic approaches for adaptation of grapevine to climate change. In Kole C (ed.), *Genomic Designing of Climate-Smart Fruit Crops*. Cham: Springer, pp. 157–270.
- Depmeier W (2009) Some examples of temperature and time resolved studies of the dehydration and hydration behavior of zeolites and clathrates. *Particle & Particle Systems Characterization* **26**, 138–150.

- Di Iorio JR, Hoffman AJ, Nimlos CT, Nystrom S, Hibbitts D and Gounder R** (2019) Mechanistic origins of the high-pressure inhibition of methanol dehydration rates in small-pore acidic zeolites. *Journal of Catalysis* **380**, 161–177.
- Doni S, Gispert M, Peruzzi E, Macci C, Mattii GB, Manzi D, Masini CM and Grazia M** (2021) Impact of natural zeolite on chemical and biochemical properties of vineyard soils. *Soil Use and Management* **37**, 832–842.
- Elemike EE, Uzoh IM, Onwuodiwe DC and Babalola OO** (2019) The role of nanotechnology in the fortification of plant nutrients and improvement of crop production. *Applied Sciences* **9**, 499.
- Erlat E and Türkerş M** (2012) Analysis of observed variability and trends in numbers of frost days in Turkey for the period 1950–2010. *International Journal of Climatology* **32**, 1889–1898.
- Eroglu N, Emekci M and Athanassiou CG** (2017) Applications of natural zeolites on agriculture and food production. *Journal of the Science of Food and Agriculture* **97**, 3487–3499.
- Feliczak-Guzik A** (2018) Hierarchical zeolites: synthesis and catalytic properties. *Microporous and Mesoporous Materials* **259**, 33–45.
- Flexas J, Bota J, Cifre J, Mariano Escalona J, Galmés J, Guliás J, Lefi El-K, Martínez-Cannellas SF, Moreno MT, Ribas-Carbo M, Riera D, Sampol B and Medrano H** (2004) Understanding down-regulation of photosynthesis under water stress: future prospects and searching for physiological tools for irrigation management. *Annals of Applied Biology* **144**, 273–283.
- Garrido J and Borges F** (2013) Wine and grape polyphenols – a chemical perspective. *Food Research International* **54**, 1844–1858.
- Gasparrini A, Guo Y, Sera F, Vicedo-Cabrera AM, Huber V, Tong S, Coelho S, Saldiva PHN, Lavigne E, Correa PM, Ortega NV, Kan H, Osorio S, Kysely J, Urban A, Jaakkola JJK, Ryti NRI, Pascal M, Goodman PG, Zeka A and Armstrong B** (2017) Projections of temperature-related excess mortality under climate change scenarios. *The Lancet Planetary Health* **1**, e360–e367.
- Gupta A, Sharma V, Sharma K, Kumar V, Choudhary S, Mankotia P, Kumar B, Mishra H, Moullick A, Ekielski A and Mishra PK** (2021) A review of adsorbents for heavy metal decontamination: growing approach to wastewater treatment. *Materials* **14**, 4702.
- Gutiérrez-Gamboa G, Zheng W and Martínez de Toda F** (2021) Strategies in vineyard establishment to face global warming in viticulture: a mini review. *Journal of the Science of Food and Agriculture* **101**, 1261–1269.
- Haj-Amor Z and Bouri S** (2020) *Climate Change Impacts on Coastal Soil and Water Management*. Boca Raton, FL: CRC Press.
- Hamid Y, Tang L, Sohail MI, Cao X, Hussain B, Aziz MZ, Usman M, He Z and Yang X** (2019) An explanation of soil amendments to reduce cadmium phytoavailability and transfer to food chain. *Science of the Total Environment* **660**, 80–96.
- Hazrati S, Tahmasebi-Sarvestani Z, Mokhtassi-Bidgoli A, Modarres-Sanavy SAM, Mohammadi H and Nicola S** (2017) Effects of zeolite and water stress on growth, yield and chemical compositions of *Aloe vera* L. *Agricultural Water Management* **181**, 66–72.
- Heo S, Bell ML and Lee JT** (2019) Comparison of health risks by heat wave definition: applicability of wet-bulb globe temperature for heat wave criteria. *Environmental Research* **168**, 158–170.
- Hidayat A, Mukti NIF, Handoko B and Sutrisno B** (2018) Biodiesel production from rice bran oil over modified natural zeolite catalyst. *International Journal of Technology* **9**, 400–411.
- Imlay JA and Linn S** (1988) DNA damage and oxygen radical toxicity. *Science* **240**, 1302–1309.
- Irimia LM, Patriche CV, Quenol H, Sfić L and Foss C** (2018) Shifts in climate suitability for wine production as a result of climate change in a temperate climate wine region of Romania. *Theoretical and Applied Climatology* **131**, 1069–1081.
- Jones GV and Alves F** (2012) Impact of climate change on wine production: a global overview and regional assessment in the Douro Valley of Portugal. *International Journal of Global Warming* **4**, 383–406.
- Jones GV and Davis RE** (2000) Climate influences on grapevine phenology, grape composition, and wine production and quality for Bordeaux, France. *American Journal of Enology and Viticulture* **51**, 249–261.
- Jones GV, White MA, Cooper OR and Storchmann K** (2005) Climate change and global wine quality. *Climatic Change* **73**, 319–343.
- Kontoudakis N, Esteruelas M, Fort F, Canals JM and Zamora F** (2010) Comparison of methods for estimating phenolic maturity in grapes: correlation between predicted and obtained parameters. *Analytica Chimica Acta* **660**, 127–133.
- Kumar CP** (2012) Climate change and its impact on groundwater resources. *International Journal of Engineering and Science* **1**, 43–60.
- Lee S, Sriutaisuk S and Kim H** (2020) Using the Tidyverse Package in R for simulation studies in SEM. *Structural Equation Modeling: A Multidisciplinary Journal* **27**, 468–482.
- Li G and Pidko EA** (2019) The nature and catalytic function of cation sites in zeolites: a computational perspective. *ChemCatChem* **11**, 134–156.
- Lu HC, Hu L, Liu Y, Cheng CF, Chen W, He F, Duan CQ and Wang J** (2022) Reducing the source/sink ratio of grapevine to face global warming in a semi-arid climate: effects on volatile composition of Cabernet Sauvignon grapes and wines. *Food Chemistry* **15**, 100449.
- Moriondo M, Jones GV, Bois B, Dibari C, Ferrise R, Trombi G and Bindu M** (2013) Projected shifts of wine regions in response to climate change. *Climatic Change* **119**, 825–839.
- OIV** (2021) Official methods for the analysis of musts and wines of the international organization of vine and wine (OIV). Methods of analysis of wines and musts. (OIV-MA-INT-00-2012). <http://www.oiv.int/oiv/info/enmethodesinternationalesvin> (Accessed 15 December 2022).
- Pirie A and Mullins MG** (1977) Interrelationships of sugars, anthocyanins, total phenols and dry weight in the skin of grape berries during ripening. *American Journal of Enology and Viticulture* **28**, 204–209.
- Poni S, Merli MC, Magnanini E, Galbignani M, Bernizzoni F, Vercesi A and Gatti M** (2014) An improved multichamber gas exchange system for determining whole-canopy water-use efficiency in grapevine. *American Journal of Enology and Viticulture* **65**, 268–276.
- Quan LJ, Zhang B, Shi WW and Li HY** (2008) Hydrogen peroxide in plants: a versatile molecule of the reactive ox-species network. *Journal of Integrative Plant Biology* **50**, 2–18.
- Rastogi A, Tripathi DK, Yadav S, Chauhan DK, Živčák M, Ghorbanpour M, El-Sheer NI and Brestic M** (2019) Application of silicon nanoparticles in agriculture. *3 Biotech* **9**, 1–11.
- Raymond C, Matthews T and Horton RM** (2020) The emergence of heat and humidity too severe for human tolerance. *Science Advances* **6**, eaaw1838.
- Reddy AR, Chaitanya KV and Vivekanandan M** (2004) Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology* **161**, 1189–1202.
- Rey G, Jougle E, Fouillet A, Pavillon G, Bessemoulin P, Frayssinet P, Clavel J and Hémon D** (2007) The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *International Archives of Occupational and Environmental Health* **80**, 615–626.
- Richter SH, Garner JP and Würbel H** (2009) Environmental standardization: cure or cause of poor reproducibility in animal experiments? *Nature Methods* **6**, 257–261.
- Sacerdoti M** (2007) The crystal structure of zeolite barrerite dehydrated in air at 400–450 °C. *Microporous and Mesoporous Materials* **102**, 299–303.
- Sadok W, Lopez JR and Smith KP** (2021) Transpiration increases under high-temperature stress: potential mechanisms, trade-offs and prospects for crop resilience in a warming world. *Plant, Cell & Environment* **44**, 2102–2116.
- Sairam RK and Saxena DC** (2000) Oxidative stress and antioxidants in wheat genotypes: possible mechanism of water stress tolerance. *Journal of Agronomy and Crop Science* **184**, 55–61.
- Salvi L, Brunetti C, Cataldo E, Storchi P and Mattii GB** (2020) Eco-physiological traits and phenylpropanoid profiling on potted *Vitis vinifera* L. cv Pinot noir subjected to *Ascohyllum nodosum* treatments under post-veraison low water availability. *Applied Sciences* **10**, 4473.
- Sánchez-Gómez R, Pérez-Álvarez EP, Salinas R, Gonzalo-Diago A, Zalacain A and Garde-Cerdán T** (2020) Effect of vine-shoot and oak extract foliar grapevine applications on oenological parameters, phenolic acids and glutathione content of white musts and wines. *Oeno One* **54**, 145–156.
- Schoedl K, Schuhmacher R and Forneck A** (2013) Correlating physiological parameters with biomarkers for UV-B stress indicators in leaves of grapevine cultivars Pinot noir and Riesling. *The Journal of Agricultural Science* **151**, 189–200.

- Shamili M, Dehghanpour S and Atrash S** (2020) Zeolite alleviates defense responses in drought stressed carrot (*Daucus carota* L.). *Journal of Plant Process and Function* **9**, 27–36.
- Sharma P, Jha AB, Dubey RS and Pessaraki M** (2012) Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany* **2012**, 217037.
- Singh G, Ramadass K, Sooriyakumar P, Hettithanthri O, Vithange M, Bolan N, Tavakkoli E, Van Zeowineieten L and Vinu A** (2022) Nanoporous materials for pesticide formulation and delivery in the agricultural sector. *Journal of Controlled Release* **343**, 187–206.
- Smith P, Fang C, Dawson JJ and Moncrieff JB** (2008) Impact of global warming on soil organic carbon. *Advances in Agronomy* **97**, 1–43.
- Sun J, Liu W, Zhang M, Geng P, Shan Y, Li G, Zhao Y and Chen P** (2018) The analysis of phenolic compounds in daylily using UHPLC-HRMSn and evaluation of drying processing method by fingerprinting and metabolomic approaches. *Journal of Food Processing and Preservation* **42**, e13325.
- Suriñach Ros A** (2017) Method validation of enzymatic ammonia method and colorimetric free amino nitrogen method applied to wine industry.
- Suter B, Triolo R, Pernet D, Dai Z and Van Leeuwen C** (2019) Modeling stem water potential by separating the effects of soil water availability and climatic conditions on water status in grapevine (*Vitis vinifera* L.). *Frontiers in Plant Science* **10**, 1485.
- Torres N, Yu R, Martinez-Luscher J, Girardello RC, Kostaki E, Oberholster A and Kurtural SK** (2022) Shifts in the phenolic composition and aromatic profiles of Cabernet Sauvignon (*Vitis vinifera* L.) wines are driven by different irrigation amounts in a hot climate. *Food Chemistry* **371**, 131163.
- Tóth JA, Lajtha K, Kotrocó Z, Krakomperger Z, Caldwell B, Bowden R and Papp M** (2007) The effect of climate change on soil organic matter decomposition. *Acta Silvatica et Lignaria Hungarica* **3**, 75–85.
- Van Leeuwen C and Darriet P** (2016) The impact of climate change on viticulture and wine quality. *Journal of Wine Economics* **11**, 150–167.
- Van Leeuwen C and Destrac-Irvine A** (2017) Modified grape composition under climate change conditions requires adaptations in the vineyard. *Oeno One* **51**, 147–154.
- Vaz S Jr, de Souza APR and Baeta BEL** (2022) Technologies for carbon dioxide capture: a review applied to energy sectors. *Cleaner Engineering and Technology* **8**, 100456.
- Wang ZP, Deloire A, Carbonneau A, Federspiel B and Lopez F** (2003) An *in vivo* experimental system to study sugar phloem unloading in ripening grape berries during water deficiency stress. *Annals of Botany* **92**, 523–528.
- Waqas M, Nizami AS, Aburiazaiza AS, Barakat MA, Asam ZZ, Khattak B and Rashid MI** (2019) Untapped potential of zeolites in optimization of food waste composting. *Journal of Environmental Management* **241**, 99–112.
- Williams LE and Araujo FJ** (2002) Correlations among predawn leaf, midday leaf, and midday stem water potential and their correlations with other measures of soil and plant water status in *Vitis vinifera*. *Journal of the American Society for Horticultural Science* **127**, 448–454.
- Williams LE and Baeza P** (2007) Relationships among ambient temperature and vapor pressure deficit and leaf and stem water potentials of fully irrigated, field-grown grapevines. *American Journal of Enology and Viticulture* **58**, 173–181.
- Wu Q, Chi D, Xia G, Chen T, Sun Y and Song Y** (2019) Effects of zeolite on drought resistance and water–nitrogen use efficiency in paddy rice. *Journal of Irrigation and Drainage Engineering* **145**, 04019024.
- Xiubin H and Zhanbin H** (2001) Zeolite application for enhancing water infiltration and retention in loess soil. *Resources, Conservation and Recycling* **34**, 45–52.
- Ye H, Song L, Schapaugh WT, Ali ML, Sinclair TR, Riar MK, Mutava RN, Li Y, Vuong T, Valliyodan B, Neto AP, Klepadlo M, Song Q, Shannon JG, Chen P and Nguyen HT** (2020) The importance of slow canopy wilting in drought tolerance in soybean. *Journal of Experimental Botany* **71**, 642–652.
- Zheng J, Chen T, Wu Q, Yu J, Chen W, Chen Y, Siddique KHM, Meng W, Chi D and Xia G** (2018) Effect of zeolite application on phenology, grain yield and grain quality in rice under water stress. *Agricultural Water Management* **206**, 241–251.
- Zijun Z, Effenev G, Millar GJ and Stephen M** (2021) Synthesis and cation exchange capacity of zeolite W from ultra-fine natural zeolite waste. *Environmental Technology & Innovation* **23**, 101595.
- Zufferey V, Spring JL, Verdenal T, Dienes A, Belcher S, Lorenzini F, Koestel C, Rosti J, Gindro K, Spangenberg J and Viret O** (2017) The influence of water stress on plant hydraulics, gas exchange, berry composition and quality of Pinot Noir wines in Switzerland. *Oeno One* **51**. <https://doi.org/10.20870/oenoone.2017.51.1.1314>