

CRACK NUCLEATION IN SHELLS WITH THROUGH-THE-THICKNESS MICROSTRUCTURE*

PAOLO MARIA MARIANO[†] AND DOMENICO MUCCI[‡]

Abstract. We refer to the common low-dimensional description of shells and thin films: surfaces endowed with out-of-surface directors satisfying a nondegeneracy condition under large strain. We consider in addition through-the-thickness material microstructure described by elements of a complete and intrinsic Riemannian manifold. We look at brittle materials. Among all possible cracked and uncracked admissible configurations, the one obtained under Dirichlet-type boundary conditions realizes the minimum of a regularized Griffith’s energy that includes curvature terms. For it, we prove the existence of minimizers for different constitutive functional choices and geometric structures with or without active through-the-thickness microstructure. Deformations are taken as *SBV* maps with a jump set included in the support of a curvature varifold with boundary. Through-the-thickness descriptors of the material microstructure are taken first as manifold-valued Sobolev maps. Then we consider pertinent *SBV* versions.

Key words. fracture, varifolds, ground states, shells, microstructures, calculus of variations

MSC codes. 74K25, 49J45, 49Q15, 49Q20, 74R05, 74E99

DOI. 10.1137/22M1490004

1. Introduction. Three topics in continuum mechanics interact in the analysis that we propose here:

- the representation of films such as thin shells modeled as surfaces endowed with directors;
- the variational description of crack nucleation;
- the multifield (at times called the phase field to adopt a common terminology in solid-state physics, although with more restricted meaning) depiction of microstructural effects on macroscopic motion.

Pertinent geometric and functional elements enter the energy. A requirement of minimality selects equilibrium configurations among those with all possible crack paths, depicted as rectifiable subsets of a domain Ω describing the shell middle surface. Fields describing macroscopic and microstructural through-the-thickness morphology and crack path are pertinent unknowns under large strain regime and Dirichlet or Dirichlet-type (so-called strong anchoring) conditions, assuming a local strong nondegeneracy requirement.

The energy considered is surface polyconvex with respect to the shell middle surface deformation gradient and convex with respect to the derivative of thickness and microstructure descriptor field. It also incorporates surface energy depending

*Received by the editors April 12, 2022; accepted for publication (in revised form) March 29, 2023; published electronically September 27, 2023.

<https://doi.org/10.1137/22M1490004>

Funding: This work has been developed within programs of the research group in “Theoretical Mechanics” of the “Centro di Ricerca Matematica Ennio De Giorgi” of the Scuola Normale Superiore at Pisa. Moreover, we acknowledge support of the Italian groups of Mathematical Physics (GNFM-INDAM) and of Analysis, Probability, and Applications (GNAMPA-INDAM).

[†]DICEA, Università di Firenze, via Santa Marta 3, I-50139 Firenze, Italy (paolomaria.mariano@unifi.it).

[‡]DSMFI, Università di Parma, Parco Area delle Scienze 53/A, I-43100 Parma, Italy (domenico.mucci@unipr.it).

on crack area and a generalized curvature tensor. This last term has a regularizing nature: It accounts for surface nonlocal interactions (precisely of second-neighbor type and thus of curvature type) and bending effects in material bonds (for the latter ones, see [53]). A term taking into account possible tip energy (it appears prominently in cohesive schemes) completes the energy. Deformations and microstructural fields are only constrained to have their jump set *contained* in the crack path *but not coinciding necessarily* with it.

Our analyses apply to several circumstances, ranging from shells considered as structural elements to thin films or to the description of biological tissues in which one dimension is definitely smaller than the other two, and a through-the-thickness microstructure implies actions hardly representable in terms of standard stresses.

1.1. Structural shells and thin films as surfaces with directors. In 1958, J. L. Ericksen and C. A. Truesdell adopted E. and F. Cosserat's 1909 scheme (which rests on a 1893 proposal by P. Duhem) to model structural elements as rods or shells as one-dimensional or two-dimensional bodies endowed, respectively, with an out-of-line or out-of-middle-surface vector field, with values, so-called *directors*, selected in the unit sphere [17]. They describe the behavior of cross sections, considered as rigid bodies. An additional scalar field may be used to represent variable thickness.

A massive body of pertinent work followed (see, for example, [6], [50], [51], [52], [43], [44], and references therein; this is an incomplete list limited just by the need to make a choice).

We may look just at bending—a curvature effect, indeed—when we constrain the director to be coincident with the normal of the shell middle surface. The plate and shell theories of G. R. Kirchhoff, T. von Karman, P. Naghdi, W. Koiter, and P. Ciarlet are based, in fact, on different curvature-dependent energies [13], [56] (see also [55], [54]).

These models imply the question of their justification in terms of dimensional reduction from three-dimensional space. A natural way to tackle the problem is via E. De Giorgi's Γ -convergence. In this view, in 1999, K. Bhattacharya and R. D. James argued that if one considers a portion of an elastic cylinder between two cross sections at reciprocal finite distance and computes the Γ -limit of the energy (and related minimizers) as the thickness goes to zero, one may obtain the energy pertinent to a membrane or to a Cosserat's surface, depending on whether the cylinder is made of a simple elastic material or a second-grade one, i.e., one including in the list of energy entries the second gradient of deformation. In the latter case, the limit generates a vector field over the surface on which we shrink the cylinder portion (see [49], [21], [22]).

Roughly speaking, membrane energy of different plate models arises from separate scaling assumptions for the energy with respect to the plate thickness. This result comes from a top-to-bottom approach. On the other hand, if we reverse the view [20] and construct a film by superposing an atomic layer over the other, we get at the continuum level a surface endowed at each point with as many directors as the number of atomic layers considered. When we consider films constituted by a mixture of atomic layers, the sequence of vectors at each point is known to within a permutation [40].

1.2. Crack paths as rectifiable sets supporting varifolds. A variational approach to the elastic-brittle behavior has been suggested in 1998 [19], based on minimality required for A. A. Griffith's energy [29]. Crack evolution is then considered by partitioning the time interval and presuming that minimality governs transitions

between subsequent minimizers at discrete times. This scheme has its roots in De Giorgi's idea of minimizing movements [16].

At first, there is distinction between deformations and crack path, although they are connected because the deformation jumps when crack margins detach. In two dimensions, we can control sequences of curves, so that we can maintain the previous distinction. To allow jumps across the crack path, deformations can be naturally considered as special bounded variation maps (*SBV*). Their distributional derivative is a finite measure with an \mathcal{H}^{n-1} -measurable jump set, where n is the ambient space dimension and \mathcal{H}^{n-1} the $(n-1)$ -dimensional Hausdorff measure (see the treatise [5]). In addition, considering a cracking process in the time-step sense mentioned above, in a two-dimensional environment, we can describe steps in which the crack margins come back at least partially in a contact without shear with respect to the original uncracked configuration. Material bonds are, however, not restored, and a crack persists. The approach requires to keep track of the previous steps (see [15] and subsequent work of the same authors).

However, in general, in three-dimensional space, we cannot always control sequences of surfaces. A way to overcome such a difficulty is to look for minimizing *SBV* deformations and to consider the crack to be coincident with the deformation jump set (see once again [15] and subsequent works of the same authors). Also, progressively onward in the course of cracking, appropriate choices in looking at a crack as coinciding with the jump set of an *SBV* deformation might allow us to describe steps in which the crack margins come back at least partially in a contact without shear with respect to the original uncracked configuration. The material bonds are, however, not restored, so the crack persists. In so doing, we should look at a time-step process such that if at a time t_i the crack margins restore the original contact, they were detached at time t_j with $j < i$, according to the *SBV* setting. However, there are cases in which material bonds break to allow the matter to reach a lower energy level but the margins of pertinent cracks do not detach as it occurs in some cracking events in windscreens of cars. Besides these specific cases, if we do not look at processes and consider a unique minimizing step—as we do here—we do not have at our disposal a “previous step” in which crack margins are detached before coming back in contact.

To maintain the distinctions between deformations and crack paths in three-dimensional space, we could consider sequences of surfaces with bounded curvature. They can be described by means of *rectifiable varifolds*, i.e., Radon measures on the Grassmannian constructed by using the tangent planes to \mathcal{H}^2 -rectifiable sets (see for a general treatment [1], [2], [3], [36]). They admit a generalized notion of *curvature* related to their support. The approach (proposed first in [27], [38], [25]) implies the introduction of an energy which is a regularization of Griffith's one, as already mentioned above.

Here we adopt for shells the varifold-based description of crack paths. The underlying two-dimensional setting notwithstanding (at least in terms of the reference domain where fields are defined) allows us to consider in a unique step cases in which crack margins remain or even “come back” at least partially in contact.

1.3. Through-the-thickness microstructures. Thin films show their own peculiar microstructures emerging from their thinness. Tents and tunnels appear in film deposition [30] on a substrate as a consequence of martensite-austenite transitions [9], [34], [35], [32]. In polycrystalline thin films, there may be grain growth larger than the film thickness in a way that grains traverse the film. In this case, film surface and grain boundary energies become comparable. Their interplay may pin the boundaries

against further migration or may enhance them [23]. However, the presence of further active material microstructure may influence drastically the film behavior, as in the presence of magnetization [8]. Besides thin films, in general, shells may be made of materials with active microstructure that determine effects hardly representable in terms of standard stresses. To account directly for them, in representing the body morphology, we not only exploit a fit region in the physical point space but also introduce descriptors of the low-spatial-scale texture of the matter and consider them to be observable. With the aim of having a general model-building framework, we take such descriptors (say, phase fields) as elements of a finite-dimensional complete Riemannian manifold \mathcal{M} not embedded into a linear space (see [11], [37], [48], [39]). The picture of shells as surfaces with directors not necessarily coinciding with the normal vector field falls within the above-mentioned general scheme. When we consider structural shells and thin films endowed with active microstructures in the sense specified above, we have at least a pertinent descriptor, ν , which complements the vector representing the cross-section behavior. This is the scheme that we investigate here accounting for the energy dependence on ν and its gradient. Such an approach is also a way to approximately describe fractures. In fact, we can choose ν to be as a scalar damage indicator, which localizes in a small neighborhood of the crack path [42], [10]. Phenomena of localization, in fact, appear also for vector choices of ν and even in linear elasticity setting as harbingers (or precursors) of fracture [41].

1.4. The main energy under investigation. With progressive extensions, in this paper, we end up with the following energy functional:

$$\begin{aligned} \mathcal{F}(u, \zeta, V, \nu) := & \int_{\Omega} (\tilde{e}(x, u, \zeta, \nu, \nabla u) + \beta_1 |\nabla \zeta|^q + f(x, \nu, \nabla \nu)) \, dx \\ & + \gamma \|V\| + \int_{\mathcal{G}_1(\Omega)} \|A\|^p \, dV + \alpha \|\partial V\|, \end{aligned}$$

where Ω is a domain in the plane; ζ is an out-of-plane vector field describing the shell cross section (cf. (3.1)); ν is the morphological descriptor of a through-the-thickness (active) material microstructure (cf. section 6); \tilde{e} is a surface polyconvex function with respect to ∇u (cf. (6.3)); f is a nonnegative, continuous, and quasiconvex $W^{1,s}(\Omega, \mathcal{M})$ function (namely, f is bounded and such that $f(x, \nu, N)$, with N a linear map from \mathbb{R}^2 onto the tangent space of \mathcal{M} at $\nu \in \mathcal{M}$, is lesser than or equal to the integral over unit square of $f(x, \nu, N + d\varphi)$, with φ a compactly supported smooth function; for a precise general definition, see Definition 6.1); V is a one-dimensional varifold with mass $\|V\|$ (the term including $\|V\|$ corresponds to the area term in Griffith's energy) and generalized curvature tensor $A := A(V)$. The boundary mass term $\|\partial V\|$ accounts for possible tip energy (cf. section 2.3 for the notation concerning curvature varifolds with boundary). We prove the existence of minimizers in terms of deformations u , vector fields ζ , descriptors ν , and varifolds V .

Specifically, we take u in the space of weak diffeomorphisms constructed on *SBV* maps—a space defined in [25]—which satisfy a local strong nondegeneracy condition. Also, ν is taken as a Sobolev map valued on an intrinsic, finite-dimensional, geodesic-complete, differentiable Riemannian manifold. When ν is itself in *SBV* with values in the same intrinsic manifold, in the list of energy entries we replace the distributional derivative of ν with its total variation (or seminorm).

2. Background material.

2.1. Special functions of bounded variation. A summable function $u \in L^1(\Omega)$, defined on a bounded domain $\Omega \subset \mathbb{R}^2$, is said to be *of bounded variation* if

the distributional derivative Du is a finite measure in Ω . In this case, the function u is approximately differentiable \mathcal{L}^2 -a.e. in Ω , and the approximate gradient ∇u agrees with the density of the Radon–Nikodym derivative of Du with respect to the Lebesgue measure \mathcal{L}^2 . Therefore, the decomposition $Du = \nabla u \mathcal{L}^2 + D^s u$ holds true, where the component $D^s u$ is singular with respect to \mathcal{L}^2 . Also, denoting by \mathcal{H}^k the k -dimensional Hausdorff measure, the *jump set* $S(u)$ of u is a countably 1-rectifiable subset of Ω that agrees \mathcal{H}^1 -essentially with the complement of u Lebesgue’s set. If, in addition, the singular component $D^s u$ is concentrated on the jump set $S(u)$, we say that u is a *special function of bounded variation* and write in short $u \in SBV(\Omega)$.

A vector-valued function $u : \Omega \rightarrow \mathbb{R}^3$ belongs to the class $SBV(\Omega, \mathbb{R}^3)$ if its components u^j are in $SBV(\Omega)$. In that case, we get

$$|Du|(B) = \int_B |\nabla u| dx + \int_{B \cap S(u)} |u^+ - u^-| d\mathcal{H}^1$$

for each Borel set $B \subset \Omega$, where the approximate gradient $\nabla u \in L^1(\Omega, \mathbb{M}^{3 \times 2})$; the jump set $S(u)$ is defined as in the scalar case, or componentwise; and u^\pm are the one-sided limits at \mathcal{H}^1 -a.e. point $x \in S(u)$ [5].

2.2. Currents carried by approximately differentiable maps. Let Ω be a bounded domain in \mathbb{R}^2 . For $u : \Omega \rightarrow \mathbb{R}^3$, an a.e. approximately differentiable map, we denote by ∇u its approximate gradient. The map u has a Lusin representative on the subset $\tilde{\Omega}$ of Lebesgue points pertaining to both u and ∇u . Also, we have $\mathcal{L}^2(\Omega \setminus \tilde{\Omega}) = 0$. We shall thus denote $M(F) := (F, \text{adj}_2 F) \in \mathbb{M}^{3 \times 2} \times \mathbb{R}^3$, where $\text{adj}_2 F$ is the 3-vector given by the 2×2 minors of the matrix $F \in \mathbb{M}^{3 \times 2}$. We say that $u \in \mathcal{A}^1(\Omega, \mathbb{R}^3)$ if $\nabla u \in L^1(\Omega, \mathbb{M}^{3 \times 2})$ and the adjoint vector $\text{adj}_2 \nabla u \in L^1(\Omega, \mathbb{R}^3)$.

The *graph* of a map $u \in \mathcal{A}^1(\Omega, \mathbb{R}^3)$ is defined by

$$\mathcal{G}_u := \left\{ (x, y) \in \Omega \times \mathbb{R}^3 \mid x \in \tilde{\Omega}, y = \tilde{u}(x) \right\},$$

where $\tilde{u}(x)$ is the Lebesgue value of u . It turns out that \mathcal{G}_u is a countably 2-rectifiable set of $\Omega \times \mathbb{R}^3$, with $\mathcal{H}^2(\mathcal{G}_u) < \infty$. The approximate tangent plane at $(x, u(x))$ is generated by the vectors $\mathbf{t}_1(x) = (1, 0, \partial_1 u(x))$ and $\mathbf{t}_2(x) = (0, 1, \partial_2 u(x))$ in \mathbb{R}^5 , where the partial derivatives are the column vectors of the gradient ∇u , and we take $\nabla u(x)$ as the Lebesgue value of ∇u at $x \in \tilde{\Omega}$. Therefore, the 2-vector

$$\xi(x) := \frac{\mathbf{t}_1(x) \wedge \mathbf{t}_2(x)}{|\mathbf{t}_1(x) \wedge \mathbf{t}_2(x)|}$$

provides an orientation to the graph \mathcal{G}_u .

The current G_u carried by the graph of u is a functional taking values

$$G_u(\omega) = \langle G_u, \omega \rangle := \int_{\mathcal{G}_u} \langle \omega, \xi \rangle d\mathcal{H}^2,$$

where ω belongs to the space $\mathcal{D}^2(\Omega \times \mathbb{R}^3)$ of compactly supported 2-forms on \mathcal{G}_u . It turns out that G_u is an *integer multiplicity* (i.m.) *rectifiable current* with finite mass $\mathbf{M}(G_u)$ equal to the area $\mathcal{H}^2(\mathcal{G}_u)$ of \mathcal{G}_u ; we then write $G_u \in \mathcal{R}_2(\Omega \times \mathbb{R}^3)$. Since the Jacobian of $x \mapsto (x, u(x))$, which is the graph map, is equal to $|\mathbf{t}_1(x) \wedge \mathbf{t}_2(x)|$, by the area formula, we get

$$\mathbf{M}(G_u) = \mathcal{H}^2(\mathcal{G}_u) = \int_{\Omega} \sqrt{1 + |\nabla u|^2 + |\text{adj}_2 \nabla u|^2} dx < \infty$$

(compare Thm. 4 at p. 225 in [28, Vol. 1, sect. 3.1.5]).

By duality, the *boundary* of G_u is the 1-current ∂G_u acting on $\mathcal{D}^1(\Omega \times \mathbb{R}^3)$, the space of compactly supported smooth 1-forms η in $\Omega \times \mathbb{R}^3$, as

$$\langle \partial G_u, \eta \rangle := \langle G_u, d\eta \rangle, \quad \eta \in \mathcal{D}^1(\Omega \times \mathbb{R}^3),$$

where $d\eta$ is the differential of η . By the Stokes theorem, we get $\partial G_u = 0$ on $\mathcal{D}^1(\Omega \times \mathbb{R}^3)$ if u is of class C^2 . Such a relation holds true also for Sobolev maps $u \in W^{1,2}(\Omega, \mathbb{R}^3)$ by approximation. However, in general, the boundary ∂G_u does not vanish and may not have finite mass in $\Omega \times \mathbb{R}^3$. On the other hand, if ∂G_u has finite mass, the boundary rectifiability theorem states that ∂G_u is an i.m. rectifiable current in $\mathcal{R}_1(\Omega \times \mathbb{R}^3)$. In particular, it turns out that u is a special function of bounded variation, $u \in SBV(\Omega, \mathbb{R}^3)$. Actually, u belongs to the class SBV_0 discussed in [5].

An extended treatment of currents is in the two-volume treatise [28].

2.3. Curvature varifolds with boundary. A 1-varifold over Ω , a domain in \mathbb{R}^2 , is a nonnegative Radon measure on the trivial bundle $\mathcal{G}_1(\Omega) := \Omega \times \mathcal{G}_{1,2}$, where $\mathcal{G}_{1,2}$ is the Grassmannian manifold of 1-planes Π (straight lines) through the origin in \mathbb{R}^2 .

If \mathfrak{b} is a 1-rectifiable subset of $\Omega \subset \mathbb{R}^2$, for $\mathcal{H}^1 \llcorner \mathfrak{b}$ a.e. $x \in \Omega$, there exists the approximate tangent 1-space $T_x \mathfrak{b}$ to \mathfrak{b} at x . We thus denote by $\Pi(x)$ the 2×2 matrix that identifies the orthogonal projection of \mathbb{R}^2 onto $T_x \mathfrak{b}$.

We define

$$(2.1) \quad V_{\mathfrak{b},\theta}(\varphi) := \int_{\mathcal{G}_1(\Omega)} \varphi(x, \Pi) dV_{\mathfrak{b},\theta}(x, \Pi) := \int_{\mathfrak{b}} \theta(x) \varphi(x, \Pi(x)) d\mathcal{H}^1(x)$$

for any $\varphi \in C_c^0(\mathcal{G}_1(\Omega))$, where $\theta \in L^1(\mathfrak{b}, \mathcal{H}^1)$ is a nonnegative density function. If θ is integer valued, $V = V_{\mathfrak{b},\theta}$ is said to be the *integer rectifiable varifold* associated with $(\mathfrak{b}, \theta, \mathcal{H}^1)$.

The *weight measure* of V is the Radon measure in Ω given by $\mu_V := \pi_{\#} V$, where $\pi : \mathcal{G}_1(\Omega) \rightarrow \Omega$ is the canonical projection. Therefore, we find $\mu_V = \theta \mathcal{H}^1 \llcorner \mathfrak{b}$, and the *mass* of V is

$$\|V\| := V(\mathcal{G}_1(\Omega)) = \mu_V(\Omega) = \int_{\mathfrak{b}} \theta d\mathcal{H}^1.$$

DEFINITION 2.1. *An integer rectifiable 1-varifold $V = V_{\mathfrak{b},\theta}$ is called a curvature 1-varifold with boundary if there exist a function $A \in L^1(\mathcal{G}_1(\Omega), \mathbb{R}^{2*} \otimes \mathbb{R}^2 \otimes \mathbb{R}^{2*})$, $A = (A_j^i)$, and a vector-valued Radon measure ∂V with finite mass $\|\partial V\|$ such that*

$$\int_{\mathcal{G}_1(\Omega)} (\Pi D_x \varphi + A D_{\Pi} \varphi + \varphi \text{tr}(AI)) dV(x, \Pi) = - \int_{\mathcal{G}_1(\Omega)} \varphi d\partial V(x, \Pi)$$

for every $\varphi \in C_c^\infty(\mathcal{G}_1(\Omega))$. We write in short $\partial V \in \mathcal{M}(\mathcal{G}_1(\Omega), \mathbb{R}^2)$. Moreover, for $p \geq 1$, the subclass of curvature 1-varifolds with boundary such that $|A| \in L^p(\mathcal{G}_1(\Omega))$ is indicated by $CV_1^p(\Omega)$ (see [27, Exs. 1, 2] for specific examples).

With respect to Allard’s approach (see [1], [2]), with definition (2.1) we gain more information. For example, if Ω is the unit disk centered at the origin and we take a one-dimensional varifold given by three half-lines from 0, which form three angles of 120° , by using Allard’s definition, we find zero mean curvature and zero boundary. At variance (compare with the results in [36]) with the view adopted here, the boundary measure is the sum of three Dirac deltas supported at the points $(0, P_i)$

in the Grassmannian $\mathcal{G}_1(\Omega)$, where P_i is the one-dimensional space determined by the i th half-line, with $i = 1, 2, 3$.

Varifolds in $CV_1^p(\Omega)$ have generalized curvature in L^p [36]. Therefore, Allard's compactness theorem applies (see [1], [2], [3]).

THEOREM 2.2. *For $1 < p < \infty$, let $\{V^{(h)}\} \subset CV_1^p(\Omega)$ be a sequence of curvature 1-varifolds $V^{(h)} = V_{\mathbf{b}_h, \theta_h}$ with boundary. Corresponding curvatures and boundaries are indicated by $A^{(h)}$ and $\partial V^{(h)}$, respectively. Assume that there exists a constant $c > 0$ such that, for every h ,*

$$\mu_{V^{(h)}}(\Omega) + \|\partial V^{(h)}\| + \int_{\mathcal{G}_1(\Omega)} |A^{(h)}|^p dV^{(h)} \leq c.$$

Then, there exists a (not relabeled) subsequence of $\{V^{(h)}\}$ and a 1-varifold $V = V_{\mathbf{b}, \theta} \in CV_1^p(\Omega)$, with curvature A and boundary ∂V , such that

$$V^{(h)} \rightharpoonup V, \quad A^{(h)} dV^{(h)} \rightharpoonup A dV, \quad \partial V^{(h)} \rightharpoonup \partial V$$

in the sense of measures. Moreover, for any convex and l.s.c. function $f : \mathbb{R}^{2} \otimes \mathbb{R}^2 \otimes \mathbb{R}^{2*} \rightarrow [0, +\infty]$, we get*

$$\int_{\mathcal{G}_1(\Omega)} f(A) dV \leq \liminf_{h \rightarrow \infty} \int_{\mathcal{G}_1(\Omega)} f(A^{(h)}) dV^{(h)}.$$

3. A skeletal model. We look first to the shell middle surface and take for it a planar reference configuration that is a two-dimensional smooth domain Ω in \mathbb{R}^2 , where Cartesian coordinates $x = (x_1, x_2)$ are fixed. A map $u : \Omega \rightarrow \mathbb{R}^3$, say, $u = (u^1, u^2, u^3)$, represents a *deformation*.

When smoothness and nonsingularity are ensured, the map u determines an immersion of Ω into \mathbb{R}^3 ; the tangent plane to the deformed film middle plane does not degenerate. Formally, it is tantamount to imposing $|\text{adj}_2 \nabla u(x)| > 0$ for any $x \in \Omega$. In other words, if $\partial_i u$ denotes a column vectors of the gradient matrix ∇u , we are imposing that the vector product $\partial_1 u \times \partial_2 u$ does not vanish at every point. Therefore, the normal to $u(\Omega)$, the deformed middle surface, is the unit vector

$$(3.1) \quad \mathbf{n}(x) = \frac{\partial_1 u(x) \times \partial_2 u(x)}{|\partial_1 u(x) \times \partial_2 u(x)|}.$$

It can be considered as a descriptor of out-of-middle-surface shell behavior. However, such information can be carried out by an \mathbb{S}^2 -valued vector field $x \mapsto \zeta(x)$ defined over Ω and constrained to be at every $x \in \Omega$ such that

$$(3.2) \quad (\partial_1 u(x) \times \partial_2 u(x)) \bullet \zeta(x) > 0,$$

where \bullet is the scalar product in \mathbb{R}^3 ; in the absence of out-of-middle-surface shear, we get $\zeta(x) = \mathbf{n}(x)$ (the scheme is standard; see [17], [50], [52], [51], [6], and references therein).

Given $q > 2$ and $r > 1$, consider the energy

$$\mathcal{G}(u, \zeta) := \int_{\Omega} \left(|\nabla u|^q + |\nabla \zeta|^r + \Phi(\nabla u, \zeta) \right) dx$$

acting on couples of Sobolev functions (u, ζ) in $[W^{1,1}(\Omega, \mathbb{R}^3)]^2$, with $|\zeta| = 1$ \mathcal{L}^2 -a.e. in Ω . Impose also the uniform bound $\|u\|_{L^\infty(\Omega)} \leq K$. Eventually, we may consider Dirichlet-type conditions for u and ζ over $\partial\Omega$ in the sense of traces.

As a matter of notation, according to [14], we define the set

$$\tilde{Y} := \left\{ (F, \zeta) \in \mathbb{M}^{3 \times 2} \times \mathbb{R}^3 \mid \det(F|\zeta) > 0 \right\},$$

where $(F|\zeta) \in \mathbb{M}^{3 \times 3}$. Therefore, according to (3.2), if $F = \nabla u$, we get

$$\det(\nabla u|\zeta) = (\partial_1 u \times \partial_2 u) \bullet \zeta.$$

DEFINITION 3.1. We shall denote by $\tilde{\mathcal{F}}$ the class of nonnegative functions $\Phi : \tilde{Y} \rightarrow [0, +\infty)$ such that

- (1) $\Phi(F, \cdot)$ is continuous in \mathbb{R}^3 for all $F \in \mathbb{M}^{3 \times 2}$;
- (2) $\Phi(\cdot, \zeta)$ is polyconvex (precisely, $\Phi(\cdot, \zeta)$ is the restriction to $\{F \in \mathbb{M}^{3 \times 2} \mid \det(F|\zeta) > 0\}$ of a polyconvex function $g : \mathbb{M}^{3 \times 2} \rightarrow \mathbb{R}$, i.e., a convex function of all the minors of F) for all $\zeta \in \mathbb{R}^3$;
- (3) $\Phi(F, \zeta) \rightarrow +\infty$ if $\det(F|\zeta) \rightarrow 0^+$.

We assume that the integrand Φ in $\mathcal{G}(u, \zeta)$ is in the class $\tilde{\mathcal{F}}$. An example is given by $\Phi(F, \zeta) = -\log(\det(F|\zeta)) \vee 0$ if $\det(F|\zeta) > 0$ and $\Phi(F, \zeta) = +\infty$ otherwise in $\mathbb{M}^{3 \times 2} \times \mathbb{R}^3$. We also fix $q > 2$ and $r > 1 \vee 2q/(3q - 4)$.

THEOREM 3.2. With the previous assumptions, minima of $\mathcal{G}(u, \zeta)$ exist among couples in $W^{1,q}(\Omega, \mathbb{R}^3) \times W^{1,r}(\Omega, \mathbb{R}^3)$, with $\|u\|_\infty \leq K$ and $|\zeta| = 1$. They are such that (3.2) holds true \mathcal{L}^2 -a.e. on Ω .

Proof. By using compactness arguments, we shall repeatedly pass to not relabeled subsequences. Taking a minimizing sequence $\{(u_h, \zeta_h)\}$, we infer that $u_h \rightharpoonup u_\infty$ weakly in $W^{1,q}(\Omega, \mathbb{R}^3)$ and $\zeta_h \rightharpoonup \zeta_\infty$ weakly $W^{1,r}(\Omega, \mathbb{R}^3)$ for some functions $(u_\infty, \zeta_\infty) \in W^{1,q}(\Omega, \mathbb{R}^3) \times W^{1,r}(\Omega, \mathbb{R}^3)$. Moreover, the a.e. convergences $u_h \rightarrow u_\infty$ and $\zeta_h \rightarrow \zeta_\infty$ imply that $\|u_\infty\|_{L^\infty(\Omega)} \leq K$ and $|\zeta_\infty| = 1$ a.e. in Ω , whereas by lower semicontinuity,

$$\int_\Omega \left(|\nabla u_\infty|^q + |\nabla \zeta_\infty|^r \right) dx \leq \liminf_{h \rightarrow \infty} \int_\Omega \left(|\nabla u_h|^q + |\nabla \zeta_h|^r \right) dx.$$

The bound $\sup_h \int_\Omega \Phi(\nabla u_h, \zeta_h) dx < \infty$ implies that

$$(3.3) \quad (\partial_1 u_h(x) \times \partial_2 u_h(x)) \bullet \zeta_h(x) > 0$$

holds true for each $h \in \mathbb{N}$ and \mathcal{L}^2 -a.e. in Ω . We claim that condition (3.3) is preserved (with possibly the equality sign instead of $>$) when passing to the limit. In fact, by the parallelogram inequality, we get for every h and for a.e. $x \in \Omega$ the bound

$$(3.4) \quad |\partial_1 u_h \times \partial_2 u_h|^{q/2} \leq C |\nabla u_h|^q, \quad q > 2,$$

for some absolute constant C . Moreover, if $1 \vee 2q/(3q - 4) < r \leq 2$, we also have strong convergence $\zeta_h \rightarrow \zeta_\infty$ in $L^p(\Omega)$ for $p = 2r/(2 - r)$, where p is greater than the conjugate exponent to $q/2$. Therefore, we infer the existence of a function $H \in L^1(\Omega)$ such that $(\partial_1 u_h \times \partial_2 u_h) \bullet \zeta_h \rightharpoonup H$ weakly in $L^1(\Omega)$. On the other hand, by (3.4) and a standard density argument, it turns out that in the distributional sense, we have, e.g.,

$$\text{Div}(u_h^1 \partial_2 u_h^2, -u_h^1 \partial_1 u_h^2) = (\partial_1 u_h^1 \partial_2 u_h^2 - \partial_2 u_h^1 \partial_1 u_h^2) \mathcal{L}^2 \llcorner \Omega \quad \forall h \in \bar{\mathbb{N}},$$

where $\bar{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$; i.e., for each test function $\varphi \in C_c^\infty(\Omega)$,

$$\int_\Omega \left(\partial_1 u_h^1 \partial_2 u_h^2 - \partial_2 u_h^1 \partial_1 u_h^2 \right) \varphi dx = \int_\Omega u_h^1 \left(\partial_1 u_h^2 \partial_2 \varphi - \partial_2 u_h^2 \partial_1 \varphi \right) dx.$$

We thus infer the weak convergence

$$\left(\partial_1 u_h^1 \partial_2 u_h^2 - \partial_2 u_h^1 \partial_1 u_h^2\right) \rightarrow \left(\partial_1 u_\infty^1 \partial_2 u_\infty^2 - \partial_2 u_\infty^1 \partial_1 u_\infty^2\right)$$

in $L^{q/2}(\Omega)$ and hence a.e. in Ω . This implies that $H = (\partial_1 u_\infty \times \partial_2 u_\infty) \bullet \zeta_\infty$ and also the pointwise convergence a.e. in Ω :

$$(\partial_1 u_h \times \partial_2 u_h) \bullet \zeta_h \rightarrow (\partial_1 u_\infty \times \partial_2 u_\infty) \bullet \zeta_\infty.$$

Therefore, the lower semicontinuity

$$\int_\Omega \Phi(\nabla u_\infty, \zeta_\infty) dx \leq \liminf_{h \rightarrow \infty} \int_\Omega \Phi(\nabla u_h, \zeta_h) dx < \infty$$

holds, whence the couple of functions (u_∞, ζ_∞) satisfies the strict inequality in (3.2) for \mathcal{L}^2 -a.e. $x \in \Omega$, as required. \square

It may happen that the deformed surface $u(\Omega)$ has a crease. This is described, e.g., when the unit normal $\mathbf{n}(x)$ is smooth outside a 1-rectifiable set \mathfrak{J} of Ω . In this case, ζ satisfies the condition $\zeta(x) \bullet \mathbf{m} > 0$ at $x \in \mathfrak{J}$ for some unit vector \mathbf{m} that lies in the cone between the one-sided limits of \mathbf{n} at $x \in \mathfrak{J}$.

Therefore, a second-order theory (yielding, e.g., to a regularity of the unit normal $\mathbf{n}(x)$) should be applied in order to describe in a precise way the above-mentioned angle condition along creases over the deformed surface.

4. Cracks and jump sets. We assume here that the components u^j of u are L^∞ -functions in the class $SBV(\Omega)$. Again, we assume a uniform bound $\|u\|_{L^\infty(\Omega)} \leq K$.

We then choose the descriptor $\zeta : \Omega \rightarrow \mathbb{S}^2 \subset \mathbb{R}^3$ of the out-of-middle-surface shell behavior as a special function of bounded variation in $SBV(\Omega, \mathbb{R}^3)$, with $|\zeta| = 1$ a.e., in such a way that the constraint (3.2) holds true \mathcal{L}^2 -a.e. in $\Omega \setminus S(u)$. Since ζ describes the behavior of shell cross sections, the continuity of matter allows us to consider the discontinuity set of ζ as included in the one of u , namely,

$$S(\zeta) \subseteq S(u).$$

The strict inclusion occurs when the crack margins are in contact across the whole shell thickness; in this case, we would have no deformation jump along the crack portion where the margins remain in contact. If one thinks of a cracking process, there could be circumstances, already mentioned above, in which margins remain, at least in part, always in contact, although the energetic content of the material bond is such that they become unstable and prefer, energetically, to break. Accounting for this aspect—i.e., a permanent contact in a portion of a crack—is a peculiarity of the present model. For analytical reasons, we replace the above set inclusion by the inequality

$$(4.1) \quad \mathcal{H}^1 \llcorner S(\zeta_h) \leq \mathcal{H}^1 \llcorner S(u_h) \quad \forall h,$$

along minimizing sequences $\{(u_h, \zeta_h)\}$.

In principle, the limit maps (u, ζ) may not satisfy condition (4.1). In fact, it may happen that the jump of u_h on $S(u_h)$ goes to zero somewhere when passing to the limit through the compactness theorem. In order to preserve inequality (4.1), we shall introduce a condition ensuring that the jump $u_h^+ - u_h^-$ on $S(u_h)$ cannot “decrease” or “disappear,” so that one has $\mathcal{H}^1(S(u_h)) \rightarrow \mathcal{H}^1(S(u))$.

Over the reference domain Ω , we consider a curvature 1-varifold with boundary $V = V_{\mathfrak{b}, \theta}$ such that the discontinuity set $S(u)$ of the deformation map u is contained in the 1-rectifiable set \mathfrak{b} . A crack in which the margins remain at least partially in contact is characterized by $\mathcal{H}^1(\mathfrak{b} \setminus S(u)) > 0$.

4.1. Membranes with cracks. At a first glance, we do not consider the out-of-middle-surface descriptor ζ .

DEFINITION 4.1. *A macroscopic configuration of a cracked membrane is a pair composed by the bounded connected open set $\Omega \subset \mathbb{R}^2$ with Lipschitz boundary and a curvature 1-varifold with boundary, namely, $V = V_{\mathbf{b},\theta} \in CV_1^p(\Omega)$ for some $p > 1$.*

We consider bounded deformation maps u in $\mathcal{A}^1(\Omega, \mathbb{R}^3)$, i.e., a.e. approximately differentiable maps u with $\nabla u \in L^1(\Omega, \mathbb{M}^{3 \times 2})$ and $|\text{adj}_2 \nabla u| \in L^1(\Omega)$, so that the current G_u carried by the rectifiable graph of u is i.m. rectifiable in $\mathcal{R}_2(\Omega \times \mathbb{R}^3)$ and with finite mass, $\mathbf{M}(G_u) < \infty$.

In general, the boundary current $\partial G_u \in \mathcal{D}_1(\Omega \times \mathbb{R}^3)$ is nonzero, and it may also have unbounded mass. We assume furthermore that fractures and holes in the graph of u are controlled by the crack V , namely,

$$(4.2) \quad \pi_{\#} |\partial G_u| \leq \mu_V,$$

where μ_V is the weight measure in Ω of the varifold V , $\pi : \Omega \times \mathbb{R}^3 \rightarrow \Omega$ is the projection onto the first two coordinates, and $|\cdot|$ is the total variation of the vector-valued measure ∂G_u , so that $\pi_{\#} |\partial G_u|(B) = |\partial G_u|(B \times \mathbb{R}^3)$ for each Borel set $B \subset \Omega$.

If $\mu_V(\Omega) < \infty$ and the bound (4.2) holds true, $|\partial G_u|$ is a finite measure, and actually $\mathbf{M}(\partial G_u) < \infty$. Therefore, the boundary rectifiability theorem yields that ∂G_u is an i.m. rectifiable current—in short $\partial G_u \in \mathcal{R}_1(\Omega \times \mathbb{R}^3)$ —and hence u is a special function of bounded variation, namely, $u \in SBV(\Omega, \mathbb{R}^3)$.

More precisely, since $|D^j u| \leq \|u\|_{\infty} \pi_{\#} |\partial G_u|$, recalling that $\mu_V = \theta \mathcal{H}^1 \llcorner \mathbf{b}$, by assumption (4.2), it turns out that the jump set of u is contained in the support \mathbf{b} of V , where the positive multiplicity function θ is integer-valued, and we actually have

$$(4.3) \quad \mathcal{H}^1 \llcorner S(u) \leq \mathcal{H}^1 \llcorner \mathbf{b}.$$

Example 4.1. Recalling that $\mu_V = \theta \mathcal{H}^1 \llcorner \mathbf{b}$, the validity of inequality (4.2) relies on the presence of the positive integer θ , which actually accounts for the multiplicity of the projection of ∂G_u . Taking, e.g., $\Omega = B^2$, the unit open disk centered at the origin, and $u : B^2 \rightarrow \mathbb{R}^3$ given by

$$u(x_1, x_2) := \begin{cases} (x_1, x_2, 0) & \text{if } x_1 < 0, \\ (1 + x_1, x_2, 0) & \text{if } x_1 > 0, \end{cases}$$

we get $\partial G_u = \gamma_{0\#} \llbracket I \rrbracket - \gamma_{1\#} \llbracket I \rrbracket$ on $\mathcal{D}^1(B^2 \times \mathbb{R}^3)$, where $I = (-1, 1)$ and $\gamma_{\alpha} : I \rightarrow \mathbb{R}^5$ is defined by $\gamma_{\alpha}(t) := (0, t, \alpha, t, 0)$ for $\alpha = 0, 1$. Therefore, $u \in SBV(B^2, \mathbb{R}^3)$, with the jump set $S(u) = \{0\} \times I$, so that inequality (4.2) holds true with, e.g., $\theta = 2$ and $\mathbf{b} = S(u)$.

We have already defined $M(F) = (F, \text{adj}_2 F) \in \mathbb{M}^{3 \times 2} \times \mathbb{R}^3$. Here we assume that a sequence $\{u_h\}_h \subset \mathcal{A}^1(\Omega, \mathbb{R}^3)$ is such that

$$\sup_h \|u_h\|_{L^\infty(\Omega)} \leq K, \quad \sup_h \|M(\nabla u_h)\|_{L^1(\Omega)} < \infty,$$

where the sequence $\{|M(\nabla u_h)|\}$ is equi-integrable on Ω . In addition, we assume that $\pi_{\#} |\partial G_{u_h}| \leq \mu_{V_h}$ for each h . If we have $\sup_h \mu_{V_h}(\Omega) < \infty$, by compactness, possibly passing to a (not relabeled) subsequence, we have $V_h \rightharpoonup V$ weakly as measures, and we find a deformation map u as above such that G_u is i.m. rectifiable and with finite

mass, $G_{u_h} \rightharpoonup G_u$ weakly as currents, and $\pi_{\#}|\partial G_u| \leq \mu_V < \infty$, whence ∂G_u is i.m. rectifiable, too.

In order to recover the weak L^1 convergence of $M(\nabla u_h)$ to $M(\nabla u)$, the starting point is the following special case of the closure theorem proven in [24]. Because of the compactness theorem in SBV , it extends a classical result proven in [7] for Sobolev maps, where the divergence form of gradient minors is exploited.

THEOREM 4.2. *Let $\Omega \subset \mathbb{R}^2$ be a bounded domain and $\{u_h\}$ a sequence of functions from $SBV(\Omega, \mathbb{R}^3)$ converging in $L^1(\Omega, \mathbb{R}^3)$ to a summable function $u : \Omega \rightarrow \mathbb{R}^3$. Assume that for some real exponents $q > 2$ and $r > 1$,*

$$\sup_h \left\{ \|u_h\|_{\infty} + \int_{\Omega} \left(|\nabla u_h|^q + |\text{adj}_2 \nabla u_h|^r \right) dx + \mathcal{H}^1(S(u_h)) \right\} < \infty.$$

Then, $u \in SBV(\Omega, \mathbb{R}^3)$, and the sequence $\mathcal{H}^1 \llcorner S(u_h)$ weakly converges in Ω to a measure μ greater than $\mathcal{H}^1 \llcorner S(u)$. Moreover, ∇u_h weakly converges to ∇u in $L^q(\Omega, \mathbb{M}^{3 \times 2})$, and $\text{adj}_2 \nabla u_h$ weakly converges to $\text{adj}_2 \nabla u$ in $L^r(\Omega, \mathbb{R}^3)$.

4.2. Accounting for the thickness. We fix $q > 2, r, p > 1, K > 0$ and introduce the class $\mathcal{A}_{q,r,p,K}$ of triplets (u, ζ, V) , where

- (i) $u : \Omega \rightarrow \mathbb{R}^3$ is a special function of bounded variation, $u \in SBV(\Omega, \mathbb{R}^3)$, with $\mathcal{H}^1(S(u)) < \infty$, such that $\|u\|_{L^{\infty}(\Omega)} \leq K, \partial_1 u \times \partial_2 u \neq 0$ \mathcal{L}^2 -a.e. on $\Omega, \nabla u \in L^q(\Omega, \mathbb{M}^{3 \times 2})$, and $|\text{adj}_2 \nabla u| \in L^r(\Omega)$;
- (ii) $\zeta : \Omega \rightarrow \mathbb{R}^3$ is a special function of bounded variation, with $\mathcal{H}^1(S(\zeta)) < \infty$, such that $|\zeta| = 1$ \mathcal{L}^2 -a.e. on Ω , and $\nabla \zeta \in L^q(\Omega, \mathbb{M}^{3 \times 2})$;
- (iii) $V \in CV_1^p(\Omega)$; i.e., V is an integer rectifiable curvature 1-varifold with boundary and generalized curvature tensor $A \in L^p(\mathcal{G}_1(\Omega), \mathbb{R}^{2*} \otimes \mathbb{R}^2 \otimes \mathbb{R}^{2*})$, with $\mu_V = \theta \mathcal{H}^1 \llcorner \mathbf{b}$;
- (iv) $\pi_{\#}|\partial G_u| \leq \mu_V$;
- (v) $\mathcal{H}^1 \llcorner S(\zeta) \leq \pi_{\#}|\partial G_u|$;
- (vi) $(\partial_1 u(x) \times \partial_2 u(x)) \bullet \zeta(x) > 0$ for \mathcal{L}^2 -a.e. $x \in \Omega \setminus \mathbf{b}$.

Then we consider on the class of triplets $(u, \zeta, V) \in \mathcal{A}_{q,r,p,K}$ the energy

$$(4.4) \quad \mathcal{F}(u, \zeta, V) := \int_{\Omega} e(x, u, \zeta, \nabla u, \nabla \zeta) dx + \|V\| + \int_{\mathcal{G}_1(\Omega)} \|A\|^p dV + \|\partial V\|,$$

where $e : \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2} \rightarrow \mathbb{R} \cup \{+\infty\}$ is the sum

$$e(x, u, \zeta, F, G) = \tilde{e}(x, u, \zeta, F) + \beta_1 |G|^q$$

for every $(x, u, \zeta, F, G) \in \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2}$, with $\beta_1 > 0$ and the first addendum a nonnegative Carathéodory function satisfying the following properties:

- (a) $\tilde{e}(x, u, \zeta, F)$ is polyconvex with respect to F , namely,

$$\tilde{e}(x, u, \zeta, F) = g(x, u, \zeta, M(F)) \quad \forall F \in \mathbb{M}^{3 \times 2},$$

where $g : \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times (\mathbb{M}^{3 \times 2} \times \mathbb{R}^3) \rightarrow [0, +\infty]$ is a Carathéodory function, with $g(x, u, \zeta, \cdot)$ convex for \mathcal{L}^2 -a.e. $x \in \Omega$ and for all $(u, \zeta) \in \mathbb{R}^3 \times \mathbb{R}^3$;

- (b) for \mathcal{L}^2 -a.e. $x \in \Omega$ and every (u, ζ, F) such that $\det(F|\zeta) > 0$,

$$\tilde{e}(x, u, \zeta, F) \geq \beta_2 (|F|^q + |\text{adj}_2 F|^r) + \Phi(F, \zeta),$$

where $\beta_2 > 0$ and $\Phi \in \tilde{\mathcal{F}}$; see Definition 3.1.

THEOREM 4.3. *The energy minimum of $\mathcal{F}(u, V, \zeta)$ is attained in the class $\mathcal{A}_{q,r,p,K}$.*

Proof. As before, we repeatedly pass to not relabeled subsequences. Choose a minimizing sequence $\{(u_h, \zeta_h, V^{(h)})\} \subset \mathcal{A}_{q,r,p,K}$. Because of assumptions (iv) and (v), by the energy lower bounds, we infer that Ambrosio’s compactness theorem in *SBV* [5, Thm. 4.8] applies to both sequences $\{u_h\}$ and $\{\zeta_h\}$. Therefore, $u_h \rightarrow u$ in $L^1(\Omega, \mathbb{R}^3)$ to some function $u \in SBV(\Omega, \mathbb{R}^3)$ with $\|u\|_{L^\infty(\Omega)} \leq K$, whereas ∇u_h weakly converges to ∇u in $L^q(\Omega, \mathbb{M}^{3 \times 2})$, and $\mathcal{H}^1 \llcorner S(u_h)$ weakly converges in Ω to a measure greater than $\mathcal{H}^1 \llcorner S(u)$. In a similar way, we prove the existence of a function $\zeta \in SBV(\Omega, \mathbb{R}^3)$ satisfying $|\zeta| = 1$ a.e. in Ω such that $\zeta_h \rightarrow \zeta$ in $L^1(\Omega, \mathbb{R}^3)$, $\nabla \zeta_h$ weakly converges to $\nabla \zeta$ in $L^q(\Omega, \mathbb{M}^{3 \times 2})$, and $\mathcal{H}^1 \llcorner S(\zeta_h)$ weakly converges in Ω to a measure greater than $\mathcal{H}^1 \llcorner S(\zeta)$. Furthermore, Allard’s compactness theorem 2.2 applies to the sequence $\{V^{(h)}\}$ in $CV^p(\Omega)$. Also, the Federer–Fleming closure theorem applies to the sequence G_{u_h} (see [28]).

Because of Theorem 4.2, we obtain a triplet (u, ζ, V) satisfying properties (i)–(v). In fact, using (a), the lower-semicontinuity result in [24] yields

$$\int_{\Omega} \tilde{e}(x, u, \nabla u, \zeta) \, dx \leq \liminf_{h \rightarrow \infty} \int_{\Omega} \tilde{e}(x, u_h, \nabla u_h, \zeta_h) \, dx$$

and

$$\int_{\Omega} |\nabla \zeta|^q \, dx \leq \liminf_{h \rightarrow \infty} \int_{\Omega} |\nabla \zeta_h|^q \, dx,$$

whereas

$$\|V\| + \int_{\mathcal{G}_1(\Omega)} \|A\|^p \, dV + \|\partial V\| \leq \liminf_{h \rightarrow \infty} \left(\|V^{(h)}\| + \int_{\mathcal{G}_1(\Omega)} \|A^{(h)}\|^p \, dV^{(h)} + \|\partial V^{(h)}\| \right),$$

so that by (4.4), we get

$$\mathcal{F}(u, \zeta, V) \leq \liminf_{h \rightarrow \infty} \mathcal{F}(u_h, \zeta_h, V^{(h)}).$$

Also, by the weak convergence $\text{adj}_2 \nabla u_h \rightharpoonup \text{adj}_2 \nabla u$ in $L^1(\Omega, \mathbb{R}^3)$, using the bound $\sup_h \int_{\Omega} |\nabla \zeta_h|^q \, dx < \infty$ for $q > 2$ and the embedding theorem, we obtain the weak convergence $(\partial_1 u_h \times \partial_2 u_h) \bullet \zeta_h \rightharpoonup (\partial_1 u \times \partial_2 u) \bullet \zeta$ in $L^1(\Omega)$. As a consequence, the lower-semicontinuity inequality

$$\int_{\Omega} \Phi(\nabla u, \zeta) \, dx \leq \liminf_{h \rightarrow \infty} \int_{\Omega} \Phi(\nabla u_h, \zeta_h) \, dx$$

holds true. Hence, by the lower bound (b), property (vi) is satisfied by the triplet (u, ζ, V) , whence $(u, \zeta, V) \in \mathcal{A}_{q,r,p,K}$, as required. \square

In fact, we could follow the view suggested in [14], considering energy functionals $\tilde{\mathcal{F}}(u, \zeta, V)$, in which we mix together the gradients of u and ζ . Namely, in (4.4), we could assume that $e(x, u, \zeta, F, G) = \tilde{e}(x, h, H)$, where $h := (u, \zeta)$, $H \in \mathbb{M}^{6 \times 2}$ is the matrix with rows (F, G) , while $\tilde{e}(x, h, H)$ is a nonnegative Carathéodory function, which is polyconvex with respect to H and such that for \mathcal{L}^2 -a.e. $x \in \Omega$ and every (h, H) , with $\det(F|\zeta) > 0$,

$$\tilde{e}(x, h, H) \geq \beta_2 \left(|H|^q + |\text{adj}_2 H|^r \right) + \Phi(F, \zeta),$$

where $\beta_2 > 0$ and $\Phi \in \tilde{\mathcal{F}}$.

Since the closure and semicontinuity properties continue to hold, the minimum of $\tilde{\mathcal{F}}(u, \zeta, V)$ is attained in the class $\mathcal{A}_{q,r,p,K}$ of triplets (u, ζ, V) in $\mathcal{A}_{q,r,p,K}$, which satisfy the additional condition $|\text{adj}_2 \nabla(u, \zeta)| \in L^r(\Omega)$, so that we should presume in this case what follows:

1. $u \in SBV(\Omega, \mathbb{R}^3)$, with $\mathcal{H}^1(S(u)) < \infty$, $\|u\|_{L^\infty(\Omega)} \leq K$, $\partial_1 u \times \partial_2 u \neq 0$ \mathcal{L}^2 -a.e. on Ω , and $\nabla u \in L^q(\Omega, \mathbb{M}^{3 \times 2})$;
2. $\zeta \in SBV(\Omega, \mathbb{R}^3)$, with $\mathcal{H}^1(S(\zeta)) < \infty$, $|\zeta| = 1$ \mathcal{L}^2 -a.e. on Ω , and $\nabla \zeta \in L^q(\Omega, \mathbb{M}^{3 \times 2})$;
3. $|\text{adj}_2 \nabla(u, \zeta)| \in L^r(\Omega)$;
4. $V \in CV_1^p(\Omega)$, with $V \in CV_1^p(\Omega)$, with $\mu_V = \theta \mathcal{H}^1 \llcorner \mathfrak{b}$;
5. $\pi_\# |\partial G_u| \leq \mu_V$;
6. $\mathcal{H}^1 \llcorner S(\zeta) \leq \pi_\# |\partial G_u|$;
7. $(\partial_1 u(x) \times \partial_2 u(x)) \bullet \zeta(x) > 0$ for \mathcal{L}^2 -a.e. $x \in \Omega \setminus \mathfrak{b}$.

However, we prefer to maintain separate membrane and out-of-middle-surface behavior, so that we refer to surface polyconvexity as considered in [40].

5. Boundary conditions. We consider two types of boundary conditions. As a first choice, we prescribe Dirichlet-type data, namely,

$$(5.1) \quad u = u_0, \quad \zeta = \frac{\partial_1 u_0 \times \partial_2 u_0}{|\partial_1 u_0 \times \partial_2 u_0|} \quad \text{on } \partial\Omega,$$

in the sense of traces, respectively, for some given a.e. injective function $u_0 \in W^{1,q}(\Omega, \mathbb{R}^3)$ with $\|u_0\|_\infty \leq K$ and $|\text{adj}_2 \nabla u_0| \in L^r(\Omega, \mathbb{R}^+)$.

Then we consider those that we call *strong anchoring conditions*, determined by assigning the boundary current, i.e., in terms of smooth and bounded 1-forms in $\Omega \times \mathbb{R}^3$,

$$\partial G_u \llcorner (\partial\Omega \times \mathbb{R}^3) = \partial G_{u_0} \llcorner (\partial\Omega \times \mathbb{R}^3),$$

a condition that clearly implies the trace equality $u = u_0$. As already mentioned, from a physical viewpoint, such a condition means that we are assigning the work performed in all possible strain modes, all considered at first to be independent and then reconciled in the limit to be compatible (see [26]). This boundary condition is generally not preserved in the minimization process because along the boundary, open cracks may have optimal placement.

A confinement condition prescribing the existence of a compact set \mathcal{C} contained in the open set Ω and such that

$$(5.2) \quad \text{spt } \mu_V \subset \mathcal{C} \quad \forall (u, \zeta, V) \in \mathcal{A}_{q,r,p,K}$$

avoids the problem.

The chain of inequalities $\mathcal{H}^1 \llcorner S(\zeta) \leq \pi_\# |\partial G_u| \leq \mu_V$ implies that the restriction to $\Omega \setminus \mathcal{C}$ of both u and ζ belongs to the Sobolev class $W^{1,q}(\Omega \setminus \mathcal{C}, \mathbb{M}^{3 \times 2})$. Therefore, the prescribed Dirichlet or strong anchoring conditions are preserved in the limit process due to the weak convergence in $W^{1,q}(\Omega \setminus \mathcal{C}, \mathbb{M}^{3 \times 2})$ of both $\{u_h\}$ and $\{\zeta_h\}$.

COROLLARY 5.1. *Let \mathcal{C} and u_0 be as above. Assume that competitors (u, V, ζ) in $\mathcal{A}_{q,r,p,K}$ have finite energy $\mathcal{F}(u, V, \zeta)$ and satisfy the prescribed boundary, clamping, and confinement conditions, with \mathcal{C} being a nonempty set. Then, the energy minimum of $\mathcal{F}(u, V, \zeta)$ is attained in the same class.*

Remark 5.1. Condition (5.2) excludes circumstances in which a crack path may go to the boundary, breaking part of a link with the environment.

6. Shells made of complex materials. To account for active microstructures (e.g., polarization in ferroelectric films), we introduce over Ω another map ν taking values on a connected, geodesic-complete, n -dimensional Riemannian differentiable manifold \mathcal{M} of class C^2 . At every $x \in \Omega$, the map ν summarizes at gross scale geometric information on material microstructure in the film thickness. For this reason, we call $\mathcal{M} = (\mathcal{M}^n, g)$ the *manifold of microstructural shapes*. The generality adopted in choosing \mathcal{M} is a way to furnish unified results, which are independent of specific microstructural features.

First, we assume the layer-descriptor map $\nu : \Omega \rightarrow \mathcal{M}$ to be an *intrinsic Sobolev map* in $W^{1,s}(\Omega, \mathcal{M})$ for some real exponent $s > 1$. Then we discuss the case in which ν can be considered as a special function of bounded variation.

The essential point is that we do not use any embedding of \mathcal{M} in some Euclidean space. Even choosing it to be isometric, it would not be unique (as Nash's theorems indicate). Thus, its choice would become part of the model, while the common effort is to offer a description of the phenomenological world as much as possible free of nonintrinsic elements.

6.1. The Sobolev case. Under previous assumptions, with $d_{\mathcal{M}}$ the geodesic distance in \mathcal{M} , by the Hopf–Rinow theorem, $(\mathcal{M}, d_{\mathcal{M}})$ is a complete metric space. Consequently, we keep referring to results in such spaces, summarizing those aspects that we need for the analysis developed here from essential references [4], [45], [46], [47] on this topic.

Let $\Omega \subset \mathbb{R}^m$ be a bounded domain and $s \geq 1$. A Borel map $\nu : \Omega \rightarrow \mathcal{M}$ is said to be an intrinsic Sobolev map in $W^{1,s}(\Omega, \mathcal{M})$ if there exists a nonnegative function $\phi \in L^s(\Omega)$ such that for every $\nu_0 \in \mathcal{M}$,

1. $x \mapsto d_{\mathcal{M}}(\nu(x), \nu_0)$ is in $L^s(\Omega)$, i.e., $\nu \in L^s(\Omega, \mathcal{M})$;
2. the distributional gradient map $x \mapsto Dd_{\mathcal{M}}(\nu(x), \nu_0)$ satisfies the inequality $|Dd_{\mathcal{M}}(\nu(x), \nu_0)| \leq \phi(x)$ for \mathcal{L}^m -a.e. $x \in \Omega$.

In this setting, the “norm” $|D\nu|(x)$, which is, in essence, the optimal function $\phi \in L^s(\Omega)$ satisfying the inequality $|Dd_{\mathcal{M}}(\nu(x), \nu_0)| \leq \phi(x)$, is well-defined for \mathcal{L}^m -a.e. $x \in \Omega$ by

$$|D\nu|(x) := \sup_{k \in \mathbb{N}} |D(d_{\mathcal{M}}(\nu(x), \nu_k))|,$$

where $\{\nu_k\}_{k \in \mathbb{N}}$ forms a dense and enumerable set in \mathcal{M} .

Weak convergence in $W^{1,s}$ of a sequence $\{\nu_h\} \subset W^{1,s}(\Omega, \mathcal{M})$ to some map $\nu \in W^{1,s}(\Omega, \mathcal{M})$ when $s > 1$ is defined by requiring that $\|d_{\mathcal{M}}(\nu_h, \nu)\|_{L^s(\Omega)} \rightarrow 0$ as $h \rightarrow \infty$ and $\sup_h \| |D\nu_h| \|_{L^s(\Omega)} < \infty$. When $s = 1$, one assumes in addition that the sequence $\{|D\nu_h|\}$ is equi-integrable.

Also, the trace operator $\text{Tr} : W^{1,s}(\Omega, \mathcal{M}) \rightarrow L^s(\partial\Omega, \mathcal{M})$ is well-defined in such a way that for continuous maps $\nu \in W^{1,s}(\Omega, \mathcal{M}) \cap C^0(\bar{\Omega}, \mathcal{M})$, it agrees with the restriction $\nu|_{\partial\Omega}$. Moreover, if $\{\nu_h\} \subset W^{1,s}(\Omega, \mathcal{M})$ weakly converges to some map $\nu \in W^{1,s}(\Omega, \mathcal{M})$, then $\text{Tr}(\nu_h)$ converges to $\text{Tr}(\nu)$ strongly in $L^s(\partial\Omega, \mathcal{M})$. Finally, traces of maps in $W^{1,s}(\Omega, \mathcal{M})$ have a $W^{1-1/s, s}$ -regularity when $s > 1$ (see [33] and [12]).

We endow the tangent bundle $T\mathcal{M}$ with the metric $d_{T\mathcal{M}}$ induced by $d_{\mathcal{M}}$ and consider the vector bundle with base space \mathcal{M} and typical fiber the space of linear homomorphisms $\text{Hom}(\mathbb{R}^m, T\mathcal{M})$. Points of such a bundle are couples (ν, N) , where $\nu \in \mathcal{M}$ and $N : \mathbb{R}^m \rightarrow T_{\nu}\mathcal{M}$ is a linear map. For any fixed $\nu \in \mathcal{M}$, we can identify $N \in \text{Hom}(\mathbb{R}^m, T_{\nu}\mathcal{M})$ with the m -tuple $(v_1, \dots, v_m) \in (T_{\nu}\mathcal{M})^m$, where $v_i = Ne_i$ and (e_1, \dots, e_m) is the canonical basis in \mathbb{R}^m . Therefore, a metric structure on the vector

bundle $\text{Hom}(\mathbb{R}^m, T\mathcal{M})$ is defined through the distance

$$D((\nu, N), (\tilde{\nu}, \tilde{N})) := \left\{ \sum_{i=1}^m d_{T\mathcal{M}}((\nu, v_i), (\tilde{\nu}, w_i))^2 \right\}^{1/2}$$

if $\tilde{N} \in \text{Hom}(\mathbb{R}^m, T_{\tilde{\nu}}\mathcal{M})$ and $w_i = \tilde{N}e_i$ for each i .

A nonnegative and continuous integrand $f : \Omega \times \text{Hom}(\mathbb{R}^m, T\mathcal{M}) \rightarrow [0, +\infty)$ is said to be *admissible* in $W^{1,s}(\Omega, \mathcal{M})$ if for some fixed point $\nu_0 \in \mathcal{M}$ and some positive constant C , the bounds

$$0 \leq f(x, \nu, N) \leq C \left(1 + d_{\mathcal{M}}(\nu, \nu_0)^s + \|N\|_{g(\nu)}^s \right)$$

hold true for all $(x, \nu, N) \in \Omega \times \text{Hom}(\mathbb{R}^m, T_{\nu}\mathcal{M})$, where $\|\cdot\|_{g(\nu)}$ is the operatorial norm. A functional

$$(6.1) \quad \nu \mapsto \int_{\Omega} f(x, \nu(x), d\nu_x) dx$$

with an admissible integrand f is well-defined on maps $\nu \in W^{1,s}(\Omega, \mathcal{M})$, where $d\nu_x$ is the approximate differential. Precisely, if $\nu : \Omega \rightarrow \mathcal{M}$ is a Borel map and $x \in \mathcal{M}$ is a point of approximate continuity of ν , a linear map $N \in \text{Hom}(\mathbb{R}^m, T_{\nu(x)}\mathcal{M})$ is said to be an *approximate differential* of ν at x if for all $\epsilon > 0$,

$$\lim_{\rho \rightarrow 0^+} \frac{1}{\rho^m} \mathcal{L}^m \left(\left\{ y \in B_{\rho}^m(x) \mid d_{\mathcal{M}}(\nu(y), \exp_{\nu(x)}(N(y-x))) \geq \epsilon |y-x| \right\} \right) = 0.$$

When it exists, the approximate differential of ν at x is unique. For smooth maps, it agrees with the classical differential (in the geometric sense). For this reason, the notation $d\nu_x$ is used for it. A Sobolev map ν in $W^{1,s}(\Omega, \mathcal{M})$ is approximately differentiable at \mathcal{L}^m -a.e. $x \in \mathcal{M}$, whence the functional (6.1) makes sense.

We ask for sequential lower semicontinuity. Thus, we need a suitable notion of quasiconvexity.

DEFINITION 6.1. *A locally bounded function $f : \Omega \times \text{Hom}(\mathbb{R}^m, T\mathcal{M}) \rightarrow \mathbb{R}$ is said to be quasiconvex if for every $(x, \nu, N) \in \Omega \times \text{Hom}(\mathbb{R}^m, T_{\nu}\mathcal{M})$ and every test function $\varphi \in C_c^{\infty}(Q_1, T_{\nu}\mathcal{M})$, where $Q_1 := [-1/2, 1/2]^m$, by looking at $d\varphi_y$ as an element of $\text{Hom}(\mathbb{R}^m, T_{\nu}\mathcal{M})$, the following inequality holds true:*

$$f(x, \nu, N) \leq \int_{Q_1} f(x, \nu, N + d\varphi_y) dy.$$

THEOREM 6.2 (see [18]). *Take $s \geq 1$. Let $f : \Omega \times \text{Hom}(\mathbb{R}^m, T\mathcal{M}) \rightarrow [0, +\infty)$ be a nonnegative and continuous admissible integrand in $W^{1,s}(\Omega, \mathcal{M})$. Then, the functional (6.1) is sequentially weakly lower semicontinuous in $W^{1,s}(\Omega, \mathcal{M})$ if and only if f is quasiconvex.*

Coming back to the physical dimension $m = 2$, a shell with out-of-middle-surface vector field ζ , a crack represented by a 1-varifold V , a descriptor ν of the through-the-thickness material morphology, and subjected to a deformation u , is modeled by a quadruplet (u, ζ, V, ν) belonging to the class $\mathcal{A}_{q,r,p,K,s}(\mathcal{M})$ constructed as follows: For $q > 2$, $r, p, s > 1$, $K > 0$, we assume that the triplet (u, ζ, V) belongs to the class $\mathcal{A}_{q,r,p,K}$ introduced above, while we let $\nu \in W^{1,s}(\Omega, \mathcal{M})$, where $\mathcal{M} = (\mathcal{M}^n, g)$ is as above.

On the class $\mathcal{A}_{q,r,p,K,s}(\mathcal{M})$, we consider the energy functional

$$(6.2) \quad \mathcal{F}(u, \zeta, V, \nu) := \int_{\Omega} e(x, u, \zeta, \nu, \nabla u, \nabla \zeta, d\nu) dx + \|V\| + \int_{\mathcal{G}_1(\Omega)} \|A\|^p dV + \|\partial V\|.$$

The energy density $e : \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M} \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2} \times \text{Hom}(\mathbb{R}^2, T\mathcal{M}) \rightarrow \mathbb{R} \cup \{+\infty\}$ is presumed to be

$$e(x, u, \zeta, \nu, F, G, N) = \tilde{e}(x, u, \zeta, \nu, F) + \beta_1 |G|^q + f(x, \nu, N)$$

for every $(x, u, \zeta, \nu, F, G, N) \in \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M} \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2} \times \text{Hom}(\mathbb{R}^2, T\mathcal{M})$, where $\beta_1 > 0$ and the first addendum is a nonnegative Carathéodory function with the following properties:

(a') $\tilde{e}(x, u, \zeta, \nu, F)$ is polyconvex with respect to F , namely,

$$(6.3) \quad \tilde{e}(x, u, \zeta, \nu, F) = g(x, u, \zeta, \nu, M(F)) \quad \forall F \in \mathbb{M}^{3 \times 2},$$

where $g : \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M} \times (\mathbb{M}^{3 \times 2} \times \mathbb{R}^3) \rightarrow [0, +\infty]$ is a Carathéodory function, with $g(x, u, \zeta, \nu, \cdot)$ convex for \mathcal{L}^2 -a.e. $x \in \Omega$ and for all $(u, \zeta, \nu) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M}$;

(b') for \mathcal{L}^2 -a.e. $x \in \Omega$ and every (u, ζ, ν, F) such that $\det(F|\zeta) > 0$,

$$\tilde{e}(x, u, \zeta, \nu, F) \geq \beta_2 (|F|^q + |\text{adj}_2 F|^r) + \Phi(F, \zeta),$$

where $\beta_2 > 0$ and $\Phi \in \tilde{\mathcal{F}}$, a function class specified in Definition 3.1.

Moreover, the third addendum $f : \Omega \times \text{Hom}(\mathbb{R}^2, T\mathcal{M}) \rightarrow [0, +\infty]$ is a nonnegative, continuous, and quasiconvex admissible integrand in $W^{1,s}(\Omega, \mathcal{M})$. We presume it is such that for positive constants $C_1, C_2 > 0$, the inequality

$$C_1 \|N\|_{g(\nu)}^s \leq f(x, \nu, N) \leq C_2 \|N\|_{g(\nu)}^s$$

holds true for every $(x, \nu, N) \in \Omega \times \text{Hom}(\mathbb{R}^2, T\mathcal{M})$.

THEOREM 6.3. *Under previous assumptions, the energy functional $\mathcal{F}(u, V, \zeta, \nu)$ attains a minimum in the class $\mathcal{A}_{q,r,p,K,s}(\mathcal{M})$.*

Proof. Choose a minimizing sequence $\{(u_h, \zeta_h, V^{(h)}, \nu_h)\} \subset \mathcal{A}_{q,r,p,K,s}(\mathcal{M})$. The growth assumptions on f imply that

$$\sup_h \int_{\Omega} \|d(\nu_h)_x\|_{g(\nu_h(x))}^s dx < \infty.$$

By compactness (see [33]), possibly passing to a (not relabeled) subsequence, $\{\nu_h\}$ weakly converges in $W^{1,s}$ to some $\nu \in W^{1,s}(\Omega, \mathcal{M})$, which satisfies $d_{\mathcal{M}}(\nu_h(x), \nu(x)) \rightarrow 0$ for \mathcal{L}^2 -a.e. $x \in \Omega$. Moreover, by exploiting Theorem 6.1, we get

$$\int_{\Omega} f(x, \nu(x), d\nu_x) dx \leq \liminf_{h \rightarrow \infty} \int_{\Omega} f(x, \nu_h(x), d(\nu_h)_x) dx.$$

Then we follow the same path adopted for Theorem 4.3. We omit further details. \square

As to the boundary conditions on the quadruplets (u, ζ, V, ν) in $\mathcal{A}_{q,r,p,K,s}(\mathcal{M})$, we first assume that

$$u = u_0, \quad \zeta = \frac{\partial_1 u_0 \times \partial_2 u_0}{|\partial_1 u_0 \times \partial_2 u_0|}, \quad \nu = \nu_0 \quad \text{on } \partial\Omega,$$

in the sense of traces, for given maps $\nu_0 \in W^{1,s}(\Omega, \mathcal{M})$ and $u_0 \in W^{1,q}(\Omega, \mathbb{R}^3)$, with $\|u_0\|_\infty \leq K$ and $|\text{adj}_2 \nabla u_0| \in L^r(\Omega)$.

Then, we also prescribe a compact set \mathcal{C} contained in Ω such that

$$\text{spt } \mu_\nu \subset \mathcal{C} \quad \forall (u, \zeta, V, \nu) \in \mathcal{A}_{q,r,p,K,s}(\mathcal{M}).$$

In this case, strong anchoring conditions are correspondingly defined in terms of the graph boundary current ∂G_u as in the previous section.

COROLLARY 6.4. *Let \mathcal{C} , u_0 , and ν_0 as above. Assume that competitors (u, V, ζ, ν) in $\mathcal{A}_{q,r,p,K,s}(\mathcal{M})$ have finite energy $\mathcal{F}(u, V, \zeta, \nu)$ and satisfy prescribed boundary, clamping, and confinement conditions, with \mathcal{C} a nonempty set. Then the energy minimum of $\mathcal{F}(u, V, \zeta, \nu)$ is attained in the same class.*

6.2. The SBV case. The through-the-thickness descriptor of the material microstructure ν might jump across the crack margins. So it could be natural to consider $\nu : \Omega \rightarrow \mathcal{M}$ as a special function of bounded variation. However, in this case, we do not have at our disposal a lower-semicontinuity result for manifold-valued BV -maps. Then we restrict ourselves to consider just the total variation of ν distributional derivative among the energy entries and base the pertinent analyses on the closure-compactness theorem proven in [4].

The class of Sobolev maps $\nu \in W^{1,s}(\Omega, \mathcal{M})$ can be equivalently defined in terms of postcomposition with Lipschitz functions $\varphi : \mathcal{M} \rightarrow \mathbb{R}$. By letting

$$\widehat{\mathcal{F}} := \{ \varphi \in \text{Lip}(\mathcal{M}, \mathbb{R}) \mid \text{Lip}(\varphi) \leq 1 \},$$

one requires $\nu \in L^s(\Omega, \mathcal{M})$ and the existence of a nonnegative function $\phi \in L^s(\Omega)$ such that $\varphi \circ \nu \in W^{1,s}(\Omega)$ and $\|\nabla(\varphi \circ \nu)\|_{L^s(\Omega)} \leq \|\phi\|_{L^s(\Omega)}$ for every $\varphi \in \widehat{\mathcal{F}}$. The optimal function $\phi \in L^s(\Omega)$, which satisfies the previous inequality independently of $\varphi \in \widehat{\mathcal{F}}$, agrees \mathcal{L}^m -essentially with the function $|D\nu|$ previously considered.

A summable map $\nu \in L^1(\Omega, \mathcal{M})$ is said to be a function of bounded variation in $BV(\Omega, \mathcal{M})$ if there exists a finite Borel measure μ in Ω such that the total variation of the distributional derivative $D(\varphi \circ \nu)$ is bounded by μ , i.e., $|D(\varphi \circ \nu)|(B) \leq \mu(B)$ for every Borel set $B \subset \Omega$ and every $\varphi \in \widehat{\mathcal{F}}$. The least measure μ satisfying such a property is denoted by $|D\nu|$ (see [4] for further details).

If $\nu \in BV(\Omega, \mathcal{M})$, the countably $(m-1)$ -rectifiable jump set $S(\nu)$ is defined in terms of the jump sets of the BV functions $\varphi \circ \nu$ for a suitable countable set of functions $\varphi \in \widehat{\mathcal{F}}$. The one-sided limits $\nu^\pm(x)$ are correspondingly defined at \mathcal{H}^{m-1} -a.e. $x \in S(\nu)$, once the \mathcal{H}^{m-1} -measurable unit normal $\mathbf{n}_\nu(x)$ to $S(\nu)$ is fixed, as the manifold elements $z^\pm \in \mathcal{M}$ such that the set $\{y \in \Omega \mid d_{\mathcal{M}}(\nu(y), z^\pm) > \epsilon, \langle y - x, \pm \mathbf{n}_\nu(x) \rangle > 0\}$ has 0-density at x for any $\epsilon > 0$.

In a similar way, we define a nonnegative function $|\nabla \nu| \in L^1(\Omega)$ in terms of $\sup |\nabla(\varphi \circ \nu)|$, where $\nabla(\varphi \circ \nu)$ denotes the approximate gradient of the BV function $\varphi \circ \nu$. Then $|\nabla \nu|$ agrees with the Radon–Nikodym derivative of $|D\nu|$ with respect to $\mathcal{L}^m \llcorner \Omega$, so that for \mathcal{L}^m -a.e. $x \in \Omega$, we compute the approximate limit

$$\text{ap } \lim_{y \rightarrow x} \frac{d_{\mathcal{M}}(\nu(y), \nu(x))}{|y - x|} = |\nabla \nu|(x).$$

A function $\nu \in BV(\Omega, \mathcal{M})$ is said to be a special function of bounded variation if $\varphi \circ \nu \in SBV(\Omega)$ for every $\varphi \in \widehat{\mathcal{F}}$. If $\nu \in SBV(\Omega, \mathcal{M})$, for every Borel set $B \subset \Omega$, we get the decomposition formula

$$|D\nu|(B) = \int_B |\nabla \nu| dx + \int_{B \cap S(\nu)} d_{\mathcal{M}}(\nu^+, \nu^-) d\mathcal{H}^{m-1}.$$

THEOREM 6.5 (see [4]). For $\mathcal{K} \subset \mathcal{M}$ a compact set, consider $\{\nu_h\} \subset SBV(\Omega, \mathcal{M})$ such that $\nu_h(\Omega) \subset \mathcal{K}$ for every h . If for some exponent $s > 1$

$$(6.4) \quad \sup_h \left(\int_{\Omega} |\nabla \nu_h|^s dx + \int_{S(\nu_h)} (1 + d_{\mathcal{M}}(\nu_h^+, \nu_h^-)) d\mathcal{H}^{m-1} \right) < \infty,$$

there exists a (not relabeled) subsequence of $\{\nu_h\}$ converging \mathcal{L}^m -a.e. in Ω to some $\nu \in SBV(\Omega, \mathcal{M})$ and such that

$$\int_{\Omega} |\nabla \nu|^s dx \leq \liminf_{h \rightarrow \infty} \int_{\Omega} |\nabla \nu_h|^s dx$$

$$\int_{S(\nu)} d_{\mathcal{M}}(\nu^+, \nu^-) d\mathcal{H}^{m-1} \leq \liminf_{h \rightarrow \infty} \int_{S(\nu_h)} d_{\mathcal{M}}(\nu_h^+, \nu_h^-) d\mathcal{H}^{m-1}.$$

In the case considered here, we take first $q > 2, r, p, s > 1, K > 0$. Then we consider a compact set \mathcal{K} in \mathcal{M} , where $\mathcal{M} = (\mathcal{M}^n, g)$ is given as above. We say that $(u, \zeta, V, \nu) \in \tilde{\mathcal{A}}_{q,r,p,K,s,\mathcal{K}}(\mathcal{M})$ if the triplet (u, ζ, V) belongs to the class $\mathcal{A}_{q,r,p,K}$ previously introduced and $\nu \in SBV(\Omega, \mathcal{M})$ satisfies the following relations:

1. $\nu(\Omega) \subset \mathcal{K}$ with $|\nabla \nu| \in L^s(\Omega)$;
2. $\mathcal{H}^1 \llcorner S(\nu) \leq \mu_V$.

Assumption 1 is of a technical nature, whereas assumption 2 means that ν jumps only over the fracture.

Thus, we consider the energy to be

$$(6.5) \quad \tilde{\mathcal{F}}(u, \zeta, V, \nu) := \int_{\Omega} e(x, u, \zeta, \nu, \nabla u, \nabla \zeta, |\nabla \nu|) dx + \|V\| + \int_{\mathcal{G}_1(\Omega)} \|A\|^p dV + \|\partial V\|.$$

The density $e : \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M} \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2} \times \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ is such that

$$(6.6) \quad e(x, u, \zeta, \nu, F, G, N) = \tilde{e}(x, u, \zeta, \nu, F) + \beta_1 |G|^q + \beta_3 |N|^s$$

for every $(x, u, \zeta, \nu, F, G, N) \in \Omega \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathcal{M} \times \mathbb{M}^{3 \times 2} \times \mathbb{M}^{3 \times 2} \times \mathbb{R}$, where $\beta_1 > 0$, $\beta_3 > 0$, and the first addendum is a nonnegative Carathéodory function satisfying the properties (a') and (b') above introduced when $\nu \in W^{1,s}(\Omega, \mathcal{M})$.

THEOREM 6.6. The minimum of the energy functional $\tilde{\mathcal{F}}(u, \zeta, V, \nu)$ given by (6.5) is attained in the class $\tilde{\mathcal{A}}_{q,r,p,K,s,\mathcal{K}}(\mathcal{M})$.

Proof. Take a minimizing sequence $\{(u_h, \zeta_h, V^{(h)}, \nu_h)\} \subset \tilde{\mathcal{A}}_{q,r,p,K,s,\mathcal{K}}(\mathcal{M})$. Since $\mathcal{H}^1 \llcorner S(\nu_h) \leq \mu_{V^{(h)}}$ and $\nu_h(\Omega) \subset \mathcal{K}$ for each h , we have $d_{\mathcal{M}}(\nu_h^+, \nu_h^-) \leq c(\mathcal{K}) < \infty$ for \mathcal{H}^1 -a.e. $x \in S(\nu_h)$ and for each h , where $c(\mathcal{K})$ is a real constant. Therefore, since $\beta_3 > 0$ in formula (6.6), the uniform bound (6.4) holds, and we can apply Theorem 6.2. The proof then goes along the same path followed in Theorem 4.3. Also, the limit point $\nu \in SBV(\Omega, \mathcal{M})$ satisfies conditions 1–2. We omit further details. \square

Under prescribed Dirichlet or strong anchoring conditions for the triplets (u, V, ζ) as given previously, if the corresponding class of competitors (u, V, ζ, ν) in $\tilde{\mathcal{A}}_{q,r,p,K,s,\mathcal{K}}(\mathcal{M})$ with finite energy (6.5) is nonempty, the energy attains its minimum in the same class. Notice that a Dirichlet condition on the through-the-thickness descriptor ν is not preserved by the weak convergence in Theorem 6.2.

Remark 6.1. In order to consider cohesive effects instead of looking at brittle fracture, as we have done here so far, we could add to the energy $\tilde{\mathcal{F}}(u, \zeta, V, \nu)$, with density (6.6), a term of the type

$$\nu \mapsto \beta_4 \int_{S(\nu)} \delta_{\mathcal{M}}(\nu^+, \nu^-) d\mathcal{H}^1, \quad \beta_4 \geq 0.$$

On the other hand, in general a term of the type

$$\int_{S(\nu)} g(\nu^+, \nu^-, \mathbf{n}_\nu) d\mathcal{H}^{m-1},$$

with $g: \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{S}^{m-1} \rightarrow [0, +\infty)$ a suitable “jointly convex” function, is presently hard to be treated when we consider \mathcal{M} as intrinsic manifold not necessarily coinciding with a linear space.

Remark 6.2. Finally, a lower-semicontinuity result necessary to analyze the *SBV* case when $D\nu$ enters the energy density instead of its total variation could perhaps be reached by adopting techniques presented in [31]. However, we do not tackle the problem here, leaving it open.

Remark 6.3. Our analysis does not exclude that distant portions of the shell might interpenetrate. Avoiding it would require to account for the shell thickness and to prescribe a pertinent bound involving it. We leave the analysis to a further ongoing work.

Acknowledgments. We thank Emilio Acerbi for very useful discussions. We also thank the Editor and two anonymous reviewers for their constructive feedback.

REFERENCES

- [1] W. K. ALLARD, *On the first variation of a varifold*, Ann. Math., 95 (1972), pp. 417–491.
- [2] W. K. ALLARD, *On the first variation of a varifold: Boundary behavior*, Ann. Math., 101 (1975), pp. 418–446.
- [3] F. J. ALMGREN, JR., *Theory of Varifolds*, mimeographed notes, Princeton University, 1965.
- [4] L. AMBROSIO, *Metric space valued functions of bounded variation*, Ann. Sc. Norm. Sup. Pisa Cl. Sci., 17 (1990), pp. 439–478.
- [5] L. AMBROSIO, N. FUSCO, AND D. PALLARA, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, Oxford, 2000.
- [6] S. S. ANTMAN, *Nonlinear Problems of Elasticity*, Springer-Verlag, Berlin, 1995.
- [7] J. M. BALL, *Convexity conditions and existence theorems in nonlinear elasticity*, Arch. Ration. Mech. Anal., 63 (1976/77), pp. 337–403.
- [8] M. BARHAM, D. J. STEIGMANN, AND D. WHITE, *Magnetoelasticity of highly deformable thin films: Theory and simulation*, Int. J. Non-Linear Mech., 47 (2012), pp. 185–196.
- [9] K. BHATTACHARYA AND R. D. JAMES, *A theory of thin films of martensitic materials with applications to microactuators*, J. Mech. Phys. Solids, 47 (1999), pp. 531–576.
- [10] B. BOURDIN, G. A. FRANCFORT, AND J.-J. MARIGO, *Numerical experiments in revisited brittle fracture*, J. Mech. Phys. Solids, 47 (2000), pp. 797–826.
- [11] G. CAPRIZ, *Continua with Microstructure*, Springer-Verlag, Berlin, 1989.
- [12] D. CHIRON, *On the definitions of Sobolev and BV spaces into singular spaces and the trace problem*, Commun. Contemp. Math., 9 (2007), pp. 473–513.
- [13] P. G. CIARLET, *Mathematical Elasticity II. Theory of Plates*, North-Holland, Amsterdam, 1997.
- [14] P. CIARLET, R. GOGU, AND C. MARDARE, *Orientation-preserving condition and polyconvexity on a surface: Application to nonlinear shell theory*, J. Math. Pures Appl., 99 (2013), pp. 704–725.
- [15] G. DAL MASO AND R. TOADER, *A model for the quasi-static growth of brittle fractures: Existence and approximation results*, Arch. Ration. Mech. Anal., 162 (2002), pp. 101–135.
- [16] E. DE GIORGI, *New problems on minimizing movements*, in Ennio De Giorgi—Selected Papers, L. Ambrosio, G. Dal Maso, M. Forti, M. Miranda, and S. Spagnolo, eds., Springer-Verlag, Berlin, 2006, pp. 699–713.
- [17] J. L. ERICKSEN AND C. A. TRUESDELL, *Exact theory of stress and strain in rods and shells*, Arch. Ration. Mech. Anal., 1 (1958), pp. 295–323.

- [18] M. FOCARDI AND E. N. SPADARO, *An intrinsic approach to manifold constrained variational problems*, Ann. Mat. Pura Appl., 192 (2013), pp. 145–163.
- [19] G. A. FRANCFORT AND J. J. MARIGO, *Revisiting brittle fracture as an energy minimization problem*, J. Mech. Phys. Solids, 46 (1998), pp. 1319–1342.
- [20] G. FRIESECKE AND R. D. JAMES, *A scheme for the passage from atomic to continuum theory of thin films, nanotubes and nanorods*, J. Mech. Phys. Solids, 48 (2000), pp. 1519–1540.
- [21] G. FRIESECKE, R. D. JAMES, AND S. MÜLLER, *A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity*, Comm. Pure Appl. Math., 55 (2002), pp. 1461–1506.
- [22] G. FRIESECKE, R. D. JAMES, AND S. MÜLLER, *A hierarchy of plate models derived from nonlinear elasticity by gamma-convergence*, Arch. Ration. Mech. Anal., 180 (2006), pp. 183–236.
- [23] H. J. FROST, *Microstructural evolution in thin films*, Mater. Charact., 4 (1994), pp. 257–273.
- [24] N. FUSCO, C. LEONE, R. MARCH, AND A. VERDE, *A lower semi-continuity result for polyconvex functionals in SBV*, Proc. R. Soc. Edinburgh Sect. A, 136 (2006), pp. 321–336.
- [25] M. GIAQUINTA, P. M. MARIANO, AND G. MODICA, *A variational problem in the mechanics of complex materials*, Discrete Contin. Dyn. Syst., 28 (2010), pp. 519–537.
- [26] M. GIAQUINTA, P. M. MARIANO, AND G. MODICA, *Stress constraints in simple bodies undergoing large strain: A variational approach*, Proc. R. Soc. Edinburgh Sect. A, 145 (2015), pp. 1–32.
- [27] M. GIAQUINTA, P. M. MARIANO, G. MODICA, AND D. MUCCI, *Ground states of simple bodies that may undergo brittle fracture*, Phys. D, 239 (2010), pp. 1485–1502.
- [28] M. GIAQUINTA, G. MODICA, AND J. SOUČEK, *Cartesian Currents in the Calculus of Variations*, Vols. 1 and 2, Springer-Verlag, Berlin, 1998.
- [29] A. A. GRIFFITH, *The phenomena of rupture and flow in solids*, Philos. Trans. R. Soc. A, 221 (1920), pp. 163–198.
- [30] M. GROSSER, H. SEIDEL, AND U. SCHMID, *Microstructure and mechanical properties of sputter deposited tantalum nitride thin films after high temperature loading*, Thin Solid Films, 629 (2017), pp. 69–78.
- [31] L. IGNAT AND X. LAMY, *Lifting of RP^{d-1} -valued maps in BV and applications to uniaxial Q -tensors. With an appendix on an intrinsic BV-energy for manifold-valued maps*, Calc. Var. Partial Differential Equations, 58 (2019), 68.
- [32] R. D. JAMES AND R. RIZZONI, *Pressurized shape memory thin films*, J. Elasticity, 59 (2000), pp. 399–436.
- [33] N. J. KOREVAAR AND R. M. SCHOEN, *Global existence theorems for harmonic maps to non-locally compact spaces*, Comm. Anal. Geom., 5 (1997), pp. 333–387.
- [34] M. KRUŽIK AND J. ZIMMER, *Modelling of thin martensitic films with nonpolynomial stored energies*, Math. Stat., 35 (2013), pp. 587–608.
- [35] C. H. LEI, Y. LIU, AND W. CHEN, *Formations and evolutions of martensitic tents and tunnels in shape memory alloy thin films*, Mech. Mater., 134 (2019), pp. 9–17.
- [36] C. MANTEGAZZA, *Curvature varifolds with boundary*, J. Differential Geom., 43 (1996), pp. 807–843.
- [37] P. M. MARIANO, *Multifield theories in mechanics of solids*, Adv. Appl. Mech., 38 (2002), pp. 1–93.
- [38] P. M. MARIANO, *Physical significance of the curvature varifold-based description of crack nucleation*, Rend. Lincei, 21 (2010), pp. 215–233.
- [39] P. M. MARIANO, *Trends and challenges in the mechanics of complex materials: A view*, Philos. Trans. Roy. Soc. A, 374 (2016), 20150341.
- [40] P. M. MARIANO, *Equilibrium configurations of mixture thin films undergoing large strains*, Math. Methods Appl. Sci., 41 (2018), pp. 479–489.
- [41] P. M. MARIANO AND F. L. STAZI, *Strain localization in elastic microcracked bodies*, Comput. Methods Appl. Mech. Engrg., 190 (2001), pp. 5657–5677.
- [42] C. MIEHE, F. ALDAKHEEL, AND S. TEICHTMEISTER, *Phase-field modeling of ductile fracture at finite strains: A robust variational-based numerical implementation of a gradient-extended theory by micromorphic regularization*, Int. J. Numer. Methods Eng., 111 (2017), pp. 816–863.
- [43] P. NEFF, *A geometrically exact Cosserat shell-model including size effects, avoiding degeneracy in the thin shell limit. Part I: Formal dimensional reduction for elastic plates and existence of minimizers for positive Cosserat couple modulus*, Contin. Mech. Thermodyn., 16 (2004), pp. 577–628.
- [44] P. NEFF, *Local existence and uniqueness for a geometrically exact membrane-plate with viscoelastic transverse shear resistance*, Math. Methods Appl. Sci., 28 (2005), pp. 1031–1060.

- [45] Y. G. RESHETNYAK, *Sobolev classes of functions with values in a metric space*, Siberian Math. J., 38 (1997), pp. 567–583.
- [46] Y. G. RESHETNYAK, *Sobolev classes of functions with values in a metric space. II*, Siberian Math. J., 45 (2004), pp. 709–721.
- [47] Y. G. RESHETNYAK, *On the theory of Sobolev classes of functions with values in a metric space*, Siberian Math. J., 47 (2006), pp. 117–134.
- [48] R. SEGEV, *A geometrical framework for the static of materials with microstructure*, Math. Models Methods Appl. Sci., 4 (1994), pp. 871–897.
- [49] Y. C. SHU, *Heterogeneous thin films of martensitic materials*, Arch. Ration. Mech. Anal., 153 (2000), pp. 39–90.
- [50] J. C. SIMO AND D. D. FOX, *On a stress resultant geometrically exact shell model. I. Formulation and optimal parametrization*, Comput. Methods Appl. Mech. Engrg., 72 (1989), pp. 267–304.
- [51] J. C. SIMO AND N. TARNOW, *A new energy and momentum conserving algorithm for the non-linear dynamics of shells*, Int. J. Numer. Methods Eng., 37 (1994), pp. 2527–2549.
- [52] J. C. SIMO, J. E. MARSDEN, AND P. S. KRISHNAPRASAD, *The Hamiltonian structure of non-linear elasticity: The material and convective representations of solids, rods, and plates*, Arch. Ration. Mech. Anal., 104 (1988), pp. 125–183.
- [53] R. SPATSCHEK AND E. A. BRENER, *Grinfeld instability on crack surfaces*, Phys. Rev. E, 64, (2001), 046120.
- [54] D. J. STEIGMANN, *Fluid films with curvature elasticity*, Arch. Ration. Mech. Anal., 150 (1999), pp. 127–152.
- [55] D. J. STEIGMANN, *Mechanics of materially uniform thin films*, Mat. Mech. Solids, 20 (2015), pp. 309–326.
- [56] P. VILLAGGIO, *Mathematical Models for Elastic Structures*, Cambridge University Press, Cambridge, 1997.