



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Heterocoumarins Are Selective Carbonic Anhydrase IX and XII Inhibitors with Cytotoxic Effects against Cancer Cells Lines

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Heterocoumarins Are Selective Carbonic Anhydrase IX and XII Inhibitors with Cytotoxic Effects against Cancer Cells Lines / Angeli, Andrea; Trallori, Elena; Carta, Fabrizio; Di Cesare Mannelli, Lorenzo; Ghelardini, Carla; Supuran, Claudiu T.*. - In: ACS MEDICINAL CHEMISTRY LETTERS. - ISSN 1948-5875. - ELETTRONICO. - 9:(2018), pp. 947-951. [10.1021/acsmchemlett.8b00362]

Availability:

This version is available at: 2158/1141181 since: 2018-11-13T10:31:47Z

Published version:

DOI: 10.1021/acsmchemlett.8b00362

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Heterocoumarins Are Selective Carbonic Anhydrase IX and XII Inhibitors with Cytotoxic Effects against Cancer Cells Lines

Andrea Angeli,[†] Elena Trallori,[‡] Fabrizio Carta,[†] Lorenzo Di Cesare Mannelli,[‡] Carla Ghelardini,[‡] and Claudiu T. Supuran^{*,†}


[†]University of Florence, NEUROFARBA Dept., Sezione di Scienze Farmaceutiche, Via Ugo Schiff 6, 50019 Sesto Fiorentino, Florence, Italy

[‡]NEUROFARBA Department, Section of Pharmacology and Toxicology, Università degli Studi di Firenze, Viale Pieraccini 6, 50139 Florence, Italy

Supporting Information

ABSTRACT: We have synthesized a new series of coumarin-based compounds demonstrating high selectivity and potent effects with low nanomolar affinity against the tumor associated carbonic anhydrase (CA, EC 4.2.1.1) isoforms hCA IX and XII. A number of these compounds were evaluated *ex vivo* against human prostate (PC3) and breast (MDA-MB-231) cancer cell lines. Compounds **4b** and **15** revealed effective cytotoxic effects after 48 h of incubation in both normoxic and hypoxic conditions with PC3 cancer cell line. However, compound **3** showed selective cytotoxic effects against MDA-MB-231 in hypoxic condition. These results may be of particular importance for the choice of future drug candidates targeting hypoxic tumors and metastases, considering the fact that a selective carbonic anhydrase CA IX inhibitor (SLC-0111) is presently in phase II clinical trials.

KEYWORDS: Carbonic anhydrase inhibitors (CAIs), selenium, metalloenzymes, coumarin, tumor, tellurium



The coumarin scaffold is found in various natural products and bioactive compounds. For this reason, it is a biologically important and highly privileged structure also in the pharmaceutical field.¹ Coumarin derivatives showed a large number of biological activities, including anticoagulant,² antibacterial,³ antiviral,⁴ and recently many different derivatives have been reported as antitumor with interesting cytotoxic activity *in vitro* and *in vivo*.^{5,6,23} The interesting heterocyclic system of coumarin derivatives, with their wide range of biological functions, has made them an excellent starting point for further chemical derivatization to identify novel therapeutic agents. In this particular contest, recently, several coumarin derivatives have been shown to constitute a new class of selective inhibitors against the human tumor-associated carbonic anhydrase (CA, EC 4.2.1.1) isoforms hCA IX and hCA XII.^{7,8} These two isoforms are active extracellular enzymes involved in metabolism of CO₂, and they have been implicated in acidification of extracellular microenvironment and, at the same time, in protection of cancer cells from the acidosis.⁹ For this reason, we developed potent and selective inhibitors from synthetic source with the aim to identify new CAIs structurally related to natural products as coumarins. In the last years, we have been engaged in the development of new synthetic methodologies involving different seleno functionalization of privileged structures^{10,11} including their possible medical applications.^{12,13} In continuation of our research on the preparation of potentially useful chalcogen derivatives, herein

we report a facile and convenient access to synthetic strategies for the replacement of the endocyclic oxygen atom by selenium provides new heterocoumarins. The selenophenyl cinnamate (**2a–b**) needed for the synthesis of selenocoumarins was prepared by reduction of diselenides **1a–b** with NaBH₄ to the corresponding selenolate, which was treated *in situ* with cinnamoyl chloride, affording the compounds **2a–b** in excellent yield. Thus, in order to synthesize selenocoumarins **3a–b**, derivatives **2a–b** were treated with anhydrous AlCl₃ in chlorobenzene at 100 °C for 30 min to afford a red reaction mixture, which, after workup, furnished compounds **3a–b** in rather good yield,¹⁴ as reported in Scheme 1.

In order to access further heterocoumarins, we evaluated the possibility to replace the carbonyl group with an isosteric thiocarbonylic moiety to employ an excess of Lawesson's reagent (2 equiv) in refluxing toluene for 4 h. Compounds **4a–b** were obtained in excellent yields as reported in Scheme 2.


We continued the possibility to replace exocyclic oxygen atom by selenium, but this time, our effort did not give the desired results. Thus, our study has been extended to the nitrogen endocyclic system. Cinnamoyl amides, differently substituted, **6a–f** were obtained from the corresponding amines **5a–f** with cinnamoyl chloride in dichloromethane at

Received: August 10, 2018


Accepted: August 29, 2018

Published: August 29, 2018

Scheme 1. Synthesis of Selenocoumarins 3a–b




Scheme 2. Synthesis of Thioseleno Coumarins 4a–b



room temperature in quantitative yields. Treatment of compounds **6a–f** with anhydrous AlCl₃ in chlorobenzene at 100 °C for 5 h afforded the corresponding quinolin-2(1H)-one derivatives **7a–f** with similar mechanistic pathway of compounds **3a–b**. Moreover, the reaction proceeded efficiently with substituents on the nitrogen (**7b**) or on the aryl group (**7a,c–f**) as reported in Scheme 3.¹⁵


Scheme 3. Synthesis of Different Quinolin-2(1H)-one Derivatives 7a–f



In order to access an exocyclic selenium, this time, the isosteric replacement of oxygen was possible in a one-pot selenation reaction with Wollins' reagent. Compounds **7a–b** were readily converted to seleno quinolin-2(1H)-one derivatives **8a–b** with an excess of WR (1.5 equiv) in refluxing toluene for 4 h affording in good yield the desiderate compounds as outlined in Scheme 4.


As a further investigation, we tried to synthesize with a similar mechanism of reaction also the endocyclic replacement of oxygen atom by tellurium; however, it proved to be difficult because telluro esters were more instable. In order to propose an alternative way to access tellurocoumarin, we sought to

Scheme 4. Synthesis of Seleno Quinolin-2(1H)-one Derivatives 8a–b



achieve it from cinnamic acid derivative **14**, thus avoiding the synthesis of telluro esters. The aldehyde group of compound **9** was protected by acetalization with triethyl orthoformate in refluxing ethanolic solution. Diaryl ditelluride **11** was prepared from aryl bromide **10** via lithiation, tellurium insertion, and oxidation in analogy with a literature procedure.¹⁶ The ditelluride acetal derivative **11** was then hydrolyzed with concentrated HCl, and the resulting aldehyde **13** was condensed with a Knoevenagel reaction with malonic acid to afford the corresponding cinnamoyl acid derivative **14**. Finally, cyclization into tellurocoumarin **15** was obtained by converting compound **14** *in situ* in the corresponding acyl chloride and subsequently treated with AlCl₃ in dichloromethyl methyl ether as reported previously in the literature¹⁷ and outlined in Scheme 5.

Scheme 5. Synthesis of Tellurocoumarin 15



All synthesized compounds, **3a–b**, **4a–b**, **7a–f**, **8a–b** and **15**, were tested *in vitro* for their inhibitory properties against the physiologically relevant hCA isoforms (I, II, IX, and XII) by means of a stopped-flow carbon dioxide hydration assay¹⁸ after a period of 6 h of incubation of the enzyme and inhibitor solutions.^{19–21} Their activities were compared to the standard carbonic anhydrase inhibitor (CAI) acetazolamide (AAZ) (Table 1).

From the analysis of constants of inhibition values reported in Table 1, we observed that all compounds, according to previous reports,^{19–21} were ineffective inhibitors of two dominant cytosolic hCA I and hCA II showing high selectivity against tumor-associated isoforms hCA IX and hCA XII. In terms of structure–activity relationships, it could be observed that methyl moiety in position 6 of seleno-coumarins **3a–b** and **4a–b** played a crucial role for activity. This moiety increased near two times the potency against hCA IX and over three times for hCA XII. Moreover, the isosteric substitution of carbonyl with a thiocarbonyl group did not influence significantly the activity. Methyl substituent on nitrogen of quinolin-2(1H)-one **7b** led to an increase in the activity two-fold against hCA IX compared to compound **7a** (K_i 44 nM to 82 nM). However, different substituents in position 8 (**7c–e**) increase the potency for both tumor-associated isoforms. The replacement of carbonyl group with selenocarbonyl moiety (**8a–b**), this time, influenced the activity especially for compound **8b**, showing a decrease of near four times that of compound **7b** (K_i 44 nM to 172 nM). Finally, tellurocoumarin **15** showed an interesting inhibition pattern compared to

Table 1. Inhibition Data of Human CA Isoforms I, II, IX, and XII with Compounds 3a–b, 4a–b, 7a–f, 8a–b, 15, and AAZ by a Stopped Flow CO₂ Hydrase Assay¹⁸

compd	K_i (nM) ^a			
	hCA I	hCAII	hCA IX	hCA XII
3a	>10000	>10000	26.3	22.9
3b	>10000	>10000	56.1	7.6
4a	>10000	>10000	39.5	25.8
4b	>10000	>10000	59.9	7.4
7a	>10000	>10000	82.0	93.0
7b	>10000	>10000	44.6	83.5
7c	>10000	>10000	23.0	6.8
7d	>10000	>10000	58.8	6.8
7e	>10000	>10000	23.4	8.4
7f	>10000	>10000	123.1	8.7
8a	>10000	>10000	54.4	7.6
8b	>10000	>10000	172.0	86.4
15	>10000	>10000	59.0	8.2
AAZ	250.0	12.1	25.8	5.7

^aMean from three different assays, by a stopped flow technique (errors were in the range of ± 5 –10% of the reported values).

selenocoumarin **3a**. The biggest chalcogen atom for compound **15** led to a decrease in potency near two times **3a** for hCA IX (K_i 59 nM to 26.3 nM), but for the other membrane isoform hCA XII, the activity increased near three times that of selenocoumarin (K_i 8.2 nM to 22.9 nM), thus showing tellurocoumarin is more selective against hCA XII.

The high isoform selectivity and upregulated expression of hCA IX and XII in a wide selection of hypoxic tumors make these compounds a desirable feature for compounds designed to target the tumor-associated enzymes. We focused our attention on the *ex vivo* activity of compounds **3a**, **4b**, **8a**, and **15**, which were evaluated for their effects on cell viability against the human prostatic (PC3) and breast (MDA-MB-231) cancer cell lines. All compounds were highly selective hCA IX and XII inhibitors and were used at different concentrations, being incubated for 48 h in both normoxic and hypoxic conditions, when overexpression of high amounts of CA IX occurs.²² In PC3 cells, selenoquinolin-2(1*H*)-one **8a** did not show any cytotoxic activity in normoxic and hypoxic conditions. However, selenocoumarin **3a** started to reduce the cell viability only at higher concentration (300 μ M) to 68% in both conditions. Its efficacy increased significantly when selenium endocyclic was replaced by tellurium; this isosteric substitution, in normoxic condition, led to a reduced viability of 54% already at 30 μ M and increased to 19% at higher concentrations (Figure 1). In the hypoxic condition, indeed, compound **15** showed less effect on cytotoxicity, which reached 71% at 30 μ M and 38% at 100–300 μ M. Moreover, the thioselenocoumarin derivative **4b** showed at 30 μ M similar cytotoxicity effect to compound **15** in normoxic and hypoxic conditions (51% and 79%, respectively).

Derivative **8a** also in the MDA-MB231 cell line did not show any activity in these *ex vivo* normoxia and hypoxia assays. However, seleno-coumarin **3a** showed only at 300 μ M a good cytotoxic activity in normoxic condition (28%). The potency against this cancer cell line increased drastically when compound **3a** was used in hypoxic condition. Indeed, this compound reduced cell viability already by more than 50% at 30 μ M and arrived to kill over 90% at 300 μ M (Figure 2), showing, thus, an interesting selectivity against this specific




Figure 1. Effects of newly synthesized compounds **3a**, **4b**, **8a**, and **15** on viability of the human prostatic cancer cell line PC3 following 48 h treatment in normoxic and hypoxic (1% O₂) conditions. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ versus control.

cancer cell line in hypoxic condition. Also this time, tellurocoumarin **15** exhibited a strong cytotoxicity in normoxic conditions, already at 30 μ M (15.9%). The potency decreased over three times in hypoxic condition, reducing the cell viability to 22% with a concentration of 300 μ M. A reduced cell viability (61%) was observed also for compound **4b** only at 30 μ M. In the hypoxic condition, this compound did not show any significant activity.

In conclusion, we report a new series of different chalcogenocoumarins and quinolin-1(2*H*)-one. These compounds were evaluated for their inhibitory properties against the two dominant cytosolic isoforms hCA I and II and the tumor associated isoforms hCA IX and XII. All derivatives, here investigated, were ineffective against the off-target cytosolic hCA I and II, whereas they showed interesting selective and potent inhibition profiles (in the low nanomolar range) against the tumor associated isoforms hCA IX and XII. For this reason, different compounds were evaluated *ex vivo* against two cancer cell lines PC3 and MDA-MB231. In particular, compounds **3a** and **15** revealed an interesting cytotoxic effect after 48 h of incubation. These results may be of particular importance for the choice of future drug candidates targeting hypoxic tumors.




Figure 2. Effects of the newly synthesized compounds **3a**, **4b**, **8a**, and **15** on viability of the human adenocarcinoma breast cell line MDA-MB231 following 48 h treatment in normoxic and hypoxic (1% O₂) conditions. ** $p < 0.01$, *** $p < 0.001$ versus control.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acsmchemlett.8b00362](https://doi.org/10.1021/acsmchemlett.8b00362).

Synthetic procedures, characterization of compounds, in vitro kinetic procedure, and biological assay (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*(C.T.S.) Tel/Fax: +39-055-4573729. E-mail: claudiu.supuran@unifi.it.

ORCID

Fabrizio Carta: 0000-0002-1141-6146

Claudio T. Supuran: 0000-0003-4262-0323

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ABBREVIATIONS

CA, carbonic anhydrases; AAZ, acetazolamide

■ REFERENCES

- (1) Kontogiorgis, C.; Detsi, A.; Hadjipavlou-Litina, D. Coumarin-based drugs: a patent review (2008 – present). *Expert Opin. Ther. Pat.* **2012**, *22*, 437–454.
- (2) Kidane, A. G.; Salacinski, H.; Tiwari, A.; Bruckdorfer, K. R.; Seifalian, A. M. Anticoagulant and Antiplatelet Agents: Their Clinical and Device Application(s) Together with Usages to Engineer Surfaces. *Biomacromolecules* **2004**, *5*, 798–813.
- (3) Appendino, G.; Mercalli, E.; Fuzzati, N.; Arnoldi, L.; Stavri, M.; Gibbons, S.; Ballero, M.; Maxia, A. Antimycobacterial Coumarins from the Sardinian Giant Fennel (*Ferula communis*). *J. Nat. Prod.* **2004**, *67*, 2108–2110.
- (4) Ma, T.; Liu, L.; Xue, H.; Li, L.; Han, C.; Wang, L.; Chen, Z.; Liu, G. Chemical Library and Structure–Activity Relationships of 11-Demethyl-12-oxo Calanolide A Analogues as Anti-HIV-1 Agents. *J. Med. Chem.* **2008**, *51*, 1432–1446.
- (5) Kaur, M.; Kohli, S.; Sandhu, S.; Bansal, Y.; Bansal, G. Coumarin: a promising scaffold for anticancer agents. *Anti-Cancer Agents Med. Chem.* **2015**, *15*, 1032–1048.
- (6) Lai, Y.; Long, Y.; Lei, Y.; Deng, X.; He, B.; Sheng, M.; Li, M.; Gu, Z. A novel micelle of coumarin derivative monoend-functionalized PEG for anti-tumor drug delivery: in vitro and in vivo study. *J. Drug Target* **2012**, *3*, 246–254.
- (7) Maresca, A.; Temperini, C.; Vu, H.; Pham, N. B.; Poulsen, S. A.; Scozzafava, A.; Quinn, R. J.; Supuran, C. T. Non-zinc mediated inhibition of carbonic anhydrases: coumarins are a new class of suicide inhibitors. *J. Am. Chem. Soc.* **2009**, *131*, 3057–62.
- (8) Bonardi, A.; Falsini, M.; Catarzi, D.; Varano, F.; Di Cesare Mannelli, L.; Tenci, B.; Ghelardini, C.; Angeli, A.; Supuran, C. T.; Colotta, V. Structural investigations on coumarins leading to chromeno[4,3-c]pyrazol-4-ones and pyrano[4,3-c]pyrazol-4-ones: New scaffolds for the design of the tumor-associated carbonic anhydrase isoforms IX and XII. *Eur. J. Med. Chem.* **2018**, *146*, 47–59.
- (9) Supuran, C. T. Inhibition of carbonic anhydrase IX as a novel anticancer mechanism. *World J. Clin. Oncol* **2012**, *7*, 98–103.
- (10) Angeli, A.; Tanini, D.; Peat, T. S.; Di Cesare Mannelli, L.; Bartolucci, G.; Capperucci, A.; Ghelardini, C.; Supuran, C. T.; Carta, F. Discovery of New Selenoureido Analogues of 4-(4-Fluorophenylureido)benzenesulfonamide as Carbonic Anhydrase Inhibitors. *ACS Med. Chem. Lett.* **2017**, *8*, 963–968.
- (11) Angeli, A.; Tanini, D.; Viglianisi, C.; Panzella, L.; Capperucci, A.; Menichetti, S.; Supuran, C. T. Evaluation of selenide, diselenide and selenoheterocycle derivatives as carbonic anhydrase I, II, IV, VII and IX inhibitors. *Bioorg. Med. Chem.* **2017**, *25*, 2518–2523.
- (12) Angeli, A.; Di Cesare Mannelli, L.; Trallori, E.; Peat, T. S.; Ghelardini, C.; Carta, F.; Supuran, C. T. Design, Synthesis, and X-ray of Selenides as New Class of Agents for Prevention of Diabetic Cerebrovascular Pathology. *ACS Med. Chem. Lett.* **2018**, *9*, 462–467.
- (13) Angeli, A.; Di Cesare Mannelli, L.; Lucarini, E.; Peat, T. S.; Ghelardini, C.; Supuran, C. T. Design, synthesis and X-ray crystallography of selenides bearing benzenesulfonamide moiety with neuropathic pain modulating effects. *Eur. J. Med. Chem.* **2018**, *154*, 210–219.
- (14) Jayachandran, T.; Manimaran, T.; Ramakrishnan, V. T. Synthesis of Selenacoumarins. *Indian J. Chem.* **1984**, *23B*, 328–330.
- (15) Natarajan, M.; Ramakrishnan, V. T. A new route for the synthesis of Coumarins, Thiocoumarins and Carbostyryls. *Indian J. Chem.* **1984**, *23B*, 720–727.
- (16) Poon, J.; Yan, J.; Singh, V. P.; Gates, P. J.; Engman, L. Alkyltelluro Substitution Improves the Radical-Trapping Capacity of Aromatic Amines. *Chem. - Eur. J.* **2016**, *22*, 12891–12903.
- (17) Christiaens, L.; Piette, J. L.; Luxen, A.; Renson, M. 2H-[1]-Benzotellurinnone-2 (telluro-1 coumarine) et Dihydro-3,4-chalcogeno-1 coumarines. *J. Heterocyclic Chem.* **1984**, *21*, 1281.

(18) Khalifah, R. G. The carbon dioxide hydration activity of carbonic anhydrase. I. Stop flow kinetic studies on the native human isoenzymes B and C. *J. Biol. Chem.* **1971**, *246*, 2561.

(19) De Luca, L.; Mancuso, F.; Ferro, S.; Buemi, M. R.; Angeli, A.; Del Prete, S.; Capasso, C.; Supuran, C. T.; Gitto, R. Inhibitory effects and structural insights for a novel series of coumarin-based compounds that selectively target human CA IX and CA XII carbonic anhydrases. *Eur. J. Med. Chem.* **2018**, *143*, 276–282.

(20) Angapelly, S.; Sri Ramya, P. V.; Angeli, A.; Supuran, C. T.; Arifuddin, M. Sulfo-coumarin-, Coumarin-, 4-Sulfamoylphenyl-Bearing Indazole-3-carboxamide Hybrids: Synthesis and Selective Inhibition of Tumor-Associated Carbonic Anhydrase Isozymes IX and XII. *ChemMedChem* **2017**, *19*, 1578–1584.

(21) Zengin Kurt, B.; Sonmez, F.; Durdagi, S.; Aksoydan, B.; Ekhteiri Salmas, R.; Angeli, A.; Kucukislamoglu, M.; Supuran, C. T. Synthesis, biological activity and multiscale molecular modeling studies for coumaryl-carboxamide derivatives as selective carbonic anhydrase IX inhibitors. *J. Enzyme Inhib. Med. Chem.* **2017**, *1*, 1042–1052.

(22) Pacchiano, F.; Carta, F.; McDonald, P. C.; Lou, Y.; Vullo, D.; Scozzafava, A.; Dedhar, S.; Supuran, C. T. Ureido-substituted benzenesulfonamides potently inhibit carbonic anhydrase IX and show antimetastatic activity in a model of breast cancer metastasis. *J. Med. Chem.* **2011**, *54*, 1896–902.

(23) Carradori, S. Selective carbonic anhydrase IX inhibitors based on coumarin scaffold as promising antimetastatic agents. WO2012070024, 2013.