



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

# FLORE

## Repository istituzionale dell'Università degli Studi di Firenze

### **The Porto Azzurro unit (Mt. Calamita promontori, southeastern Elba island, Tuscany): stratigraphic, tectonic and metamorphic evolution**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

The Porto Azzurro unit (Mt. Calamita promontori, southeastern Elba island, Tuscany): stratigraphic, tectonic and metamorphic evolution / F.GARFAGNOLI; F.MENNA; E.PANDELI; G.PRINCIPI. - In: BOLLETTINO DELLA SOCIETÀ GEOLOGICA ITALIANA. - ISSN 0037-8763. - STAMPA. - spec. n.3 (2005):(2005), pp. 119-138.

*Availability:*

This version is available at: 2158/255214 since:

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze  
(<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

## The Porto Azzurro Unit (Mt. Calamita promontory, south-eastern Elba Island, Tuscany): stratigraphic, tectonic and metamorphic evolution

FRANCESCA GARGAGNOLI (\*), FRANCESCO MENNA (\*),  
ENRICO PANDELI (\*), (\*\*\*) & GIANFRANCO PRINCIPI (\*), (\*\*)

### ABSTRACT

Elba Island has a key role in the reconstructions of the stratigraphic, tectonic, metamorphic and magmatic evolution of the Northern Tyrrhenian Sea and the inner part of the Northern Apennines chain. The Porto Azzurro Unit, cropping out in the south eastern part of Elba Island, is the deepest tectonic unit of the central-eastern Elba structural pile of Tuscan, Ligurian and Ligurian-Piemontese Nappes, which were intruded by Late Tortonian-Lower Pliocene granitoids and mainly acidic dykes. Moreover, in this part of the island, the relationships between the emplacement of the plutonic bodies and the final deformations of the tectonic stack are easily detectable. To improve our geological knowledge of south eastern Elba, the authors carried out 1:10,000 geological mapping of the Calamita promontory (mostly composed of the Porto Azzurro Unit) and performed petrographical and structural studies on its rocks. The Porto Azzurro Unit consists of a Paleozoic, probably pre-Carboniferous basement (Mt. Calamita Metamorphic Complex), which is unconformably overlain by the ?Triassic Verrucano metasiliciclastics (Barabacca Quartzites) and ?Upper Triassic-?Hettangian metacarbonates. In the Mt. Calamita Metamorphic Complex, five main lithofacies were recognized and mapped. In particular, garnet-bearing-, albite mica-schists (lithofacies a) geometrically underlie a phyllitic-quartzitic unit (lithofacies b); porphyroids (lithofacies e), metabasite bodies (lithofacies d) and graphite-rich siliciclastics (lithofacies c) are also present. The rocks of lithofacies a are similar to those of the ?pre-Paleozoic-?Paleozoic Micaschist Complex of the Larderello geothermal field, whereas the other lithofacies can probably be correlated with the ?Ordovician formations of the Tuscan Metamorphic Succession (e.g. Apuan Alps). The complex deformation-metamorphic evolution of the Porto Azzurro Unit consists of the following events: a) a Variscan tectono-metamorphic event ( $D_x$ ), recognized in the Mt. Calamita Metamorphic Complex, which is defined by a pre-Alpine foliation and mineral relicts (garnet); b) two Alpine tectono-metamorphic folding events ( $D_1$  and  $D_2$ ) in the greenschist facies, which also deformed the Mesozoic cover; c) a later folding event ( $D_3$ ) which probably occurred during or immediately after the thermometamorphic imprint (including the magnetite-rich skarn bodies), caused by Neogene magmatic intrusions; d) subsequently, the uplift of the magmatic bodies caused low-angle detachments within the Porto Azzurro Unit (between the Mt. Calamita Metamorphic Complex and the Mesozoic cover) and between the latter and the overlying tectonic units (e.g. Zuccale Fault between the Porto Azzurro Unit and the Cretaceous Flysch Unit); e) a final weak antiformal folding ( $D_4$ ) of the whole promontory took place before the development of NW-SE and NE-SW trending high-angle normal fault systems, locally sealed by hydrothermal, sometimes Fe-rich mineralizations. The lithostratigraphical, tectonic, metamorphic and magmatic evolution of the Porto Azzurro Unit is similar to that detected for the Larderello geothermal region. Thus, the Mt. Calamita area can be considered as an older, but similar geological

model for all future interpretations of the deep structure of southern Tuscany crossed by the CROP 18 profile.

**KEY WORDS:** *Northern Apennines, south-eastern Elba Island, Paleozoic-Mesozoic successions, tectonic and metamorphic evolution, granitoid emplacement.*

### RIASSUNTO

**L'Unità di Porto Azzurro (Monte Calamita, Isola d'Elba sud-orientale, Toscana): evoluzione stratigrafica, tattonica e metamorfica.**

L'Isola d'Elba ha un ruolo chiave nella ricostruzione stratigrafica, tattonica, metamorfica del Tirreno Settentrionale e delle porzioni più interne della catena Appenninica settentrionale. L'Unità di Porto Azzurro, che affiora nella parte dell'Isola sud-orientale, è l'unità più profonda della pila tattonica dell'Elba centro-orientale, che è costituita da unità toscane, liguri e liguri-piemontesi, intruse da granitoidi e filoni, per lo più acidi, del Tortoniano sup.-Pliocene inferiore. In questa parte dell'isola, inoltre, sono ben esposte le relazioni tra corpi magmatici e deformazioni finali subite dalle varie unità. Al fine di migliorare le conoscenze geologiche relative all'Elba sud-orientale, gli autori hanno effettuato un rilevamento geologico alla scala 1:10.000 del Promontorio del Calamita (costituito per la maggior parte dall'Unità di Porto Azzurro) ed hanno eseguito uno studio petrografico e strutturale sui litotipi affioranti. L'Unità di Porto Azzurro è costituita da una successione paleozoica, probabilmente di età pre-carbonifera (Complesso Metamorfico di Monte Calamita), sormontata in discordanza stratigrafica dai meta-sedimenti silicoclastici del Verrucano triassico (Quarziti di Barabacca) e da successioni carbonatiche del ?Triassico superiore-?Hettangiano. Nel Complesso Metamorfico di Monte Calamita sono state distinte e cartografate cinque principali litofacie. In particolare, sui micaschisti a granato ed albite (lithofacies a), giace geometricamente un'unità filladico-quarzitica (lithofacies b). Sono inoltre presenti porfiroidi (lithofacies e), rocce silicoclastiche ricche in grafite (lithofacies c) e livelli di metabasiti (lithofacies d). Le rocce della litofacies a sono simili a quelle, di età ?pre-paleozoica-?paleozoica del Complesso dei Micaschisti del campo geotermico di Larderello, mentre le altre litofacie possono essere correlate con formazioni ?Ordoviciane della successione metamorfica toscana (es. quella esposta sulle Alpi Apuane). La complessa evoluzione metamorfico-deformativa dell'Unità di Porto Azzurro può essere così schematizzata: a) evento tattono-metamorfico variabile ( $D_x$ ), riconosciuto nel Complesso Metamorfico di Monte Calamita, testimoniato dalla presenza di relitti tessiturali (scistosità pre-alpina  $S_x$ ) e mineralogici (granati); b) due eventi plicativi tattono-metamorfici alpini, con metamorfismo in facies di scisti verdi, che interessano anche le coperture mesozoiche; c) evento plicativo successivo ( $D_3$ ), avvenuto probabilmente durante o immediatamente dopo la ricristallizzazione termometamorfica, dovuta ai granitoidi neogenici, alla quale sono legati anche i corpi di skarn a magnetite; d) successivamente la risalita dei corpi magmatici ha causato la formazione di detachment a basso angolo all'interno dell'Unità di Porto Azzurro (in particolare tra il Complesso Metamorfico di Monte Calamita e le coperture mesozoiche) e tra quest'ultima e le unità tattonomiche sovrastanti (es. la faglia di Zuccale tra l'Unità di Porto Azzurro e l'Unità del Flysch Cretaceo); e) è seguito l'evento plicativo  $D_4$ , che ha

(\*) Dipartimento di Scienze della Terra, Università di Firenze; via G. La Pira, 4 - 50121 Florence, Italy.

(\*\*\*) CNR (National Research Council of Italy) - Istituto di Geoscienze e Georisorse - Sezione di Firenze - Via G. La Pira, 4 - 50121 Florence (Italy). [pandeli@geo.unifi.it](mailto:pandeli@geo.unifi.it)

prodotto una blanda mega-antiforme, alla scala dell'intero promontorio. Infine si sviluppano faglie normali ad alto angolo ad andamento prevalentemente NW-SE e NE-SW, localmente interessate da mineralizzazioni a ossidi/idrossidi di ferro. L'assetto stratigrafico e l'evoluzione tectonica, metamorfica e magmatica dell'Unità di Porto Azzurro è molto simile a quella del campo geotermico di Larderello. Pertanto, il promontorio di M. Calamita può costituire un modello di riferimento per l'interpretazione della struttura profonda della Toscana meridionale attraversata dal profilo CROP 18.

**TERMINI CHIAVE:** Appennino settentrionale, Isola d'Elba sud-orientale, successioni paleozoiche-Mesozoiche, evoluzione tectonica e metamorfica, messa in posto di granitoidi.

## INTRODUCTION

Elba Island, the biggest island of the Tuscan Archipelago, represents the westernmost outcrop of the Northern Apennines chain. The geological peculiarities of this island are not only linked to its complex tectonic pile of nappes and to the well-known Fe-rich ores, but also to the well-exposed interactions between Neogene magmatic intrusions and tectonics (TREVISAN, 1952; BARBERI *et alii*, 1969a; KELLER & PIALLI, 1990; PERTUSATI *et alii*, 1993;

BORTOLOTTI *et alii*, 2001; TANELLI *et alii*, 2001). Moreover, its geographical location, midway between Tuscany and Corsica, makes Elba Island a key place for the reconstruction of the extensional processes in the hinterland of the Northern Apennines (e.g. the opening of the Northern Tyrrhenian Sea) from the Early Miocene (BARTOLE *et alii*, 1991; BARTOLE, 1995). In this framework, the metamorphic rocks of the lowest tectonic unit (Complex I of TREVISAN, 1951 and BARBERI *et alii*, 1969; Porto Azzurro Unit of BORTOLOTTI *et alii*, 2001), exposed in the Mt. Calamita promontory (SE Elba: location in fig. 1) are poorly studied, perhaps because of their complex deformation and metamorphic evolution. In fact, such mostly pre-Alpine rocks (Calamita Schists *Auctt.*) suffered strong thermometamorphism and hydrothermalism related to Messinian-Pliocene magmatism, which locally obliterate the previous polyphase tectono-metamorphic framework. A similar younger geological framework is present at depth in the Larderello geothermal field (cfr. BATINI *et alii*, 1983; ELTER & PANDELI, 1990, 1996) located in southern Tuscany about 40 km distant from Elba. To improve the geological model of the geothermal field, crossed by the CROP-18 profile, the authors carried out 1:10,000 geological mapping, and lithostratigraphical,




Fig. 1 - Tectonic sketch map of Elba Island according to BARBERI *et alii* (1969a) and location of the studied area (squared area).  
- Carta tectonica schematica dell'Isola d'Elba secondo BARBERI *et alii* (1969a) e localizzazione dell'area studiata (nel riquadro).

petrographical and structural studies on the Porto Azzurro Unit to reconstruct its tectonic and metamorphic evolution and its relationships with the Neogene magmatic intrusions.

### GEOLOGICAL FRAMEWORK

According to BARBERI *et alii* (1967a), the tectonic building of the Elba Island is composed of a structural pile of at least five main tectonic units («Complexes» of TREVISAN, 1950, fig. 1) of which the lowermost three belong to the Tuscan Domain, while the uppermost two relate to the Ligurian Domain.

In last ten years several papers have contributed to improve the tectono-stratigraphic framework of the island. In particular, BORTOLOTTI *et alii* (2001) performed 1:10,000 mapping of central-eastern Elba and proposed a new stratigraphic and tectonic model in which the five «Complexes» of TREVISAN were reinterpreted and renamed. BORTOLOTTI *et alii* (2001) recognised nine main tectonic units which are (fig. 2), from bottom to top of the tectonic stack:

1) *Porto Azzurro Unit* (TREVISAN's Complex I): this consists of ?Paleozoic mica-schists, phyllites and quartzites with local amphibolitic horizons (?Carboniferous-?Permian Calamita Gneiss: TREVISAN, 1950; BARBERI *et alii*, 1967a, 1969a,b; «Complesso scistoso basale di Capo Calamita»: BARBERI *et alii*, 1967b; Calamita Schists: PERRIN, 1975; Mt. Calamita Fm.: BORTOLOTTI *et alii*, 2001) and Triassic to Hettangian metasiliciclastics and metacarbonates. Particularly in the Porto Azzurro zone and in the eastern areas of the Calamita promontory, the Mt. Calamita Fm. is typically cross-cut by the aplitic and microgranitic dyke swarm of the La Serra-Porto Azzurro monzogranitic pluton (MARINELLI, 1959) of 5.1-6.2 Ma (SAUPÉ *et alii*, 1982; FERRARA & TONARINI, 1985), which produced a strong thermometamorphic imprint in the host rocks (MARINELLI, 1959; BARBERI *et alii*, 1967b). The contact with the overlying imbricated tectonic units is a cataclastic horizon, up to ten metres-thick, locally mineralised, linked to the well-known low-angle Zuccale fault, which post-dates the magmatic intrusions (KELLER & PIAZZI, 1990; DANIEL & JOLIVET, 1995; PERTUSATI *et alii*, 1993; BORTOLOTTI *et alii*, 2001; COLLETTINI & BARCHI, 2004).

2) *Ortano Unit* (lower part of TREVISAN's Complex II): this comprises phyllites and quartzites (Capo d'Arco Schists, intruded by rare aplitic and more feric dykes), porphyroids and porphyritic schists, and phyllites, quartzitic metasandstones and metaconglomerates at the top. This succession has been correlated with the Ordovician formations of central Sardinia and of Apuan Alps (PANDELI & PUXEDDU, 1990). Also this unit suffered thermometamorphic recrystallization due to the Serra Porto Azzurro intrusion (DURANTI *et alii*, 1992). A vacuolar carbonate cataclasite, with clasts of phyllites and porphyritic schists at the base, marks the contact with the overlying unit.

3) *Acquadolce Unit* (upper part of TREVISAN's Complex II). The basal part is made up of locally dolomitic massive marbles, grading upwards to calcschists. The latter are capped by a thick siliciclastic succession, which includes some layers of calcschists containing a Lower Cretaceous microfauna (DURANTI *et alii*, 1992). BOR-




Fig. 2 - Central Eastern Elba tectonic sketch according to BORTOLOTTI *et alii* (2001). PU=Porto Azzurro Unit; UO=Ortano Unit; AU=Acquadolce Unit (a - Porticciolo sub-unit, b - S. Filomena sub-unit); MU=Monticano Roccastrada Unit; TN=Tuscan Nappe; GU=Grassera Unit; OU=Ophiolitic Unit (a - Acquaviva sub-unit, b - M. Serra sub-unit., b<sub>1</sub> - Capo Vita sub-unit, c - Sassi Turchini sub-unit, d - Volterraio sub-unit, e - Magazzini sub-unit, f - Bagnaia sub-unit); EU=Unità del Flysch Paleogenico; CU=Unità del Flysch Cretaceo.

- Schema tettonico dell'Elba centro orientale secondo BORTOLOTTI *et alii* (2001). PU=Unità di Porto Azzurro; UO=Unità di Ortano; AU=Unità dell'Acquadolce (a - sub-unità del Porticciolo, b - sub-unità di S. Filomena); MU=Unità Monticano Roccastrada; TN=Falda Toscana; GU=Unità del Grassera; OU=Unità Ophiolitica (a - sub-unità dell'Acquaviva, b - sub-unità di M. Serra, b<sub>1</sub> - sub-unità di Capo Vita, c - sub-unità dei Sassi Turchini, d - sub-unità del Volterraio, e - sub-unità di Magazzini, f - sub-unità di Bagnaia); EU=Unità del Flysch Paleogenico; CU=Unità del Flysch Cretaceo.

TOLOTTI *et alii* (2001) and PANDELI *et alii* (2001a) consider this succession to be a metamorphosed oceanic Ligurian-Piemontese unit similar to the *Schistes Lustrés* of Corsica and Gorgona islands. The presence of a meta-serpentinite slice at its top is in agreement with this interpretation. The Mio-Pliocene thermometamorphic imprint is also evident in these rocks.

4) *Monticano-Roccastrada Unit* (lower part of TREVISAN's Complex III). This includes basal fossiliferous graphitic metasediments of Late Carboniferous-Early Permian age (Rio Marina Fm.), unconformably overlain by the detrital Verrucano succession (?Middle-Late Triassic). The epimetamorphic succession of Capo Castello, consisting of Upper Jurassic siliceous metalimestones to Oligocene metagreywackes, and the ?Mesozoic phyllitic-carbonate rocks of Capo Pero have been considered as part of the original cover of Verrucano (PANDELI *et alii*, 1995; BORTOLOTTI *et alii*, 2001).

5) *Tuscan Nappe Unit* (central part of TREVISAN's Complex III). This is represented by calcareous-dolomitic breccias (Calcare Cavernoso Auctt.) cropping out between Porto Azzurro and Rio Marina, and near Norsi-Valdana. Northwards, in the Cavo area, the overlying carbonatic (Pania di Corfino Fm., Cetona Fm. and «Calcare Massiccio», Late Triassic-Hettangian), carbonatic-cherty (Limano and Grotta Giusti Limestones and «Rosso Ammonitico», Middle-Late Liassic) and marly-carbonatic (Posidonia Marlstones, Dogger) non-metamorphic formations also crop out.

6) *Grassera Unit* (upper part of TREVISAN's Complex III). Most of the unit is made up of varicoloured slates and siltstones with rare metalimestone or meta-chert intercalations; basal calcschists also occur. These rocks, previously included in the «Posidonia Marlstones» of the Tuscan Nappe (BARBERI *et alii*, 1969a,b), differ from both




Fig. 3a

Tuscan and Ligurian lithotypes, show a slight metamorphism (up to the anchizone/epizone boundary) and tectonically lie on different formations of the Tuscan Nappe (PANDELI *et alii*, 2001b). Therefore, BORTOLOTTI *et alii* (2001) tentatively attributed also the Grassera Unit to Ligurian-Piemontese oceanic domain.

7) *Ophiolitic Unit* (TREVISAN's Complex IV). Within this Ligurian unit, BORTOLOTTI *et alii* (2001) recognized four main tectonic sub-units with Jurassic to Cretaceous formations (from bottom to top): *a*) Acquaviva sub-unit with serpentinites-ophicalcites, basalts, Mt. Alpe Cherts and Palombini Shales; *b*) Sassi Turchini sub-unit with serpentinites; *c*) Mt. Serra sub-unit with ophicalcites, Mt. Alpe Cherts, Nisportino Fm., Calpionella Limestones and Palombini Shales; *d*) Volterraio sub-unit with serpentinite, basalts, Mt. Alpe Cherts, Nisportino Fm. and Calpionella Limestones.

8) *Paleogene Flysch Unit* (lower part of TREVISAN's Complex V). This mainly consists of shales, marls with limestone, sandstone and ophiolitic breccia intercalations including fossils of Paleocene-Eocene age. KELLER & PIAZZI (1990) and BORTOLOTTI *et alii* (2001) correlated this unit with the Epiligurian Lanciaia Fm. of southern Tuscany.

9) *Cretaceous Flysch Unit* (upper part of TREVISAN's Complex V). This Ligurian, Lower to Upper Cretaceous unit, consists of basal Palombini Shales and varicoloured shales, which vertically pass to turbiditic siliciclastic sandstones and conglomerates (Ghiaieto Sandstones) and these latter to alternating marlstones and marly limestones (Marina di Campo Fm.).

Both Flysch Units were intruded by aplitic and porphyritic dykes and laccoliths at about 7-8 Ma (DINI *et alii*, 2002).

The evolution of deformation of Elba Island can be dated back to the Variscan Orogeny, which affected the Mt. Calamita Fm. and the Ortano Unit (PANDELI & PUXEDDU, 1990). During the Upper Cretaceous-Eocene closure of the Ligurian-Piemontese Ocean, intraoceanic deformations of the Ligurian and the «*Schistes Lustrès*»-like units took place. The subsequent collision and ensialic shortening of the Adriatic margin occurred in the Upper Eocene/Oligocene to Lower Miocene time interval, during which the «*Schistes Lustrès*»-like, Ligurian and Tuscan Units were deformed and piled up. During ?Early-Middle Miocene times, extensional tectonics began and led to the thinning of the structural pile, the opening of the Tyrrhenian Sea and the uplift of magmatic bodies, such as the Mt. Capanne (about 6.8 Ma: JUTEAU *et alii*, 1984; FERRARA & TONARINI, 1993) and the Serra-Porto Azzurro (5.1-6.2 Ma: SAUPÉ, 1982; FERRARA & TONARINI, 1993; 5.9 Ma: MAINERI *et alii*, 2003) granitoids in Elba Island.

The emplacement of the magmatic bodies determined both a thermometamorphic and metasomatic imprint (e.g. skarn bodies: DIMANCHE, 1971; TANELLI, 1977) in the country rocks and gravitational detachments within the pile of nappes (TREVISAN, 1951; PERTUSATI *et alii*, 1993; DANIEL & JOLIVET, 1995; BORTOLOTTI *et alii*, 2001). A final high-angle, mainly N-S trending normal faulting dissected the Elba tectonic stack (BORTOLOTTI *et alii*, 2001) and was locally sealed by hematite-rich ores (DESCHAMPS *et alii*, 1983; LIPPOLT *et alii*, 1995).




Fig. 3 - a) Geological sketch map and geological cross-sections of the Mt. Calamita promontory; b) Stratigraphic and structural sketch of the units cropping out in the Mt. Calamita promontory.

- a) Carta geologica schematica e sezioni geologiche del promontorio del Monte Calamita; b) schema stratigrafico e strutturale delle unità affioranti nel promontorio del Monte Calamita.

## GEOLOGY OF THE CALAMITA PROMONTORY

Most of the Mt. Calamita promontory is made up of polymetamorphic rocks of the Porto Azzurro Unit. In fact, the Cretaceous Flysch Unit (Marina di Campo Fm.) tectonically lies, through the Zuccale Fault (KELLER & PIAZZI, 1990; PERTUSATI *et alii*, 1993; DANIEL & JOLIVET, 1995; KELLER & COWARD, 1996; BORTOLOTTI *et alii*, 2001; COLLETTINI & BARCHI, 2004) on top of the Porto Azzurro Unit only in the Capoliveri-Zuccale-Lido di Capoliveri area. Another minor outcrop of the Cretaceous Flysch Unit (Ghiaieto Ss.) is that of the Scoglio Corbella (southwest of the studied area). The Ophiolitic Unit crops out only in the outermost of the two Gemini Islands (Serpentinites and Gabbros) and in some emerging rocks (marls and marly limestones of the Nisportino Fm.) between the Scoglio Corbella and the Mt. Calamita promontory (fig. 3a). Quaternary deposits, mostly consisting of cemented sands locally characterized by large-scale cross-stratification (eolian sands and beach rocks: MAZZANTI, 1983; CENTAMORE *et alii*, 1988), are present in the western coast of the promontory and in the Calagrande-Calanova area.

The stratigraphic and structural study of the unfossiliferous Porto Azzurro Unit has always been difficult because of the Messinian-Lower Pliocene thermometamorphic overprint and skarn mineralizations, which locally completely obliterate previous textures and mineralogical associations of these polymetamorphic rocks. Apart from the ?Triassic metasiliclastics (Verrucano Auctt.; Barabacca Quartzites of PERRIN, 1975) and ?Triassic-?Hettangian metacarbonates (e.g. Calanchiole marble and crystalline dolostone), the underlying Mt. Calamita Fm. (BORTOLOTTI *et alii*, 2001) has been considered as an undifferentiated unit both in the 1:25,000 geological map (BARBERI *et alii*, 1967a) and in the Sheet 126 of the 1:100,000 Geological Map of Italy by BARBERI *et alii* (1969a,b). Only in the 1:25,000 map did BARBERI and co-workers distinguish amphibolite horizons. In the same

TABLE 1

Lithological, petrographical and textural features of the Porto Azzurro Unit in the Calamita promontory.  
– Caratteristiche litologiche, petrografiche e tessiturale dell'Unità di Porto Azzurro nel promontorio del Monte Calamita.

UNITS	LITHOLOGY	TEXTURE	MINERALOGICAL ASSOCIATIONS	
			PRE/SYN ALPINE TECTONOMETAMORPHIC MINERALS	THERMOMETAMORPHIC MINERALS (HT/LP events)
Lithofacies a	micaschists and phyllites	granolepidoblastic to porphyroblastic	Ms + Ab + Qtz + Chl ± Bt local pre-Alpine Grit, frequent Ab porphyroblasts with helicitic inclusion trails (Op ± Qtz ± Ms) Acc. Min.: Ap, Ox, Py, Ttn, Tur, Zrn	And + Bt + Ms or Bt ± And
Lithofacies b	quartzitic phyllites and quartzites	granolepidoblastic to granoblastic/blastopsmamic	Qtz + Ms + Ser ± Chl (± Ab) millimetric levels with abundant Hem pigment, rare Ab; Acc. Min.: Ap, Hem, Py, Tur, Zrn	Crd + And + Bt (Ginevro and Sassi Neri area), And + Bt + Ms or Bt + Chl ± And
Lithofacies c	black quartzites and phyllites	granolepidoblastic to granoblastic	Qtz + Ms + Op ± Ab + Chl abundant Opaque and organic matter pigment Acc. Min.: Ap, Ox, Tur	Qtz + Chl + Ep skarn (Ginevro and Sassi Neri); FeAct + FePrgr + Hbl + Mag + Py ± Lm
Lithofacies d	amphibolites	granolepidoblastic to nematoblastic	Hbl + Qtz + acid Pl (+Pl) + Ms Acc. Min.: Ap, Ox, Ttn, Tur, Zrn	Bt or Bt + And
Lithofacies e	porphyroids	granoblastic/porphyroblastic	Qtz + Pl + Kfs + Bt porphyroblasts and aggregates of: Qtz, Pl, Kfs Acc. Min.: Ap, Ox, Ttn, Tur, Zrn	Ep + Tr-Act
Barabarca Quartzite (?Late Ladinian- ?Carian)	quartzites	blastopsmamic, granoblastic to granolepidoblastic	Qtz + Ser (± Ms) ± Chl. Acc. Min.: Ap, Hem, Tur, Ox (Fe)	Bt + And
Tocchi Fm. (?Carian)	phyllites, calcschists and impure marbles	diablastic, nematoblastic to lepidoblastic	carbonate beds: Cal (or Cal ± Mg) ± Qtz ± Ab + Ox phyllitic beds: Ser + Chl + Qtz ± Ms + Ox Acc. Min.: Hem, Py, Mag, Ccp, Mrc, Tur	Bt + Act ± Tr skarn (Calamita Mine); Tr-Act + Adr + Hd + Hem + Mag + Py
Calanchiole Marble and Cristalline Dolostone (?Rhaetian- ?Hettangian)	marbles and crystalline dolostones	granoblastic to xenoblastic	Cal + Dol ± Qtz ± Ser. Acc. Min.: Ap, Hem, Ox (Fe).	Cal, Cal + Hem skarn (Calamita Mine); Adr + Hd + Hem + Mag + Py

year, BARBERI *et alii* (1967b) recognized in the «Complesso scistoso basale di Capo Calamita» three petrofacies: «Scisti muscovitico-biotitici ad andalusite» and «Scisti muscovitico-biotitici con metablastesi plagioclasica» of possibly Paleozoic age, and «Scisti quarzoso-muscovitico-biotitici con intercalazioni di quarziti nella parte alta», which they correlated to the Triassic Verrucano. Successively, PUXEDDU *et alii* (1984) distinguished, on the evidence of petrography-geochemistry, four groups of rocks, which could be tentatively attributed to the Carboniferous (e.g. S. Lorenzo Group) and Lower Paleozoic (Lower Quartzite and Phyllite Fm. and Upper Quartzite and Phyllite Fm., and Boccheggiano Fm.) units of Tuscany (cfr. PANDELI *et alii*, 1994). The presence of Variscan rocks in the Mt. Calamita Fm. was also suggested by the local discovery of pre-Alpine foliation relics (PANDELI & PUXEDDU, 1990; FASANO *et alii*, 2001). As it regards the protolith of the thermometamorphosed amphibolite horizons within the Mt. Calamita Fm., in the Pareti-Morcone and in the Remaiolo areas (fig. 3a), BARBERI & INNOCENTI (1965) and BARBERI *et alii* (1967b) suggested a basic magmatic origin (Triassic volcanics? or tectonic slices of Ligurian basalts?); the same horizons were correlated to the within-plate Paleozoic metabasites of southern Tuscany by PUXEDDU *et alii* (1984). In any case, Ca-Mg-hornfels, locally associated with the amphibolite horizon of the Pareti-Morcone area, was interpreted by BARBERI *et alii* (1967b) as metamorphosed dolostone. A carbonatic protolith of Paleozoic age (e.g. Silurian dolostones of the Apuan Alps) was suggested also by DIMANCHE (1971) for the skarn intercalations of the Ginevro mining area (fig. 3a).

The data from our 1:10,000 geological map of the Mt. Calamita promontory, and from meso-/microstructural and petrographical studies on its polymetamorphic rocks, are discussed in the following paragraphs.

#### LITOLOGICAL AND PETROGRAPHICAL DATA

The non-fossiliferous Porto Azzurro Unit shows an apparent thickness of at least 800 m and consists of a recrystallized ?Triassic-Hettangian? cover succession which lies on the Mt. Calamita Metamorphic Complex (*sensu* INTERNATIONAL SUBCOMMISSION ON STRATIGRAPHIC CLASSIFICATION, 1987), made up of pre-Paleozoic? to Paleozoic formations.

From bottom to top, the following units were distinguished (see fig. 3a,b; see also table 1 for detailed petrographical data, mineral abbreviations as KRETZ, 1983, and Adularia = Adu, Marcasite = Mrc, Opaques = Op, Sericite = Ser, Fe-pargasite = Fe-prg):

– *Mt. Calamita Metamorphic Complex*. This metamorphic complex represents most of the outcrops of the Porto Azzurro Unit (fig. 4) and lies unconformably below the Verrucano metasediments (Barabarca Quartzites). In spite of the local strong thermometamorphic imprint (e.g. in the eastern part of the promontory) and magnetite-rich skarn mineralizations (e.g. the Ginevro and Sassi Neri areas; fig. 3a), at least four lithofacies can be distinguished.

– *Lithofacies a*. It mainly consists of silver grey, grey-brown to grey-greenish, quartzitic mica-schists and phyllites with local ochreous alteration. The muscovite and muscovite-biotite mica-schists typically include syntectonic albite porphyroblasts (fig. 5a). Rare garnet porphy-



Fig. 4 - A typical outcrop of the phyllites and quartzites of the Mt. Calamita Metamorphic Complex (Cala di Mola area) affected by Alpine folding ( $S_1$  and  $S_2$  = Alpine foliations) and intruded by aplitic dykes (A); the latter are dissected by low-angle (white arrow) and, successively, by high-angle (black arrows) faults.

– Tipico affioramento delle fillidi e quarziti del Complesso Metamorfico di Monte Calamita (presso Cala di Mola) caratterizzate da deformazioni alpine ( $S_1$  and  $S_2$  = foliazioni alpine) e intruse da dicchi aplitici (A); questi ultimi sono tagliati da faglie a basso (frecce bianche) e, successivamente, ad alto angolo (frecce nere).

roblasts also occur (fig. 5b). These latter are broken, chloritized and wrapped by the Alpine foliations (see below) and are sometimes recrystallized into thermometamorphic biotite pseudomorphic aggregates. These lithotypes crop out in the Poggio Turco-Remaiolo-Costa dei Gabbiani area (fig. 3a) and correspond to the «Scisti muscovitico-biotitici con metablastesi plagioclasica» of BARBERI *et alii* (1967b). Centimetre- to 2m-thick, dark grey to grey-brown quartzites are locally present as intercalations or, in the Pareti-Innamorata area (fig. 3a), as the dominant lithology. The contact with Lithofacies b is tentatively represented in the geological map (fig. 3a) as dotted lines.

– *Lithofacies b*. This is represented by centimetric to decimetric alternations of granolepidoblastic, often quartzose grey-greenish phyllites and grey granoblastic to blastopsammitic metasandstones. In particular, they are fine- to coarse-grained quartzitic metasandstones and chlorite-poor phyllites (fig. 5c), which locally include millimetre-scale alignments of Fe-oxides (e.g. in the areas west and north-west of Mt. Calamita). In any case, some samples collected NE of Mt. Calamita are albite-bearing metasandstones and phyllites which were probably rich in chlorite, now mostly transformed into thermometamorphic biotite. Lenticular bodies of quartzitic metasandstones, meters to tens of metres thick, have been distinguished in the Capoliveri-Morcone area (fig. 3a).

– *Lithofacies c*. This consists of alternating black phyllites and grano-lepidoblastic quartzites which are rich in organic matter (fig. 5d). They form decimetric to decametric lenticular bodies within lithofacies a in the Poggio Turco-Costa dei Gabbiani area (the thickest bodies are shown in fig. 3a). The contact with the mica-schists is locally gradual. Similar lithotypes (alternating grey to light grey quartzites and quartzose microconglomerates with layers of black phyllites) also occur in the Punta del Praticciolo (fig. 3a) coast, and were inter-




Fig. 5a




Fig. 5b




Fig. 5c




Fig. 5d




Fig. 5e




Fig. 5f

**Fig. 5 - Photomicrographs of the lithofacies within the Mt. Calamita Metamorphic Complex:** a) albite porphyroblasts (Ab) in the mica-schists of lithofacies a (crossed polars, 15 $\times$ ); b) Pre- $S_1$ , fractured and chloritized garnet porphyroblasts (Grt) in the mica-schists of lithofacies a, Bt=static thermometamorphic biotite (parallel polars, 25 $\times$ ); c) quartzitic phyllite of lithofacies b (crossed polars, 10 $\times$ ); d) graphite-rich phyllite of lithofacies c (crossed polars, 10 $\times$ ) including post- $S_2$  static andalusite porphyroblasts (And); e) passage from mica-schists (M) of lithofacies a to amphibolite (A) of the lithofacies d (crossed polars, 15 $\times$ ); f) zoned and twinned, acidic plagioclase composite porphyroblast within the porphyroids of lithofacies e (crossed polars, 10 $\times$ ).  $S_1$ ,  $S_2$  = Alpine foliations.

- Microfotografia delle litofacie all'interno del Complesso Metamorfico di Monte Calamita: a) porfiroblasti di albite (Ab) nei micascisti della litofacie a (nicols incrociati, 15 $\times$ ); b) porfiroblasti di granato pre- $S_1$ , fratturati e cloritizzati (Grt) nei micascisti della litofacie a, Bt=biotite statica termometamorfica (nicols paralleli, 25 $\times$ ); c) filladi quarzitiche della litofacie b (nicols incrociati, 10 $\times$ ); d) filladi grafitose della litofacie c (nicols incrociati, 10 $\times$ ), che includono porfiroblasti di andalusite statica post- $S_2$ ; e) passaggio tra micascisti (M) della litofacie a alle anfiboliti (A) della litofacie d (nicols incrociati, 15 $\times$ ); f) porfiroblasto zonato e geminato di plagioclasio acido composito nei porfiroidi della litofacie e (nicols incrociati, 10 $\times$ ).  $S_1$ ,  $S_2$  = foliazioni alpine.

interpreted as Carboniferous metasediments by PUXEDDU *et alii* (1984).

– *Lithofacies d.* This consists of green, banded light and dark-green to black amphibolites, which occur as lenses tens of meters thick in western (Pareti-Morcone), south-western (north of Punta del Praticciolo) and south-eastern (Remaiolo) areas of the promontory. The amphibolite to the east of the Remaiolo area shows transitional boundaries to the mica-schists by millimetre-scale alternations of the two lithologies (fig. 5e). At the microscale, the less thermometamorphosed samples are characterized by a granonematoblastic texture with albite-oligoclase+tremolite-actinolite (after hornblende)+ titanite; a post-tectonic metasomatic assemblage of garnet (grossular-andradite) ± hedembergite also occur, for example NE of Morcone (i.e.: Ca-Mg hornfels of BARBERI *et alii*, 1967b)

– *Lithofacies e.* Grey to dark grey rocks, which show a typical microaugen texture at the mesoscale and form discontinuous (mostly due to faulting), decimetres to some metres-thick horizons within the phyllitic-quartzitic lithotypes of the Capo Calvo area (east coast of the promontory). The porphyroclastic texture is due to pre-kinematic (magmatic) blasts, up to 1 cm in size, of feldspars (mostly acidic plagioclases), quartz and minor granoblastic/granular aggregates of quartz+plagioclase+k-feldspar in a quartzose-feldspathic-micaceous groundmass (fig. 5f). The latter is affected by the same pervasive foliation as the surrounding quartzose metapelites; this foliation appears parallel to the lithological contact and deformed by the D2 folding.

It is still difficult to define the geometric-stratigraphical setting of the Mt. Calamita Metamorphic Complex lithofacies, because of the polyphase folding (see later) and the presence of possible tectonic slices (see MARINELLI, 1959), but we hypothesize that Lithofacies a is at the geometric base of the succession as shown by the geological cross-sections in fig. 3a. The stratigraphical contact with the overlying Barabarca quartzite was locally affected by the Alpine syn-metamorphic folding.

In the ?Triassic-?Hettangian cover we distinguish the following informal formations:

– *Barabarca quartzite.* This unit widely crops out along the Zuccale-Barabarca coast; other minor exposures are located at Calanchiole (south of Spiaggia del Lido, see fig. 3a), south of the Innamorata Gulf, east of the Punta del Praticciolo and around the Remaiolo Gulf. Its maximum apparent thickness is about 50 m. The main lithotype is poorly-sorted-, quartzitic metaconglomerates («anageniti» Auctt.) and quartzites with typical white and pink quartz pebbles in a quartzose-micaceous matrix (fig. 6a). Quartzitic and phyllitic lithics, as well as black quartz+tourmaline clasts («tourmalinites», fig. 6b), are also present; these latter lithics characterize the ?Late Ladinian-Carnian Verrucano basal siliciclastics at the regional scale (CAVARRETTA *et alii*, 1989, 1992; PANDELI, 2002 and references therein). These blasto-psefitic/-psammitic rocks, which are medium- to thick-bedded (max. 1.5-2m) and locally normally graded, sometimes show centimetric to decimetric ±quartzose metapelitic intercalations, wine-red, grey or green in colour. Erosional contacts are frequently seen at the base of coarser beds. In the Barabarca and Calanchiole areas, the «anageniti»



Fig. 6a - Quartzose metaconglomerates («Anageniti») of the Barabarca Quartzite Fm. at Barabarca (T=tourmalinite clast).

– *Metaconglomerati quarzitici («Anageniti») della Formazione delle Quarziti di Barabarca presso Barabarca (T=clasto di tourmalinolite).*




Fig. 6b - Photomicrograph of the contact (dashed line) between a tourmalinitic clast (T) and the surrounding low-sorted quartzitic matrix in a meta-psophite of the Barabarca Quartzites Fm. (crossed polars, 15×).

– *Microfotografia del contatto (linea tratteggiata) tra un clasto di tourmalina (T) e la circostante matrice quarzitica ben classata in una meta-psofite della Formazione delle Quarziti di Barabarca (nichols incrociati, 15×).*




Fig. 7a - Stratigraphic passage from alternating phyllites and metacarbonate beds of the Tocchi Fm. (T) and the overlying Calanchiole marble and crystalline dolostone (M) at the Calanchiole beach.  
– Passaggio stratigrafico tra l’alternanza di livelli filladici e carbonatici della Formazione di Tocchi (T) e i sovrastanti Marmi e dolomie cristalline delle Calanchiole (M) presso la spiaggia delle Calanchiole.

include and grade upwards into alternating fine- to very coarse-grained quartzites and grey, greenish and pinkish-violet, more or less quartzitic phyllites and metasiltstones. The contact with the overlying Tocchi Fm. is gradual (appearance of carbonate intercalations), but it is frequently tectonized.

– *Tocchi Fm.* This occurs at Calanchiole (south of Spiaggia del Lido), Madonna delle Grazie, and in the Calamita mining area (fig. 3) with a maximum apparent thickness of about 150 m. Where it is not affected by tectonism (it generally represents a preferential detachment horizon) or by mineralization (e.g. the Punta della Calamita skarn), it consists of alternating light grey to orange-yellow, impure, more or less dolomitic metadolostone and calcschist beds. They are up to 60 cm thick, green, grey-green and brownish sericitic-chloritic phyllites. The stratigraphical position and its lithological association is similar to the Carnian Tocchi Fm. of southern Tuscany (cfr. COSTANTINI *et alii*, 1980). At Madonna delle Grazie beach, this unit is strongly tectonized (Madonna



Fig. 7b - Photomicrograph of a calcschist of the Tocchi Fm. (crossed polars, 15 $\times$ );  $S_1$  and  $S_2$  = Alpine foliations.  
– Microfotografia dei calcescisti della Formazione di Tocchi (nicols incrociati, 15 $\times$ );  $S_1$  e  $S_2$  = foliazioni alpine.

delle Grazie cataclasite). The transition to the overlying carbonate unit (fig. 7a) is gradual (decrease in pelitic lithotypes).

– *Calanchiole marble and crystalline dolostone.* This unit, with a maximum apparent thickness of 170 m, crops out south of Capoliveri-Madonna delle Grazie, in the north-eastern Gemini Isle, in the Calamita-Punta Rossa mining area (fig. 8) (where it is largely affected by skarn mineralization, e.g. in the Macei Alti Mine) and at the Scoglio del Remaiolo (the little island south of Remaiolo in fig. 3a). The lower part generally consists of up to 1.5 m-thick beds of stratified, white to greyish (sometimes yellowish to reddish, because of Fe-oxides or hydroxides) marbles, with local millimetre to centimetre-scale phyllite and calcschist intercalations (e.g. between Calanchiole and Spiaggia del Lido and south of Madonna delle Grazie beach). The upper part consists of whitish massive to irregularly bedded, often saccoidal marbles and crystalline dolostones (e.g. at Punta Calamita, figs. 3a and 8). A whitish and greyish banding can be locally present within the carbonate rocks. Correlation with the Upper Triassic (Grezzoni Fm.) and Hettangian (Marbles Fm.) metacarbonates has been suggested by all the authors (BARBERI *et alii*, 1967a, b; BARBERI *et alii*, 1969a,b).

The Porto Azzurro Unit is separated from the overlying Cretaceous Flysch Unit (Marina di Campo Fm.) by the Zuccale cataclasite (Zuccale Fault *Auctt.*: see below) (KELLER & PIALLI, 1990; PERTUSATI *et alii*, 1993; DANIEL



Fig. 8 - Whitish marbles in the upper part of the Calanchiole marble and crystalline dolostone at Punta Calamita; SK=skarn bodies.  
– Marmi biancastri nella parte alta dei Marmi e dolomie cristalline delle Calanchiole presso Punta Calamita; SK=mineralizzazioni a skarn.

& JOLIVET, 1995; KELLER & COWARD, 1996; BORTOLOTTI et alii, 2001; COLLETTINI & BARCHI, 2004).

TABLE 2

Blastesis evolution of the Porto Azzurro Unit. The Variscan evolution characterized only the units of Mt. Calamita Metamorphic Complex.  
– Evoluzione della blastesi nell'Unità di Porto Azzurro. L'evoluzione varisica ha interessato solo le unità del Complesso Metamorfico di Monte Calamita.

METAMORPHIC EVENT DEFORMATIONAL EVENT	VARISICAN EVOLUTION		ALPINE EVOLUTION			
	M <sub>x</sub>	D <sub>x</sub>	MA <sub>1</sub>	MA <sub>2</sub>	MAT	MA <sub>3</sub>
DA <sub>1</sub>	DA <sub>2</sub>	DA <sub>3</sub>				
Qtz	====	====	====	====		
Ms	====	====	====	====		
Ser			====	====		
Chl			====	====		
Ab			====	====		
Pli	====		====	====		
Bt	==?==		====	====		
Grt	====					
Op			====	====		
Cal			====	====		
Dol			====	====		
And			====	====		
Crd			====	====		
Kfs			====	====		
Hbl	==?==		====	====		
Tr-Act			====	====		

#### STRUCTURAL DATA

All the formations of the Porto Azzurro Unit were affected by Alpine deformations and metamorphism (see also table 2 for details). In particular, the following plastic deformation events can be distinguished:

**D<sub>1</sub>** - This is defined by the main foliation at the meso- (fig. 9) and microscale ( $S_1 = \text{Ser}/\text{Ms} + \text{Qtz} + \text{Chl} + \text{Ab} + \text{Cc}$ ) (e.g.: figs. 5a and 11), which is continuous or sub-millimetre spaced and generally sub-parallel to the lithological parititions (e.g. Mt. Calamita Metamorphic Complex) or sedimentary bedding (e.g. Verrucano beds). This folia-




Fig. 9 - Alpine deformations within the quartzites of the Mt. Calamita Metamorphic Complex at the Innamorata beach; the main foliation ( $S_1$ ) is isoclinally folded into  $F_2$  ( $AP_2 = F_2$  axial plane); then, the  $F_2$  folds are deformed into open  $D_3$  folds, characterized by a subvertical axial plane ( $AP_3$ ).

– Deformazioni alpine nelle quarziti del Complesso Metamorfico di Monte Calamita presso la spiaggia dell'Iammorata; la foliazione principale ( $S_1$ ) è piegata a formare le pieghe isoclinali  $F_2$  ( $AP_2 =$  piano assiale  $F_2$ ); successivamente le pieghe  $F_2$  sono deformate in pieghe  $F_3$  aperte e caratterizzate da piano assiale sub-verticale ( $AP_3$ ).

tion is the axial plane foliation of isoclinal folds, which are rarely recognizable, because of the the strong transposition by the following  $D_2$  event. During the  $D_1$  event, the syntectonic growth of albite porphyroblasts and the breakage and chloritization of pre- $S_1$  garnets occurred.

For the southern and southwestern areas, the stereonets in fig. 10 show for the rare  $D_1$  folds N-S trending axes and a western dip of the axial surface, while at Cala Nova, probable  $D_1$  folds are characterized by ENE-WSW trending axes and NE axial plane dips.

**D<sub>2</sub>** - The  $D_2$  event is represented by tight to isoclinal folds with sharp or rarely rounded hinges. These are the most common folds recognizable at the mesoscale (tens of centimetres to tens of metres) all over the promontory (e.g. in the Innamorata area; fig. 9). Syntectonic quartz veins, folded by  $D_2$  into close to tight folds (e.g. rootless  $F_2$  hinges) are locally present. The associated axial plane foliation ( $S_2$ ) is represented by a millimetre- to centimetre-spaced, zonal to discrete crenulation cleavage (figs. 5a and 11) made up of the alignment of opaque minerals and Ser/Ms blasts; locally, especially in the metapelitic lithotypes, this foliation is more pervasive ( $S_2 = \text{Op} + \text{Ms} / \text{Ser} \pm \text{Chl} \pm \text{Qtz}$  in fig. 7b).

Also in the case of the attitude of  $D_2$  structures, significant variations are recognizable going around the promontory (fig. 10), probably due to later deformation events. At Barabbarca and Innamorata, the  $F_2$  axial planes dip towards the west or the south, and the axial plunge is about 10-20° towards the SW (also towards the NE in the Barabbarca area). In the Remaiolo and Poggio Turco areas,  $F_2$  have westward, 60-70° dipping axial planes and axes plunging at 50-60° to the SW. At Capo Calvo a north-eastern  $F_2$  axial plane dip and axes plunge is recognizable, whereas at Cala Nova they are both directed to the north. The vergence of  $F_2$  is towards the southeast.

**D<sub>3</sub>** -  $D_3$  folds are symmetric, mostly open, tens of centimetres to metres in size, and characterized by rounded hinges and generally steeply-dipping to subvertical axial planes (fig. 9). The  $S_3$  zonal, spaced crenulation ( $S_3$  in fig. 5b) is locally marked by the alignment of opaque minerals. The  $D_3$  structural elements are well exposed only in




Fig. 10 - Distribution of the structural elements of the  $D_1$ ,  $D_2$ ,  $D_3$  and  $D_4$  folding events in the Calamita promontory (Schmidt nets, lower hemisphere).

- Distribuzione degli elementi strutturali degli eventi plicativi  $D_1$ ,  $D_2$ ,  $D_3$  e  $D_4$  nel promontorio del Monte Calamita (reticolo di Schmidt, emisfero inferiore).

the western and southern areas (fig. 10). In particular, in the Innamorata and Remaiolo-Poggio Turco areas the strike of the steep axial surface is about NE-SW and the axes plunge towards the SW. The latter is instead towards the NW in the Barabarca area and mostly towards N-NW in the Mt. Calamita area. It is noteworthy that the growth of thermometamorphic minerals (e.g. andalusite±biotite: see later fig. 14b) occurred before and after the  $D_3$  crenulation event.

**$D_4$**  - The structures related to a possible  $D_4$  folding event can be recognized only at the map scale. In particular, the areal distribution of the attitude of the main foliation and the orientation of the axes of the major folds (i.e.  $F_2$  folds in fig. 10) outline an overall large-scale antiform in the Porto Azzurro Unit in the Mt. Calamita promontory, with a NW-SE trending axis (fig. 10). The presence of this megafold at the scale of the whole promontory was also suggested also by BARTOLE *et alii* (1991).

In some samples of the Mt. Calamita Metamorphic Complex, a pre- $D_1$  tectono-metamorphic event ( $D_x$ ) can be also recognised. This is indicated by local intrafoliar relicts of pre- $S_1$  foliation ( $S_x = \text{Ms} \pm \text{Chl} \pm \text{Qtz} \pm \text{Bt}$ ) (fig. 11), by the opaque-rich inclusion trails within zoned albite porphyroblasts which are discordant or truncated by  $S_1$ , and by relict garnet porphyroblasts within  $S_1$  (fig. 5b). In the Paleozoic succession of the Northern Apennines, these structures and minerals are associated with the Variscan Orogeny, in particular the tectono-metamorphic Sudetic event (ELTER & PANDELI, 1990; CONTI *et alii*, 1991; PANDELI *et alii*, 1994).

Two main brittle deformations were also recognized:

*Low-angle detachment faults.* These structures are locally present within the Mt. Calamita Metamorphic Complex associated with metres to tens of metres-thick cataclastic horizons (mylonites and ultramylonites of MARINELLI, 1959), affecting the leucocratic dykes (e.g. top-to-east displacements along the north-eastern coast of the promontory: see fig. 7 in KELLER & PIAZZI, 1990). In particular, the cataclasites are generally medium- to fine-grained cataclastic breccias with phyllitic-quartzitic, or white quartz (fig. 12), or quartz+feldspar+biotite (brecciated dykes?) clasts in a quartzitic or quartzitic-micaceous matrix. Some samples reveal that the thermometamorphism occurred before the cataclastic event. Moreover, tourmalinization processes locally affected such rocks before and/or after the cataclastic phenomena. In any case predate the youngest Fe-mineralization (e.g. at Punta di Cala Nova and in the Capo Calvo area, where the tourmalinized cataclasites are cut by high-angle fractures filled by Fe-minerals).

Other low-angle faults detach the Mesozoic cover formations (including the skarn bodies: e.g. Calamita mining area) from the original basement (Mt. Calamita Metamorphic Complex) or the carbonate units from the Verrucano (e.g.: the Madonna delle Grazie cataclasite). The tectonic superposition of the overturned  $F_2$  Spiaggia del Lido-Barabarca-Madonna delle Grazie megafold, formed by the Mesozoic formations (Barabarca quartzite at the base), on the Paleozoic phyllites probably belongs to this same event.

The best known example of detachment in this area is the Zuccale fault (KELLER & PIAZZI, 1990; PERTUSATI *et alii*, 1993; DANIEL & JOLIVET, 1995; KELLER & COWARD, 1996; BORTOLOTTI *et alii*, 2001; COLLETTINI & BARCHI, 2004). This tectonically complex horizon, up to 7 m thick,




Fig. 11 - Photomicrograph of a  $F_2$  fold, which deforms the main foliation  $S_1$  and the previous (?Variscan)  $S_x$  foliation (parallel polars, 2,5 $\times$ ). - Microfotografia di una piega  $F_2$ , che deforma la foliazione principale  $S_1$  e la foliazione  $S_x$  precedente (?varisica) (nicols paralleli, 2,5 $\times$ ).





Fig. 12 - Photomicrograph of a quartz cataclasite associated with a low-angle fault, crossed by a quartz vein (V) in the Mt. Calamita Metamorphic Complex along the panoramic road from the Calamita mining area to Poggio Turco (crossed polars, 15 $\times$ ). - Microfotografia di una cataclasite quarzitica associata ad una faglia a basso angolo, tagliata da una vena di quarzo (V) nel Complejo Metamorfico di Monte Calamita, lungo la strada panoramica tra la miniera di Calamita e Poggio Turco (nicols incrociati, 15 $\times$ ).

is well exposed along the coast between Calanchiole and Zuccale, where it has a shallow (max 15°) dip to the west, and has two parts:

a) the *upper part* is a Fe-carbonate-rich breccia, brown in colour and at times vacuolar, made up of millimetre to 10 cm-sized, recrystallized and at times silicified clasts of mainly carbonate and minor pelitic and arenaceous rocks, derived from the overlying Cretaceous Flysch. S-C, C and C'-type fabric (see also PERTUSATI *et alii*, 1993 and COLLETTINI & BARCHI, 2004) suggest a «top-to-ENE» sense of shear (fig. 13a);

b) the *lower part* is a polymictic breccia with up to 5 cm sized thermometamorphosed quartzitic and phyllitic



*Fig. 13a - S-C-type fabric showing a «top-to-ENE» sense of shear in the upper part of the Zuccale cataclasite at Punta di Zuccale.*  
- *Fabric di tipo S-C, che indica un senso di taglio «top-to-ENE» nella parte alta della cataclasite associata alla Faglia di Zuccale presso Punta Zuccale.*



*Fig. 13b - The Zuccale cataclasite at Punta di Zuccale. Zl=Lower part, Zu=upper part, B=Barabarca Quartzites, CF=Cretaceous Flysch Unit. Zl includes Verrucano and Tocchi Fm. slices (S).*  
- *Cataclasite di Zuccale presso Punta Zuccale. Zl=parte inferiore, Zu=parte superiore, B=Quarziti di Barabarca, CF=Unità del Flysch Cretaceo. Zl include scaglie di Verrucano e della Formazione di Tocchi (S).*

elements (from the Verrucano and Tocchi Fm.) in a foliated phyllosilicate and phyllosilicate-carbonatic matrix. Moreover, decimetre to metre-sized tectonic slices of Verrucano siliciclastics and of phyllite-carbonate alternations of the Tocchi Fm. are often present (fig. 13b). Mylonitic fabrics locally occur in the basal portion. Some tens of metres north of Punta Zuccale, where the brecciated footwall (Verrucano anagenites and Tocchi Fm.) of the Zuccale shear horizon is well exposed, W-E trending slicken-side striae and calcite fibre veins are present at the base of the cataclastic horizon, which are deformed by weak N-S trending mesofolds.

*High-angle faults.* This is the last deformation event that affected the studied tectonic units, and it is represented by NW-SE and NE-SW-trending joints and high-angle normal faults, which cut both magmatic dykes and the low-angle locally tourmalinized detachment faults and are locally filled by Fe-oxides and hydroxides (e.g. at Punta di Cala Nova and Capo Calvo in fig. 3a). An evident system of NW-SE trending normal faults is present in the Innamorata-Praticciolo area which downthrow the Meso-

zoic covers (north-eastern Gemini Isle) towards the SW with respect to the Calamita Fm. and the Ophiolitic Unit (south-western Gemini Isle) with respect to the formers (fig. 3a).

#### CONTACT METAMORPHISM AND HYDROTHERMALISM

The Messinian-Lower Pliocene magmatism produced a wide metamorphic aureole in the country rocks of the studied area (MARINELLI, 1959; BARBERI *et alii*, 1967b). Our studies, in agreement with MARINELLI (1959), reveal that the contact metamorphism which imprinted the Mt. Calamita Metamorphic Complex (see tables 1 and 2) is generally ascribed to the albite-epidote hornfels/hornblende hornfels facies and made up of green to brown Bt (after regional Chl) $\pm$ And( $\pm$ Ms). The skarn bodies of the Calamita mine (Tr-Act + Adr + Hd + Hem + Mag + Py) have a similar metamorphic grade. In the eastern areas, which are characterized by a complex network of aplitic to microgranitic dykes and skarn bodies with Fac+Feprg + Hbl + Mag + Py  $\pm$  Lm (Sassi Neri and Ginevro mines in

fig. 3a: DIMANCHE, 1971; TANELLI *et alii*, 2001), the static recrystallization is stronger (hornblende hornfels facies) and obliterates previous textures and mineralogical associations. In these areas, the finding of Crd (sometimes transformed into pinnite) associated with Bt, And (fig. 14a) and Ms is frequent. However, local higher temperature conditions are recorded in the Ginevro Mine where a Qtz+Kfs+Pl+Crd(+Bt) and rare Kfs+Crd+Pl+Crn were described by DIMANCHE (1971) in agreement with MARINELLI (1959). In the surroundings of Mt. Calamita and generally towards the north-west, there is an evident decrease in static recrystallization (regional chlorite is less transformed in static biotite; andalusite is present as discontinuous spots instead of sub-idiomorphic porphyroblasts).

Most of the hydrothermal veins hosted within the thermometamorphosed rocks of the Porto Azzurro Unit show a rather homogeneous composition (Qtz+FeCal+FeOx/idrox±Chl, Qtz+Chl+Adu±Tr-Act±Ti, Qtz+Adu+Chl+Ep, Qtz+Ep+Cal, Adu±Ep) over the whole promontory, although veins of Bt, Tur±Qtz, Qtz+Cc+Ab, Qtz+Ti+Bt+Pl and Cpx+Ti were seen locally in eastern areas.

## DISCUSSION

The data collected during this study improve our knowledge of the stratigraphic and magmatic-structural history of the Mt. Calamita promontory. From a stratigraphic point of view, in the polymetamorphic Porto Azzurro Unit we distinguish a Paleozoic basement (Mt. Calamita Metamorphic Complex) which is overlain by ?Triassic-?Hettangian cover units (Barabarca Quartzites, Tocchi Fm. and Grezzoni/Marmi) in agreement with previous authors (BARBERI *et alii*, 1967a,b; BARBERI *et alii*, 1969b). In spite of the lack of fossils, the reconstructed lithostratigraphy of the Porto Azzurro Unit strongly recalls that of the typical successions of the Tuscan Metamorphic Units, and in particular those of southern Tuscany (e.g. Monticiano-Roccastrada Unit: BERTINI *et alii*, 1991 and references therein).

The stratigraphic attribution of the lithofacies defined within the Mt. Calamita Metamorphic Complex is more difficult. The presence of pre-S<sub>1</sub> foliation (S<sub>x</sub>) and minerals (garnet) only in the Mt. Calamita Metamorphic Complex, and their lack in the overlying Mesozoic formations, suggests a pre-Triassic age for the former lithodemic unit. In particular, at the regional scale, the pre-S<sub>1</sub> foliation and minerals are related to the Sudetian tectono-metamorphic event of the Variscan orogeny (CONTI *et alii*, 1991; PANDELI *et alii*, 1994 and references therein). Therefore the Mt. Calamita Metamorphic Complex is probably made up of pre-Carboniferous rocks.

In particular, *lithofacies a*, at the geometric base of the succession, shows similar petrographical and microstructural features to the Micaschist Complex (?Early Paleozoic or ?pre-Paleozoic), present at the base of the Monticiano-Roccastrada Unit, in the sub-surface of the Larderello Geothermal Region (ELTER & PANDELI, 1990, 1996; BERTINI *et alii*, 1991). In fact, acidic plagioclase porphyroblasts and the presence of pre-Alpine almandine are distinctive features in both units. The scarcity of these latter porphyroblasts in the Calamita mica-schists is in part due to their thermometamorphic transformation into biotite and/or can be related to the different chem-




Fig. 14a - Sub-idiomorphic composite andalusite porphyroblasts in the Mt. Calamita Metamorphic Complex (crossed polars, 2,5×).  
- Porfiroblasti compositi sub-idiomorfi di andalusite nel Complejo Metamorfico di Monte Calamita (nolci incrociati, 2,5×).




Fig. 14b - Pre- to syn-D<sub>3</sub> andalusite porphyroblasts (And) in the Mt. Calamita Metamorphic Complex. S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>=Alpine foliations (crossed polars, 15×).  
- Porfiroblasti di andalusite da pre- a sin-D<sub>3</sub> nel Complejo Metamorfico di Monte Calamita. S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub>=foliazioni alpine (nolci incrociati, 15×).

istry of the protolith with respect to that of the Larderello mica-schists.

The metasiliciclastics of *lithofacies b* are characterized by a variable chlorite content (generally low, possibly due to its transformation into thermometamorphic biotite) and a lack or low content of acidic plagioclase blasts in the quartz-rich metasandstones. These features are also typical of the phyllitic-quartzitic units which underlie (Capo Arco Schists) and overlie the Porphyroids in the Ortano Unit of Eastern Elba (see figs. 1 and 2). BORTOLOTTI *et alii* (2001) attributed these rocks to the Late Ordovician metasiliciclastics of Tuscany (e.g. Upper Quartzite and Phyllite of the Apuan Alps; Buti banded quartzite and phyllite of the Mts. Pisani: PANDELI *et alii*, 1994 and references therein), which the authors generally correlate with the transgressive Caradocian sediments of central Sardinia (cfr. CARMIGNANI coord., 2001). The local presence of concentrations and bands of Fe-oxide agrees with this correlation (see also PUXEDDU *et alii*,

1984). The presence of the chlorite- and albite-rich Lower Phyllite and Quartzite (PUXEDDU *et alii*, 1984) has not been definitely confirmed by our sampling.

The graphite-rich *lithofacies c* contains frequent black quartzites which are known as lydites in the Silurian-Early Devonian units of Tuscany, such as the Orthoceras-bearing dolostone and Lydian stones of the Apuan Alps (CONTI *et alii*, 1991) and the Risanguigno Fm. of the Monticiano Roccastrada Unit (COSTANTINI *et alii*, 1987/88) and central Sardinia (CARMIGNANI coord., 2001). On the other hand, similar lithotypes mark the passage from Porphyroids/Porphyritic Schists to the overlying metasiliciclastics in the Ortano Unit succession (DURANTI *et alii*, 1992; BORTOLOTTI *et alii*, 2001) of probable ?Ordovician age.

The analysed amphibolites of *lithofacies d* of the Pareti-Morcone to Remaiolo areas, because of the high content of Ti minerals, are related to a basic magmatic protolith in agreement with BARBERI & INNOCENTI (1965), BARBERI *et alii* (1967b) and PUXEDDU *et alii* (1984). These metabasite horizons appear intercalated both in the micaschists (with a gradual transition in the Remaiolo area) and in lithofacies b (Pareti-Morcone). On the basis of Ti/100-Zr-Y and Ti content, PUXEDDU *et alii* (1984) related all these amphibolites to within-plate basalts as well as the ?Lower Paleozoic Valle del Giardino metabasites of the Apuan Alps and of Niccioleta Mine in southern Tuscany. Therefore, the amphibolitic horizons within the mica-schists of the Remaiolo area are different from those intercalated in the Micaschist complex of the Larderello subsurface and exposed in the Cerreto Pass area (north-west of the Apuan Alps), which were attributed to ocean floor basalts (PUXEDDU *et alii*, 1984; MOLLI *et alii*, 2002).

The orthometamorphic rocks of *lithofacies e* are affected by the D<sub>1</sub> and D<sub>2</sub> tectono-metamorphic events together with the surrounding phyllitic-quartzitic rocks, which are cut by non-foliated aplitic to microgranitic dykes dated to 5.2 to 6.1 Ma by SAUPÉ *et alii* (1982). These data suggest a pre-Miocene age for the protolith. Moreover, the thermometamorphic minerals grow onto S<sub>1</sub>-S<sub>2</sub>, and the radiometric age of the corresponding foliation in the metamorphic units of Tuscany is Late Oligocene-Middle Miocene (KLIGFIELD *et alii*, 1986; DEINO *et alii*, 1992; BRUNET *et alii*, 2000), and therefore rules out that these rocks could be considered foliated Messinian-Lower Pliocene dykes. Therefore, it is likely that lithofacies e correlates with the Ordovician Porphyroids of the Ortano Unit and the Apuan Alps. In any case, the Capo Calvo porphyroids are different from the Ordovician metavolcanic rocks because of the presence of intrusive or sub-intrusive textures (e.g. granular aggregates of Or+Qtz+Pl, lack of corroded quartz phenocrysts).

The lithostratigraphical and petrographical data of the Mt. Calamita Metamorphic Complex strongly recall those of the Variscan succession of the Northern Apennines and, in particular, of southern Tuscany. In fact, in the subsurface of the Larderello-Travale geothermal region, below the Triassic rocks, the lower part of the Monticiano-Roccastrada Unit consists of phyllite-quartzite units with local metabasite levels (e.g.: the Phyllitic-Quartzitic Complex) which overlie a deep Micaschist Complex (ELTER & PANDELI, 1990, 1994; 1996a; BERTINI *et alii*, 1991). Moreover, both the Larderello Paleozoic units suffered static high-temperature metamorphism due to Pliocene granitoid intrusion (BATINI *et alii*, 1983; MUSUMECI *et alii*, 2002; GIANELLI & RUGGIERI, 2002 and references therein)

The deformation evolution of the Calamita promontory and its relationships with the magmatic intrusions are more complex than previously hypothesized (see the synthetic sketch in fig. 15). The oldest tectono-metamorphic event (D<sub>x</sub>), which is ascribed to the Sudetic event of the Variscan Orogeny, is only recognizable in the Mt. Calamita Metamorphic Complex. The Alpine deformations affected both the pre-Triassic rocks and the overlying Mesozoic cover formations. In particular, two main ductile events in the greenschist metamorphic facies (D<sub>1</sub> and D<sub>2</sub>) firstly affected the Porto Azzurro Unit and show strong similarities with the tectono-metamorphic events recognized in the Tuscan Metamorphic Units (CAMPETTI *et alii*, 1999; CARMIGNANI & KLIGFIELD, 1990; ELTER & PANDELI, 1994b; CAROSI *et alii*, 1997; CORSI *et alii*, 2001; COSTANTINI *et alii*, 1987/88). In the studied units, D<sub>1</sub> is the strongest shortening event («Tangential or Syn-Nappe phase»: CARMIGNANI & KLIGFIELD, 1990; CARMIGNANI *et alii*, 1994; ELTER & SANDRELLI, 1995; DECANDIA *et alii*, 2001), dated to about 27 Ma in the Apuan Alps by KLIGFIELD *et alii*, 1986), during which isoclinal folding and the associated very penetrative S<sub>1</sub> (main foliation at the meso-scale) was produced. During D<sub>2</sub> (dated to about 14-12 Ma in the Apuan Alps by KLIGFIELD *et alii*, 1986), the main foliation was isoclinally or tightly folded, and generally the D<sub>1</sub> structures were transposed. The asymmetries of the D<sub>2</sub> folds show an eastern (SE) vergence in agreement with the D<sub>2</sub> folds in the Ortano Unit and the Monticiano-Roccastrada Unit of eastern Elba (BORTOLOTTI *et alii*, 2001; ELTER & PANDELI, 2001). We have no data to attribute D<sub>2</sub> to the post-orogenic regional extension of the Northern Apennines chain («Post-Nappe phase», according to a «Core Complex» extensional model: CARMIGNANI & KLIGFIELD, 1990; CARMIGNANI *et alii*, 1994; ELTER & SANDRELLI, 1995; DECANDIA *et alii*, 2001) or to a final «serrage» event, possibly coupled with syn-orogenic extension (FAZZUOLI *et alii*, 1994; CAROSI *et alii*, 1995; JOLIVET *et alii*, 1998; BORTOLOTTI *et alii*, 2001; CAROSI *et alii*, 2002). We think that an Adriatic vergence of the D<sub>2</sub> folds in Elba does not fit well with the first «Core Complex»-like model. In fact, the latter suggests a centrifugal discharge (westwards in southern Tuscany) of the units from the so-called Tuscan Mid-Ridge (i.e. the Apuan Alps to Mt. Leoni regional structural lineament; the «paleo-apenninic antiform» of ELTER & SANDRELLI, 1994), which represents the main positive structure of the Tuscan Metamorphic Units in Tuscany.

The D<sub>3</sub> weak folding event recognized in the Porto Azzurro Unit probably occurred during the emplacement of the magmatic bodies (5.1 to 6.2 Ma according to SAUPÉ *et alii*, 1982). In fact, the petrographical and microstructural data reveal that the development of S<sub>3</sub> probably in part overlapped with the blastesis of the thermometamorphic minerals and the skarn mineralizations.

The preliminary data, which show an overall decrease in thermal recrystallization from the Ginevro mine towards the north-west of Mt. Calamita, mean it is difficult to make a direct linkage between the La Serra-Porto Azzurro pluton (cropping out northwards of the Mt. Calamita promontory) and the magmatic body present at shallow depths in the studied area, which could be related to an apophysis of the main La Serra-Porto Azzurro pluton. In any case, our data suggest that the top of the Mt. Calamita magmatic body is probably localized east of the Ginevro-Sassi Neri area.




AGE	DEFORMATIONS (D) and METAMORPHIC (M) EVENTS	FOLIATIONS	STRUCTURES
Carb. Inf.	D <sub>x</sub> /M <sub>x</sub>	S <sub>x</sub>	
27-19 Ma (?)	D <sub>1</sub> /M <sub>1</sub>	S <sub>1</sub>	
14-12 Ma (?)	D <sub>2</sub> /M <sub>2</sub>	S <sub>2</sub>	
5.1-6.2 Ma	HT/LP event D <sub>3</sub>	---	
<5.1-6.2 Ma	D <sub>4</sub> Late D <sub>4</sub>	---	
<5.1 Ma	D <sub>5</sub>	---	

Fig. 15 - Sketch of the deformation-metamorphic evolution of the Porto Azzurro Unit (chronological data after KLINGFIELD *et alii*, 1986; SAUPÉ *et alii*, 1982; BRUNET *et alii*, 2000; DEINO *et alii*, 1992)  
– Schema dell’evoluzione tettono-metamorfica dell’Unità di Porto Azzurro (dati cronologici da KLINGFIELD *et alii*, 1986; SAUPÉ *et alii*, 1982; BRUNET *et alii*, 2000; DEINO *et alii*, 1992).

After D<sub>3</sub> deformation and thermometamorphism, low-angle shearing (D<sub>4</sub> event) in Mt. Calamita and detachments between the Mt. Calamita Metamorphic Complex and the overlying Mesozoic units, possibly triggered by

the uplift of the magmatic bodies, locally occurred (e.g. in the Calamita Mining area). The Zuccale cataclasite was also formed, after the magmatic intrusions and thermometamorphism, in a possibly fluid-rich, top-to-the-east

shear zone (KELLER & PIALLI, 1990; PERTUSATI *et alii*, 1993; DANIEL & JOLIVET, 1995; KELLER & COWARD, 1996; BORTOLOTTI *et alii*, 2001; COLLETTINI & BARCHI, 2004). It is noteworthy of that the kinematic data collected in the Zuccale area suggest a western provenance for the Cretaceous Flysch, in particular from the Golfo Stella. Therefore, the Zuccale Fault in the studied area cannot be directly related to the uplift of the Calamita promontory or of the La Serra-Porto Azzurro pluton. The Cretaceous Flysch Unit probably came from an at-present collapsed high, to the west of the Calamita promontory (e.g.: an uplifted magmatic body in the Golfo Stella area?). In the last stages of the D<sub>4</sub> event all the previous structures were deformed into a major mega-antiform with a NW-SE axis, which is recognizable at the scale of the whole promontory (see geological sections in fig. 10). Also the Zuccale fault shows local weak folding and an overall antiformal attitude, from Zuccale to Porto Azzurro through Capoliveri areas. We think that this mega-structure can be related to the final uplift of the consolidated pluton. Finally, the NW-SE and NE-SW-trending high-angle normal faulting (D<sub>5</sub> event) occurred and modelled the western (e.g. the west-downthrown system of the Praticciolo-Innamorata-Gemini islands), southern (e.g. Remaiolo-Punta dei Ripalti) and eastern coasts (e.g. Capo Calvo-Forte Focardo alignment). These locally mineralized faults are probably coeval with the high-angle normal faults of eastern Elba (cfr. BORTOLOTTI *et alii*, 2001), which host Adu+Hem ores dated about 5 Ma (LIPPOLT *et alii*, 1995).

## CONCLUSIONS

The 1:10,000 scale geological mapping of the Mt. Calamita promontory and the stratigraphical, petrographical and structural studies allow the following conclusions:

1) The Porto Azzurro Unit consists of Paleozoic, probably pre-Carboniferous successions (Mt. Calamita Metamorphic Complex), which are unconformably overlain by the Triassic Verrucano metasiliciclastics (Barabacca Quartzites) and Upper Triassic-?Hettangian metacarbonates. The Mt. Calamita Metamorphic Complex mostly consists of a deep garnet-bearing, albite micaschist unit (lithofacies *a*), similar to the Micaschist complex of the Larderello subsurface, which underlies a phyllitic-quartzitic unit (lithofacies *b*), probably correlatable with the Ordovician formations of the Tuscan Mid-Ridge (e.g. Buti banded quartzite and phyllite in RAU & TONGIORGI, 1974) and the Ortano Unit in the eastern Elba. This correlation is also strengthened by the existence of local porphyroid-like horizons. The analysed amphibolite horizons of the Mt. Calamita Metamorphic Complex probably derive from the metamorphism of within-plate basalts.

2) The deformation and metamorphic evolution of the Porto Azzurro Unit is more complex than previously hypothesized. Apart from the relicts of the Variscan tectono-metamorphism recognized in the Mt. Calamita Metamorphic Complex, all the formations of the Porto Azzurro Unit suffered two Alpine tectono-metamorphic events (D<sub>1</sub> and D<sub>2</sub>) under greenschist facies metamorphic conditions similar to most of the other Tuscan metamorphic units in Tuscany. In Messinian-Lower Pliocene times, the emplacement of anatetic granitoids produced

thermometamorphism in the host rocks and weak D<sub>3</sub> folding, followed by low-angle detachments. The uplift of the granitic body (D<sub>4</sub> event) produced the Apennine-trending mega-antiform at the scale of the whole promontory. Finally, this complex structural and magmatic assemblage was dissected by NW-SE and NE-SW-trending fault systems.

3) The stratigraphic and most of the structural and magmatic-hydrothermal evolution of the Calamita promontory show close analogies with that of the buried metamorphic units of the Larderello geothermal region (cfr. BATINI *et alii*, 1983; ELTER & PANDELI, 1990, 1993; PANDELI *et alii*, 1994; BERTINI *et alii*, 1991; DECANDIA *et alii*, 2001; MUSUMECI *et alii*, 2002; GIANELLI & RUGGIERI, 2002 and references therein). Thus, the Mt. Calamita area can be considered an analogous geological model for the studies of the deep structure of southern Tuscany, and in particular of the Larderello-Travale geothermal region, crossed by the CROP 18 profile.

## ACKNOWLEDGEMENTS

We thank the referees (Prof. C.A. Ricci, University of Siena and Prof. R. Carosi, University of Pisa) for the critical reading and improvement of the paper. This research work was supported by the Projet COFIN MURST 2003 - Tectonic Process in Accretionary Prism (coordinator Prof. G. Principi), by the ex 60% Grant (Prof. G. Principi) and CNR-IGG grants.

## REFERENCES

- BARBERI F. & INNOCENTI F. (1965) - Studio chimico-petrografico di un'anfibolite rinvenuta negli scisti termometamorfici di Capo Calamita (Isola d'Elba). Boll. Soc. Geol. It., **84**, 41-92.
- BARBERI F., GIGLIA G., INNOCENTI F., MARINELLI G., RAGGI G., RICCI C.A., SQUARCI P., TAFFI L. & TREVISAN L. (1967a) - Carta geologica dell'isola d'Elba. Scala 1:25.000. C.N.R. Roma.
- BARBERI F., INNOCENTI F. & RICCI C.A. (1967b) - Il Complesso Sciustoso di Capo Calamita (Isola d'Elba). Atti Soc. Toscana Sci. Nat. Pisa, Mem., P.V., Ser. A, **72** (2), 579-617.
- BARBERI F., DALLAN L., FRANZINI M., GIGLIA G., INNOCENTI F., MARINELLI G., RAGGI R., RICCI C.A., SQUARCI P., TAFFI L. & TREVISAN L. (1969a) - Isola d'Elba, Foglio 126. Carta Geologica d'Italia. Serv. Geol. d'It., E.I.R.A., Firenze.
- BARBERI F., DALLAN L., FRANZINI M., GIGLIA G., INNOCENTI F., MARINELLI G., RAGGI R., RICCI C.A., SQUARCI P., TAFFI L. & TREVISAN L. (1969b) - Foglio 126 (Isola d'Elba). Note illustrative della Carta Geologica d'Italia alla scala 1:100.000. Serv. Geol. d'It., 32 pp.
- BARTOLE R. (1995) - The North Tyrrhenian-Northern Apennines post-collisional system: constraints for a geodynamic model. Terra Nova, **7**, 7-30.
- BARTOLE R., TORELLI L., MATTEI G., PEIS D. & BRANCIOLINI G. (1991) - Assetto stratigrafico-strutturale del Tirreno settentrionale: stato dell'arte. Studi Geologici Camerti, vol. spec. 1991/1, 115-140.
- BATINI F., BERTINI G., GIANELLI G. & PANDELI E. (1983) - Deep structure of the Larderello field. Mem. Soc. Geol. Ital., **25**, 219-235.
- BERTINI G., CAMELI G.M., COSTANTINI A., DECANDIA F.A., DI FILIPPO M., DINI I., ELTER F.M., LAZZAROTTO A., LIOTTA D., PANDELI E., SANDRELLI F. & TORO B. (1991) - Struttura geologica fra i Monti di Campiglia e Rapolano Terme (Toscana meridionale): stato attuale delle conoscenze e problematiche. Studi Geologici Camerti, vol. spec. 1991/1, 155-178.
- BORTOLOTTI V., FAZZUOLI M., PANDELI E., PRINCIPI G., BABBINI A. & CORTI S. (2001) - Geology of Central and Eastern Elba Island, Italy. Ofioliti, **26** (2a), 97-150.
- BRUNET C., MONIÉ P., JOLIVET L. & CADET J.P. (2000) - Migration of compression and extension in the Tyrrhenian Sea, insights from <sup>40</sup>Ar/<sup>39</sup>Ar ages on micas along a transect from Corsica to Tuscany. Tectonophysics, **33** (21), 127-155.

- CAMPETTI A., CAROSI R., DECANDIA F.A., ELTER P., LAZZAROTTO A., MONTOMOLI C., PERTUSATI P.C. & POLVEROSI B. (1999) - *Assetto ed evoluzione strutturale dei Monti dell'Uccellina (Toscana Meridonale)*. Boll. Soc. Geol. It., **108**, 149-166.
- CARMIGNANI L. Coord. (2001) - *Note illustrative della Carta Geologica della Sardegna alla scala 1:200.000. Memorie descrittive della Carta Geologica d'Italia*, **60**, 283 pp.
- CARMIGNANI L. & KLINGFIELD R. (1990) - *Crustal extension in the northern Apennines: the transition from compression to extension in the Alpi Apuane core complex*. Tectonics, **9**, 1275-1303.
- CARMIGNANI L., DECANDIA F.A., FANTOZZI P.L., LAZZAROTTO A., LIOTTA D. & OGGIANO G. (1994a) - *Tertiary extensional tectonics in Tuscany (Northern Apennines, Italy)*. Tectonophysics, **238**, 295-315.
- CAROSI R., CERBAI N. & MONTOMOLI C. (1995) - *Deformation history of the Verrucano of Pisani Mts. (Northern Apennines, Italy)*. Annales Tectonicae, **9**, 55-74.
- CAROSI R., MONTOMOLI C., BERTUCCELLI N. & PROFETI M. (2002) - *The structural evolution of the southern Apuan Alps: new constraints on the tectonic evolution of the Northern Apennines (Italy)*. Comptes Rendus Geoscience, **334**, 339-346.
- CAVARETTA G., PUDEXDU M., FRANCESCHELLI M., PANDELLI E. & VALORI A. (1989) - *Tourmalinates from hydrothermal systems related to Permian rift magmatism of the Hercynian Saalian phase in Tuscany (Italy)*. In: D.L. Miles (Ed.) - Proceedings of the Sixth International Symposium on Water-Rock Interaction held in Malvern (GB), 3-8 August 1989. A.A. Balkema/Rotterdam/Brookfield/1989, 141-144.
- CAVARETTA G., PUDEXDU M., FRANCESCHELLI M., PANDELLI E. & VALORI A. (1992) - *Tourmalinates from the Triassic Verrucano of the Northern Apennines, Italy*. In: L. Carmignani & F.P. Sassi (Eds.) - «Contribution to the Geology of Italy with special regard to the Paleozoic basements». A volume dedicated to Tommaso Cocozza. IGCP No. 276, Newsletter, Vol. 5, spec. issue, Centrooffset, Siena 1992, 335-338.
- COLLETTINI C. & BARCHI M. (2004) - *A comparison of structural data and seismic images for low-angle normal faults in the Northern Apennines (Central Italy): constraints on activity*. In: Alsop G.I. & Holdsworth R.E. (Eds.) - Flow processes in faults and shear zones. Geol. Soc. London, Spec. Pubbl., **224**, 95-112.
- CONTI P., COSTANTINI A., DECANDIA F.A., ELTER F.M., GATTIGLIO M., LAZZAROTTO A., MECCHERI M., PANDELLI E., RAU A., SANDRELLI F., TONGIORGI M. & DI PISA A. (1991) - *Structural frame of the Tuscan Paleozoic: a review*. Boll. Soc. Geol. It., **110**, 523-541.
- CORSI B., ELTER F.M., PANDELLI E. & SANDRELLI F. (2001) - *Caratteri strutturali del Gruppo del Verrucano (Unità di Monticiano-Roccastrada) nella Toscana meridionale ed insulare*. Atti Tic. Sc. Terra, **42**, 47-58.
- COSTANTINI A., DECANDIA F.A., LAZZAROTTO A. & SANDRELLI F. (1987/88) - *L'Unità di Monticiano Roccastrada tra la Montagnola Senese e il Monte Leoni (Toscana meridionale)*. Atti Tic. Sci. Terra, **31**, 382-420.
- COSTANTINI A., GANDIN A., MATTIAS P.P., SANDRELLI F. & TURI B. (1980) - *Un'ipotesi per l'interpretazione paleogeografica della Formazione di Tocchi*. Mem. Soc. Geol. It., **21**, 203-216.
- DANIEL & JOLIVET (1995) - *Detachment faults and pluton emplacement: Elba Island (Tyrrhenian Sea)*. Bull. Soc. Géol. France, **166**, 341-354.
- DECANDIA F.A., LAZZAROTTO A. & LOTTA D. (2001) - *Structural features of southern Tuscany*. Ofioliti, **26**, 287-300.
- DEINO A., KELLER J.V.A., MONELLI G. & PIALLI G. (1992) - *Datazioni  $^{40}\text{Ar}/^{39}\text{Ar}$  del metamorfismo dell'Unità di Ortano-Rio Marina (Isola d'Elba): risultati preliminari*. Studi Geologici Camerti, vol. spec. 1992/2, CROP 1-1A, 187-192.
- DESCHAMPS Y., DAGALLIER G., MACAUDIERE L., MARIGNAC C., MOINE B. & SAUPÉ F. (1983) - *Le gisement de pyrite-hématite de Valle Giove (Rio Marina Ile d'Elbe, Italie)*. Schweiz. Mineral. Petrogr. Mitt., **63**, 149-165.
- DIMANCHE F. (1971) - *Les minéraux de magnetite et les skarns du Ginevro (Ile d'Elbe - Italie)*. Mineral. Dep., **6**, 356-379.
- DINI A., INNOCENTI F., ROCCHI S., TONARINI S. & WESTERMAN D.S. (2002) - *The magmatic evolution of the late Miocene laccolith-pluton-dyke granitic complex of Elba Island, Italy*. Geol. Mag., **139**, 257-279.
- DURANTI S., PALMERI, PERTUSATI P.C. & RICCI C.A. (1992) - *Geological evolution and metamorphic petrology of the basal sequences of Eastern Elba (Complex II)*. Acta Vulca., Marinelli, **2**, 213-229.
- ELTER F.M. & PANDELLI E. (1990) - *Alpine and Hercynian orogenic phases in the basement rocks of the Northern Apennines (Larderello geothermal field, southern Tuscany, Italy)*. Eclogae geol. Helv., **83** (2), 241-264.
- ELTER F.M. & PANDELLI E. (1993) - *Alpine tectono-metamorphic framework of the Tuscan Paleozoic (Southern Tuscany, Italy)*. Annales Tectonicae, **7**, 71-84.
- ELTER F.M. & PANDELLI E. (1994) - *Caratteri petrografici e strutturali del basamento metamorfico nel sottosuolo dell'area geotermica di Travale-Radicondoli (Toscana meridionale, Italia)*. Studi Geologici Camerti, Vol. spec. 1994/1, 103-112.
- ELTER F.M. & PANDELLI E. (1996) - *Structural setting of the Paleozoic Crystalline Basement of the Northern Apennines (Italy)*. In: O. Oncken & C. Janssen (Eds.) - Proceedings of the Eleventh International Conference on Basement Tectonics «Basement Tectonics 11-Europe and other regions» held in Potsdam, Germany, July 1994-Kluwer Academic Publishers Dordrecht/Boston/London, 79-90.
- ELTER F.M. & PANDELLI E. (2001) - *Structural evolution of anchiz/epimetamorphic units of Central and Eastern Elba (Ortano, Acquadolce, Monticiano-Roccastrada and Grassera Units)*. Ofioliti, **26** (2a), 219-228.
- ELTER F.M. & SANDRELLI F. (1994) - *La fase post-Nappe nella Toscana Meridionale: nuova interpretazione nell'evoluzione dell'Appennino Settentrionale*. Atti Tic. Sc. Terra, **37**, 173-193.
- FASANO F., LIOTTA D. & RICCI C.A. (2001) - *Hercynian and Apennine deformations superimpositions in Southern Tuscany: the case of Mt. Calamita-Elba Island*. In: Stratigraphic and structural evolution on the Late carboniferous to Triassic continental and marine successions in Tuscany (Italy). Regional reports and general correlation. Abstract volume, 36-37. April 30-May 7, 2001. Siena, Italy.
- FAZZUOLI M., PANDELLI E. & SANI F. (1994) - *Considerations on the sedimentary and structural evolution of the Tuscan Domain since Early Liassic to Tortonian*. Mem. Soc. Geol. It., **48**, 31-50.
- FERRARA G. & TONARINI S. (1993) - *L'Isola d'Elba: un laboratorio di geocronologia*. Mem. Soc. Geol. It., **49**, 227-232.
- GIANELLI G. & RUGGIERI G. (2002) - *Evidence of a contact metamorphic aureole with high-temperature metasomatism in the deepest part of the active geothermal system of Larderello*. Geothermics, **31**, 443-474.
- INTERNATIONAL SUBCOMMISSION ON STRATIGRAPHIC CLASSIFICATION (AMOS SALVADOR, CHAIRMAN) (1987) - *Stratigraphic classification and nomenclature of igneous and metamorphic rock bodies*. Geological Society of America Bulletin, **99**, 440-442.
- JUTEAU M., MICHAUD A., ZIMMERMANN J.L. & ALBAREDE F. (1984) - *Isotopic heterogeneities in the granitic intrusion of Monte Capanne (Elba Island, Italy) and dating concepts*. J. Petrol., **25**, 532-545.
- KELLER J.V.A. & PIALLI G. (1990) - *Tectonics of the Island of Elba: a reappraisal*. Boll. Soc. Geol. It., **109**, 413-425.
- KELLER J.V.A. & COWARD M.P. (1996) - *The structure and evolution of the Northern Tirrenian sea*. Geol. Mag., **133**, 1-16.
- KLINGFIELD R., HUNZIKER J., DALLMEYER R.D. & SCHAMEL S. (1986) - *Dating of deformation phases using K-Ar and  $^{40}\text{Ar}/^{39}\text{Ar}$  techniques: result from the Northern Apennines*. J. Struct. Geol., **8** (7), 781-798.
- KRETZ R. (1983) - *Symbols for rock-forming minerals*. Amer. Mineral., **68**, 277-279.
- LIPPOLT H., WERNICKE R.S. & BÄHR R. (1995) - *Paragenetic specularite and adularia (Elba, Italy): Concordant (U+Th)-He and K-Ar ages*. Earth Plan. Sci. Lett., **132**, 43-51.
- MARINELLI G. (1959) - *Le intrusioni terziarie dell'Isola d'Elba*. Atti Soc. Tosc. Sci. Nat., Mem., Ser. A, **66**, 50-223.
- MAINERI C., BENVENUTI M., COSTAGLIELA P., DINI A., LATTANTI P., RUGGIERI G. & VILLA I.M. (2003) - *Sericitic alteration at the La Crocetta deposits (Elba Island, Italy): interplay between magmatism, tectonic and idrothermal activity*. Mineralium Deposita, **38**, 67-86.

- MAZZANTI R. (1983) - *Il punto sul Quaternario della fascia costiera e dell'arcipelago di Toscana*. Boll. Soc. Geol., It., **102**, 419-556.
- MOLLI G., MONTANINI A. & FRANK W. (2002) - *MORB-derived Variscan amphibolites in the northern Apennine Basement: The Cerreto metamorphic slices (Tuscan-Emilian Apennine, NW Italy)*. Ophioliti, **27**, 17-30.
- MUSUMECI G., BOCINI L. & CORSI R. (2002) - *Alpine tectonothermal evolution of the Tuscan Metamorphic Complex in the Larderello geothermal field (northern Apennines, Italy)*. Journal of the Geological Society, London, **159**, 443-456.
- PANDELI E. (2002) - *Sedimentary-tectonic evolution of the Tuscan area (Northern Apennines, Italy) from Late «Autunian» to Carnian*. Boll. Soc. Geol. It., Vol. Spec. **1** (2002), 251-262.
- PANDELI E., GIANELLI G., PUXEDDU M. & ELTER F.M. (1994) - *The paleozoic basement of the Northern Apennines: stratigraphy, tectono-metamorphic evolution and Alpine hydrothermal processes*. Mem. Soc. Geol. It., **48**, 627-654.
- PANDELI E., BORTOLOTTI V. & PRINCIPI (1995) - *La successione toscana epimetamorfica di Capo Castello (Cavo, isola d'Elba nord-orientale)*. Atti Tic. Sc. Terra, **38**, 171-191.
- PANDELI E., CORTI S., FRANCESCHELLI M. & PECHIONI E. (2001a) - *The Varicoloured Slates of the Grassera Unit (Central-Eastern Elba, Tuscany): petrological-mineralogical data comparisions with other Tuscan and Ligurian-Piemontese Units*. Ophioliti, **26** (2a), 197-206.
- PANDELI E. & PUXEDDU M. (1990) - *Paleozoic age for the Tuscan upper metamorphic sequences of Elba and its implications for the geology of the Northern Apennines (Italy)*. Ecl. Geol. Helv., **83** (1), 123-142.
- PANDELI E., PUXEDDU M. & RUGGIERI G. (2001b) - *The metasilicic-clastic-carbonate sequence of the Acquadolce Unit (Eastern Elba Island): new petrographic data and paleogeographic interpretation*. Ophioliti, **26** (2a), 207-218.
- PERRIN M. (1975) - *L'Ile d'Elbe et la limite alpes-Apennin: données sur la structure géologique et l'évolution tectogénétique de l'Elbe alpine et de l'Elbe apennine*. Boll. Soc. Geol. It., **94**, 1929-1955.
- PERTUSATI P.C., RAGGI G., RICCI C.A., DURANTI S. & PALMERI R. (1993) - *Evoluzione post-collisionale dell'Elba centro-orientale*. Mem. Soc. Geol. It., **49**, 297-312.
- PUXEDDU M., SAUPÉ F., DÉCHOMETS R., GIANELLI G. & MOINE B. (1984) - *Geochemistry and stratigraphic correlations-application to the investigation of geothermal and mineral resources of Tuscany, Italy*. Chem. Geol., **43**, 77-113.
- SAUPÉ F., MARIGNAC C., MOINE B., SONET J. & ZIMMERMANN J.L. (1982) - *Datation par les méthodes K/Ar et Rb/Sr de quelques roches de la partie orientale de l'Ile d'Elbe (Province de Livourne, Italie)*. Bull. Mineral., **105**, 236-245.
- RAU A. & TONGIORGI M. (1974) - *Geologia dei Monti Pisani a Sud-Est della Valle del Guappero*. Mem. Soc. Geol. It., **13**, 227-408.
- TANELLI G. (1977) - *I giacimenti a skarn della Toscana*. Rend. Soc. It. Min. Petr., **33** (2), 875-903.
- TANELLI G., BENVENUTI M., COSTAGLIOLA P., DINI A., LATTANZI P., MANIERI C., MASCARO I. & RUGGIERI G. (2001) - *The iron mineral deposits of Elba island: state of the art*. Ophioliti, **26** (2a), 239-248.
- TREVISAN L. (1950) - *L'Elba orientale e la sua tettonica di scivolamento per gravità*. Mem. Ist. Geol. Univ. Padova, **16**, 5-39.
- TREVISAN L. (1951) - *La 55<sup>a</sup> Riunione Estiva della Società Geologica Italiana. Isola d'Elba, 18-23 Settembre 1951*. Boll. Soc. Geol. It., **70**, 435-472.

*Manoscritto pervenuto il 5 Maggio 2004; testo approvato per la stampa l'11 Novembre 2004; ultime bozze restituite il 7 Aprile 2005.*