



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Cardiac and vascular remodeling in older adults with borderline isolated systolic hypertension: The ICARe Dicomano Study.**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Cardiac and vascular remodeling in older adults with borderline isolated systolic hypertension: The ICARe Dicomano Study / R. PINI; M.C. CAVALLINI; F. BENCINI; L. STAGLIANO; E. TONON; F. INNOCENTI; G. BALDERESCHI; N. MARCHIONNI; M. DI BARI; R.B. DEVEREUX; G. MASOTTI; MJ. ROMAN. - In: HYPERTENSION. - ISSN 0194-911X. - STAMPA. - 38:(2001), pp. 1372-1376.

*Availability:*

This version is available at: 2158/309664 since:

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

# Cardiac and Vascular Remodeling in Older Adults With Borderline Isolated Systolic Hypertension

## The ICARe Dicomano Study

Riccardo Pini, M. Chiara Cavallini, Francesca Bencini, Loredana Staglianò, Elisabetta Tonon, Francesca Innocenti, Giorgio Baldereschi, Niccolò Marchionni, Mauro Di Bari, Richard B. Devereux, Giulio Masotti, Mary J. Roman

**Abstract**—Although borderline isolated systolic hypertension (ISH), defined as a blood pressure of 140 to 159/ $<90$  mm Hg, is a proven cardiovascular risk factor, the major clinical trials on treatment of ISH have used a cutoff of 160 mm Hg. Moreover, no data exist on the cardiovascular modifications associated with borderline ISH. Therefore, we compared subjects with borderline ISH to subjects with diastolic hypertension (diastolic blood pressure  $\geq 90$  mm Hg) or ISH. Community-dwelling residents (age  $\geq 65$  years) of a small town in Italy (Dicomano) underwent extensive clinical examination, echocardiography, carotid ultrasonography, and applanation tonometry. Only untreated subjects were included in this analysis: 95 with diastolic hypertension, 87 with borderline ISH, and 43 with ISH. Despite lower systolic and mean pressures in borderline ISH, left ventricular mass was similar to that in diastolic hypertension. In univariate and multivariate analysis, pulse pressure but not systolic pressure was related to left ventricular mass. Borderline ISH subjects had a tendency to greater carotid cross-sectional area and stiffness index than did diastolic hypertensive subjects despite lower mean carotid pressure, whereas the number of atherosclerotic plaques was similar in the 2 groups. Pulse pressure but not systolic pressure was independently related to carotid remodeling. In our community-based, older population, individuals with borderline ISH had a similar prevalence of left ventricular hypertrophy and carotid atherosclerosis as that of subjects with diastolic hypertension, despite lower systolic and mean pressures. Among blood pressure values, pulse pressure was the single or strongest independent predictor of cardiovascular remodeling. (*Hypertension*. 2001;38:1372-1376.)

**Key Words:** age ■ carotid arteries ■ atherosclerosis ■ arterial compliance ■ hypertrophy

The prevalence of isolated systolic hypertension (ISH), defined as a systolic blood pressure (SBP) of  $\geq 160$  mm Hg with a diastolic blood pressure (DBP)  $<90$  mm Hg, increases with age and becomes particularly high among subjects  $>60$  years of age.<sup>1</sup> In 1993, the Fifth Joint National Committee on High Blood Pressure changed the definition of ISH by reducing the SBP threshold from 160 to 140 mm Hg.<sup>2</sup> This change was based on the observation that subjects with borderline ISH, defined as a SBP of 140 to 159 mm Hg with a DBP  $<90$  mm Hg, suffered a higher incidence of cardiovascular death than did normotensive subjects.<sup>3</sup> However, despite this evidence, the major clinical trials on the effects of antihypertensive treatment of ISH have used a cutoff of 160 mm Hg.<sup>4,5</sup> Moreover, no data are available regarding the cardiac and vascular modifications associated with borderline ISH. Therefore, the present study was undertaken to analyze the cardiac and vascular characteristics of subjects with borderline ISH compared with subjects with diastolic (essential) hypertension (HTN) or ISH.

## Methods

### Study Population

All community-dwelling residents aged  $\geq 65$  years of a small town in Italy (Dicomano) underwent an extensive assessment consisting of home interview, laboratory testing, and clinical examination. Further clinical information was gathered from primary care physicians of the participants with a structured questionnaire regarding the history and treatment of clinical conditions, including hypertension. The general design of this survey (ICARe Dicomano Study) has been previously published in detail.<sup>6</sup>

Blood pressure was measured by the first and fifth Korotkoff phases using a cuff sized for arm circumference. Three measures were obtained 1 to 2 minutes apart; the second and the third blood pressure measurements were averaged, and mean values were considered as the reference SBP and DBP. Mean blood pressure was calculated as  $DBP + [0.33 \times \text{pulse pressure}]$ .

Subjects receiving antihypertensive medication were excluded from the present study. Enrolled subjects were grouped according to the following criteria: HTN is  $DBP \geq 90$  mm Hg; borderline ISH is  $SBP 140$  to  $159$  mm Hg and  $DBP < 90$  mm Hg, and ISH is  $SBP \geq 160$  mm Hg and  $DBP < 90$  mm Hg. Subjects with blood pressure

Received March 14, 2001; first decision April 6, 2001; revision accepted July 10, 2001.

From the Department of Critical Care Medicine and Surgery-Unit of Gerontology and Geriatrics, University of Firenze, Firenze, Italy, the Azienda Ospedaliera Careggi, Firenze, Italy; and the Department of Medicine, New York Presbyterian Hospital-Weill Medical College of Cornell University (R.B.D., M.J.R.).

Correspondence to Riccardo Pini, MD, Unit of Gerontology and Geriatrics, Via delle Oblate 4, 50141 Firenze, Italy. E-mail: rpini@unifi.it  
© 2001 American Heart Association, Inc.

*Hypertension* is available at <http://www.hypertensionaha.org>



TABLE 2. Echocardiographic Characteristics of Hypertensive Subjects

Variable	HTN (n 94)	P(1 vs 2)	Borderline ISH (n 85)	P(2 vs 3)	ISH (n 43)	P(1 vs 3)	P
IVS <sub>d</sub> , mm	8.4 1.7		8.5 1.9		9.0 2.2		NS
PWT <sub>d</sub> , mm	8.1 1.4		7.8 1.3		8.4 1.4		NS
LVID <sub>diastolic</sub> , mm	51.6 6.9		52.3 6.0		53.0 4.9		NS
LVID <sub>systolic</sub> , mm	32.4 7.0		32.2 6.4		32.6 6.4		NS
LV mass, g	153 49		153 47		170 53		NS
LV mass index, g/m <sup>2</sup>	89 26	NS	94 27	NS	102 26	0.048	0.047
LV relative WT	0.32 0.09		0.30 0.06		0.32 0.05		NS
LV hypertrophy (LVMI)	13 (14%)	NS	19 (22%)		13 (30%)		NS 0.049
Fractional shortening, %	38		39 7		39 9		NS
E wave, m/sec	0.58 0.16		0.62 0.17		0.61 0.19		NS
A wave, m/sec	0.78 0.16	NS	0.77 0.17	0.016	0.87 0.19	0.042	0.012
E wave/A wave	0.74 0.18	0.031	0.84 0.33	0.010	0.69 0.18	NS	0.003
Relaxation time, msec	99		98 16		95 21		NS
Deceleration time, msec	262	NS	221 54	NS	248 71	NS	0.021
Significant valvular disease	13 (14%)		15 (18%)		7 (16%)		NS

IVS<sub>d</sub> indicates inter-ventricular septum thickness in diastole; PWT<sub>d</sub>, left ventricular posterior wall thickness in diastole; LVID, left ventricular internal diameter; WT, wall thickness; and LVMI, left ventricular mass index.

multiple linear regression analysis. Two-tailed 0.05 was considered significant.

An expanded Methods section can be found in an online data supplement available at <http://www.hypertensionaha.org>.

## Results

### Clinical Features of Hypertensive Patients

Demographic features, laboratory data, and pertinent cardiovascular history of the 3 groups are compared in Table 1. Subjects with ISH were older and had lower body mass index than patients with HTN, but no differences were found in gender distribution, height, or body surface area. Compared with HTN subjects, borderline ISH subjects had lower weight and body surface area. By definition, borderline ISH subjects had lower blood pressure values than did ISH patients; however, when compared with HTN patients, borderline ISH patients had lower SBPs, DBPs, and mean blood pressures with similar PP. Lipid profile tended to be most favorable in the ISH, but differences did not reach statistical significance; no differences were found between the 3 groups regarding previous history of stroke, transient ischemic attack, and coronary artery or peripheral vascular diseases. Diabetes tended to be more prevalent in the older ISH subjects than in borderline ISH or HTN patients. The percentage of current or former smokers was similar in the 3 groups.

### Echocardiographic Findings in the HTN Group and ISH Groups

Echocardiographic findings in the 3 groups are compared in Table 2. LV mass and LV mass indexed for body surface area in borderline ISH were similar to those in HTN subjects despite lower systolic and mean pressures. ISH subjects tended to have higher LV mass than HTN subjects, and this difference became significant when indexing for body surface area. Borderline ISH patients had increased LV mass index compared with that of previously reported normotensive

subjects (83 23 g/m<sup>2</sup>, P 0.023), whereas HTN patients did not. When this index was used to identify patients with LV hypertrophy, borderline ISH and ISH patients exhibited a 1.5- to 2-fold increase in the prevalence of LV hypertrophy (22% and 30%, respectively) compared with that of HTN patients (14%), whereas normotensive subjects had a lower prevalence (9%). LV mass exhibited a positive correlation with PP (r 0.182, P 0.010), whereas LV mass index correlated positively with PP (r 0.230, P 0.001) and negatively with DBP (r -0.169, P 0.016). When all blood pressure measures (SBP, DBP, mean blood pressure, and PP) were included in a multivariate analysis, both LV mass and LV mass index were independently related only to PP (r 0.182, P 0.010 and 0.230, P 0.001, respectively). LV fractional shortening, an index of systolic function, was similar in the 3 groups. Borderline ISH patients had a significantly greater E/A velocity ratio and a tendency to a shorter deceleration time than did the other 3 groups.

### Comparison of Carotid Artery Structure and Stiffness in Hypertensive Groups

In the borderline ISH group, the carotid intimal-medial thickness and internal diameters were intermediate between the values observed in the HTN and ISH groups, despite having lower DBPs and mean blood pressures than those of the HTN patients (Table 3), but the differences between the 3 groups did not reach statistical significance. The carotid cross-sectional area, an index of vascular remodeling similar to LV mass, tended to increase from HTN to borderline ISH to ISH patients. A similar trend was present when carotid cross-sectional area was indexed for body surface area. When included in a multivariate analysis model, only PP was independently related to carotid cross-sectional area, both absolute (r 0.270, P 0.0001) and indexed (r 0.289,

TABLE 3. Carotid Artery Pressure, Structure, and Stiffness in Hypertensive Subjects

Variable	HTN (n=95)	P (1 vs 2)	Borderline ISH (n=87)	P (2 vs 3)	ISH (n=42)	P (1 vs 3)	P
SBP, mm Hg	142±19	NS	137±17	<0.0001	154±23	0.009	<0.0001
DBP, mm Hg	83±11	<0.0001	75±9	NS	73±9	<0.0001	<0.0001
MBP, mm Hg	103±13	0.001	95±10	NS	100±11	NS	0.001
PP, mm Hg	59±14	NS	62±15	<0.0001	81±21	<0.0001	<0.0001
IMT, mm	0.83±0.16		0.84±0.19		0.89±0.17		NS
Diastolic diameter, mm	6.09±0.96		6.16±1.04		6.44±1.07		NS
Systolic diameter, mm	6.85±0.96		7.01±1.09		7.35±1.17		NS
Systolic expansion, %	13±4		14±5		14±5		NS
RWT	0.28±0.07		0.28±0.07		0.28±0.07		NS
Cross-sectional area, mm <sup>2</sup>	17.98±4.19	NS	18.68±5.72	NS	20.60±5.48	0.040	0.043
CSA index, mm <sup>2</sup> /m <sup>2</sup>	10.60±2.55	NS	11.46±3.41	NS	12.46±3.51	0.013	0.012
Subjects with plaques, %	72 (76%)		68 (78%)		36 (86%)		NS
Plaque number	2±2	NS	2±2	0.007	3±2	0.001	0.001
Plaque maximum thickness, mm	2.14±0.62	NS	2.33±0.66	NS	2.47±0.66	0.048	0.041
Beta stiffness index	4.99±2.33	NS	5.19±2.95	NS	6.42±3.03	NS	0.030
Reflected wave, mm Hg	18±10	NS	21±11	NS	25±14	0.014	0.009
AI (corrected for PP)	0.30±0.12		0.32±0.13		0.30±0.15		NS
AI (corrected for MBP)	0.17±0.08	0.035	0.22±0.11	NS	0.25±0.13	0.002	0.001

IMT indicates intimal-medial thickness; RWT, carotid relative wall thickness; CSA index, cross-sectional area/body surface area; and AI, augmentation index.

P<0.0001). Number and thickness of carotid plaques and carotid artery Beta stiffness index exhibited an increase from HTN to borderline ISH to ISH group; however, these values were not statistically different between borderline ISH and HTN subjects. In univariate analysis, the number of carotid plaques was independently related to PP (0.273, P<0.01) but not to SBP. Both borderline ISH and HTN patients had stiffer carotid arteries than did normal subjects (Beta stiffness index=3.97±1.93, P=0.006 and P=0.041, respectively). In multivariate analysis, only carotid PP (β=0.590, P<0.0001) and SBP (β=-0.448, P<0.0001) were related to the stiffness index. The negative correlation coefficient of SBP is an indication of a strong collinearity between PP and SBP; in fact, when PP was removed from the model, SBP exhibited a positive coefficient β=0.228, P=0.016). Despite similar carotid SBP, borderline ISH patients had higher augmentation index, adjusted for mean blood pressure, than did HTN or normotensive subjects (0.15±0.09, P<0.0001).

### Discussion

In our population aged ≥65 years, subjects with borderline ISH demonstrated the same cardiac and vascular abnormalities as those of patients with HTN, despite lower values of systolic and mean blood pressure. Moreover, borderline ISH patients exhibited comparable prevalences of ischemic heart disease and cerebrovascular events as those of patients with HTN or ISH.

Recently, several observational studies have yielded evidence to suggest that in older subjects, PP compared with SBP may be a better predictor of cardiovascular events. Despite these data, the classification of hypertension is still based on SBP and DBP without special consideration of PP.

The present study supports previous observations that PP is a stronger predictor of cardiovascular disease than SBP. In fact, when comparing borderline ISH to HTN subjects, the 2 groups exhibited a similar prevalence of established cardiac and cerebrovascular diseases despite the lower values of SBP and mean blood pressure that were found in the borderline ISH group. Moreover, this study indicates possible underlying mechanisms to explain the strong association of PP with cardiac and cerebrovascular events. In fact, borderline ISH patients exhibited cardiac and carotid remodeling that was similar to ISH patients, an observation that has never previously been documented in a geriatric population. Thus, based on our observations and previous studies, older patients with borderline ISH may be especially likely to benefit from antihypertensive treatment.

### Potential Study Limitations

Our data were collected from a population of older white subjects living in a small town in Italy; therefore, the conclusions derived from the results presented in this study may not be directly applicable to subjects in other geographical areas or with a different racial composition. However, the prevalence of borderline ISH found in our population (17%) is similar to that of previous studies (20 to 26%). In this study, subjects were classified as borderline ISH based on the study blood pressure measurements and may include individuals with white-coat hypertension. Recently, Fagard et al found that 24% of 700 ISH patients had SBP <140 mm Hg on ambulatory blood pressure monitoring. However, because white-coat hypertensive patients have less cardiac and vascular remodeling than do sustained hypertensive patients, the exclusion of these patients might increase



the association of PP with cardiovascular remodeling and its importance when compared with SBP.

### Acknowledgments

Sponsored in part by the Italian Ministry of Scientific and Technological Research (National Special Project on Heart Failure), by the Regional Government of Tuscany, and by the Azienda Ospedaliera Careggi-Firenze, Italy.

### References

- Black HR. Isolated systolic hypertension in the elderly: lessons from clinical trials and future directions. *J Hypertens* 1999;17(suppl 5):S49-S54.
- The fifth report of the Joint National Committee on detection, evaluation, and treatment of high blood pressure (JNC V). *Arch Intern Med* 1993;153:154-183.
- Sagie A, Larson MG, Levy D. The natural history of borderline isolated systolic hypertension. *N Engl J Med* 1993;329:1912-1917.
- Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991;265:3253-3264.
- Staessen JA, Fagard R, Thijs L, Celis H, Arabadzisz G, Birkenhager WH, Bulpitt CJ, de Leeuw PW, Dollery CT, Fletcher AE, Forette F, Leonetti G, Nachev C, O'Brien ET, Rosenfeld J, Rodicio JL, Tuomilehto J, Zanchetti A. Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension: the Systolic Hypertension in Europe (Syst-Eur) Trial Investigators. *Lancet* 1997;350:757-764.
- Di Bari M, Marchionni N, Ferrucci L, Pini R, Antonini E, Chiarlone M, Marsilli A, De Alfieri W, Fumagalli S, Masotti G. Heart failure in community-dwelling older persons: aims, design and adherence rate of the ICARE Dicomano project: an epidemiologic study. *Insufficienza Cardiaca negli Anziani Residenti a Dicomano*. *Am Geriatr Soc* 1999;47:664-671.
- Pini R, Cavallini MC, Bencini F, Silvestrini G, Tonon E, Marchionni N, Di Bari M, Devereux RB, Masotti G, Roman MJ. Cardiac and vascular remodeling in older adults with isolated systolic hypertension: the ICARE Dicomano study. *Circulation* 2000;102(suppl II):II-698. Abstract.
- Di Bari M, Salti F, Nardi M, Pahor M, De Fusco C, Tonon E, Ungar A, Pini R, Masotti G, Marchionni N. Undertreatment of hypertension in community-dwelling older adults: a drug-utilization study in Dicomano, Italy. *J Hypertens* 1999;17:1633-1640.
- Devereux RB, Roman MJ. Evaluation of cardiac and vascular structures by echocardiography and other non-invasive techniques. In: Laragh JH, Brenner BM, eds. *Hypertension: Pathophysiology, Diagnosis, Treatment*. 2nd ed. New York, NY: Raven Press; 1995:196-198.
- Sahn D, DeMaria A, Kisslo J, Weyman A, for the Committee on M-mode Standardization of the American Society of Echocardiography. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* 1978;58:1072-1083.
- Schiller NB, Shah PM, Crawford M, DeMaria A, Devereux RB, Feigenbaum H, Gutgesell H, Reichek N, Sahn D, Schnittger I, Silverman NH, Tajik AJ. Recommendations for quantitation of the left ventricle by two-dimensional echocardiography: American Society of Echocardiography Committee on Standards, Subcommittee on Quantitation of Two-Dimensional Echocardiograms. *Am Soc Echocardiogr* 1989;2:358-367.
- Appleton CP, Jensen JL, Hatle LK, Oh JK. Doppler evaluation of left and right ventricular diastolic function: a technical guide for obtaining optimal flow velocity recordings. *J Am Soc Echocardiogr* 1997;10:271-292.
- Palmieri V, Dahlof B, DeQuattro V, Sharpe N, Bella JN, de Simone G, Paranicas M, Fishman D, Devereux RB. Reliability of echocardiographic assessment of left ventricular structure and function: the PRESERVE study. *J Am Coll Cardiol* 1999;34:1625-1632.
- Roman MJ, Pickering TG, Schwartz JE, Pini R, Devereux RB. Association of carotid atherosclerosis and left ventricular hypertrophy. *Am Coll Cardiol* 1995;25:8390.
- Roman MJ, Saba PS, Pini R, Spitzer M, Pickering TG, Rosen S, Alderman MH, Devereux RB. Parallel cardiac and vascular adaptation in hypertension. *Circulation* 1992;86:1909-1918.
- Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R. Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation* 1986;74:1399-1406.
- Lemne C, Jogestrand T, de Faire U. Carotid intima-media thickness and plaque in borderline hypertension. *Stroke* 1995;26:34-39.
- Roman MJ, Pickering TG, Pini R, Schwartz JE, Devereux RB. Prevalence and determinants of cardiac and vascular hypertrophy in hypertension. *Hypertension* 1995;26:369-373.
- Roman MJ, Pini R, Pickering TG, Devereux RB. Non-invasive measurements of arterial compliance in hypertensive compared with normotensive adults. *J Hypertens* 1992;10:S115-S118.
- Saba PS, Roman MJ, Pini R, Spitzer M, Ganau A, Devereux RB. Relation of arterial pressure waveform to left ventricular and carotid anatomy in normotensive subjects. *Am Coll Cardiol* 1993;22:1873-1880.
- Saba PS, Cavallini MC, Scorzoni D, Longhini C, Pini R, Ganau A. Arterial tonometry: principles and clinical applications in hypertension. *High Blood Press* 1996;5:244-250.
- Kawasaki T, Sasayama S, Yagi S, Asakawa T, Hirai T. Non-invasive assessment of the age related changes in stiffness of major branches of the human arteries. *Cardiovasc Res* 1987;21:678-687.
- Hirai T, Sasayama S, Kawasaki T, Yagi S. Stiffness of systemic arteries in patients with myocardial infarction. *Circulation* 1989;80:78-86.
- Kelly R, Daley J, Avolio A, O'Rourke M. Arterial dilation and reduced wave reflection. Benefit of diltiazem in hypertension. *Hypertension* 1989;14:14-21.
- Murgo JP, Westerhof N, Giolma JP, Altobelli SA. Aortic input impedance in normal man: relationship to pressure wave form. *Circulation* 1980;62:1051-116.
- Khattar RS, Acharya DU, Kinsey C, Senior R, Lahiri A. Longitudinal association of ambulatory pulse pressure with left ventricular mass and vascular hypertrophy in essential hypertension. *Hypertension* 1997;15:737-743.
- Domanski MJ, Davis BR, Pfeffer MA, Kastantin M, Mitchell GF. Isolated systolic hypertension: prognostic information provided by pulse pressure. *Hypertension* 1999;34:375-380.
- Staessen JA, Gasowski J, Wang JG, Thijs L, Den Hond E, Boissel JP, Coope J, Ekblom T, Gueyffier F, Liu L, Kerlikowske K, Pocock S, Fagard RH. Risks of untreated and treated isolated systolic hypertension in the elderly: meta-analysis of outcome trials. *Lancet* 2000;355:865-872.
- Blacher J, Staessen JA, Girerd X, Gasowski J, Thijs L, Liu L, Wang JG, Fagard RH, Safar ME. Pulse pressure not mean pressure determines cardiovascular risk in older hypertensive patients. *Arch Intern Med* 2000;160:1085-1089.
- Fagard RH, Staessen JA, Thijs L, Gasowski J, Bulpitt CJ, Clement D, de Leeuw PW, Dobovisek J, Jaaskivi M, Leonetti G, Ben E, Palatini P, Parati G, Rodicio JL, Vanhanen H, Webster J. Response to antihypertensive therapy in older patients with sustained and nonsustained systolic hypertension. *Circulation* 2000;102:1139-1144.
- The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 1997;157:2413-2446.
- Glynn RJ, Chae CU, Guralnik JM, Taylor JO, Hennekens CH. Pulse pressure and mortality in older people. *Arch Intern Med* 2000;160:2765-2772.
- Vaccarino V, Holford TR, Krumholz HM. Pulse pressure and risk for myocardial infarction and heart failure in the elderly. *Am Coll Cardiol* 2000;36:130-138.
- Cavallini MC, Roman MJ, Pickering TG, Schwartz JE, Pini R, Devereux RB. Is white coat hypertension associated with arterial disease or left ventricular hypertrophy? *Hypertension* 1995;26:413-419.