



FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Extracellular calcium negatively modulates tyrosine phosphorylation and tyrosine kinase activity during capacitation of human

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Extracellular calcium negatively modulates tyrosine phosphorylation and tyrosine kinase activity during capacitation of human spermatozoa / M. Luconi; C. Krausz; G. Forti; E. Baldi. - In: BIOLOGY OF REPRODUCTION. - ISSN 0006-3363. - STAMPA. - 55:(1996), pp. 207-216. [10.1095/biolreprod55.1.207]

Availability:

This version is available at: 2158/336110 since: 2017-05-21T19:52:20Z

Published version: DOI: 10.1095/biolreprod55.1.207

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf)

Publisher copyright claim:

(Article begins on next page)

Extracellular Calcium Negatively Modulates Tyrosine Phosphorylation and Tyrosine Kinase Activity during Capacitation of Human Spermatozoa¹

Michaela Luconi, Csilla Krausz, Gianni Forti, and Elisabetta Baldi²

Dipartimento di Fisiopatologia Clinica, Unita' di Andrologia, Universita' di Firenze, 50139 Florence, Italy

ABSTRACT

Capacitation of spermatozoa, a complex process occurring after sperm ejaculation, is required to obtain fertilization of the oocyte in vivo and in vitro. Although most of the biochemical/ biophysical events that occur during capacitation in vitro have been characterized, the molecular mechanisms underlying these complex events are still obscure. Increases of intracellular free Ca²⁺ concentrations ([Ca²⁺],) and protein tyrosine phosphorylation have previously been demonstrated during in vitro capacitation of human spermatozoa. In the present study we investigated the relationship between extracellular/intracellular Ca2+, protein tyrosine phosphorylation, and tyrosine kinase and phosphatase activities during sperm capacitation. We report that the increase in tyrosine phosphorylation of several protein bands that occurs during sperm capacitation is independent of the presence of Ca2+ in the external medium and, at least partially, of the increase in [Ca2+], occurring during the process. Indeed, the spontaneous increase in phosphorylation was still present in Ca2+-free/EGTA-containing-medium and in the presence of the intracellular Ca2+ chelator BAPTA/AM. Moreover, phosphorylation of proteins and protein tyrosine kinase (PTK) activity was enhanced if spermatozoa were incubated in Ca2+-free medium, suggesting the presence of Ca2+-inhibited tyrosine kinase(s) in human sperm. This hypothesis is further substantiated by the lower tyrosine phosphorylation observed after incubation with the ionophore A23187 and the endoplasmic Ca2+-ATPase inhibitor thapsigargin, which promote Ca2+ influx in human sperm. The ability of the cells to undergo acrosome reaction in response to progesterone, which can be considered a functional endpoint of capacitation, was highly compromised when spermatozoa were incubated in Ca2+-free medium or in the presence of EGTA, confirming that Ca2+ is required for sperm capacitation. Conversely, in the presence of erbstatin, a inhibitor of tyrosine kinase activity, which blunts tyrosine phosphorylation during capacitation, response to progesterone was maintained, suggesting that tyrosine phosphorylation must be kept at a low level (physiologically by the presence of Ca2+ in the external medium, or pharmacologically by the presence of erbstatin) in order to obtain response to progesterone. This mechanism may be important in vivo during sperm transit in the female genital tract to ensure appropriate timing of full capacitation in the proximity of the oocyte.

INTRODUCTION

The process of capacitation is an event that is essential in order for mammalian spermatozoa to acquire their ability to fertilize the oocyte (for review see [1]). During this process, spermatozoa become actively motile (expressing a distinct motility pattern called hyperactivation) and responsive to physiological stimuli of the acrosome reaction, such as zona pellucida proteins and progesterone [1]. Although physiological capacitation occurs in the female genital tract, the process can be reproduced in vitro by incubation in defined media (whose composition is similar to oviductal fluid) and thus appropriately studied [1]. To date, it is known that capacitation is characterized by a series of complex biochemical/biophysical events involving regional modifications of sperm membrane sterols and phospholipids, changes in ionic movements, and other events that for the most part have been well documented [1]. However, the precise molecular mechanisms underlying these complex events are still far from being completely clear. In addition, the lack of a defined assay that can identify capacitated spermatozoa (for review see [2]) limits the interpretation of in vitro studies. Increasing evidence indicates that Ca²⁺ ions are of fundamental importance in this process: an increase of intracellular free Ca^{2+} concentrations ([Ca^{2+}]_i) during capacitation has been indeed demonstrated in spermatozoa from several mammalian species [3-6]. Such increase is accompanied by an enhancement of sperm responsiveness to physiological stimuli of the acrosome reaction such as progesterone [6–9] and zona pellucida proteins [10], which have been shown to increase $[Ca^{2+}]_i$ [6, 11–13] in capacitated spermatozoa only. In addition to an increase in $[Ca^{2+}]_{i}$, capacitation is also characterized by an increase in phosphorylation of sperm proteins, particularly in tyrosine residues [14-17], indicating activation of sperm tyrosine kinases during this process. Recent evidence suggests a role for tyrosine kinase(s) in the process of fertilization [17, 18]. Moreover, a role of tyrosine kinase(s) in the initiation of flagellar movement has been demonstrated in the rainbow trout sperm [19]. These studies indicate that these enzymes, in addition to their well-characterized role in cell proliferation and differentiation [20], may also play an important role in the biology of highly specialized, mature cells such as spermatozoa.

However, little is known about the signals that promote activation of tyrosine kinase during capacitation in human sperm, or about the role, if any, of tyrosine kinase activation for capacitation of spermatozoa. In a recent paper, Aitken et al. [21] reported that the ionophore A23187 enhances tyrosine phosphorylation of human sperm during capacitation, suggesting a positive correlation between the two phenomena.

The increases in $[Ca^{2+}]_i$ and protein tyrosine phosphorylation during in vitro capacitation of human spermatozoa show a similar time course [6, 16, 22], suggesting the possible existence of a close relationship between the two events.

In the present study we investigated whether the increase in tyrosine phosphorylation is dependent on the increase in $[Ca^{2+}]_i$ by using an experimental approach involving Western blot analysis of tyrosine phosphorylation and assays of sperm tyrosine kinase and phosphatase activities in sper-

Accepted March 12, 1996.

Received November 8, 1995.

¹This work was supported by grants from Consiglio Nazionale delle Ricerche (CNR, Rome, targeted project FATMA, contract n. 94.00548-PF41), Ministero dell'Universita' e della Ricerca Scientifica (Rome) and Regione Toscana (Florence).

²Correspondence: Elisabetta Baldi, Ph.D., Dipartimento di Fisiopatologia Clinica, Unita' di Andrologia, Universita' di Firenze, viale Pieraccini 6, I-50139 Florence, Italy. FAX: 39 55 4221 848.

matozoa incubated in the absence of extracellular Ca^{2+} and/ or in the presence of various Ca^{2+} chelators. In addition, in order to verify the occurrence of capacitation in the different conditions, we evaluated progesterone-stimulated acrosome reaction. We report that 1) extracellular Ca^{2+} is not required for the increase in tyrosine phosphorylation during capacitation, 2) extracellular calcium negatively modulates tyrosine kinase activity in human sperm, and 3) progesterone can stimulate the acrosome reaction of human sperm when capacitation is obtained in the presence of an inhibitor of tyrosine kinase activity.

MATERIALS AND METHODS

Chemicals

Sodium orthovanadate (Na₃VO₄), progesterone, fatty acid-free BSA, gelatine, glycine, trizma base, Temed, ammoniumpersulfate, Triton X-100, PMSF, soybean trypsin inhibitor, O-phospho-DL-tyrosine, Ponceau S and Coomassie R250, EGTA, and EDTA were from Sigma Chemical Company (St. Louis, MO). Thapsigargin, erbstatin, genistein, tyrphostin A47, 1,2-bis(o-aminophenoxy)-ethane-N, N, N', N'-tetra-(acetoxymethyl)-ester (BAPTA/AM), ionomycin, fura-2/AM, A23187, and fluorescein isothiocyanate (FITC)-labeled Arachis hypogea (peanut) lectin were from Calbiochem (La Jolla, CA). Unconjugated and peroxidaseconjugated monoclonal (PY20) antiphosphotyrosine antibodies were from ICN (Costa Mesa, CA). Reagents for SDS-PAGE were from Bio-Rad Laboratories (Hercules, CA). Protein molecular weight standards, Tween 20, and Nonidet P-40 (NP-40) were from FLUKA Chemie AG (Bucks, Switzerland). Percoll was obtained from Pharmacia LKB (Uppsala, Sweden). Enhanced chemiluminescence (ECL) Western blotting detection reagents were from Amersham (Buckinghamshire, UK). Medi-Cult (Copenhagen, Denmark) was the supplier of human serum albumin (HSA)-free sperm preparation medium (229.5 mg/L CaCl₂·2H₂O, 346.55 mg/L KCl, 173.27 mg/L MgSO4.7H2O, 5891.28 mg/L NaCl, 2200 mg/L NaHCO3, 137.15 mg/L NaH2PO4.2H2O, 866.36 mg/L D-glucose, 8.66 mg/L phenol red, 97.0 mg/L Na pyruvate, 1000 mg/L Hepes sodium salt, 3010 mg/L Hepes acid, 1.0 ml/L SSR2 (comp. A), 1.0 ml/L SSR2 (comp. B), 50 mg/L streptomycin, 50 000 IU/L penicillin), and Ca²⁺-free sperm preparation medium.

Preparation and Incubation of Spermatozoa

Human semen was obtained from normospermic men undergoing semen analysis for couple infertility in our lab. Semen was collected according to the procedure recommended by WHO [23] via masturbation after 3-4 days of sexual abstinence. Samples with a linear progressive motility of less than 50% at 60 min and with leukocytes and/ or an immature germ cell concentration greater than 10⁶/ ml were not included in the study. Semen samples were processed as previously described [6, 24] with the exception that sperm preparation medium (SPM) was used instead of Biggers, Whitten and Whittingham (BWW) medium [6]. Briefly, spermatozoa were separated on 40% and 80% Percoll gradients, combined, washed in BSA-free SPM, and resuspended in a small volume of the same medium. All these initial steps were performed in complete absence of BSA in order to avoid initial capacitation of the samples. Indeed, we have previously demonstrated that spermatozoa prepared in these conditions are not able to respond to progesterone, indicating absence of capacitation, and that they can acquire this ability after reincubation in BSA-containing medium [6]. Spermatozoa were counted and resuspended in BSA-free SPM (noncapacitated samples) and in SPM with added BSA (3 mg/ml; capacitating samples) at the concentration of 3×10^6 /ml in the different conditions tested. Samples were incubated at 37° C in a CO₂ atmosphere, and aliquots were collected at different incubation times and processed as described below. For experiments in which the effect of progesterone was to be evaluated, spermatozoa were capacitated for 2 h in complete SPM medium and then stimulated with progesterone (4 µg/ ml, final concentration) for 5 min.

Evaluation of Sperm Motility and Acrosome Reaction

Sperm motility was evaluated with use of a fully automated computer-assisted semen analyzer (Hamilton-Thorne motion analyzer: Hamilton-Thorne, Danvers, MA) as previously described [25]. Sperm acrosome reaction was evaluated through use of the fluorescent probe FITC-labeled Arachis hypogea (peanut) lectin as previously described [25, 26]. Briefly, spermatozoa (10×10^{6} /ml), after capacitation, were stimulated for 1 h with progesterone (4 μ g/ml), centrifuged at 400 \times g for 10 min, incubated in 0.5 ml of hyposmotic swelling medium for 1 h at 37°C, centrifuged, and finally resuspended in 50 µl of ice-cold methanol. The sperm suspension was layered on a slide, air-dried at room temperature, and stored at -20° C. Fluorescence was observed under a fluorescent microscope (Leitz, type 307-148002; Wetzlar, Germany), and the acrosome reaction was evaluated on a total of 100 spermatozoa per slide. As outlined by Aitken et al. [27], only curly-tailed sperm were considered viable and thus scored.

SDS-PAGE and Western Blot Analysis

After the different incubations, the samples were processed for SDS-electrophoresis as previously described [16]. Briefly, sperm samples, containing 3×10^6 cells/ml, were centrifuged at 400 \times g at 4°C for 10 min, washed in HSA-free SPM, and resuspended in 10 µl lysis buffer (20 mM Tris [pH 7.4], 150 mM NaCl, 0.25% NP-40, 1 mM Na_3VO_4 , 1 mM PMSF). After measurement of proteins, the sperm extracts, containing approximately 20 µg of proteins, were diluted in equal volume of double-strength concentrated Laemmli sample buffer (double-strength sample buffer = 62.5 mM Tris [pH 6.8] containing 10% glycerol, 4% SDS, 2.5% pyronin, and 200 mM dithiothreitol), vor-texed, incubated at 95°C for 5 min, and then loaded onto 7.5% polyacrylamide/bisacrylamide gels. After SDS-PAGE, proteins were transferred to nitrocellulose (Sigma) and stained with Ponceau to verify equal protein loading. In some experiments, equivalent protein loading was verified by staining parallel gels with Coomassie. If protein loading was not equivalent in each lane, sperm proteins were newly run. The nitrocellulose was immunostained with peroxidase-conjugated monoclonal antiphosphotyrosine antibody (PY20), and the phosphotyrosine antibodyreacted proteins were revealed by the ECL system (Amersham). To obtain immunoabsorption, the antibody was reacted with 40 mM O-phospho-DL-tyrosine for 1 h at 4°C. Quantification of the bands was made directly on the films by image analysis with a C3077/01 video camera connected to the video frame grabber M4477 (Hamamatsu Photonics, Hamamatsu City, Japan). This video frame grabber is a plug-in board used in a Macintosh Ilsi PC (Apple, Carpinteria, CA). Acquisition of image was performed by Image Quest IQ Base software (Hamamatsu Photonics). Image processing and analysis were obtained with IMAGE free software, kindly provided by Wayne Rasband (NIMH, Bethesda, MD). For technical reasons, quantifications were performed in ECL low-exposed films.

Evaluation of Tyrosine Phosphatase and Kinase Activities

Tyrosine phosphatase and kinase activities were evaluated with an ELISA (tyrosine phosphatase assay kit, tyrosine kinase assay kit; Boehringer Mannheim GmbH, Mannheim, Germany). After 1 h of incubation in complete medium or in Ca²⁺-free, or A23187 (10 µM)-containing medium, sperm extracts were washed and resuspended in modified TN buffer [18, 28] (150 mM NaCl, 20 mM Tris [pH 7.2], 0.5% NP-40, 1 mM PMSF, and 1 mM Na₃VO₄ for protein tyrosine kinase [PTK] or 50 µg/ml erbstatin for protein tyrosine phosphatase [PTP] assay) and then vortexed for 2 min to obtain membrane preparations [18, 28]. Since both methods may be subjected to interferences due to tyrosine kinase activity (for PTP) and phosphatase activity (for PTK) present in the extracts, the assays were conducted in the presence of 50 μ g/ml of the tyrosine kinase inhibitor erbstatin (for PTP) or 1 mM Na₃VO₄ (for PTK). To evaluate PTP activity, the enzyme reaction was performed in duplicate at 30°C for 15 min in the presence of 1 nM tyrosine phosphatase substrate (corresponding to amino acids 1-17 of human gastrin phosphorylated on tyrosine 12). The reaction was stopped with the tyrosine phosphatase inhibitor Na₃VO₄ (100 μ M); the unmetabolized substrate, immobilized on microtiter plates, was determined immunochemically with ABTS (2-2'-azino-di-[3-ethylbenthiazoline sulfonate]) via antiphosphotyrosine antibody, directly conjugated to peroxidase. The rate of dephosphorylation was determined with a phosphopeptide standard curve. For the tyrosine kinase assay, the enzyme reaction was conducted in duplicate at 30°C for 30 min in an assay buffer (20 mM Tris-HCl [pH 7.2], 0.1% 2-mercaptoethanol, 1 mM ATP, 10 mM MgCl₂, 1 mg/ml BSA, and 1 nM enzyme substrate [corresponding to the amino acid sequence 1–17 of gastrin biotinylated at the N-terminal glutamate]). The reaction was stopped with 0.5 mM of the tyrosine kinase inhibitor piceatannol (provided in the kit from Boehringer Mannheim GmbH); then tyrosine phosphorylated and unphosphorylated substrates were immobilized on microtiter plates, and tyrosine phosphorylated substrate was determined as described above for the PTP assay. The rate of phosphorylation was quantitated with a phosphopeptide standard curve.

Measurement of [Ca²⁺], Concentration

 $[Ca^{2+}]_i$ was measured in fura-2-loaded spermatozoa after 2-h incubation in complete and Ca²⁺-free medium by a fluorimetric method. Spermatozoa were loaded with fura-2/ AM as previously described [6, 26], and fluorescence was measured with use of a single-wavelength spectrofluorometer (University of Pennsylvania Biomedical Group, Philadelphia, PA) set at 340 nm excitation with emission at 510 nm. Basal $[Ca^{2+}]_i$ was calculated according to Grynkiewicz et al. [29], using ionomycin (60 μ M) for evaluation of maximal fluorescence (F_{max}) followed by EGTA (7.5 mM) for evaluation of minimal fluorescence (F_{min}), and assuming a K_d of fura-2 for Ca²⁺ of 224 nM [29]. In spermatozoa incubated in Ca²⁺-free medium, F_{min} was evaluated by adding ionomycin directly to the sample or after incubation with

Measurement of Proteins

Protein concentrations were evaluated by the Bio-Rad protein assay reagent exactly as indicated by the manufacturer, with BSA used as standard.

Statistical Analysis

Data are expressed as mean \pm SEM. Statistical comparisons were made with Student's *t*-test for paired data.

RESULTS

Role of Extracellular Calcium in Time-Dependent Changes in Tyrosine Phosphorylation of Sperm Proteins during Capacitation

As expected, after incubation of human spermatozoa for 1 and 2 h in complete, Ca²⁺-containing, capacitating medium, tyrosine phosphorylation of several sperm protein bands of molecular mass ranging between 40 and 100 kDa was increased at 2 h as compared to 1 h (compare lane 1 with lane 5 in Figs. 1 and 4). In most of the experiments, the higher degree of tyrosine phosphorylation was observed in two protein bands with apparent molecular masses of 75 and 97 kDa. Sperm proteins were specifically phosphorylated on tyrosine residues, since preimmunoabsorption of the PY20 antibody with O-phospho-DL-tyrosine completely abolished immunoreactivity (data not shown). These data confirm previous results obtained by our group showing progressive increase of tyrosine phosphorylation of sperm proteins at different times of capacitation [16, 22]. When spermatozoa were incubated in a medium devoid of Ca²⁺ (nominally Ca²⁺-free medium) for 1 and 2 h, tyrosine phosphorylation of sperm proteins was enhanced as compared to that seen in complete medium at both incubation times (compare lane 2 with lane 1, and lane 6 with lane 5, in Figs. 1 and 4; compare also lane 2 with lane 1 in Fig. 3B); and the spontaneous increase of phosphorylation at 2 h vs. 1 h was still present (compare lane 6 with lane 2 in Figs. 1 and 4). The average percentage increases of tyrosine phosphorylation in Ca^{2+} -free vs. complete medium for the 97- and the 75-kDa bands were, respectively, 50.71 ± 3.4 (n = 7) and 151.3 ± 64.8 (n = 7) at 1-h incubation and 51 ± 10.1 (n = 8) and 205 \pm 74.6 (n = 5) at 2-h incubation. In contrast, the presence of EDTA (3 mM) in the incubation medium abolished tyrosine phosphorylation of proteins at both the 1- and 2-h time points (compare lane 3 with lane 2, and lane 7 with lane 6, in Fig. 1). Since EDTA chelates both Mg^{2+} and Ca^{2+} , we tested the effect of EGTA, which predominantly chelates Ca²⁺. As shown in Figures 1 and 2, EGTA (3 mM) only slightly affected tyrosine phosphorylation of sperm proteins (compare lane 4 with lane 2 in Fig. 1; lane 8 with lane 6 in Fig. 2). The concentration of free Mg2+ in the EDTA- and EGTA-containing media, measured according to Fabiato [30], was, respectively, 230 µM and 370 µM, while the concentration of Ca^{2+} (assumed to be 10 μM due to contamination) was found to be less than 10^{-8} M with both chelating agents; thus the decrease in tyrosine phosphorylation observed in the presence of EDTA was likely to be due to chelation of Mg^{2+} . This result is in agreement with findings of Hayashi et al. [19], who showed that cAMP-dependent tyrosine phosphorylation of a rainbow trout sperm protein of 15 kDa was completely inhibited in EDTA-containing medium but



FIG. 1. Western blot analysis of the reactivity of antiphosphotyrosine antibody with human sperm proteins at 1 and 2 h of in vitro incubation in the presence or absence of added Ca²⁺, EDTA (3 mM), and EGTA (3 mM). Spermatozoa were incubated for the indicated times at 37°C in an atmosphere of CO₂, collected, and run on 7.5% acrylamide/bisacrylamide gel; tyrosine phosphorylation was evaluated by Western blot analysis using a peroxidase-conjugated antiphosphotyrosine antibody (PY20) followed by the ECL detection system. Densitometric analysis (arbitrary units) for the 97 (D-97)- and 75 (D-75)-kDa tyrosine phosphorylated bands is reported below each lane. Molecular weight markers (×10³) are indicated to the right of the blot; Rf = running front. Representative of three similar experiments.

was only slightly affected by EGTA, indicating the role of Mg²⁺ in PTK activation in rainbow trout sperm. Furthermore, we found that the spontaneous increase in tyrosine phosphorylation observed during capacitation in complete medium was still present when EGTA was included in the Ca^{2+} -free medium (compare lane 8 with lane 4 in Fig. 1); there was an increase in relation to that seen at 1 h that was similar to the increase found in complete medium (compare densitometric analysis for lanes 8 vs. 4 and lanes 5 vs. 1 in Fig. 1). To evaluate whether incubation in Ca^{2+} free medium was consistent with a decrease in $[Ca^{2+}]_{i}$, spermatozoa were incubated for 2 h in capacitating conditions (complete medium) or in Ca^{2+} -free medium, and basal $[Ca^{2+}]_i$ was then evaluated by a spectrofluorometric method. We found that when incubation was conducted in Ca²⁺free medium, basal $[Ca^{2+}]_i$ was reduced by 44.7 $\pm~2.3\%$ (n = 3) from the value obtained after incubation in complete medium. When $[Ca^{2+}]_i$ measurement was conducted in the presence of 0.5 mM EGTA to chelate residual calcium present in the medium, basal $[Ca^{2+}]_i$ was reduced by $81.4 \pm 1.8\%$ (n = 3).

To further investigate the relationship between extracellular Ca²⁺ and sperm tyrosine phosphorylation during capacitation, spermatozoa were incubated for 2 h with various concentrations of added Ca²⁺ (0.03, 0.15, 0.3, 1.5 [Ca²⁺ concentration in SPM], and 3 mM). As shown in Figure 2, incubation of sperm with increasing concentrations of Ca²⁺ progressively blunted sperm tyrosine phosphorylation of proteins in a concentration-dependent manner, suggesting that Ca²⁺ ions negatively modulate tyrosine phosphorylation of sperm proteins during capacitation. To explore this possibility further, we evaluated the effect of two known stimulators of Ca²⁺ influx and acrosome reaction, the Ca²⁺ ionophore A23187 and the endoplasmic Ca²⁺-ATPase in-



FIG. 2. Effect of different Ca²⁺ concentrations in the capacitation medium on sperm tyrosine phosphorylation. Spermatozoa were incubated at 37°C for 2 h without Ca²⁺ in the presence or absence of EGTA (3 mM), or with increasing Ca²⁺ concentrations; they were then collected and processed as described for Figure 1 for detection of tyrosine phosphorylation of proteins. Molecular weight markers (×10³) are indicated to the right of the blot; Rf = running front. Representative of two similar experiments.

hibitor thapsigargin [31, 32]. As shown in Figure 3A, the presence, during the 2-h incubation, of the Ca²⁺ ionophore A23187 (10 μ M) in complete medium inhibited tyrosine phosphorylation of 97- and 75-kDa proteins (compare lane 3 with lane 1 in Fig. 3A) as well as of other proteins (not shown). The average percentage decrease in tyrosine phosphorylation after A23187 incubation was $47 \pm 4.9\%$ for the 97- and 76.6 \pm 9.2% for the 75-kDa (n = 3) protein band. In this respect, it must be mentioned that in a recent paper, Aitken et al. [21] showed that incubation with the ionophore A23187 leads to an enhancement of tyrosine phosphorylation of proteins in human sperm. However, the pattern of tyrosine phosphorylated proteins shown in that study [21] differs from that reported by several other groups ([15–17]; present paper), thus pointing to the possibility that methodological differences might be responsible for the conflicting results. In addition, a preliminary report by Moos et al. [33] indicates that the treatment of human sperm with A23187 decreases sperm phosphorylation. Results similar to those obtained with A23187 were observed with thapsigargin (10 μ M), which inhibited tyrosine phosphorylation of the 97- and 75-kDa protein bands and of other bands (not shown) in complete medium (compare lane 3 with lane 1 in Fig. 3B) but was less effective in Ca^{2+} -free medium (compare lane 4 with lane 2 in Fig. 3B). Since thapsigargin has been shown to be effective in increasing sperm $[Ca^{2+}]_i$ only when Ca^{2+} is present in the extracellular medium [31], the latter result is not surprising. We also evaluated the effect of A23187 in spermatozoa incubated in the presence of the tyrosine phosphatase inhibitor Na₃VO₄ (1 mM) in order to examine whether tyrosine phosphatases may contribute to the inhibitory effect of the ionophore. The inhibitory effect of A23187 was still evident when Na₃VO₄ was present in the medium (compare lane 4 with lane 2 in Fig. 3A). Although Na_3VO_4 is not a specific inhibitor of tyrosine phosphatases, this result suggests that PTP activity does not contribute to the inhibition of tyrosine phosphorylation induced by A23187. Indeed, if the inhibitory effect of A23187 on tyrosine phosphorylation was due to activation of tyrosine phosphatases, it should not have persisted in the presence of Na₃VO₄.

Effect of the Intracellular Ca²⁺ Chelator BAPTA on Sperm Tyrosine Phosphorylation

To further investigate whether Ca^{2+} increase during capacitation was involved in the enhancement of tyrosine



FIG. 3. **A**) Effect of the Ca²⁺ ionophore A23187 on tyrosine phosphorylation of sperm proteins. Spermatozoa were capacitated for 2 h at 37°C in an atmosphere of CO₂ in Ca²⁺-containing medium with or without A23187 (10 μ M). The effect of A23187 was also tested in the presence of the tyrosine phosphatase inhibitor Na₃VO₄ (1 mM). Representative of three similar experiments. **B**) Effect of the endoplasmic Ca²⁺-ATPase inhibitor thapsigargin (T; 10 μ M) on tyrosine phosphorylation of 97-kDa and 75-kDa protein bands of spermatozoa. Spermatozoa were allowed to capacitate for 2 h at 37°C in an atmosphere of CO₂ with or without added Ca²⁺ and in the presence or absence of thapsigargin for 1 h, and tyrosine phosphorylation was evaluated. Representative of two similar experiments. Tyrosine phosphorylated proteins were revealed as described for Figure 1. Molecular weight markers (×10³) are indicated to the right (**A**) or to the left (**B**) of the blot.

phosphorylation, spermatozoa were incubated with the intracellular Ca²⁺ chelator BAPTA/AM (10 µM) in either the presence or absence of Ca2+ in the external medium, and protein tyrosine phosphorylation was evaluated at 1 and 2 h. In preliminary experiments, loading of spermatozoa with 10 µM BAPTA/AM was found to completely abolish the [Ca²⁺], increase in response to progesterone in fura 2-loaded sperm (results not shown), indicating that BAPTA effectively chelates intracellular Ca²⁺ in these conditions. As shown in Figure 4, when spermatozoa were incubated in the presence of BAPTA, sperm tyrosine phosphorylation was reduced both at 1- and 2-h incubation either in the presence (compare lane 3 with lane 1 for 1-h incubation; lane 7 with lane 5 for 2-h incubation) or in the absence (compare lane 4 with lane 2 for 1-h incubation; lane 8 with lane 6 for 2-h incubation) of Ca^{2+} in the external medium; this result, consistently observed, might suggest a partial Ca²⁺-dependence of sperm tyrosine kinase activity during capacitation. Alternatively, the result might be explained,



FIG. 4. Effect of the intracellular Ca²⁺ chelator BAPTA (10 μ M) on sperm protein tyrosine phosphorylation at 1- and 2-h in vitro incubation in the presence or absence of added Ca²⁺ in the medium. Spermatozoa were incubated in the different conditions for the indicated times at 37°C in an atmosphere of CO₂, collected, and run on 7.5% acrylamide/bisacrylamide gel; tyrosine phosphorylation was evaluated by Western blot analysis as described for Figure 1. Densitometric analysis (arbitrary units) for the 97 (D-97)- and 75 (D-75)-kDa tyrosine phosphorylated bands is reported below each lane. Molecular weight markers (×10³) are indicated to the right of the blots; Rf = running front. Representative of two similar experiments.

as in the case of EGTA incubation (Fig. 1), by a possible chelating effect of BAPTA on intracellular Mg^{2+} . Although the affinity of BAPTA for Mg^{2+} is much lower than that for Ca²⁺, the possibility of a chelating effect of BAPTA on intracellular $\hat{M}g^{2+}$ cannot be excluded. On the other hand, in the presence of BAPTA, the spontaneous increase in tyrosine phosphorylation at 2-h incubation (compared to 1 h) was still present both in complete (compare lane 7 with lane 3 in Fig. 4) and in Ca^{2+} -free (compare lane 8 with lane 4 in Fig. 4) medium. Moreover, densitometric analysis of the 97- and 75-kDa phosphoprotein bands indicates that the relative increase in tyrosine phosphorylation of this protein at 2-h incubation in the presence of BAPTA in Ca²⁺free medium was similar to, if not higher than, that observed in complete medium (compare densitometric analysis in lane 8 vs. 4 for BAPTA and lane 5 vs. 1 for complete medium in Fig. 4).

The results presented so far indicate that Ca²⁺ ions negatively modulate tyrosine phosphorylation in human sperm. It has been shown that an influx of Ca²⁺ decreases PTP activity [34]. We therefore investigated the possible role of tyrosine phosphatase(s) activity in the tyrosine phosphorylation increase during capacitation. Spermatozoa were incubated for 2 h with the tyrosine phosphatase inhibitor Na₃VO₄ (1 mM) in the presence or absence of Ca²⁺. As shown in Figures 3A and 5, Na₃VO₄ increased tyrosine phosphorylation of the 97- and 75-kDa as well as (not shown) other protein bands, in complete medium (compare lane 3 with lane 1 in Fig. 5; lane 2 and lane 1 in Fig. 3A), whereas no apparent effect was observed in Ca2+-free medium (compare lane 4 with lane 2 in Fig. 5). This result suggests that the increase of tyrosine phosphorylation observed in Ca²⁺-free medium is not mediated by inhibition of tyrosine phosphatase(s).

To investigate further the relative contributions of sperm tyrosine phosphatase and tyrosine kinase activities in the



FIG. 5. Effect of incubation for 2 h with the tyrosine phosphatase inhibitor Na₃VO₄ (1 mM) on tyrosine phosphorylation of 97-kDa and 75-kDa protein bands in human sperm. Spermatozoa were incubated in complete medium with or without added Ca²⁺ in the external medium at 37°C in an atmosphere of CO₂ in the presence or absence of Na₃VO₄. Tyrosine phosphorylation of the two proteins at 2-h capacitation with or without added Ca²⁺ is shown in lanes 1 and 2, respectively. Molecular weight markers (×10³) are indicated to the right of the blots; Rf = running front. Representative of three similar experiments.

increase in tyrosine phosphorylation observed in the absence of extracellular Ca²⁺, we evaluated these activities in spermatozoa incubated for 1 h in complete medium, in the absence of Ca²⁺, and in the presence of A23187. The activity of the two enzymes was evaluated by measuring the rate of phosphorylation (for kinase) and dephosphorylation (for phosphatase) of peptide substrates by a photometric assay. Tyrosine phosphatase activity at 1-h incubation in complete medium was 2.24 ± 0.04 pmol/µg protein/min (n = 3) and was not modified in the absence of Ca^{2+} in the external medium or in the presence of A23187 (10 μ M) (Table 1), suggesting that these enzymes are not involved in the increase in tyrosine phosphorylation observed in Ca²⁺-free medium. Tyrosine phosphatase activity was inhibited, in all the conditions tested, by the concomitant presence of the tyrosine phosphatase inhibitor Na_3VO_4 (not shown). Sperm tyrosine kinase activity after 1-h incubation in complete medium was 85 ± 10.7 fmol/µg protein/min (n = 4). When spermatozoa were incubated in a medium devoid of Ca²⁺, the rate of substrate phosphorylation was significantly increased (Table 1), thus suggesting higher tyrosine kinase activity in this condition that may be responsible for the higher tyrosine phosphorylation observed. However, the results obtained with Å23187 (10 μ M) were conflicting, since in 2 of 4 experiments the tyrosine kinase activity was reduced by about 30%, while a slight (10-20%) increase was observed in the other 2 experiments, with an average percentage inhibition of about 20% that was not significant (Table 1). At present we have no explanation for these conflicting results; they may be attributable, however, to variability among the different subjects studied.

Effect of Different Incubation Conditions on Motility and on Basal and Stimulated Acrosome Reaction of Spermatozoa

The effect of the different incubation conditions on motility parameters (and, for comparison, on protein tyrosine phosphorylation) of human spermatozoa are reported in Table 2. Total and progressive motilities were not significantly affected by incubation in Ca^{2+} -free medium both at 1- and 2-h incubation, while tyrosine phosphorylation was increased. Incubation in BAPTA-containing Ca^{2+} -free medi-

TABLE 1. Effect of incubation in Ca^{2+} -free medium and in presence of A23187 on PTP and PTK activity during capacitation of human spermatozoa.^a

| Activity ^b | n | Ca²+-free medium | A23187 | | |
|-----------------------|---|------------------|----------|--|--|
| РТР | 3 | 109 ± 6 | 74 ± 9.2 | | |
| РТК | 4 | $154 \pm 14^{*}$ | 81 ± 29 | | |

^a Spermatozoa were capacitated for 1 hr in complete medium without added Ca²⁺ and in complete medium in the presence of A23187 (10 μ M). ^b Total enzyme activities, measured with use of a photometric method as described in the text, are expressed as percentage of control (capacitation in complete medium = 100%). Values represent means ± SEM of the indicated number of experiments.

* p < 0.05 vs. values in complete medium.

um (which slightly decreased tyrosine phosphorylation over that in Ca^{2+} -free medium alone) did not affect motility after 1 h and slightly decreased it after 2 h. In contrast, treatment with Na₃VO₄ reduced all motility parameters while increasing tyrosine phosphorylation. EGTA inhibited motility and tyrosine phosphorylation, whereas the addition of EDTA (not shown in the table) totally inhibited motility and tyrosine phosphorylation. Incubation with erbstatin, which dose-dependently blunted tyrosine phosphorylation during capacitation in both the presence [16] and absence (Fig. 6) of Ca^{2+} , had a variable effect on sperm motility at 1 h and inhibited it at 2 h. Sperm viability, evaluated by staining with eosin [23], was not affected by incubation with these agents, with the exception of EDTA, which reduced viability by about 20–30%.

We next evaluated spontaneous and progesterone-stimulated acrosome reaction in spermatozoa incubated in the different conditions used to detect tyrosine phosphorylation. As shown in Table 3, spontaneous acrosome reaction was not modified in spermatozoa capacitated in Ca²⁺-free, BAPTA-, EGTA-, and Na₃VO₄-containing media. In contrast, incubation with erbstatin (25 μ g/ml) consistently induced an increase in spontaneous acrosome reaction (Table 3). Similar results were also obtained with two different tyrosine kinase inhibitors, genistein and tyrphostin A47 (results not shown), which have been shown to decrease sperm tyrosine phosphorylation [18, 35]. This finding is difficult to explain, since erbstatin does not affect sperm viability and had a slight inhibitory effect on sperm motility (Table

TABLE 2. Effect of incubation in different conditions on total and progressive motility of human spermatozoa.*

| | To mot Capac tir | tal ility: itation ne | Progr mot Capac tir | Tyrosine phos- phory- lation: Capacita- | |
|---|--|--|---|--|--|
| Treatment | 1 h | 2 h | 1 h | 2 h | tion time 1 h/2 h |
| Ca ²⁺ -free medium NA ₃ VO ₄ 1 mM BAPTA 10 μM EGTA 3 mM Erbstatin 25 μg/ml | $\begin{array}{c} \leftrightarrow \\ \downarrow \\ \leftrightarrow \\ \downarrow \\ \leftrightarrow \\ \downarrow \end{array}$ | $\begin{array}{c} \leftrightarrow \\ \downarrow \\ \downarrow \downarrow \\ \downarrow \downarrow \\ \downarrow \end{array}$ | $\begin{array}{c} \leftrightarrow \\ \downarrow \\ \leftrightarrow \\ \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \end{array}$ | $\begin{array}{c} \leftrightarrow \\ \leftrightarrow \\ \rightarrow \\$ | $\uparrow\uparrow\\\downarrow\\\downarrow\downarrow\\\downarrow\downarrow$ |

* Spermatozoa were incubated for 1 or 2 h in the indicated conditions, and motility was evaluated by using a fully automated motion analyzer. Arrows indicate changes in sperm motility and tyrosine phosphorylation of proteins in comparison with values obtained after incubation in complete medium or, in the case of BAPTA and EGTA, in Ca²⁺-free medium. Results shown represent the average effect of motility in at least 5 different experiments for each condition. \leftrightarrow , No change; \uparrow , increase; \downarrow , decrease; $\leftrightarrow \downarrow$, variable results.



FIG. 6. Effect of erbstatin (E) on the levels of tyrosine phosphorylation of sperm proteins at 2-h incubation in the absence of added Ca²⁺. Spermatozoa were incubated at 37°C in an atmosphere of CO₂ with or without Ca²⁺ in the presence of 5, 25, and 50 µg/ml erbstatin. Tyrosine phosphorylated proteins were revealed as described for Figure 1. Molecular weight markers (×10³) are indicated to the right of the blot. Rf = running front. Representative of two similar experiments.

2). Moreover, the acrosome reaction was evaluated by scoring only curly-tailed (and thus viable) spermatozoa [27], further implying that the acrosome reaction-inducing activity of tyrosine kinase inhibitors is not attributable to toxic effects. In addition, tyrosine kinase inhibition with genistein did not affect sperm [Ca2+]i [35], excluding the involvement of this pathway in the acrosome reaction-inducing activity of these compounds. As for response to progesterone, when spermatozoa were incubated in capacitating, complete medium, progesterone increased the acrosome reaction (Table 3), as indicated by the difference between percentage progesterone-induced and percentage spontaneously acrosome-reacted spermatozoa (considered to be the percentage of spermatozoa in the population capable of responding to progesterone-thus showing acrosome reaction following progesterone challenge [ARPC]). In contrast, the steroid was ineffective when preincubation was conducted in Ca²⁺-free medium or in the presence of EGTA (Table 3), confirming that the presence of Ca^{2+} in the external medium is required to obtain capacitation of human spermatozoa [1, 36]. When spermatozoa were incubated in the presence of the phosphatase inhibitor Na_3VO_4 , the acrosome reaction in response to progesterone was lower than that obtained in complete medium (Table 3). On the other hand, the presence of erbstatin during capacitation did not apparently modify the ability of progesterone to induce the acrosome reaction (Table 3).



FIG. 7. Effect of progesterone (P; 4 μ g/ml, 5-min incubation) on tyrosine phosphorylation of sperm proteins. Spermatozoa were capacitated for 2 h in Ca²⁺-containing medium, washed, reincubated in Ca²⁺-free medium, and stimulated with progesterone in the presence or absence of EGTA (E; 3 mM). Tyrosine phosphorylated proteins were revealed as described for Figure 1. Molecular weight markers (×10³) are indicated to the right of the blot. Representative of two similar experiments.

Effect of Progesterone on Sperm Protein Tyrosine Phosphorylation in Absence of Extracellular Ca²⁺

To test the role of Ca^{2+} in the progesterone-mediated increase in tyrosine phosphorylation of sperm proteins, spermatozoa that had been capacitated for 2 h in complete medium were incubated with progesterone for 5 min in Ca^{2+} -free medium in the presence or absence of EGTA (3 mM). As shown in Figure 7, in both conditions progesterone was able to induce an increase in tyrosine phosphorylation of the 97-kDa protein band that was similar in magnitude (1.5-fold over basal) to that observed in Ca^{2+} -containing medium [16, 37].

DISCUSSION

In the present study we investigated the interrelationship between the well-documented increases of $[Ca^{2+}]_i$ [3–6] and protein tyrosine phosphorylation [14–17] occurring during the process of capacitation of spermatozoa. We clearly demonstrate that the spontaneous increase in tyrosine phosphorylation of several protein bands during in vitro capacitation of human sperm was still present when the cells were incubated in Ca²⁺-free/EGTA-containing medium or in the presence of the intracellular Ca²⁺ chelator BAPTA, thus implying that extracellular Ca²⁺ and the spontaneous increase in $[Ca^{2+}]_i$ during capacitation are not necessary for the increase in tyrosine phosphorylation. At the same time, we show that incubation in (nominally) Ca²⁺-free medium resulted in an increase in sperm tyrosine phosphorylation and tyrosine kinase activity. Although this

TABLE 3. Effect of capacitation in different conditions on basal acrosome reaction and in response to progesterone.^a

| | | Condition of capacitation ⁶ | | | | | | | | | | | | | |
|----------|-----------------|--|--------------------------|----|---------------------------------|------|-------------|-----|------|-----------------------------------|----|------|---|----|------|
| - Exp | Complete medium | | Without Ca ²⁺ | | Without Ca ²⁺ + EGTA | | + Erbstatin | | | + Na ₃ VO ₄ | | | | | |
| | С | Р | ARPC ^b | С | Р | ARPC | С | Р | ARPC | С | Р | ARPC | С | Р | ARPC |
| 1 | 8 | 21 | 13 | 7 | 10 | 3 | 4 | 8.5 | 4.5 | 27 | 42 | 15 | 6 | 11 | 5 |
| 2 | 5 | 15 | 10 | 9 | 10 | 1 | 5 | 6 | 1 | _ | | | 7 | 14 | 7 |
| 3 | 7 | 25 | 18 | 9 | 9 | 0 | 7 | 9 | 2 | 33 | 51 | 18 | 7 | 9 | 2 |
| 4 | 14 | 26.5 | 12.5 | 16 | 18 | 2 | 6 | 13 | 7 | 38 | 65 | 27 | _ | _ | |
| 5 | 17 | 42 | 25 | 17 | 29 | 12 | 10 | 11 | 1 | 50 | 66 | 16 | | | |

^a Spermatozoa were capacitated for 2 h in the different conditions, centrifuged, resuspended in complete medium, and incubated for 1 h in presence (P) or absence (C) or progesterone (4 μ g/ml). Results of single experiments obtained in different sperm samples are shown. Values represent percentage acrosome-reacted sperm.

^b ARPC = acrosome reaction following progesterone challenge (difference between percentage progesterone-induced and percentage spontaneous acrosome-reacted spermatozoa).

condition cannot be considered an absolute absence of Ca^{2+} , since contamination from water and other sources (including extrusion of Ca²⁺ from spermatozoa) cannot be excluded, this result, consistently observed in several experiments conducted in different sperm samples, leads to the hypothesis that Ca²⁺ ions restrain the increase in tyrosine phosphorylation during capacitation. Indeed, in the virtual absence of Ca^{2+} in the external medium, the uptake of the ion that occurs during sperm capacitation is apparently reduced as demonstrated by the decreased [Ca²⁺]_i observed in this condition. Thus, the higher tyrosine phosphorylation observed in Ca^{2+} -free medium is likely to be due to the lower levels of Ca²⁺ in the cell. This hypothesis is further substantiated by our experimental results showing a concentration-dependent decrease in tyrosine phosphorylation in the presence of increasing concentrations of added Ca²⁺ as well as after incubation with agents that promote an influx of Ca^{2+} in human sperm, such as the Ca^{2+} ionophore A23187 and the endoplasmic Ca²⁺-ATPase inhibitor thapsigargin.

Moreover, since tyrosine kinase activity during capacitation was higher in Ca²⁺-free medium than in Ca²⁺-containing medium (Table 1), the presence of Ca²⁺-inhibited tyrosine kinase in human sperm is suggested. It is worth noting that in two recent studies conducted in epididymal mouse spermatozoa, Visconti et al. [38, 39] reported that the increase in tyrosine phosphorylation during capacitation in this species is dependent on the presence of Ca^{2+} in the external medium [38] and is mainly regulated by cAMPdependent protein kinase [39]. However, the pattern of tyrosine phosphorylated proteins that increase during capacitation in this species is quite different from the pattern for human sperm. In particular, in human sperm, an increase in tyrosine phosphorylation has been reported in a protein band in the 95-97-kDa molecular mass range, among others ([15-17]; present paper). Such protein, which has been recently sequenced [17], has been shown to be the sperm receptor for ZP3, the zona pellucida protein responsible for inducing the acrosome reaction [14, 17, 18]. Conversely, in the studies reported by Visconti et al. [38, 39], a 97-kDa protein band had a similar tyrosine phosphorylation intensity in both noncapacitated and capacitated sperms and appeared to be different from the ZP3 receptor [17]. It is possible that epididymal spermatozoa have a different mode of capacitation than ejaculated sperm and/or that important species-related differences exist among mammals in this process.

So far, the possible role of Ca^{2+} ions in phosphorylation of sperm proteins in processes other than capacitation has been investigated in few studies [14, 19, 32, 40, 41]. These studies suggest that sperm kinase activity is poorly modulated or, as in the case of bovine sperm [40], even inhibited by an increase in $[Ca^{2+}]_i$, in agreement with our findings. In that report, however, total sperm kinase activity was evaluated. Tesarik et al. [37] reported that the increase in tyrosine phosphorylation induced by progesterone in human sperm was independent of the increase in Ca^{2+} . In the present paper we confirm these studies, showing that the increase in tyrosine phosphorylation in response to progesterone also occurs in Ca²⁺-free/EGTA-containing medium—thus suggesting that progesterone affects $[Ca^{2+}]_i$ and protein phosphorylation by two distinct pathways. In agreement with this hypothesis, recent data obtained by our group indicate an involvement of tyrosine kinase in the generation of the sustained phase of $[Ca^{2+}]_i$ increase induced by progesterone [35].

An important, yet unresolved, question regarding sperm capacitation is which event, among all the events described to occur during this process, is absolutely required to achieve capacitation. In the past, Ca²⁺ has been indicated to be essential for capacitation, motility, and the acrosome reaction of human sperm [1, 36]; however, recent data demonstrate that in particular conditions, the acrosome reaction can also be obtained in Ca²⁺-free medium [25, 42, 43] and that the capacitative effect of the phorbol ester TPA (tetradecanoylphorbol 13-acetate) can be demonstrated also in the absence of Ca^{2+} [44]. More recently, Bielfeld et al. [45] have reported that the acrosome reaction in response to zona pellucida proteins necessitates both $[Ca^{2+}]_i$ increase and kinase activation. We evaluated the acrosome reaction in response to progesterone as an index of capacitation. Indeed, although a clear assay for identifying capacitation has not been defined (for review see [2]), we [6] and others [7-9] have shown that response to progesterone does not occur in uncapacitated sperm and that thus this parameter can be used as a measure of the occurrence of capacitation. Our findings on progesterone-stimulated acrosome reaction clearly confirm that Ca²⁺ is essential to achieve sperm capacitation in vitro. Indeed, if sperm are preincubated in Ca²⁺-free medium or in the presence of EGTA, the acrosome reaction in response to progesterone is absent or highly compromised, suggesting lack of capacitation in these conditions, as already indicated in previous studies [6-9]. Conversely, inhibition of tyrosine kinase during capacitation does not prevent the response to progesterone, and conditions that increase tyrosine phosphorylation (presence of Na_3VO_4 , absence of Ca^{2+}) impair or prevent the response to progesterone, indicating that capacitation does not likely occur in these conditions. It appears that tyrosine phosphorylation must be kept at a low level (by the presence of Ca^{2+} in the medium or by pharmacological inhibition of its activity with erbstatin) to produce a response to progesterone. In view of the fact that tyrosine kinase activation is necessary for exocytosis in response to physiological stimuli such as zona proteins [14, 18] and progesterone [16, 37], it is conceivable that low levels of tyrosine phosphorylation are required in order to obtain a response to these stimuli. An intriguing possibility is that Ca2+-inhibited tyrosine kinase may provide a mechanism for the control of tyrosine phosphorylation during capacitation. Although the in vitro condition is much different from the in vivo one, our results suggest that the Ca²⁺-dependent decrease in tyrosine phosphorylation might be important in vivo during sperm transit in the female genital tract for maintaining low levels of tyrosine phosphorylation in spermatozoa in order to ensure full responsiveness to the physiological agents (progesterone, zona pellucida proteins) at the moment of interaction with the oocyte.

Another important issue concerns the role of Ca^{2+} in maintenance of sperm motility. Recent data indicate a negative correlation between Ca^{2+} and motility [39, 41, 46]. Furthermore, the increase in motility induced by the phorbol ester, TPA, in human sperm was found to be independent of extracellular Ca^{2+} [47]. In our study, we found that motility was not affected by incubation in (nominally) Ca^{2+} -free medium (which resulted in increased tyrosine phosphorylation), was slightly affected by the intracellular Ca^{2+} chelator BAPTA, and was blunted by incubation with EGTA after 2-h incubation, suggesting that Ca^{2+} ions are at least partially involved in maintenance of sperm motility in vitro. Conversely, no apparent correlation appears to exist between motility and level of phosphorylation in tyrosine (see Table 2), although a 2-h incubation with erbstatin decreased motility. However, factors controlling sperm motility and capacitation in vivo are probably multiple and are unknown, thus making it difficult to compare the in vitro condition with the physiological one.

Spermatozoa do not appear to be the only cell type in which Ca²⁺ negatively affects phosphorylation. Indeed, an increase in intracellular Ca2+ has been shown to affect phosphorylation of proteins and phosphatase activity in other cell systems [34, 48-50]. In particular, A23187 has been shown to inhibit epidermal growth factor-induced tyrosine phosphorylation in HER14 cells [50]. However, the ionophore has also been shown to inhibit PTP activity in the same cells [51], suggesting that the inhibition of epidermal growth factor-mediated phosphorylation by A23187 in HER14 cells is due to an inhibitory action of Ca²⁺ on tyrosine kinase as in our system. The possible molecular mechanism(s) involved in Ca2+-mediated inhibition of sperm tyrosine kinase activity is at present under investigation in our lab. In particular, we are studying the possible involvement of Ca²⁺-regulated kinases in this phenomenon.

In conclusion, our results show that tyrosine phosphorylation in human sperm during capacitation in vitro is, at least partially, independent of the concomitant increase of Ca^{2+} , and that this ion negatively modulates tyrosine phosphorylation and tyrosine kinase activity. This mechanism might be relevant in vivo during physiological capacitation of sperm in the female genital tract.

ACKNOWLEDGMENTS

We thank Dr. Maria G. Torcia (Unita' di Endocrinologia, Universita' di Firenze) and Dr. Fabio Marra (Istituto di Medicina Interna, Universita' di Firenze) for helpful advice, and Dr. Mario Maggi and Prof. Mario Serio (Unita' di Endocrinologia, Universita' di Firenze) for critical reading of the manuscript. We wish to thank Dr. Andrea Romani (Dept. of Physiology and Biophysics, CWRU, Cleveland, OH) for calculating concentrations of Ca^{2+} and Mg^{2+} .

REFERENCES

- Yanagimachi R. Mammalian Fertilization. In: Knobil E, Neill JD (eds.), The Physiology of Reproduction, second edition. New York: Raven Press; 1994: 189–317.
- Cohen-Dayag A, Eisenbach M. Potential assays for sperm capacitation in mammals. Am J Physiol 1994; 267:C1167-C1176.
- Coronel CE, Lardy HA. Characterization of Ca²⁺ uptake by guinea pig epididymal spermatozoa. Biol Reprod 1987; 37:1097–1107.
- White DR, Aitken RJ. Relationship between calcium, cyclic AMP, ATP, and intracellular pH and the capacity of hamster spermatozoa to express hyperactivated motility. Gamete Res 1989; 22:163–177.
- Zhou R, Shi B, Chou KCK, Oswalt MD, Hang A. Changes in intracellular calcium of porcine sperm during in vitro incubation with seminal plasma and a capacitating medium. Biochem Biophys Res Commun 1990; 172:47-53.
- Baldi E, Casano R, Falsetti C, Krausz Cs, Maggi M, Forti G. Intracellular calcium accumulation and responsiveness to progesterone in capacitating human spermatozoa. J Androl 1991; 12:323–330.
- Mendoza C, Tesarik J. A plasma membrane progesterone receptor is switched on by increasing intracellular free calcium. FEBS Lett 1993; 330:57-60.
- Shi Q-X, Roldan ERS. Evidence that a GABA_A-like receptor is involved in progesterone-induced acrossomal exocytosis in mouse spermatozoa. Biol Reprod 1995; 52:373–381.
- Meyers SA, Overstreet JW, Liu IKM, Drobnis EZ. Capacitation in vitro of stallion spermatozoa: comparison of progesterone-induced acrosome reaction in fertile and subfertile males. J Androl 1995; 16:47– 54.
- Florman HM. Sequential focal and global elevations of sperm intracellular Ca²⁺ are initiated by the zona pellucida during acrosomal exocytosis. Dev Biol 1994; 165:152–164.
- 11. Florman HM, Tombes RM, First NL, Babcock DF. An adhesion-as-

sociated agonist from the zona pellucida activates G protein-promoted elevations of internal Ca^{2+} and pH that mediate mammalian sperm acrosomal exocytosis. Dev Biol 1989; 135:133–146.

- Thomas P, Meizel S. Phosphatidylinositol 4,5,-bisphosphate hydrolysis in human sperm stimulated with follicular fluid or progesterone is dependent upon Ca²⁺ influx. Biochem J 1989; 264:539–564.
- 13. Blackmore PF, Beebe SJ, Danforth DR, Alexander N. Progesterone and 17α hydroxyprogesterone: novel stimulators of calcium influx in human sperm. J Biol Chem 1990; 265:1367–1380.
- Leyton L, Saling PM. 95 kd sperm proteins bind ZP3 and serve as tyrosine kinase substrates in response to zona binding. Cell 1989; 51: 1123-1130.
- Naz RK, Ahmad K, Kumar R. Role of membrane phosphotyrosine proteins in human spermatozoal function. J Cell Sci 1991; 99:157– 165.
- Luconi M, Bonaccorsi L, Krausz Cs, Gervasi G, Forti G, Baldi E. Stimulation of protein tyrosine phosphorylation by platelet-activating factor and progesterone in human spermatozoa. Mol Cell Endocrinol 1995; 108:35-42.
- Burks DJ, Carballada R, Moore HDM, Saling PM. Interaction of a tyrosine kinase from human sperm with the zona pellucida at fertilization. Science 1995; 269:83–86.
- Leyton L, LeGuen P, Bunch DO, Saling PM. Regulation of mouse gamete interaction by a sperm tyrosine kinase. Proc Natl Acad Sci USA 1992; 89:11692–11695.
- Hayashi H, Yamamoto K, Yonekawa H, Morisawa M. Involvement of tyrosine protein kinase in the initiation of flagellar movement in rainbow trout spermatozoa. J Biol Chem 1987; 262:16692–16698.
- Puoyssegur J, Seuwen K. Transmembrane receptors and intracellular pathways that control cell proliferation. Annu Rev Physiol 1992; 54: 195-210.
- Aitken RJ, Paterson M, Fisher H, Buckingham DW, van Duin M. Redox regulation of tyrosine phosphorylation in human spermatozoa and its role in the control of human sperm function. J Cell Sci 1995; 108:2017–2025.
- Baldi E, Krausz Cs, Luconi M, Bonaccorsi L, Maggi M, Forti G. Actions of progesterone on human sperm: a model of non-genomic effects of steroids. J Steroid Biochem Mol Biol 1995; 53:199-203.
- 23. World Health Organization. WHO Laboratory Manual for the Examination of Human Semen and Semen-Cervical Mucus Interaction, Third Edition. Cambridge: University Press; 1992.
- Baldi E, Falsetti C, Krausz Cs, Gervasi G, Carloni V, Casano R, Forti G. Stimulation of platelet-activating factor synthesis by progesterone and A23187 in human spermatozoa. Biochem J 1993; 292:209–216.
- Krausz Cs, Gervasi G, Forti G, Baldi E. Effect of platelet-activating factor on motility and acrosome reaction of human spermatozoa. Hum Reprod 1994; 9:471–476.
- Falsetti C, Baldi E, Krausz Cs, Casano R, Failli P, Forti G, Baldi E. Decreased responsiveness to progesterone of spermatozoa in oligozoospermic patients. J Androl 1993; 14:17–22.
- Aitken RJ, Buckingham DW, Fang HG. Analysis of the response of human spermatozoa to A23187 employing a novel technique for assessing acrosome reaction. J Androl 1993; 14:132–141.
- Bunch DO, Saling PM. Generation of mouse sperm membrane fraction with zona receptor activity. Biol Reprod 1991; 44:672–680.
- Grynkiewicz G, Poenie M, Tsien RY. A generation of calcium indicators with greatly improved fluorescence properties. J Biol Chem 1985; 260:3440-3450.
- Fabiato A. Computer program for calculating total from specified free or free from specified total ionic concentrations in aqueous solutions containing multiple metals and ligands. Methods Enzymol 1988; 157: 378-417.
- Blackmore PF. Thapsigargin elevates and potentiates the ability of progesterone to increase intracellular free calcium in human sperm: possible role of perinuclear calcium. Cell Calcium 1992; 14:53–60.
- Meizel S, Turner KO. Initiation of the human sperm acrosome reaction by thapsigargin. J Exp Zool 1993; 267:350–355.
- Moos J, Carrera A, Tesarik J, Moss SB, Gerton GL, Kopf GS. Regulation, localization, and identity of phosphotyrosine-containing proteins in human sperm. Biol Reprod 1995; 52(suppl 1):168 (abstract 445).
- Ostergaard HL, Trowbridge IS. Negative regulation of CD45 protein tyrosine phosphatase activity by ionomycin in T cells. Science 1991; 253:1423-1425.
- Bonaccorsi L, Luconi M, Forti G, Baldi E. Tyrosine kinase inhibition reduces the plateau fase of calcium increase in response to progesterone in human sperm. FEBS Lett 1995; 364:83–86.

- DasGupta S, Mills CL, Fraser LR. Ca²⁺-related changes in the capacitation state of human spermatozoa assessed by a chlortetracycline fluorescence assay. J Reprod Fertil 1993; 99:135–143.
- Tesarik J, Moos J, Mendoza C. Stimulation of protein tyrosine phosphorylation by a progesterone receptor on the cell surface of human sperm. Endocrinology 1993; 33:328–335.
- Visconti PE, Bailey JL, Moore GD, Pan D, Olds-Clarke P, Kopf GS. Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation. Development 1995; 121:1129-1137.
- 39. Visconti PE, Moore GD, Bailey JL, Leclerc P, Connors SA, Pan D, Olds-Clarke P, Kopf GS. Capacitation of mouse spermatozoa. II. Protein tyrosine phosphorylation and capacitation are regulated by a cAMP-dependent pathway. Development 1995; 121:1139–1150.
- Carr DW, Acott TS. The phosphorylation of a putative sperm microtubule-associated protein 2 (MAP2) is uniquely sensitive to regulation. Biol Reprod 1990; 43:795–805.
- Duncan AE, Fraser LR. Cyclic AMP-dependent phosphorylation of epididymal mouse sperm proteins during capacitation in vitro: identification of an Mr 95 000 phosphotyrosine-containing protein. J Reprod Fertil 1993; 97:287-299.
- Rotem R, Paz GF, Homonnai ZT, Kalina M, Lax J, Breitbart H, Naor Z. Ca²⁺-Independent induction of acrosome reaction by protein kinase C in human sperm. Endocrinology 1992; 131:2235–2243.
- Foresta C, Rossato M, Di Virgilio F. Extracellular ATP is a trigger for the acrosome reaction in human spermatozoa. J Biol Chem 1992; 267: 19443–19447.

- Furuya S, Endo Y, Osumi K, Oba M, Suzuki S. Effects of modulators of protein kinase C on human sperm capacitation. Fertil Steril 1993; 59:1285–1290.
- 45. Bielfeld P, Anderson RA, Mack SR, De Jonge CJ, Zaneveld LJD. Are capacitation or calcium ion influx required for the human sperm acrosome reaction? Fertil Steril 1994; 62:1255–1261.
- McLaughlin EA, Ford WCL. Effects of cryopreservation on the intracellular calcium concentration of human spermatozoa and its response to progesterone. Mol Reprod Dev 1994; 37:241–246.
- Rotem R, Paz GF, Homonnai ZT, Kalina M, Naor Z. Further studies on the involvement of protein kinase C in human sperm flagellar motility. Endocrinology 1990; 127:2571–2577.
- Sihra TS, Bogonez E, Nicholls DG. Localized Ca²⁺ entry preferentially effects protein dephosphorylation, phosphorylation, and glutamate release. J Biol Chem 1992; 267:1983–1989.
- Katoh S, Funajama A, Kohno H, Ohkubo Y. Dephosphorylation of tyrosine of epidermal growth factor receptor is inhibited by Ca²⁺ pretreatment in isolated liver membrane. Arch Biochem Biophys 1993; 307:52–56.
- Errasfa M, Stern A. Melittin inhibits epidermal growth factor-induced protein tyrosine phosphorylation: comparison with phorbol myristate acetate and calcium ionophore A23187. Biochim Biophys Acta 1994; 1222:471-476.
- Errasfa M, Stern A. Inhibition of protein tyrosine phosphatase activity in HER14 cells by melittin and Ca²⁺ ionophore A23187. Eur J Pharmacol 1993; 247:73–80.