



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Neurotransmitter role of endogenous morphine in CNS.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Neurotransmitter role of endogenous morphine in CNS / Massimo Guarna; Carla Ghelardini; Nicoletta Galeotti; George B Stefano; Enrica Bianchi. - In: MEDICAL SCIENCE MONITOR. - ISSN 1234-1010. - STAMPA. - 11:(2005), pp. RA190-RA193.

Availability:

This version is available at: 2158/326946 since: 2016-11-09T11:06:20Z

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Received: 2005.05.08
Accepted: 2005.05.12
Published: 2005.06.01

Neurotransmitter role of endogenous morphine in CNS

**Massimo Guarna¹, Carla Ghelardini², Nicoletta Galeotti²,
George B. Stefano³, Enrica Bianchi⁴**

¹ Department of Anatomical and Biomedical Sciences, University of Siena, Italy

² Department of Clinical and Preclinical Pharmacology, University of Florence, Italy

³ Neuroscience Research Institute, State University of New York, U.S.A.

⁴ Department of Neuroscience, University of Siena, Italy

Source of support: Departmental sources

Summary

Endogenous morphine is present in the mammalian brain as determined by gas chromatography/mass spectrometry. The criteria essential for satisfying the definition of morphine as a neurotransmitter are examined. The detection of endogenous morphine-like compounds inside brain neurons by immunocytochemistry and the Ca⁺ dependent release of endogenous morphine from rat brain slices provide evidence for its transmitter status. Indirect evidence that endogenous morphine modulates thermnociception and weakens memory through mu opioid receptors again supports a neurotransmitter role for this chemical messenger. Evidence has been found for its endogenous synthesis in animal tissues as well. These findings indicate that endogenous morphine might function as neuromodulator/neurotransmitter agent in the CNS.

key words:

endogenous morphine • neurotransmitter • opiate receptor • mu3 opiate receptor subtype • nitric oxide • amygdala • hippocampus • memory

Full-text PDF: http://www.MedSciMonit.com/pub/vol_11/no_6/7467.pdf

Word count: 1542

Tables: –

Figures: 1

References: 39

Author's address: Enrica Bianchi, Dipartimento di Neuroscienze, Via A. Moro 6, 53100 Siena, Italy, e-mail: bianchi16@unisi.it

BACKGROUND

A morphine-like material in animal tissues was originally demonstrated by immunological recognition [1]. Subsequently, endogenous morphine-like compounds have been identified in mouse and calf brain [2,3]. The molecular structure of the HPLC purified compound was confirmed as morphine by liquid and gas chromatographic retention times and mass spectrometry in various tissues such as bovine brain, rat and mouse brain, hypothalamus and adrenal glands, mammalian lung, invertebrate and human tissues, human cerebrospinal fluid and human plasma [4-16]. Codeine and morphine also have been identified in human cerebrospinal fluid [14] and non human primate brain [17]. Morphine precursors, reticuline [18], thebaine [19] and codeine [20-22] are some of the main intermediates of morphine biosynthesis in the poppy plant and have been found in mammals.

The pathway for morphine biosynthesis has been established in the opium poppy and animal tissue. In this regard, radiolabeled reticuline was transformed into salutaridine by rat liver microsomes *in vitro* and a similar conversion has been observed *in vivo* [23]. The conversion of salutaridine, thebaine and codeine into morphine has been demonstrated in several rat tissues, including the brain, providing evidence for the biosynthetic pathway of endogenous morphine (eM) in mammals [5,24]. Incubation of human neural cancer cell lines with L-tyrosine leads to the formation of morphine as well [25]. Addition of reticuline to intact invertebrate ganglia *in vivo* leads to the formation of morphine [26]. *Ascaris suum*, a mammalian parasite produces morphine in mammalian hosts and *in vitro*, a phenomenon that may dull the host discomfort so that the parasite can escape host surveillance [27].

IMMUNOCYTOCHEMICAL LOCALIZATION OF ENDOGENOUS MORPHINE IN MAMMALIAN BRAIN

Morphine-like immunoreactivity has been demonstrated in cell bodies, fibers and terminals of neurons in different brain areas of the rat [28], mouse [2] and man [29] by indirect immunofluorescence and unlabeled antibody enzyme procedure. The antisera revealed numerous nerve cell bodies with variable morphology, dendrites and nerve fibers throughout the cerebral cortex. In the caudate putamen numerous medium size immunoreactive perikaria were visible. Nerve cell body and fibers were stained in the hippocampal formation; dot-like immunoreactivity was visible around granule cells of dentate gyrus and dorsal hippocampus; immunoreactive nerve cell bodies were also revealed throughout the brainstem. In the cerebellum numerous granule cells in the granular layer and small neurons and fibers in the molecular layer are immunoreactive. In human brain tissue, morphine-like immunoreactivity was identified in cerebral and cerebellar cortex. Morphine positive terminals were observed in rat hippocampal formation sometimes forming synaptic contact with the soma of unlabeled cells [29]. In different regions of the brainstem positive morphine-like nerve fibers apposed to μ opiate receptor (μ OR) immunoreactive cell bodies were observed (Figure 1).

MORPHINE RELEASE

Neurotransmitters and neuromodulators are released from neurons by depolarization in a Ca^{++} dependent manner.

Neuron depolarization by high potassium concentrations caused the release of endogenous morphine from eel chromaffin cells [20]. The release of endogenous morphine from rat brain slices was observed *in vitro* in the presence of high potassium concentration; this effect disappeared when Ca^{++} was omitted from the superfusion medium [7]. Furthermore, exogenously applied morphine can affect the release of other neurotransmitters, i.e., dopamine, in nervous tissues suggesting a similar role for endogenous morphine [30].

MORPHINE UPTAKE

The slow i.c.v. infusion of [3H]morphine demonstrated the capacity of specific brain neurons to store the alkaloid [28]. Otherwise synaptosomes from mid-brain, cerebral cortex and cerebellum failed to take up and store any trace of the tritiated alkaloid [28]. However, this finding also suggests that an alternate degradation process may be operating. The major pathway for the detoxication of exogenous morphine is conversion of morphine to morphine 6 glucuronide and/or morphine 3 glucuronide. The presence of morphine 6-dehydrogenase in mammalian liver [31] suggests the metabolic pathway of morphine to morphinone as a route in endogenous morphine metabolism. Thus, endogenous morphine conversion may represent the way to terminate its action.

ROLE OF ENDOGENOUS MORPHINE IN THE MODULATION OF THERMAL NOCICEPTION

The depletion of endogenous morphine in mouse brain was obtained by immunoneutralization of this molecule from brain extracellular spaces. An affinity purified anti-morphine IgG was administered to mice, which then underwent the hot plate test. As shown by the licking latency in the hot plate test, the nociceptive threshold in response to thermal stimulus was significantly lowered by anti-morphine IgG with respect to controls. We attributed this effect to a decrease in endogenous morphine levels present in mouse brain extracellular spaces [8]. Endogenous morphine depletion attenuated the antinociceptive effect of the μ selective agonist DAMGO in the hot plate test whereas enkephalin or DPDE antinociceptive effect was unmodified [8]. Therefore, endogenous morphine appears to be involved in the modulation of thermonociception in mice via the μ OR receptor [8].

ROLE OF ENDOGENOUS MORPHINE IN WEAKENING MEMORY PROCESSES UNDER STRESS CONDITION

Classically, exogenous morphine administration impairs memory processes [32] and a highly significant regression coefficient between analgesia and memory impairment was found after opiate administration [33]. Twelve hour food deprivation (fasting) induced a stress condition, which does not interfere with motor coordination, spontaneous motility and exploratory activity. An anti-morphine IgG was administered to mice in order to induce morphine depletion in extracellular spaces. Fed and stressed animals were both administered with anti-morphine IgG and submitted to passive avoidance test to study the effect of endogenous morphine depletion on working memory [34]. Fed animals were used as a control. Acquisition and consolida-




Figure 1. Bright field of coronal section of mouse brainstem double stained with anti- μ OR (brown) and anti-morphine (violet) IgG. Morphine-like immunoreactive nerve fibers are closely apposed to a μ OR positive neurons.

tion of memory was significantly impaired in stressed animals with respect to control. This was demonstrated by a decrease in entry latency into the dark room in the retention session of the passive avoidance test. This effect was reversed to the basal value after endogenous morphine depletion. Therefore, endogenous morphine may be involved in weakening memory processes under stress conditions [34]. This may provide a strategy for limiting future hesitancy when action is called for.

MU RECEPTOR INTERACTION: POSTSYNAPTIC RESPONSE SYSTEM

Recently the μ_3 OR has been cloned in human neural (cell line), immune and vascular tissues [35]. This μ_3 OR splice variant is morphine selective and opioid peptide insensitive and is coupled to nitric oxide release [36]. In rat limbic tissues (hippocampus and amygdala) morphine is present as demonstrated by biochemical, including mass spectrometry, and immunocytochemical techniques [28,37,38]. Additionally, adding exogenous morphine to these tissues results in constitutive nitric oxide release, which does not occur with opioid peptides, indirectly demonstrating the μ_3 OR subtype [36]. Taken together, a highly specific postsynaptic receptor exists in these tissues for morphine.

Criteria for defining a chemical messenger as a neurotransmitter exists. These criteria are the following: 1) The chemical must be produced and found within a neuron; 2) When a neuron is depolarized, it must release the chemical in a Ca^{++} dependent manner; 3) When released, the chemical messenger must act on a post-synaptic receptor and cause a biological effect; 4) After exerting its action, the chemical messenger must be inactivated via a reuptake mechanism or by an enzyme that terminates its action; and lastly, 5) If the chemical messenger is applied on the post-synaptic membrane, it should have the same effect as when it is released by a neuron. Taken together, the earlier noted studies demonstrate that morphine functions as a neurotransmitter.

CONCLUSIONS

Reticuline [18], thebaine [19] and codeine [12,20,38] are some of the main intermediates of morphine biosynthesis in the poppy plant and these compounds have been found

in mammals and invertebrates, especially in their neural tissues. Radiolabeled reticuline was transformed into salutaridine by rat liver microsomes *in vitro* and a similar conversion has been observed *in vivo* [23]. The conversion of salutaridine, thebaine and codeine into morphine has been demonstrated in several rat tissues, including the brain, providing evidence for the biosynthetic pathway of endogenous morphine in mammals [5,24]. Reticuline exposure to invertebrate ganglia leads to morphine formation as well [26]. Presently morphine appears to be endogenously synthesized by brain microsomes even if microsomes are intracellular membrane structures belonging to different brain cell type. In a neural cancer cell line L-tyrosine exposure leads to morphine formation [25]. Morphine immunoreactivity has been demonstrated inside the cell body, fibers and terminals of neurons in different brain areas of the rat [28], mouse [2], humans [29] and invertebrate neural tissues [39]. We also have demonstrated that endogenous morphine can be released in a chemically detectable form from rat brain slices in a Ca^{2+} and K^+ dependent manner, fulfilling a criterion for endogenous morphine as a neurotransmitter [12]. When endogenous morphine was depleted from mice brain extracellular spaces, acute thermociception was increased and working memory weakness, a classic effect of exogenous morphine administration, was impaired [8,34]. Both these effects appeared to be mediated through μ OR. We failed to observe an active uptake of [^3H]morphine by rat brain synaptosomes [28]. However, its enzymatic conversion to different metabolites may satisfy this criterion. A novel μ_3 OR exists, which is coupled to nitric oxide release and only responds to morphine. This occurs in both human, mammalian and invertebrate tissues. Endogenous morphine and the μ receptor are associated throughout the phylogensis of the nerve system of mammal and lower animals, suggesting that this molecule is a physiological ligand of the μ receptor.

REFERENCES:

- Gintzler AR, Levy A, Spector S: Antibodies as a means of isolating and characterizing biologically active substances: Presence of a non-peptide morphine-like compound in the central nervous system. *Proc Natl Acad Sci USA*, 1976; 73: 2132-36
- Gintzler AR, Gershon MD, Spector S: A nonpeptide morphine-like compound: immunocytochemical localization in the mouse brain. *Science*, 1978; 199: 447-48
- Killian AK, Schuster CR, House JT et al: A non-peptide morphine-like compound from brain. *Life Sci*, 1981; 28: 811-17
- Goldstein A, Barrett RW, James IF et al: Morphine and other opiates from beef brain and adrenal. *Proc Natl Acad Sci USA*, 1985; 82: 5203-7
- Donnerer J, Oka K, Brossi A et al: Presence and formation of codeine and morphine in the rat. *Proc Natl Acad Sci USA*, 1986; 83: 4566-67
- Donnerer J, Cardinale G, Coffey J et al: Chemical characterization and regulation of endogenous morphine and codeine in the rat. *J Pharmacol Exp Ther*, 1987; 242: 583-87
- Guarna M, Neri C, Petrioli F, Bianchi E: Potassium-induced release of endogenous morphine from rat brain slices. *J Neurochem*, 1998; 70: 147-52
- Guarna M, Bianchi E, Bartolini A et al: Endogenous morphine modulates acute thermociception in mice. *J Neurochem*, 2002; 80: 271-77
- Goumon Y, Bouret S, Casares F et al: Lipopolysaccharide increases endogenous morphine levels in rat brain. *Neuroscience Letters*, 2000; 293: 135-38
- Goumon Y, Stefano GB: Identification of Morphine in the Rat Adrenal Gland. *Mol Brain Res*, 2000; 77: 267-69

11. Goumon Y, Weeks BS, Cadet P, Stefano GB: Identification of morphine in the adrenal medullary chromaffin PC-12 cell line. *Mol Brain Res*, 2000; 81: 177-80
12. Munjal ID, Minna JD, Manneckjee R et al: Possible role of endogenous morphine and codeine on growth regulation of lung tissue. *Life Sci*, 1996; 57: 517-21
13. Stefano GB, Digenis A, Spector S et al: Opiate-like substances in an invertebrate, a novel opiate receptor on invertebrate and human immunocytes, and a role in immunosuppression. *Proc Natl Acad Sci USA*, 1993; 90: 11099-103
14. Cardinale CJ, Donnerer J, Finck AD et al: Morphine and codeine are endogenous components of human cerebrospinal fluid. *Life Sci*, 1987; 40: 301-6
15. Liu Y, Bilfinger TV, Stefano GB: A rapid and sensitive quantitation method of endogenous morphine in human plasma. *Life Sci*, 1997; 60: 237-43
16. Brix-Christensen V, Tonnesen E, Sanchez RG et al: Endogenous morphine levels increase following cardiac surgery as part of the antiinflammatory response? *Int J Cardiol*, 1997; 62: 191-97
17. Neri C, Guarna M, Bianchi E et al: Endogenous morphine and codeine in the brain of non-human primate. *Med Sci Monit*, 2004; 10(6): MS1-MS5
18. Zhu W, Ma Y, Cadet P et al: Presence of reticuline in rat brain: A pathway for morphine biosynthesis. *Mol Brain Res*, 2003; 117: 83-90
19. Kodaira H, Listek CA, Jardine I et al: Identification of the convulsant opiate thebaine in the mammalian brain. *Proc Natl Acad Sci USA*, 1989; 86: 716-19
20. Epple A, Navarro I, Horak P, Spector S: Endogenous morphine and codeine: release by the chromaffin cells of the eel. *Life Sci*, 1993; 52: PL117-PL121
21. Hofmann U, Seefried S, Schweizer E et al: Highly sensitive gas chromatographic-tandem mass spectrometric method for the determination of morphine and codeine in serum and urine in the femtomolar range. *J Chromatogr B Biomed Sci Appl*, 1999; 727: 81-88
22. Stefano GB, Scharer B: Endogenous morphine and related opiates, a new class of chemical messengers. *Adv Neuroimmunol*, 1994; 4: 57-68
23. Weitz CJ, Faull KF, Goldstein A: Synthesis of the skeleton of the morphine molecule by mammalian liver. *Nature*, 1987; 330: 674-77
24. Kodaira H, Spector S: Transformation of thebaine to oripavine, codeine, and morphine by rat liver, kidney, and brain microsomes. *Proc Natl Acad Sci USA*, 1988; 85: 1267-71
25. Poeaknapo C, Schmidt J, Brandsch M et al: Endogenous formation of morphine in human cells. *Proc Natl Acad Sci USA*, 2004; 101: 14091-96
26. Zhu W, Stefano GB: Reticuline exposure to invertebrate ganglia increases endogenous morphine levels. *Neuro Endocrinol Lett*, 2004; 25: 323-30
27. Goumon Y, Casares F, Pryor S et al: *Ascaris suum*, an internal parasite, produces morphine. *J Immunol*, 2000; 165: 339-43
28. Bianchi E, Alessandrini C, Guarna M, Tagliamonte A: Endogenous codeine and morphine are stored in specific brain neurons. *Brain Research*, 1993; 627: 210-15
29. Bianchi E, Guarna M, Tagliamonte A: Immunocytochemical localization of endogenous codeine and morphine. *Adv Neuroimmunol*, 1994; 4: 83-92
30. Stefano GB, Hall B, Makman MH, Dvorkin B: Opioid inhibition of dopamine release from nervous tissue of *Mytilus edulis* and *Octopus bimaculatus*. *Science*, 1981; 213: 928-30
31. Yamano S, Takahashi A, Todaka T, Toki S: *In vivo* and *in vitro* formation of morphinone from morphine in rat. *Xenobiotica*, 1997; 27(7): 645-56
32. Izquierdo I: Effect of beta-endorphin and naloxone on acquisition, memory, and retrieval of shuttle avoidance and habituation learning in rats. *Psychopharmacology (Berl)*, 1980; 69: 111-15
33. Braida D, Gori E, Sala M: Relationship between morphine and etonitazene-induced working memory impairment and analgesia. *Eur J Pharmacol*, 1994; 271: 497-504
34. Guarna M, Ghelardini C, Galeotti N et al: Effects of endogenous morphine deprivation on memory retention of passive avoidance learning in mice. *Int J Neuropsychopharmacol*, 2004; 7: 311-19
35. Cadet P, Mantione KJ, Stefano GB: Molecular identification and functional expression of mu3, a novel alternatively spliced variant of the human mu opiate receptor gene. *Journal of Immunology*, 2003; 170: 5118-23
36. Stefano GB, Goumon Y, Casares F et al: Endogenous morphine. *Trends in Neurosciences*, 2000; 9: 436-42
37. Zhu W, Ma Y, Bell A et al: Presence of morphine in rat amygdala: Evidence for the δ opiate receptor subtype via nitric oxide release in limbic structures. *Med Sci Monit*, 2004; 10(12): BR433-BR439
38. Spector S, Munjal I, Schmidt DE: Endogenous morphine and codeine. Possible role as endogenous anticonvulsants. *Brain Res*, 2001; 915: 155-60
39. Sonetti D, Mola L, Casares F et al: Endogenous morphine levels increase in molluscan neural and immune tissues after physical trauma. *Brain Res*, 1999; 835: 137-47