



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **The killer shrimp *Dikerogammarus villosus* (Sowinsky, 1894) is spreading in Italy.**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

The killer shrimp *Dikerogammarus villosus* (Sowinsky, 1894) is spreading in Italy / E. TRICARICO; G. MAZZA; G. ORIOLI; C. ROSSANO; F. SCAPINI; F. GHERARDI. - In: AQUATIC INVASIONS. - ISSN 1818-5487. - ELETTRONICO. - 5:(2010), pp. 211-214. [10.3391/ai.2010.5.2.14]

*Availability:*

This version is available at: 2158/390890 since:

*Published version:*

DOI: 10.3391/ai.2010.5.2.14

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

## Short communication

## The killer shrimp, *Dikerogammarus villosus* (Sowinsky, 1894), is spreading in Italy

Elena Tricarico, Giuseppe Mazza, Gabriele Orioli, Claudia Rossano, Felicita Scapini and Francesca Gherardi\*

Dipartimento di Biologia Evoluzionistica "Leo Pardi", Università di Firenze, via Romana 17, 50125 Firenze, Italy

E-mail: [elena.tricarico@unifi.it](mailto:elena.tricarico@unifi.it) (ET), [m.beppe81@libero.it](mailto:m.beppe81@libero.it) (GM), [gabri.orioli@gmail.com](mailto:gabri.orioli@gmail.com) (GO), [claudia\\_rossano@libero.it](mailto:claudia_rossano@libero.it) (CR), [felicita.scapini@unifi.it](mailto:felicita.scapini@unifi.it) (FS), [francesca.gherardi@unifi.it](mailto:francesca.gherardi@unifi.it) (FG)

\* Corresponding author

Received: 23 November 2009 / Accepted: 11 January 2010 / Published online: 21 January 2009

### Abstract

In 2008, the killer shrimp, *Dikerogammarus villosus*, native to the Ponto-Caspian region, was found for the first time in Central Italy, in Bilancino, an artificial lake situated in the watershed of the River Arno (Tuscany). This new record shows that this species' range is expanding in Italy. It is thus imperative to identify the pathways and vectors of spread of this species in order to halt this invasion process.

**Key words:** *Dikerogammarus villosus*, inland waters, Italy

Because of its predatory voracity and aggressive behaviour, *Dikerogammarus villosus* (Sowinsky, 1894) is called the "killer shrimp". It is a crustacean amphipod native to the Ponto-Caspian region. After the opening of the Danube-Main-Rhine canal in 1992, as the result of both natural expansion and transportation in ballast waters (Casellato et al. 2007), the species quickly invaded central and western Europe (Van den Brink and Van der Velde 1991; Dick and Platvoet 2000; Kinzler and Maier 2003; Kley and Maier 2003; Grabowski et al. 2007b).

Its progressive spread across Europe poses a serious threat to the diversity of indigenous amphipods and to the wider indigenous communities as well. *Dikerogammarus villosus* shows a number of life history traits typical of an invasive species, such as the short duration of embryonic development, rapid growth, short generation time, early sexual maturity, long reproductive period, and large reproductive capacity (Pöckl 2007, 2009). Besides being a shredder and detritus feeder, it is a voracious predator of both macroinvertebrates and the eggs, larvae, and adults of fish species (Dick et al. 2002; Devin et al. 2003; Casellato et al. 2007; Platvoet et al. 2009). It tolerates wide ranges of temperature, oxygen concentration, and salinity (Brujijns et al. 2001; Bij de Vaate et al. 2002;

Devin et al. 2003; Brooks et al. 2009) and adapts to several types of substrate (Devin et al. 2003), favoured in this by its polymorphic pigmentation (Devin et al. 2004a). Its aggressive behaviour and voracity cause the replacement of indigenous gammarids (Dick and Platvoet 2000; Van Riel et al. 2006, 2007; Casellato et al. 2008; Leuven et al. 2009; Van der Velde et al. 2009) along with changes in the composition of the invaded communities and damages to the food webs of the recipient water bodies (Dick et al. 2002; Haas et al. 2002; Van der Velde et al. 2002). All these characteristics give this invader the potential of becoming widely distributed in inland water systems.

Since 2003, *D. villosus* has been reported in Northern Italy (Lake Garda, Po and Mincio rivers; Casellato et al. 2006, 2007, 2008). During 2008-2009, *D. villosus* was found in several samples of macrobenthos taken from the shoreline of the Lake Bilancino (40 km from Florence, Tuscany, Central Italy; 43°58'41"N, 11°16'54"E; altitude: 255 m above sea level; Figure 1).

Lake Bilancino is an artificial lake built in 1995 (but opened in 2002) to provide Florence with water and energy; it is located in an agricultural area, near a busy highway, and is an oligotrophic lake (total nitrogen TN: 0.83 mg/l;


nitric nitrogen  $\text{NO}_3^-$ : 0.5 mg/l; chlorophyll *a*: 2.3  $\text{mg/m}^3$ ), with low dissolved oxygen saturation during summer in the hypolimnion stratum (<20%) and poor water transparency (on average 3.3 m) (data from the Regional Agency for the protection on the environment, ARPAT 2009). The lake has six influent rivers (Lora, Calecchia, Tavaiano, Sieve, Sorcella, and Stura) and only one effluent (Sieve).

Due to the large number of established alien species, this lake can be considered as a hot spot of xenodiversity. Since 2002, 14 alien species have become established in the lake: the Chinese freshwater jellyfish *Craspedacusta sowerbyi* Lankaster, 1880; the Ponto-Caspian zebra mussel *Dreissena polymorpha* (Pallas, 1771); the American bladder snail *Haitia acuta* (Draparnaud, 1805); the New Zealand mudsnail *Potamopyrgus antipodarum* (Gray, 1843); the Chinese pond mussel *Sinanodonta woodiana* (Lea, 1834); the red swamp crayfish *Procambarus clarkii* (Girard, 1852); the crucian carp *Carassius carassius* (Linnaeus, 1758); the carp *Cyprinus carpio* (Linnaeus, 1758); the black bullhead *Ictalurus melas* (Rafinesque, 1820); the channel catfish *Ictalurus punctatus* (Rafinesque, 1818); the pumpkinseed sunfish *Lepomis gibbosus* (Linnaeus, 1758); the largemouth bass *Micropterus salmoides* Lacépède, 1802; the wels catfish *Silurus glanis* Linnaeus, 1758; and the coypu *Myocastor coypus* (Molina, 1782) (Lori and Cianfanelli 2006; ARPAT 2009). Among the other alien species in the lake, *D. polymorpha*, recorded in Bilancino since 2005 (Lori and Cianfanelli 2006), is causing the strongest economic impact amounting to 20,000 Euro per year to clean pipes and tanks of the water treatment plant managed by the company Publiacqua (P. Grossi, pers. comm.).

*Dikerogammarus villosus* is the only gammarid in the lake, while the indigenous species *Echinogammarus veneris* is reported in the rivers of the same basin (Ruffo and Stoch 2005). During October 2008 and April 2009, we sampled four sites (two on the Northern side, two on the Southern one) along the littoral area of the lake. Sampling was performed up to 2 m from the banks and from 0 to 60 cm depth in areas where gammarids seemed most concentrated. Individuals were collected using a hand net (0.5 mm mesh size) in a square transect of 50×50 cm randomly sorted, and preserved in vials with 75% ethanol; then, in the laboratory, under a light microscope, their sex and maturation level were analyzed. The mean density did



**Figure 1.** View of Lake Bilancino (Tuscany, Central Italy). Photo by C. Rossano.



**Figure 2.** Frequency of males, mature females, immature females, and juveniles sampled in October 2008 and April 2009 in Lake Bilancino. The numbers over the bar indicate sample size. \* denotes a significant difference after the G test with Williams' correction.



**Figure 3.** Two mating individuals of *Dikerogammarus villosus*: the male (the biggest and striped shrimp) is guarding the female. Photo by C. Rossano.

not change between October and April (independent samples Student's *t*-test:  $t=-0.81$ ,  $df=6$ ,  $P=0.45$ ; October:  $210\pm 88$  ind/m<sup>2</sup>; April:  $369\pm 175$  ind/m<sup>2</sup>). From a total of 1571 individuals, the sex ratio was biased towards females (males=335, females= 510, *G* test with Williams' correction:  $G=36.49$ ,  $df=1$ ,  $P<0.001$ ) without any significant difference between October and April ( $G=0.10$ ,  $df=1$ ,  $P>0.1$ ; Figure 2). Our findings confirm previous studies on this species in different areas (e.g. Casellato et al. 2006; Pöckl 2009). In the invasion process, female-biased sex ratio could be advantageous since it increases the reproductive capacity of the population (Devin et al. 2004b). The number of ovigerous females was higher in October (85%) than in April (31%) ( $G=164.68$ ,  $df=1$ ,  $P<0.001$ ), while the opposite was found for the juveniles (April 57% vs October 33%;  $G=220.81$ ,  $df=1$ ,  $P<0.001$ ; Figure 2). In April, we also observed several matings (Figure 3). Similar to other species of Gammaridae, *D. villosus* is a multivoltine species with three generations reported per year (Devin et al. 2004b). In Austria, Pöckl (2009) found a reproductive resting period from October to December, and three peaks of juveniles in May-June, August-October, and January. Breeding typically lasts nine months, but it can extend for more than 10 months in warmer conditions (Grabowski et al. 2007a; Pöckl 2009). For example, in Lake Garda, Casellato et al. (2006) found ovigerous females all year round, with peaks between June and October, while juveniles were the most abundant component of the population particularly in October. Further samplings are needed in Bilancino to better understand the life cycle of the species in this newly invaded area.

In Lake Bilancino, *D. villosus* mostly occupies gravelly bottoms as in Lake Garda (Casellato et al. 2007); it also inhabits the clusters of *D. polymorpha*, which confirms the strong association between the two species. Indeed, in many European countries the invasion of the killer shrimp has been facilitated by the habitat alteration caused by previous introductions of *D. polymorpha* (Gonzales and Downing 1999; Simberloff and Von Holle 1999; Jazdzewski and Konopacka 2002). The invaded habitats are in fact physically modified by zebra mussel: its high-density and dense clusters provide suitable shelter to macroinvertebrates, including *D. villosus*. Moreover, zebra mussel faeces become an additional food source for the indigenous gammarids and for other detriti-

vorous macroinvertebrates, which in turn provide prey for *D. villosus* (Ricciardi et al. 1997; Gonzáles and Downing 1999).

This is the first record of *D. villosus* for Central Italy and warns of the likely further expansion of this species' range across Italy. Similar to *D. polymorpha* (Lori and Cianfanelli 2006), *D. villosus* has the potential of colonizing the River Sieve, the effluent of Lake Bilancino, and subsequently the River Arno downstream of its confluence, whereas its dispersal upstream might be limited. *Dikerogammarus villosus* seems to have been introduced into Lake Garda by tourist boats from central Europe (Casellato et al. 2006, 2007), possibly in association with *D. polymorpha* (Quaglia et al. 2008); it might have been also released into Lake Bilancino as a contaminant of fish stocks. Obviously, further studies are needed to understand the exact pathways and vectors of this species in order to prevent and hopefully control its further spread across Central and Southern Italy.

#### Acknowledgements

We thank Prof. Sandra Casellato (University of Padoa, Italy) for having identified the species and Dr. Paolo Grossi of Publicacqua for the logistic support during sampling. We also thank Dr. Jaimie Dick and two anonymous reviewers for their helpful suggestions. The study was in part supported by the project ALT ("Atlante delle Specie Alloctone in Toscana") cofinanced by Regione Toscana, the Municipality of Scarperia, and the University of Florence.

#### References

- ARPAT (2009) Proposte per l'ecosistema di Bilancino. Regione Toscana, pp 228
- Bij de Vaate A, Jazdzewski K, Ketelaars HAM, Gollasch S, Van der Velde G (2002) Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in Europe. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1159-1174, doi:10.1139/f02-098
- Brooks SJ, Platvoet D, Mills CL (2008) Cation regulation and alteration of water permeability in the amphipod *Dikerogammarus villosus*: an indicator of invasion potential. *Fundamental and Applied Limnology* 172: 183-189, doi:10.1127/1863-9135/2008/0172-0183
- Bruijs MCM, Kelleher B, Van der Velde G, Bij de Vaate A (2001) Oxygen consumption, temperature and salinity tolerance of the invasive amphipod *Dikerogammarus villosus*: indicator of further dispersal via ballast water transport. *Archiv für Hydrobiologie* 152: 633-646
- Casellato S, La Piana G, Latella L, Ruffo S (2006) *Dikerogammarus villosus* (Sowinsky, 1894) (Crustacea, Amphipoda, Gammaridae) for the first time in Italy. *Italian Journal of Zoology* 73: 97-104, doi:10.1080/1125000500502293
- Casellato S, Visentin A, La Piana G (2007) The predatory impact of *Dikerogammarus villosus* on fish. In: Gherardi F (ed), Biological invaders in inland waters: profiles, distribution and threats. Springer, Berlin, pp 495-507, doi:10.1007/978-1-4020-6029-8\_27



- Casellato S, Masiero L, La Piana G, Gigliotti F (2008) The alien amphipod crustacean *Dikerogammarus villosus* in Lake Garda (N-Italy): the invasion continues. In: Rabitsch W, Essl F, Klingenstein F (eds), Biological Invasions – from Ecology to Conservation. NEOBIOTA 7: 115-122
- Devin S, Piscart C, Beisel JN, Moreteau JC (2003) Ecological traits of the amphipod invader *Dikerogammarus villosus* on a mesohabitat scale. *Archiv für Hydrobiologie* 158: 43-56, doi:10.1127/0003-9136/2003/0158-0043
- Devin S, Bollache L, Beisel JN, Moreteau JC, Perrot-Minnot MJ (2004a) Pigmentation polymorphism in the invasive amphipod *Dikerogammarus villosus*: some insights into its maintenance. *Journal of Zoology London* 264: 391-397, doi:10.1017/S0952836904005862
- Devin S, Piscart C, Beisel JN, Moreteau JC (2004b) Life History Traits of the Invader *Dikerogammarus villosus* (Crustacea: Amphipoda) in the Moselle River, France. *International Review of Hydrobiology* 89: 21-34, doi:10.1002/iroh.200310667
- Dick JTA, Platvoet D (2000) Invading predatory crustacean *Dikerogammarus villosus* eliminates both native and exotic species. *Proceedings of the Royal Society London B* 267: 977-983, doi:10.1098/rspb.2000.1099
- Dick JTA, Platvoet D, Kelly DW (2002) Predatory impact of the freshwater invader *Dikerogammarus villosus* (Crustacea: Amphipoda). *Canadian Journal of Fisheries and Aquatic Science* 59: 1078-1084, doi:10.1139/f02-074
- Gonzales MJ, Downing A (1999) Mechanism underlying amphipod responses to zebra mussel (*Dreissena polymorpha*) invasion and implications for fish–amphipod interactions. *Canadian Journal of Fisheries and Aquatic Sciences* 56: 679-685, doi:10.1139/cjfas-56-4-679
- Grabowski M, Bacela K, Konopacka A (2007a) How to be an invasive gammarid (Amphipoda: Gammaroidea) – comparison of life history traits. *Hydrobiologia* 590: 75-84, doi:10.1007/s10750-007-0759-6
- Grabowski M, Jazdzewski K, Konopacka A (2007b) Alien Crustacea in Polish waters – Amphipoda. *Aquatic Invasions* 2: 25-38, doi:10.3391/ai.2007.2.1.3
- Haas G, Brunke M, Streit B (2002) Fast turnover in dominance of exotic species in the Rhine River determines biodiversity and ecosystem function: an affair between amphipods and mussels. In: Leppäkoski E, Gollasch S, Olenin S (eds), Invasive aquatic species of Europe: distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 426-432
- Jazdzewski K, Konopacka A (2002) Invasive Ponto-Caspian species in waters of the Vistula and Oder basins and southern Baltic Sea. In: Leppäkoski E, Gollasch S, Olenin S (eds), Invasive aquatic species of Europe: Distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 384-398
- Kinzler W, Maier G (2003) Asymmetry in mutual predation: Possible reason for the replacement of native gammarids by invasives. *Archiv für Hydrobiologie* 157: 473-481, doi:10.1127/0003-9136/2003/0157-0473
- Kley A, Maier G (2003) Life history characteristics of the invasive freshwater gammarids *Dikerogammarus villosus* and *Echinogammarus ischnus* in the river Main and in the Main–Donau canal. *Archiv für Hydrobiologie* 156: 457-470, doi:10.1127/0003-9136/2003/0156-0457
- Leuven RSEW, Van der Velde G, Baijens I, Snijders J, van der Zwart C, Lenders HJR, Bij de Vaate A (2009) The river Rhine: a global highway for dispersal of aquatic invasive species. *Biological Invasions* 11: 1989-2008, doi:10.1007/s10530-009-9491-7
- Lori E, Cianfanelli S (2006) New records of *Dreissena polymorpha* (Pallas, 1771) (Mollusca: Bivalvia: Dreissenidae) from Central Italy. *Aquatic Invasions* 1: 281-283, doi:10.3391/ai.2006.1.4.11
- Platvoet D, Van der Velde G, Dick JTA, Shuqiang, L (2009) Flexible omnivory in *Dikerogammarus villosus* (Sowinsky, 1894) (Amphipoda). Amphipod pilot species project (AMPIS) report 5. *Crustaceana* 82: 703-720, doi:10.1163/156854009X423201
- Pöckl M (2007) Strategies of a successful new invader in European fresh waters: fecundity and reproductive potential of the Ponto-Caspian amphipod *Dikerogammarus villosus* in the Austrian Danube, compared with the indigenous *Gammarus fossarum* and *G. roeseli*. *Freshwater Biology* 52: 50-63, doi:10.1111/j.1365-2427.2006.01671.x
- Pöckl M (2009) Success of the invasive Ponto-Caspian amphipod *Dikerogammarus villosus* by life history traits and reproductive capacity. *Biological Invasions* 11: 2021-2041, doi:10.1007/s10530-009-9485-5
- Quaglia F, Lattuada L, Manteca P, Bacchetta R (2008) Zebra mussels in Italy: where do they come from? *Biological Invasions* 10: 555-560, doi:10.1007/s10530-007-9152-7
- Ricciardi A, Whoriskey FG, Rasmussen JB (1997) The role of the zebra mussel (*Dreissena polymorpha*) in structuring macroinvertebrate community on hard substrata. *Canadian Journal of Fisheries and Aquatic Sciences* 54: 2596-2608, doi:10.1139/cjfas-54-11-2596
- Ruffo S, Stoch F (eds) (2005) Checklist e distribuzione della fauna italiana. Memorie del Museo Civico di Storia Naturale di Verona, 307 pp
- Simberloff D, Von Holle B (1999) Positive interactions of non indigenous species: Invasional meltdown? *Biological Invasions* 1: 21-32, doi:10.1023/A:1010086329619
- Van den Brink FWB, Van der Velde G (1991) Amphipod invasion on the Rhine. *Nature* 352: 576, doi:10.1038/352576a0
- Van der Velde G, Nagelkerken I, Rajagopal S (2002) Invasions by alien species in inland freshwater bodies in Western Europe: the Rhine Delta. In: Leppäkoski E, Gollasch S, Olenin S (eds), Invasive species of Europe: distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 360-372
- Van der Velde G, Leuven RSEW, Platvoet D, Barcelá K, Huijbregts MAJ, Hendriks HWM, Kruijt D (2009) Environmental and morphological factors influencing predatory behaviour in invasive non-indigenous gammaridean species. *Biological Invasions* 11: 2043-2054, doi:10.1007/s10530-009-9500-x
- Van Riel MC, Van der Velde G, Rajagopal S, Marguiller S, Dehairs F, Bij de Vaate A (2006) Trophic relationships in the Rhine food web during invasion and after establishment of the Ponto-Caspian invader *Dikerogammarus villosus*. *Hydrobiologia* 565: 39-58, doi:10.1007/s10750-005-1904-8
- Van Riel MC, Healy EP, Van der Velde G, Bij de Vaate A (2007) Interference competition among native and invader amphipods. *Acta Oecologica* 31: 282-289, doi:10.1016/j.actao.2006.12.006