



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Intraoperative positive fluid balance improves tissue diffusion of ceftizoxime.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Intraoperative positive fluid balance improves tissue diffusion of ceftizoxime / Di Filippo A;Cammelli R;Novelli A;Mazzei T;Tonelli F;Fallani S;Cassetta MI;Messori D;De Gaudio AR. - In: CHEMOTHERAPY. - ISSN 0009-3157. - STAMPA. - 51:(2005), pp. 51-56. [10.1159/000085610]

Availability:

This version is available at: 2158/772066 since: 2016-09-23T17:46:34Z

Published version:

DOI: 10.1159/000085610

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Intraoperative Positive Fluid Balance Improves Tissue Diffusion of Ceftizoxime

A. Di Filippo^a R. Cammelli^a A. Novelli^b T. Mazzei^b F. Tonelli^c S. Fallani^b
M.I. Cassetta^b D. Messeri^d A.R. De Gaudio^a

^aUniversity of Florence, Department of Critical Care, Section of Anesthesiology and Intensive Care,

^bDepartment of Clinical and Preclinical Pharmacology, ^cDepartment of Clinical Physiopathology, Unit of Surgery, ^d3rd Local Health Unit, Pistoia, Infectious Diseases Unit, Florence, Italy

Key Words

Antibiotic · Ceftizoxime · Surgical infection prophylaxis · Intraoperative fluid balance · Pharmacokinetics

Abstract

Aim of Study: To demonstrate that administration of fluids and the consequent improvement of fluid balance during a surgical procedure can modify the tissue diffusion of ceftizoxime. **Methods:** Twenty-eight patients (30–79 years) undergoing major abdominal surgery of the colon were administered ceftizoxime 30 mg/kg i.v. at induction of anesthesia. A sample of arterial blood was taken before administration of the drug (t_0) and then again at the time of vascular occlusion of the colon segment to be removed (t_1). A sample of the segment of removed colon was taken. The patients were divided into two groups on the basis of the fluid balance between t_0 and t_1 : group A ($n = 17$) with a fluid balance $<1,000$ ml and group B ($n = 11$) with a fluid balance $>1,000$ ml. The parameters evaluated in each group were: weight, height and age of the patients, serum and tissue antibiotic concentration, percent ratio of serum and tissue concentration, time elapsed between t_0 and t_1 , volume of administered fluids between t_0 and t_1 , diuresis and hourly diuresis between t_0 and t_1 and body fluid distribution, obtained using a bioelectrical impedance analyzer. The

mean results obtained in the two groups were then compared using Student's *t* test. **Results:** The balance of fluids calculated up to t_1 was 675 ± 308 ml for group A and $1,411 \pm 405$ ml for group B ($p < 0.01$). The means of the recorded values that showed statistically significant differences were: mean percent concentration ratio (43.6 ± 8.4 vs. $84 \pm 16\%$; $p < 0.05$), concentration in the colonic segment (16.3 ± 7.9 vs. 37.2 ± 25.9 mg/ml; $p < 0.05$), urinary volume gathered up to t_1 (538 ± 557 vs. 169 ± 104 ml; $p < 0.05$), hourly urinary volume up to t_1 (311.1 ± 296 vs. 97.6 ± 77.9 ml/h; $p < 0.05$), percent variation of resistance (95.1 ± 5.1 vs. 89.7 ± 8.6 ; $p < 0.05$). The other means did not show any significant statistical differences. **Conclusions:** A higher tissue water level seems to facilitate the penetration of the antibiotic into the tissue according to the pharmacokinetic characteristics of ceftizoxime: high amount of free drug (not bound to plasma proteins) and high hydrosolubility.

Copyright © 2005 S. Karger AG, Basel

Introduction

The effectiveness of perioperative surgical infection prophylaxis depends on the following pharmacological aspects of the antimicrobial agent used: (a) the optimal time for antibiotic administration which is at induction of anesthesia [1]; (b) the highest concentration that must

be obtained in the tissues where bacteria grow [2]; (c) the antibiotic serum concentration that must be higher than the minimal inhibitory concentration of the possible contaminating bacteria during an intervention [3].

Intraoperative fluid therapy could interfere with these aspects by modifying body fluid compartment distribution. In particular, fluid balance, hemodilution, and interstitial fluid content [4] may influence the tissue diffusion of antimicrobial agents.

The aim of this study was to evaluate whether an intraoperative positive fluid balance can modify the tissue diffusion of a cephalosporin (ceftizoxime) with a high amount of free drug (not bound to plasma proteins) and high hydrosolubility, administered for short-term surgical infection prophylaxis.

Patients and Methods

Thirty-five consecutive ASA I-II patients undergoing major colon surgery under general anesthesia, were admitted to the study. All the patients were informed about the goals of the study and gave their signed, written consent. All the patients were submitted to the same bowel preparation and were weighed immediately before the start of the intervention. The surgical procedures were performed by the same surgeon and via the same cutaneous xiphopubic incision with removal of a segment of colonic tissue. Seven patients requiring blood transfusions, blood derivatives, vasoactive drugs administered for hemodynamic instability or diuretics up to resection of the colon segment were excluded from the study.

The anesthetic technique was the following: premedication with morphine (10 mg i.m.) followed by the induction of anesthesia with Fentanyl (1.5 µg/kg), Propofol (2 mg/kg) and Atracurium (0.6 mg/kg), Isoflurane MAC95 in O₂ and air, and continuous intravenous administration of Atracurium (0.4 mg/kg/h until reawakening).

As the administration of different volumes of fluids can modify the clearance of the antibiotic, the protocols of fluid infusion were the same during the period of the study; crystalloid solutions (Ringer acetate) were given to all the patients at the dosage of 15 ml/kg/h during the first hour and 12 ml/kg/h up to the end.

Ceftizoxime 30 mg/kg was administered over 10 min through a peripheral vein at induction of anesthesia. A 5-ml arterial blood sample was taken immediately before the induction of anesthesia (t_0) to obtain the baseline standard serum antibiotic concentration and the hematocrit values. A second 5-ml arterial blood sample was taken at the time of vascular occlusion of the colonic segment to be removed (t_1) to measure the serum antibiotic concentration and hematocrit.

A fragment of the removed segment of approximately 5 × 1 cm was washed with saline, dried with sterile gauze and conserved in a test tube at -20°C. The tissues were weighed, diluted 1:1 (wt/vol) in sterile normal saline (pH 6.3), homogenized with a Polytron PT 10-35 homogenizer (Kinematica, Lucerne, Switzerland), and centrifuged at low speed; the supernatant was used for the assay [5].

The concentration of the antibiotic in serum and tissues was determined in triplicate by a validated large-plate agar diffusion

technique, according to Good Laboratory Practice (GLP) standards [6, 7]. Ceftizoxime concentrations were determined using the Antibiotic Medium II (BBL) as the culture medium and *Escherichia coli* Sc 12,355 as the test organism, with a lower limit of sensitivity of 0.125 mg/l. Standard concentrations were prepared daily in pooled serum for blood samples and in normal saline for tissue specimens. The test organism was added by the surface layer technique. After homogeneous distribution of the culture, the excess liquid was removed with a pipette. The plates were incubated at 37°C in air overnight.

Best-fit standard curves were obtained by linear regression analysis. The linearity was $\log y = 0.102x - 1.8$ for plasma samples and $\log y = 0.103x - 1.95$ for tissues; the correlation coefficient was not less than 0.99. Intra-assay precision ranged from 4.5 to 9.8% for serum samples and from 1.3 to 7.4% for tissue samples. Inter-assay precision at a level of 1 mg/l ranged from 1.83 to 4.82% for serum and from 1.4 to 5.8% for tissues.

The ratio of serum antibiotic concentration at t_1 to tissue antibiotic concentration was calculated as percent ratio. The fluid balance between t_0 and t_1 was obtained by calculating the difference between the volume of the administered fluids and the intraoperative losses due to fasting (2 ml/kg/h), perspiratio insensibilis (8 ml/kg/h of surgery) and diuresis. A cutoff value of intraoperative fluid balance was chosen (1,000 ml) to divide the patients into two groups on the basis of the intraoperative fluid balance reported in other works [8, 9] and the time between the induction of anesthesia and the time of vascular exclusion of the colonic segment to be removed previously measured (about 90 min): group A ($n = 17$) with in a fluid balance between t_0 and $t_1 < 1,000$ ml and group B ($n = 11$) with in a fluid balance $> 1,000$ ml. The mean weight, height and age were calculated in the two groups. To evaluate the effect of fluid balance on tissue diffusion of the antibiotic, the following parameters were registered or calculated: tissue concentration of the antibiotic (mg/ml); ratio of tissue to plasma concentration (%); diuresis between t_0 and t_1 (ml); hourly diuresis between t_0 and t_1 (ml/h), hematocrit between t_0 and t_1 (expressed as percentage), and time elapsed between t_0 and t_1 (min).

To evaluate the body compartment distribution of administered fluids, an impedance analyzer (BIA/STATM Akern S.r.l. Florence, Italy) [10-12] was used to measure the variation of resistance between t_0 and t_1 (%), which expresses conductor opposition to the flow of alternate currents, and is inversely proportional to total body fluids, and the variation in capacitive reactance between t_0 and t_1 (%), which expresses condenser opposition to the flow of alternating current, and is directly proportional to cellular mass.

Statistical analysis was performed using Student's *t* test. A *p* value < 0.05 was considered significant.

Results

Group A ($n = 17$) and group B ($n = 11$) were not statistically different in terms of age (respectively: 57.7 ± 12 vs. 54.7 ± 14 years), weight (68.9 ± 8.4 vs. 72.9 ± 9.1 kg) and height (165 ± 9.9 vs. 168.2 ± 6.6 cm).

Group A patients had a mean fluid balance of 675 ± 308 ml, while group B had a higher mean fluid balance:




Fig. 1. Linear correlation between tissue concentration of ceftizoxime and intraoperative fluid balance in a series of patients submitted to major colon surgery.

Table 1. Mean (\pm standard deviation) of the values measured between t_0 (start of intervention) and t_1 (vascular exclusion of the colonic segment to be removed) in two groups of patients: group A: fluid balance $<1,000$ ml; group B: fluid balance $>1,000$ ml

	Group A (n = 17)	Group B (n = 11)	p
Fluid balance, ml	675 \pm 308	1,411 \pm 405	<0.05
Tissue /serum concentration, %	33.6 \pm 8.4	93.9 \pm 5.9	<0.05
Tissue concentration in the colonic portion, mg/ml	16.3 \pm 7.9	37.2 \pm 25.9	<0.05
Serum concentration, mg/ml	49.6 \pm 24	40.1 \pm 26.6	NS
Diuresis, ml	538 \pm 557	169.3 \pm 104	<0.05
Hourly diuresis, ml/h	311.1 \pm 296	97.6 \pm 77.9	<0.05
Hematocrit variation, %	83 \pm 7.3	81.3 \pm 10.4	NS
Resistance variation, %	95.1 \pm 5.1	89.7 \pm 8.6	<0.05
Capacitive reactance variation, %	108.8 \pm 21	106.9 \pm 27	NS
Time elapsed between t_0 and t_1 , min	79 \pm 34	92 \pm 22	NS

Statistics: Student's t test; $p < 0.05$ was considered significant.

1,411 \pm 405 ml ($p < 0.05$). Mean tissue concentration (16.3 \pm 7.9 of group A vs. 37.2 \pm 25.9 of group B, $p < 0.05$), mean ratio of tissue to serum concentration (43.6 \pm 28.4% of group A vs. 84 \pm 16% of group B; $p < 0.05$), mean diuresis (538 \pm 557 ml of group A vs. 169 \pm 104 ml of group B, $p < 0.05$), mean hourly diuresis (311.1 \pm 296 ml/h of group A vs. 97.6 \pm 77.9 of group B, $p < 0.05$)

and mean percent variation of the resistance (95.1 \pm 5.1 of group A vs 89.7 \pm 8.6 of group B, $p < 0.05$) were statistically significantly different in the two groups ($p < 0.05$). The results are reported in table 1. The linear correlation between tissue concentration versus intraoperative fluid balance is shown in figure 1.

Discussion and Conclusion

The ideal characteristics of short-term antimicrobial prophylaxis include: use of antibiotic with a narrow spectrum of activity that includes the main pathogens responsible for postoperative infections, bactericidal effect, high therapeutic index, high tolerability, favorable kinetic parameters such as intravenous administration, long half-life of elimination and good tissue penetration [1].

The tissue diffusion of the antibiotic mainly depends on the gradient existing between plasma and tissue concentrations [13]. The gradient can be reduced if the antibiotic has a high percentage of protein binding: only the drug not bound to protein diffuses into tissue.

Anesthesia and surgical trauma can interact with tissue diffusion of drugs: hemodilution and replacement of intraoperative blood loss can determine homeostatic modifications in body fluid: general anesthesia and vasoactive drugs can modify the vascular system [14–16]. In addition, the depression of sympathetic tone observed at induction of anesthesia can cause previously adequate refilling pressure to become insufficient [17, 18]. Therefore, it may be necessary to expand the circulating volume before or immediately after the induction of anesthesia and during the intervention [19, 20].

Finally, the vascular splanchnic capacity can suddenly increase when intra-abdominal pressure is rapidly reduced by a laparotomy [21]. These physiological modifications, induced by anesthesia and surgery, require administration of fluid and a fluid balance between the administered fluids and the intraoperative losses must be achieved. The imperceptible losses include the amount of fluid that must be reinfused as a basal requirement (2 ml/kg/h) to maintain equilibrium and that lost due to surgery, which depends on the type of surgery (minor: 4 ml/kg/h, intermediate: 6 ml/kg/h, major: 8 ml/kg/h) [22]. The fluid balance must be calculated during surgery because fluid overload may be responsible for the onset of interstitial edema that can lead to hypoxemia at the pulmonary level and cause prolonged postoperative intensive care in patients with reduced cardiopulmonary capacity [23]. Instead, hypovolemia may lead to oliguria which, in addition to the reduction in glomerular filtration caused by the anesthetics, reduces cardiac output (and thus oxygen transport), decreasing the ability to tolerate and adapt to significant hematocrit and hemoglobin reduction [24].

The aim, therefore, is to re-establish the volume, which must be empirically evaluated by measuring the systemic arterial pressure, maintaining a value that can assure blood

distribution to all the organs in a way that allows an adequate supply (DO_2) and consumption (VO_2) of oxygen.

The effects of hemodilution on the administered drug concentration during anesthesia and on the diffusion of these drugs in the interstitium are not well understood. Antibiotic tissue distribution during the intraoperative period and its relationship with intraoperative fluid balance has not been studied enough.

Ceftizoxime is a third-generation cephalosporin that, in recent years, has been widely used in the prophylaxis of postoperative infection for its particular efficacy against Enterobacteriaceae involved in the infections occurring in 'clean-contaminated' surgery. It has, therefore, become an alternative prophylaxis in abdominal surgery. After a bolus administration, the pharmacokinetics of ceftizoxime are well suited for prophylaxis. The concentrations reached after intravenous bolus administration of 1 g are 107–136 mg/l [25]. The volume of distribution studied in healthy volunteers is 15–28 liters [26]; the elimination half-life is 1.1–2.2 h [27]; serum clearance is 110–200 ml/min and renal clearance is 100–160 ml/min, just high enough to enable 70% of the drug to reach the urine within 2 h of administration [27]. Ceftizoxime does not pass the blood-brain barrier, and concentrates more in peripheral tissue [26]. The peak tissue concentrations are obtained 1–3 h after administration [27]. Ceftizoxime has a mean protein binding of 30% [28].

Ceftizoxime is mostly eliminated via glomerular filtration with a small component of unmodified elimination in urine through a tubular secretion mechanism [27]; its elimination is directly related to renal function and the dosage must be reduced in proportion to the decrease in creatinine clearance [29].

In clinical trials, prophylactic ceftizoxime has been shown to be able to reduce the incidence of postoperative infections just as cefazolin and cefamandole [30].

The pharmacokinetic characteristics of ceftizoxime (high water solubility, low protein binding, high renal clearance) could make its serum and tissue concentration curve particularly susceptible to fluid modifications. In fact, in a clinical trial on 53 patients undergoing emergency explorative laparotomy, Rosemurgy et al. [31] found that significant blood losses and subsequent abundant fluid replacement caused serious reductions in antibiotic serum concentrations that are significantly related to the appearance of postoperative infection.

In our study, we attempted to demonstrate that the tissue concentrations of ceftizoxime are increased by a high fluid balance.

To evaluate the body compartment distribution of administered fluids, an impedance analyzer was used to measure resistance, which expresses conductor opposition to the flow of alternating current, and is inversely proportional to total body fluids, and the capacitive reactance, which expresses condenser opposition to the flow of alternating current, and is directly proportional to cellular mass. These measurements were useful for assessing nutritional status [10] and dialysis-related modifications of body fluid distribution [11, 12].

The results of this paper demonstrate that in patients with higher fluid balance, in whom we observed reduced resistance to fluid accumulation, antibiotic diffusion and tissue concentration were higher.

On the basis of these results we hypothesize that if diuresis is reduced and the fluid balance increases during an operation, the total body fluid increases and a high fluid content at the tissue level allows the passage of a higher amount of drug in the tissues.

The differences observed in diuresis could be related, among others, to a relative preoperative dehydration (but the bowel preparation was similar in all the patients) as well as to the effect of mechanical ventilation, or intraoperative minimal variation of kidney perfusion.

The passage of a higher amount of drug in the tissues is in agreement with the pharmacokinetic characteristics

of ceftizoxime described above: i.e. highly free (not bound to plasma protein) and hydrosoluble drug [30].

Bioelectrical impedance is essential for the evaluation of the effects of fluid balance and provided significant supporting data. In a recent study, bioimpedance analysis variations demonstrated a strict correlation with fluid losses during hemodialysis. In that case, the bioimpedance analysis variation of 20% was equivalent to a body weight decreased by 2.8 ± 0.8 kg [32]. This value is comparable with the variation (7% for 1 liter of fluid balance) found in our study.

In our series of patients, the time elapsed from the administration of drug is not significantly related to the tissue concentration of ceftizoxime. This lacking correlation could mean that during a surgical intervention the tissue distribution of ceftizoxime is modified by fluid distribution more than by peak drug concentration.

Further study seems to be necessary to achieve a better understanding of the relationship between fluid administration and tissue penetration of antibiotic agents. Knowledge of the mechanisms of tissue diffusion of antimicrobial agents could help explain the infrequent but not rare failures of surgical infection prophylaxis, particularly in colon surgery [33]. For these reasons as well, the controversy over the choice of antibiotic for surgical prophylaxis is still open [34, 35].

References

- Classen DC, Scott Evans R, Pestotnik SL, Horn SD, Menlove RL, Burke JP: The timing of prophylactic administration of antibiotics and the risk of surgical wound infection. *N Engl J Med* 1992;326:281–285.
- Redington J, Ebert SC, Craig WA: Role of antimicrobial pharmacokinetics and pharmacodynamics in surgical prophylaxis. *Rev Infect Dis* 1991;13(suppl 10):790s–799s.
- Novelli A: Antimicrobial prophylaxis in surgery: The role of pharmacokinetics. *J Chemother* 1999;11:565–572.
- Holte K, Sharrock NE, Kehlet H: Pathophysiology and clinical implications of perioperative fluid excess. *Br J Anaesth* 2002;89:622–632.
- De Lalla F, Novelli A, Pellizzer G, Molocchi F, Viola R, Rigon A, Stecca C, Dal Pizzol V, Fallani S, Periti P: Regional and systemic prophylaxis with teicoplanin in monolateral and bilateral total knee replacement procedures: Study of pharmacokinetics and tissue penetration. *Antimicrobial Agents Chemother* 1993;37:2693–2697.
- Mazzei T, Tonelli F, Novelli A, et al: Penetration of cefotetan into suction skin blister fluid and tissue homogenates in patients undergoing abdominal surgery. *Antimicrob Agents Chemother* 1994;38:2221–2223.
- Periti P, Rizzo M, Novelli A, Reali EF, Dami A, Boni S, Mini E, Mazzei T: Pharmacokinetics and penetration into extravascular fluid of imipenem in patients with normal renal function; in Berkada B, Kuemmerle HP (eds): *Progress in Chemotherapy, Antimicrobial Section*. Munich, Ecomed, 1987, vol 1, pp 990–992.
- Moretti EW, Robertson KM, El-Moalem H, Gan TJ: Intraoperative colloid administration reduces postoperative nausea and vomiting and improves postoperative outcomes compared with crystalloid administration. *Anesth Analg* 2003;96:611–617.
- Gan TJ, Soppitt A, Maroof M, el-Moalem H, Robertson KM, Moretti E, Dwane P, Glass PS: Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 2002;97:820–826.
- Talluri T, Lietdke RJ, Evangelisti A, Talluri J, Maggia G: Fat-free mass qualitative assessment with bioelectric impedance analysis (BIA). *Ann N Y Acad Sci* 1999;873:94–98.
- Talluri A, Maggia G: Bioimpedance analysis (BIA) in hemodialysis: Technical aspects. *Int J Artif Organs* 1995;18:687–692.
- Scanferla F, Landini S, Fracasso A, Morachiello P, Righetto F, Toffoletto PP, Bazzato G: On-line bioelectric impedance during haemodialysis: Monitoring of body fluids and cell membrane status. *Nephrol Dial Transplant* 1990;5(suppl 1):167–170.
- Craig W: Pharmacodynamics of antimicrobial agents as a basis for determining dosage regimens. *Eur J Clin Microbiol Infect Dis* 1993;12(suppl 1):6s–8s.
- Novelli GP: Effects of enflurane and halothane on the microcirculation. *Acta Anaesthesiol Scand Suppl* 1979;71:64–68.
- Coates DP, Monk CR, Prys-Roberts C, Turtle M: Hemodynamic effects of infusions of the emulsion formulation of propofol during nitrous oxide anesthesia in humans. *Anesth Analg* 1987;66:64–70.

- 16 Banic A, Krejci V, Erni D, Wheatley AM, Sigurdsson GH: Effects of sodium nitroprusside and phenylephrine on blood flow in free musculocutaneous flaps during general anesthesia. *Anesthesiology* 1999;90:147–155.
- 17 Tibballs J, Malbezin S: Cardiovascular responses to induction of anaesthesia with thiopentone and suxamethonium in infants and children. *Anaesth Intensive Care* 1988;16:278–284.
- 18 Ismail EF, Kim SJ, Salem MR, Crystal GJ: Direct effects of propofol on myocardial contractility in in situ canine hearts. *Anesthesiology* 1992;77:964–972.
- 19 Matsukawa T, Sessler DI, Sessler AM, Schroeder M, Ozaki M, Kurz A, Cheng C: Heat flow and distribution during induction of general anesthesia. *Anesthesiology* 1995;82:662–673.
- 20 Matsukawa T, Kurz A, Sessler DI, Bjorksten AR, Merrifield B, Cheng C: Propofol linearly reduces the vasoconstriction and shivering thresholds. *Anesthesiology* 1995;82:1169–1180.
- 21 Eisele R, Dissmann W, Kotter D, Nasser M, Thimme W: Behavior of the portal vein system in man after laparotomy. *Surg Gynecol Obstet* 1975;141:870–874.
- 22 Kaye AD, Gorgono AW: Fluid and electrolyte physiology. *Anesthesia USA* 2000;45:1587–1644.
- 23 Gow KW, Phang PT, Tebbutt-Speirs SM, English JC, Allard MF, Goddard CM, Walley KR: Effect of crystalloid administration on oxygen extraction in endotoxemic pigs. *J Appl Physiol* 1998;85:1667–1675.
- 24 Van Bommel J, Siegemund M, Henny CP, Trouwborst A, Ince C: Critical hematocrit in intestinal tissue oxygenation during severe normovolemic hemodilution. *Anesthesiology* 2001;94:152–160.
- 25 Fasching CE, Peterson LR, Bettin KM, Gerding DN: High-pressure liquid chromatographic assay of ceftizoxime with an anion-exchange extraction technique. *Antimicrobial Agents Chemother* 1982;22:336–337.
- 26 Cable D, Edralin G, Overturf GD: Human cerebrospinal fluid pharmacokinetics and treatment of bacterial meningitis with ceftizoxime. *J Antimicrobial Chemother* 1982;10(suppl C):121–127.
- 27 Nakayama I, Akieda Y, Tajima K, Kawamura H, Kawaguchi H, Nishimoto A, Ishyama S: Antimicrobial activity, absorption, excretion, metabolism, distribution in tissues and the clinical use in surgical infection of a new cephalosporin antibiotic, ceftizoxime. *Chemotherapy (Tokyo)* 1980;28(suppl 5):495–510.
- 28 Murakawa T, Sakamoto H, Fukada S, Nakamoto S, Hirose T, Itoh N, Nishida M: Pharmacokinetics of ceftizoxime in animals after parenteral dosing. *Antimicrobial Agents Chemother* 1982;21:925–934.
- 29 Cutler RE, Blair AD, Burgess ED, Parks D: Pharmacokinetics of ceftizoxime. *J Antimicrobial Chem* 1982;10(suppl C):91–97.
- 30 Fainstein V, Weaver S, Bodey GP: In vitro susceptibilities of *Aeromonas hydrophila* against new antibiotics. *Antimicrobial Agents Chemother* 1982;22:513–514.
- 31 Rosemurgy AS II, Dillon KR, Kurto HA, Albrink MH: Ceftizoxime use in trauma celiotomy: Pharmacokinetics and patient outcomes. *J Clin Pharmacol* 1995;35:1046–1051.
- 32 Di Iorio BR, Scalfi L, Terracciano V, Bellizzi V: A systematic evaluation of bioelectrical impedance measurement after hemodialysis session. *Kidney Int* 2004;65:2435–2440.
- 33 Rau HG, Mittelkotter U, Zimmermann A, Lachmann A, Kohler L, Kullmann KH: Perioperative infection prophylaxis and risk factor impact in colon surgery. *Chemotherapy* 2000;46:353–363.
- 34 Dietrich ES, Bieser U, Frank U, Schwarzer G, Daschner FD: Ceftriaxone versus other cephalosporins for perioperative antibiotic prophylaxis: A meta-analysis of 43 randomized controlled trials. *Chemotherapy* 2002;48:49–56.
- 35 Bratzler DW, Houck PM, for the Surgical Infection Prevention Guidelines Writers Workgroup: Antimicrobial Prophylaxis for Surgery: An Advisory Statement from the National Surgical Infection Prevention Project. *Clin Infect Dis* 2004;38:1706–1715.