



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Pharmacological in vivo test to evaluate the bioavailability of some St. John's wort innovative oral preparations**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Pharmacological in vivo test to evaluate the bioavailability  
of some St. John's wort innovative oral preparations / A R Bilia; M C Bergonzi; B Isacchi; N Galeotti; C  
Ghelardini; F F Vincieri. - In: PLANTA MEDICA. - ISSN 0032-0943. - STAMPA. - 72:(2006), pp. 241-241.  
[10.1055/s-2006-950041]

*Availability:*

This version is available at: 2158/774259 since: 2017-10-31T11:15:24Z

*Published version:*

DOI: 10.1055/s-2006-950041

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto  
stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze  
(<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

## Pharmacological *in vivo* test to evaluate the bioavailability of some St. John's wort innovative oral preparations

AR Bilia <sup>1</sup>, MC Bergonzi <sup>1</sup>, B Isacchi <sup>1</sup>, N Galeotti <sup>2</sup>, C Ghelardini <sup>2</sup>, FF Vincieri <sup>1</sup>

- <sup>1</sup>Department of Pharmaceutical Sciences, University of Florence, 50019 Sesto Fiorentino, Florence, Italy
- <sup>2</sup>Department of Preclinical & Clinical Pharmacology, University of Florence, 50139 Florence, Italy

- [Congress Abstract](#)

---

Preparations based on extracts of St. John's wort are widely marketed for treating mild to moderately severe depressive disorders and other health conditions such as anxiety and sleep disorders [1]. Active principles are not yet discovered and flavonols, based on quercetin aglycone, naphthodianthrones (hypericin and pseudohypericin) and phloroglucinols such as hyperforin, adhyperforin seems to be related to this action. Thus, flavonols and naphthodianthrones are polyphenols, quite polar derivatives but their water solubility is very scarce; phloroglucinols are lipophilic and completely not water-soluble constituents. In addition, hypericins and hyperforins are not stable with regard to heat and light [2].

In this study the optimisation of technological and pharmaceutical aspects of dried commercial extract of St. John's wort were evaluated by the *in vivo* "Porsolt test". Solid dosage forms containing  $\beta$ -cyclodextrin and micellar systems (SDS, ASC-8) were compared in the "Porsolt test" with the extract alone. The extract showed the antidepressant activity in the mice after 60 minutes and with the dosage of 100mg/kg. The same antidepressant activity appeared in 30min with a micellar solution of SDS 40mM containing the same quantity of extract (100mg/kg), while with micelles of ASC-8 40 mM the effect appeared at 15min and with a dosage of 30mg/kg. In the case of colyophilized with  $\beta$ -cyclodextrin the best results were obtained at 30min, administering 60mg/kg of the extract.

**Acknowledgements:** The financial support of MIUR (PRIN 2004) and Ente Cassa di Risparmio di Firenze is gratefully acknowledged for financial support.

**References:** 1. Chatterjee, S.S. *et al.* (1998), *Pharmacopsychiatry* 31: 7-15. 2 Bilia A.R. *et al.* (2001), *Int. J. Pharm.* 213: 199-208.