



FLORE

Repository istituzionale dell'Università degli Studi di Firenze

When the rule becomes the exception. No evidence of gene flow between two Zerynthia cryptic butterflies suggests the emergence of

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

When the rule becomes the exception. No evidence of gene flow between two Zerynthia cryptic butterflies suggests the emergence of a new model group / Zinetti F; Dapporto L; Vovlas A; Chelazzi G; Bonelli S; Balletto E; Ciofi C. - In: PLOS ONE. - ISSN 1932-6203. - ELETTRONICO. - 8:(2013), pp. e65746-1-e65746-11. [10.1371/journal.pone.0065746]

Availability:

This version is available at: 2158/806085 since: 2019-08-13T18:57:46Z

Published version: DOI: 10.1371/journal.pone.0065746

Terms of use: Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf)

Publisher copyright claim:

(Article begins on next page)

When the Rule Becomes the Exception. No Evidence of Gene Flow between Two *Zerynthia* Cryptic Butterflies Suggests the Emergence of a New Model Group

Francesca Zinetti^{1,9}, Leonardo Dapporto²*⁹, Alessio Vovlas³, Guido Chelazzi¹, Simona Bonelli³, Emilio Balletto³, Claudio Ciofi¹

1 Dipartimento di Biologia, Università degli Studi di Firenze, Florence, Italy, 2 Istituto Comprensivo Materna Elementare Media Convenevole da Prato, Prato, Italy, 3 Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università degli Studi di Torino, Torino, Italy

Abstract

There is increasing evidence that most parapatric cryptic/sister taxa are reproductively compatible across their areas of contact. Consequently, the biological species concept, which assumes absence of interbreeding, is becoming a not so effective criterion in evolutionary ecology. Nevertheless, the few parapatric sister taxa showing complete reproductive barriers represent interesting models to study speciation processes and the evolution of reproductive isolation. In this study, we examined contact populations in northwestern Italy of two butterfly species, Zerynthia polyxena and Z. cassandra, characterized by different genitalic morphotypes. We studied levels of divergence among 21 populations distributed from Sicily to France using three genetic markers (the mitochondrial COI and ND1 genes and the nuclear wingless gene) and genitalic geometric morphometrics. Moreover, we performed species distribution modelling to estimate different climatic requirements of Z. polyxena and Z. cassandra. We projected climatic data into glacial maximum scenarios in order to verify if and to which extent glacial cycles could have contributed to speciation processes. Genetic and morphometric analyses identified two main groups. All specimens showed a concordant pattern of diversification, including those individuals sampled in the contact area. Haplotype distribution and climatic models showed that during glacial maxima both species experienced a strong range contraction and presumably remained separated into different microrefugia in southern France, in the Italian Peninsula and on the islands of Elba and Sicily. Long term separation was probably favoured by reduced dispersal ability and high phylopatry, while genitalic diversification probably favoured interbreeding avoidance. Conversely, the aposematic wing pattern remained almost identical. We compared our results with those obtained in other species and concluded that Z. polyxena and Z. cassandra represent a valuable model in the study of speciation.

Citation: Zinetti F, Dapporto L, Vovlas A, Chelazzi G, Bonelli S, et al. (2013) When the Rule Becomes the Exception. No Evidence of Gene Flow between Two Zerynthia Cryptic Butterflies Suggests the Emergence of a New Model Group. PLoS ONE 8(6): e65746. doi:10.1371/journal.pone.0065746

Editor: Dan Mishmar, Ben-Gurion University of the Negev, Israel

Received December 10, 2012; Accepted April 26, 2013; Published June 6, 2013

Copyright: © 2013 Zinetti et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was part of the program "A multitaxa approach to study the impact of climate change on the biodiversity of Italian ecosystems" funded by the Italian Ministry of University and Research, the Spanish Ministerio de Ciencia e Innovación project CGL2010-21226/BOS and the "Definizione dello status di conservazione delle falene e della malacofauna terrestre dell'Arcipelago Toscano" project. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: leonadap@gmail.com

• These authors contributed equally to this work.

Introduction

In biparental species, gene exchange across populations is strictly linked to mating and to production of fertile offspring. For this reason, the biological concept, identifying species as reproductively isolated interbreeding groups of individuals [1] received large support by ecologists and evolutionary biologists. However, molecular studies have shown that hybridization and introgression among well established species are relatively common events, mainly in parapatric sister taxa at their contact zones (e.g. [2], [3], [4], [5], [6]). Increasing evidence for a grey-scaled species delimitation made the assumption of complete reproductive isolation weaker and new species concepts were then introduced (reviewed by [7]). Recent theories suggest that species may even differentiate in the presence of hybridization, provided that some genes linked to differential fitness or to sexual interactions are exchanged at low frequencies [7], [8], [9], [10], [11]. In butterflies, on the other hand, adaptive genes can be exchanged among species million of years after speciation events [6].

Contact zones between sister and cryptic species represent excellent models for the study of speciation, particularly when species showing complete reproductive isolation and/or highly permeable reproductive barriers can be compared [4]. European butterflies are often used as model organisms for such studies [4], [12], [13]. However, comparative research has been hampered by the high proportion of sister taxa showing weak reproductive barriers. A recent revision of the entire European butterfly fauna has shown that 16% of 440 species hybridize in nature, often producing fertile hybrids, and that parapatric sister species are almost always involved in this process [4].

There are exceptions to this pattern. The reproductively isolated cryptic sister taxa of the *Leptidea* genus, for instance, are characterized by identical appearance and relatively low genetic divergence [12], [13], [14]. Complete reproductive isolation reached in a relatively short time (between 270,000 and 120,000 years) in the absence of divergent external traits [12], [13] has been considered a "perfect crime" by Descimon & Mallet [4] and represents an excellent context to study butterfly evolution [4], [11], [13]. Based on analysis of male genitalic structures, Dapporto [15] identified two completely distinct morphotypes in the broadly recognized species of the Papilionid butterfly Zerynthia polyxena. One morphotype is segregated to the Italian Appennines (\mathcal{Z} . *cassandra*) while the other is found in remnant southern and central European populations (\mathcal{Z} . *polyxena*). These morphotypes come into close contact in north-western Italy, where they also occur in sympatry in the Beigua mountain area with no evidence of morphologically intermediate individuals [15]. Genetic data are scarce, only two specimens from the Z. cassandra range were sequenced for the mitochondrial COI gene and no information was provided on their genitalic characters [16]. However, these specimens revealed about 2% divergence with respect to other European individuals.

In this study, we assessed levels of genetic divergence and introgression at the contact zone between Z. cassandra and Z. polyxena by integrating geometric morphometric comparison of male genitalia and sequence analysis of one nuclear and two mitochondrial DNA markers. Moreover, we applied distribution modelling to assess species occurrence in different climatic settings across the current range. We then projected such differences into reconstructed climatic scenarios of the last glacial period and evaluated the extent to which distributional changes driven by climatic events may have been responsible for long term genetic isolation. Considering the clear differences in genitalic morphology, identical aposematic wing colouration and well defined contact area between Z. cassandra and Z. polyxena, new evidence on genetic variation and climatic preferences may indicate whether these species are likely to be reproductively separated and which factors shaped the observed distributional patterns. Our dataset may also suggest whether the sister taxa examined in this work represent a good model for butterfly evolutionary studies.

Materials and Methods

Geometric Morphometrics

Geometric morphometrics was conducted on 217 Zerynthia polyxena and Z cassandra males from Italy and southern France. Samples included 186 individuals from a dataset published by Dapporto [15]. Permission to collect species listed under the 92/ 43/EEC Annex IV was granted by the Italian Ministry of the Environment (U.prot PNM-2011-0010400-13/05/2011). Additional specimens were obtained from public Museums and Research Institutions (Table S1). All specimens are deposited in public collections. Of particular relevance are 15 males collected in the area of sympatry on Mount Beigua, where three Z. polyxena were collected in 1997, four Z. polyxena and two Z. cassandra in 1999, three Z. cassandra in 2000 and two Z. cassandra and one Z. polyxena in 2009. Landmarks and sliding semi-landmarks [17] were identified on the lateral section of the valvae using the TPS (thinplate spline) software. Six points along the valvae rim were considered as landmarks, while 24 additional points were defined as sliding semi-landmarks which could slide along the outline trajectory ([17] and Figure S1). Digitalization and definition of sliders were carried out using TPSDIG 2.16 [18] and TPSUTIL 1.53 [19], respectively. A generalized procrustes analysis was applied to landmark data in order to remove variation in location, scale and orientation, and to superimpose the objects into a common coordinate system [17]. We calculated partial warps from the shape residuals of the generalized procrustes analysis using TPSRELW 1.49 [20]. We then obtained principal components (PCs or relative warps) by applying a principal components analysis. We used the relative warps in a partitioning around medioids (PAM) analysis implemented in the *cluster* R package in order to identify the most likely number of morphotypes in our sample as suggested by Borcard et al. [21]. We generated specimen clusters between 2 and 216 and selected the partition showing the highest silhouette width (a relative measure of interversus intra-clusters dissimilarity spread) as the most suitable number of morphotypes.

Genetic Analysis

A subset of 106 Z. polyxena and Z. cassandra individuals were used for DNA analysis (Table S1). DNA was extracted using standard phenol-chloroform procedures. Two mitochondrial and one nuclear genes were sequenced and compared. The full-length "barcode" region (657 bp) of the mitochondrial cytochrome oxidase I (COI) gene was amplified and sequenced using the light-strand primer Parn Zer F1439 (5' - ATCGCTTATACT-CAGCCATC - 3') and the heavy-strand primer Parn_Zer_R2185 (5' - GGGAAATTATTCCAAATCCTG - 3') designed on the tyrosine tRNA and the COI gene, respectively, using a consensus sequence of Z. polyxena and Parnassius bremeri COI genes [22], [23]. Primer numbers refer to the 3' end of the published Parnassius bremeri mitochondrial genome sequence [23]. A 472 bp fragment of the mitochondrial NADH dehydrogenase subunit 1 (ND1) gene was amplified and sequenced using PCR primers described in [24]. Finally, amplification and sequence of a 393 bp fragment of the nuclear DNA wingless (wg) gene was obtained using the lightstrand primer wg_F29 (5' - CAGTAAAGACTTGCTGGATGC -3') and the heavy-strand primer wg_R382 (5' - TGCACCTTT-CAACCACAAAC - 3') designed specifically for this project. The 3' base of these primers refer to the published Zerynthia polyxena partial wingless sequence [22].

Polymerase chain reaction (PCR) amplifications were conducted in a total volume of 25 μ l containing 1–5 μ l of extracted DNA, 1× PCR buffer, 1.5 mM MgCl₂, 100 µM of each dNTP, 0.5 µM of each primer and 1 unit of Taq DNA polymerase (Invitrogen). Thermal profiles for COI amplification consisted of an initial denaturation step of 5 min at 94°C, followed by 35 cycles of 30 sec at 94°C, 30 sec at 50 and 90 sec at 72°C, with a final extension step of 7 min at 72°C. PCR profiles for ND1 and wg amplification had an initial denaturation step of 5 min at 94°C, followed by 35 cycles of 30 sec at 94° C, 30 sec at 52° C and 60 sec at 72° C, with a final extension step of 7 min at 72°C. Samples that resulted in a poor PCR product were amplified using 1 µl of extracted DNA, 1× Restorase buffer, 200 µM of each dNTP, 0.5 µM of each primer and 1.25 units of Restorase DNA polymerase (Sigma-Aldrich), a blend of high quality Taq DNA polymerase and a DNA repair enzyme which has proved effective in the amplification of damaged DNA [25]. PCR mix was incubated for 15 min at 37°C and for 5 min at 72° C. Primers were then added to the mix prior to amplification. PCR cycles consisted of an initial denaturation step of 2 min at 94°C, followed by 40 cycles of 30 sec at 94°C, 30 sec at $50-52^{\circ}$ C and 60 sec at 72° C, with a final extension step of 5 min at 72°C. PCR products were cycle-sequenced using BigDye Terminator v3.1 (Applied Biosystems) according to the manufacturer's protocol. Cycle sequencing reactions were resolved on an Applied Biosystems 3100 DNA analyzer and raw sequence chromatographs from both strands were edited and aligned using CodonCode Aligner 3.0.1 (CodonCode Corporation). The resulting consensus sequences consisted of a total of 657, 438 and 348

nucleotides of COI, ND1 and wg genes, respectively (Genbank accession numbers: KC119707-KC119746).

Haplotype Networks and Phylogenetic Analysis

Sequences were checked for insertions, deletions and stop codons that would result in non-functional proteins. Mean maximum likelihood (ML) distances within and between species and among localities were calculated using MEGA 5 [26]. We inferred haplotypes networks for COI and ND1 genes using the median-joining network method [27] implemented in the program NETWORK 4.6.1.0 (http://www.fluxus-engineering.com). We set the same default weight (10) for each character/site, given the absence of hyper-variable sites. The Epsilon parameter was set to 0. We used the default "connection cost" distance calculation methods and we applied the MP option [28] on results of the median joining calculation. Eight GenBank sequences of \mathcal{Z} . *polyxena* from France, Romania and Russia [22], [29], [30] were also used for construction of haplotype networks.

Best fit of molecular evolution model to each of our dataset was assessed using JMODELTEST [31] under the Bayesian Information Criterion. Likelihood values were calculated for 88 models using a maximum likelihood optimization of tree topology implemented in Phyml [32]. Each model was tested allowing for variation in nucleotide frequencies, different substitution rates among sites and proportion of invariable sites. The models of sequence evolution that best fit the COI and ND1 datasets were the Kishino and Yano (HKY) model [33] and the Transition model (TIM) [34], respectively. The Kimura 2-parameter (K80) model [35] resulted the best fit for the wingless dataset.

Partitioned Bayesian inference was applied to the combined dataset of COI, ND1 and wg sequences. Bayesian inference was conducted using Metropolis-coupled Markov chain Monte Carlo method implemented in MRBAYES 3.1.2 [36] applying the best substitution model for each partition of the combined dataset. The TIM and the K80 models could not be implemented in MRBAYES. We applied the general time-reversible model (GTR) [37], which best matches the assumptions of the transition model, and the HKY model (instead of the K80 model) with equal stationary state frequency. Approximation of the posterior probabilities of trees was performed by two independent runs starting with default prior values and initial random trees with three heated and one cold Markov chains, which ran for 2×10^6 generations with a 1000 generation sampling interval. Stationarity of the analysis was determined by examining the standard deviation of split frequencies between the two simultaneous runs and the potential scale reduction factor [38]. The first 25% of trees were discarded as burn-in and trees were used for analysis only after the chain became stable. The remaining trees were used to construct a 50%-majority rule consensus tree. GenBank sequences of the Papilionidae Zerynthia rumina, Bhutanitis thaidina and Allancastria cretica [22], [30] were used as outgroups. The phylogenetic tree was rooted on Bhutanitis thaidina.

Distribution Modelling

Locations of 334 European occurrence sites for Z cassandra and 575 sites for Z polyxena were assessed based on known species distribution [15], [39], (Dapporto & Zinetti unpublished data). The 19 WorldClim variables [40] were used to perform maximum entropy distribution modelling. We used a resolution grid of 0.04 degrees. These variables generally show collinearity which may bias results. We therefore selected 10,000 random cells across a plot in the Mediterranean region spanning from 1°00'00" E to 35°00'00" E and from 33°00'00" N to 70°00'00" N. For each cell, we analyzed the correlation among the WorldClim variables. We then selected the most biologically meaningful variable for those cases in which two or more variables showed a Pearson correlation coefficient higher than 0.8. Biological variables were selected mostly on the known species requirements based on adults activity and/or extreme climatic conditions experienced by larvae. We predicted potential distributions based on butterfly presence data using MAXENT 3.3.2 (http://www.cs.princeton.edu/schapire/ maxent). The software implements a machine-learning algorithm based on maximum entropy to identify areas with optimal environmental conditions [41]. Considering that different sampling efforts for different areas can produce false signals of climatic preferences [41], we applied a spatial filter to select a single random specimen for each cell of 0.4×0.4 degrees using the gridSample function implemented in the R package DISMO. The filter resulted in 117 and 287 presence points for Z. cassandra and Z. polyxena, respectively. We used default parameter settings and removed hinge and likelihood features to increase prediction accuracy [42]. Each model was replicated 100 times with a crossvalidation test performed on 10% of presence data. Goodness of the model was examined using receiver operating characteristic (ROC) plots and quantified by the area under curve (AUC). AUC values higher than 0.7 indicated that predictions of the model were higher than random values [43]. Relative importance of variable contribution was assessed by Jackknife of sequential variables removal [41]. We assessed models for butterfly distributions in modern day climate and then projected expected distributions into Last Glacial Maximum (LGM) WorldClim data. We used the Community Climate System Model (CCSM) and the Model for Interdisciplinary Research on Climate (MIROC) obtained from the Paleoclimate Modelling Intercomparison Project Phase II using the same variables of the modern climate models. These models differ in the reconstruction of several climatic variables and are well known to produce different results. For instance, CCSM and MIROC tend to overestimate and underestimate, respectively, proxy evidence of LGM winter sea ice [44]. As a result, for Mediterranean butterflies, the CCSM model tends to project narrower distributions at LGM than MIROC (e.g. [45], [46]).

Results

Geometric Morphometrics

Analysis of 217 male genitalia resulted in 52 relative warps. PAM analysis showed that the partition in two clusters had the highest silhouette width (Fig. 1a). This result supported the distinction of two morphotypes. Specimen assignment by PAM confirmed that all samples originating from Italy, south to the river Po, grouped in the first cluster (\mathcal{Z} . cassandra), while the other populations were included in the second cluster (\mathcal{Z} . polyxena). Specimens from Mount Beigua were found in both groups (Fig. 1b). Shape variability along PC1 (35.16% of variance explained) and PC2 (22.63% of variance explained) indicated that most shape variance was explained by the length of the valvar apex (Fig. 1b).

Genetic Diversity and Haplotype Networks

We obtained 103 consensus sequences and 22 haplotypes characterized by 35 variable nucleotide sites for the COI gene, and 86 sequences and 14 haplotypes defined by 23 polymorphic sites for the ND1 gene. Unambiguous sequences of the nuclear DNA wingless gene were obtained for 97 individuals. A total of four wg genotypes were characterized by four polymorphic sites. All \mathcal{Z} . cassandra specimens showed the same wg genotype, while three different genotypes were found for \mathcal{Z} . polyxena. Three samples of \mathcal{Z} . polyxena were heterozygous at two polymorphic sites (Table 1).



Figure 1. Analysis of genitalic morphology. a) Partitioning Around Medioids (PAM) analysis. The red arrow indicates the solution that assumes the existence of two morphotypes having the best fit (highest average silhouette width) among all possible partitions. b) Partition in two groups obtained by PAM analysis plotted for the first two relative warps (PC1 and PC2). A high concordance with geographic origin is shown for both *Z. cassandra* (c) and *Z. polyzena* (p) morphotypes with the only exception of the Mount Beigua population (black dots), which includes individuals belonging to both species. Deformation grids of male genitalia confirm previous results on morphological diversification between the studied species [15].

doi:10.1371/journal.pone.0065746.g001

Maximum likelihood mean distances between species were consistently higher than within species values for COI, ND1 and wg genes (Table 2 and Table S1).

In the COI haplotype network, Z cassandra and Z polyxena were separated by a minimum of 13 mutations (from C2 to C14, Fig. 2a). C1 was the most common Z cassandra haplotype in Italy and the only haplotype of this species occurring north of Tuscany. The second most common haplotype was found in southern Italy and was the only sequence occurring on Elba island. Other Z.

Table 1. Nucleotide variation and genotype designation in 97 *Zerynthia cassandra* and *Z. polyxena* individuals sequenced for the wingless (wg) gene.

wg genotype	Polyr	norphi	c site		Specimens	
	57	132	165	237		
W1	А	G	С	G	Z. cassandra (N=76)	
W2	G	А	Т	А	Z. polyxena (N = 17)	
W3	G	А	С	G	Z. polyxena (N = 1)	
W4	G	А	T/C	A/G	Z. polyxena (N=3)	

doi:10.1371/journal.pone.0065746.t001

cassandra haplotypes were found in central and southern Italy. The COI haplotype network for Z. polyxena described two major groups. The first one included haplotypes from France, while the second grouped sequences from eastern Europe and northern Italy (Fig. 2a). One haplotype from the French region was also found in northern Italy. The ND1 haplotype network revealed a similar pattern with haplotypes of Z. cassandra and Z. polyxena separated by a minimum of 10 mutations (from N6-N8 to N11, Fig. 2b). N1 was the most common Z. cassandra haplotype occurring in northern and central Italy. The second most common haplotype (N2) occurred from Tuscany down south to Sicily. The ND1 network confirmed the characterization of a French and a northern Italian haplogroups for Z. polyxena. The Italian specimen showing the French COI haplotype had also a French ND1 sequence. All specimens from France and the Italian Alpine region had a Z. polyxena haplotype, corroborating geometric morphometrics results (Fig. 2c). On the other hand, all individuals from the Apennines had a Z. cassandra haplotype for all markers, with the exception of specimens from Mount Beigua where haplotypes of both taxa were found in sympatry but segregated in accordance to morphometric analyses (Fig. 2c).

Table 2. Genetic diversity of COI, ND1 and wingless genes in Zerynthia cassandra and Z. polyxena.

	Zerynthia cassandra			Zerynt	hia polyxe	na			
	N	h	A+T	N	h	A+T	ML between species	ML within species	
COI	79	12	0.71	24	10	0.71	0.014±0.005	0.006±0.002	
ND1	75	10	0.77	11	4	0.78	0.032±0.008	0.011±0.003	
wg	76	1	0.46	21	3	0.46	0.010±0.005	0.004±0.002	

Number of individuals for which unambigous consensus sequences were obtained (N), number of haplotypes or genotypes (h) and A+T mean frequencies are shown for each species. Maximum likelihood (ML) mean distances between and within species are also reported. doi:10.1371/journal.pone.0065746.t002

doi:10.1371/journal.pone.0065746.002

Phylogenetic Analyses

The topology of the Bayesian tree constructed using the combined genetic datasets revealed two monophyletic lineages, supported by 100% posterior probability values (Fig. 3). The first lineage included all specimens morphologically attributed to Z. *cassandra*, while the second lineage comprised all Z. *polyxena* samples. The first lineage was characterized by two main clades. The first one included specimens from central and southern Italy and Sicily. All but one individuals from Sicily were grouped together. The second clade comprised specimens from northern and central Italy (including those from the region of sympatry) and Elba Island. The lineage of Z. *polyxena* consisted of two well supported clades. The first clade included all specimens from France and one from northern Italy (Vercelli), while the second one consisted of all samples from northern Italy.

Species Distribution Modelling

The MAXENT modelling resulted in a good fit for both species, with an AUC score of 0.861 and 0.977 for Z. polyxena and Z. cassandra, respectively. The Jackknife evaluation of the importance of variables revealed that temperature was more important than precipitation. Maximum temperature in the warmest month (BIO5), minimum temperature in the coldest month (BIO6), mean precipitation in the driest quarter (BIO9) and mean diurnal temperature range (BIO2) were most important in the Z, polyxena model (File S1), whereas BIO6, BIO9 and precipitation in the warmest quarter (BIO18) were significant in the Z. cassandra model (File S2). The areas with a logistic response higher than 0.5 largely matched the observed species distributions. The only exception was the Italian peninsula, which was predicted to have a suitable climate for Z. polyxena despite the total absence of the species from this area (Fig. 4). In particular, the regions predicted to be potential areas of distribution of the two species largely overlapped in the Apennines and, to a lesser extent, in the Maritime Alps and southern France. Projections of the two climatic reconstructions for the last glacial maximum revealed different results between the MIROC and the CCSM models. In particular, the MIROC model projected a larger area with logistic values higher than 0.5 (Fig. 4), a pattern found in other butterfly studies (e.g. [45], [46]) In these projections, both species showed a reduced expected occurrence in the northern regions of the Mediterranean range and a fragmented distribution in southern areas. However, in both the MIROC and CCSM models, Z. polyxena was expected to occur across the Italian peninsula (Fig. 5).

Discussion

Species Divergence

The Papilionidae Zerynthia cassandra and Z, polyxena revealed a strong and consistent pattern of genetic and morphological

differentiation. Analysis of mitochondrial and nuclear DNA and genitalic morphometric data resulted in a phylogenetic reconstruction that clearly divided samples collected from Sicily to southern France into two clades. Divergence of traits was maintained in the region where *Z. cassandra* and *Z. polyxena* occur in sympatry, with no evidence for hybridization.

Levels of genetic divergence between Z. cassandra and Z. polyxena (1.5% COI, 3% ND1 and 1% wg) were similar to, or lower than, values reported for most European sister and/or cryptic species comparisons [13], [29], [47], [48], [49]. These studies also showed that most taxa hybridize at their contact areas. Hybrid zones are generally large (50-250 km) in butterfly species for which introgression occurs over their European range [13], [29], [49], [50], [51]. The cryptic butterflies Polyommatus icarus/P. celina, and Aricia agestis/A. cramera, for instance, have hybrid zones 200 and 50 km wide, respectively, and show clear phylogenetic divergence from 3% to 5% of COI sequences. Nevertheless, they show clear evidence of introgression with intermediate morphotypes and discrepancy between nuclear and mitochondrial DNA sequences [49], [52]. We sampled very close populations of \mathcal{Z} . cassandra and Z. polyxena (Vercelli, Vigevano and Alessandria) with no apparent geographical barriers and analyzed specimens from the only area where the two species have been found in sympatry (Mount Beigua). We recorded no evidence of introgression of either mitochondrial, nuclear or morphological markers despite evidence of rather long dispersal was suggested by the occurrence of a French haplotype in a specimen collected in northern Italy. The Vigevano and Vercelli sampling sites of the northern Italy contact zone, where we found only Z polyxena morphotypes and DNA sequences, are 37 and 50 km from Alessandria, respectively, where only Z. cassandra sequences and morphotypes are recorded. Although no evidence for introgression in a limited set of specimens and markers is not a definitive proof of lack of hybridization, we recovered a very different pattern from that described for most European sister taxa. Moreover, although a very narrow hybridization area may occur between these localities, the belt would be strikingly narrower than those reported for other European species.

Theory suggests that the stronger the selection of resident alleles over the two sides of a hybrid zone, the narrower the area of the hybrid zone itself (reviewed in [50]). In the absence of recognition mechanisms allowing individuals to mate with conspecifics, hybrid areas can be seen as population sinks and are unlikely to enlarge over larger portions of the areas occupied by the two different entities [53]. So far, no experiments have been conducted on preand post-copulatory mechanisms involved in intraspecific mating avoidance in *Zerynthia*, and fitness depression has yet to be demonstrated in *Z. polyxena* and *Z. cassandra* hybrids. In fact, crossing experiments have been performed between *Z. polyxena* and formerly classified *Z. polyxena cassandra* from France [4], which is



Figure 2. Median-joining haplotype network of mitochondrial DNA COI (a) and ND1 (b) sequences for *Zerynthia cassandra* and *Z. polyxena*. Eastern European haplotypes (C24–C29) are shown in the same bullet color. The lower map (c) shows distribution of haplotypes and genitalic morphotypes for each sampling location. Each pie is divided into a number of slices equal to the number of sampled butterflies for that location. Each slice is further divided into four sectors. Starting from the centre of the pie, the first two sectors show the COI and ND1 haplotype colors, respectively, found

at that location (see networks reconstruction). The third and the outer sectors of the slice show assignment of wingless genotype and genitalic morphotype, respectively, to either *Z. cassandra* (black) or *Z. polyxena* (white). A grey sector indicates absence of the corresponding marker for that individual.

doi:10.1371/journal.pone.0065746.g002

actually a population of Z polyxena. However, our results suggest that strict recognition systems, strong hybrid depression and/or interactions determining mutual exclusion among these species occur and, as a result, a form of almost complete reproductive barrier emerged in a relatively short evolutionary time. Further investigations may therefore be necessary to clarify the origin of and mechanisms involved in these barriers.

Distribution Modelling and Conclusive Remarks

The current range of Z cassandra and Z polyxena matched the distribution predicted by climatic models. The only relevant discrepancy between observed and predicted distribution was for central and southern Italy, an area with suitable habitat for Z. polyxena but with no records of occurrence. In particular, the model predicted both taxa to be present in Corsica and Sardinia, where no record of them has ever been reported. Similarly, Z polyxena was predicted to occur across most of the Italian Apennines and in Sicily where this species has never been observed.

The projection of climatic models onto glacial maximum scenarios revealed that both species, particularly Z. cassandra, probably experienced range contractions which could have favoured isolation and divergence. Modelling of species occurrence resulted in wider distributions recovered by MIROC with respect to CCSM. A similar result was found by [45] and [46]. However, Z. polyxena was predicted to occur in the Italian Peninsula during glacial maxima by both models. Therefore neither niche modelling on current distribution nor retrodictions on LGM can explain the absence of Z. polyxena from the Italian peninsula.

Morphological, genetic and climatic information and data from other butterfly groups (e.g. [4]) suggest that genitalic divergence and reproductive barriers between Z. polyxena and Z. cassandra may have developed in a relatively short time. In fact, most cryptic sister species occurring in Europe show a divergence time similar to that reported for Z. polyxena and Z. cassandra. However, no instances have been reported of both a strong diversification in genitalic structure and complete absence of introgression, as shown in our study [13], [16], [29], [49]. An important factor determining the evolution of two very distinct morphologies and genetic units is probably due to the low dispersal ability of Zerynthia [54]. Moreover, males and females of both species are highly phylopatric and their area of activity mostly depends on the distribution of the same larval host plants (Aristolochia spp.) [55]. A good indicator of such a limited dispersal capacity is the absence of both species from most Mediterranean islands, despite suitable climatic conditions and the occurrence of potential host plants. Island populations of Z. cassandra are found only on the islands of Elba and Sicily [56], which are both close to the mainland. Similarly, *Z. polyxena* is known to occur on a few islands close to the Balkan peninsula [39]. From this perspective, the presence of two distinct areas with suitable climatic conditions for Z. polyxena in northern Italy-France and Greece, respectively, described by the CCSM model for the last glacial maximum, does not necessarily imply the occurrence of this species in both areas. Haplotype networks clearly suggest that both species have strong genetic structure, indicating that several populations might have survived the LGM and previous glacial periods in separate micro-refugia. A prominent pattern in Z. polyxena, for instance, is shown by separate groups occurring in eastern Europe (Russia and Romania), northern Italy and France. Tuscany (central Italy) is rich in endemic haplotypes of Z. cassandra, suggesting that this region may have functioned as a glacial refugium, as confirmed by distribution



Figure 3. Majority rule (50%) consensus tree resulting from Bayesian analysis of the combined COI, ND1 and wingless gene datasets. Node supports inferred from Bayesian posterior probability are shown above recovered branches. doi:10.1371/journal.pone.0065746.g003



Figure 4. Representation of the logistic output of the Maxent analyses for *Zerynthia cassandra* (a) and *Z. polyxena* (b). Values >0.5 indicate the likely presence of a species. Bullets show current distributions. In the lower map (c) blue and yellow areas show logistic output >0.5 for *Z. cassandra* and *Z. polyxena*, respectively. Areas where both species are predicted to occur are reported in green. Bullet and circles show current distribution of *Z. cassandra* and *Z. polyxena*, respectively. doi:10.1371/journal.pone.0065746.g004

modelling projections. Conversely, the area north-west of Tuscany (Liguria and Piemonte) appeared to have been colonized more recently by a single Tuscan haplotype of Z. cassandra. Z. polyxena from northern Italy showed a more complex situation with most haplotypes being endemic but highly related to the eastern haplogroups and just one haplotype shared with French populations.

Nazari and Sperling [16] estimated a divergence time of 1.8 My (lower Pleistocene) between Z. polyxena and Z. cassandra, supporting our hypothesis that a series of glacial and interglacial phases could have initiated and maintained the inter and intraspecific diversification process. Our analyses suggest that the last glacial stage did not result in extensive extinctions of Zerynthia across the study area and that Italy was probably a suitable habitat for both taxa. Since no relict populations of Z. polyxena



Figure 5. Projection of the Maxent models (based on present species distribution and climate data) on climatic reconstruction for the last glacial maximum using MIROC and CCSM circulation models. Dark areas show predicted species distribution (logistic output >0.5) during the last glacial age. doi:10.1371/journal.pone.0065746.g005

were found in the Italian peninsula, it is likely that Z. polyxena and Z. cassandra were allopatric before the LGM, probably as a result of reproductive barriers and/or competition for resources. There is also evidence that Z. cassandra came into contact with Z. polyxena after a relatively recent colonization event of Liguria and Piemonte from central Italy. This body of evidence suggests that in the last glacial stages Z. cassandra probably experienced isolation in several Italian micro-refugia. Such pattern was followed by expansion and marginal contacts with Z. polyxena during interglacial periods. The presence of a single haplotype endemic to Elba Island suggests that island colonization may have been hindered by sea level rise at the end of the last Ice Age, while genetic structure of mainland populations changed following complex and recurrent colonization events, as shown for other butterfly species from the same region [57], [58].

Recent literature suggests that speciation can occur even when populations are in contact and able to hybridize (reviewed in [7]). Epistasis conferring enhanced fitness in different areas may result in limited exchange of genomic components associated to reproduction. In Zerynthia, genetic interactions affecting shape of genitalia may have played a particular role at the initial stage of diversification, when the two taxa probably came into repeated contact during glacial and interglacial periods. On the other hand, there is evidence that other adaptive and neutral variation may be more easily exchanged among species [6]. In the Mediterranean, distinct species share adaptive variation across the Maghreb and Italy [45], [57], while in Neotropical butterflies adaptive Müllerian wing colouration can be exchanged among species despite their strong genetic diversification [6]. These examples may help explaining why Z, polyxena and Z, cassandra have maintained identical aposematic wing patterns and colouration [59], [60].

The Zerynthia species considered in our study seem to follow the reproductive isolation rule required by the biological species concept. From this perspective, they represent an exception compared to most European sister/cryptic species. We therefore recommend to consider Z, polyxena and Z, cassandra as good species, also in the restrictive sense of Descimon and Mallet [4], and to promote these taxa as an important model for the study of speciation, correlation between genotypic and phenotypic traits, evolution and maintenance of adaptive patterns.

Supporting Information

Figure S1 Schematic representation of fixed landmarks and sliding semi-landmarks considered in geometric morphometric analyses.

(PDF)

Table S1 Additional information on *Zerynthia polyxena* and Z. *cassandra* samples analysed in this study. (DOC)

File S1 Supplementary results for the Maxent model of *Z. polyxena*.

(PDF)

File S2 Supplementary results for the Maxent model of *Z. cassandra*.

(PDF)

Acknowledgments

Our work was conducted in collaboration with the Arcipelago Toscano National Park, the Capanne di Marcarolo Nature Park, the Parco fluviale del Po tratto vercellese/alessandrino e Riserva Naturale del Torrente Orba, and the Migliarino, San Rossore, Massaciuccoli Nature Park. We thank Chiara Natali for assistance during laboratory work and Roger Vila and Vlad Dincă for providing three additional mitochondrial COI gene sequences and for their help during samples collection. We are also grateful to Nicola Scatassi and Gabriele Panizza for their enthusiastic support during field work. We also thank Martin Wiemers and an anonymous referee for their constructive suggestions.

Author Contributions

Conceived and designed the experiments: LD. Performed the experiments: LD FZ AV. Analyzed the data: LD FZ AV SB EB CC. Contributed

References

- Mayr E (1942) Systematics and the origin of species. New York, NY: Columbia University Press.
- Mallet J (2005) Hybridization as an invasion of the genome. Trends Ecol Evol 20: 229–237.
- Currat M, Ruedi M, Petit RJ, Excoffier L (2008) The hidden side of invasions: massive introgression by local genes. Evolution 62: 1908–1920.
- Descimon H, Mallet J (2009) Bad species. In: Settele J, Shreeve T, Konvička M, Van Dyck H, editors. Ecology of butterflies in Europe. Cambridge University Press. pp. 219–249.
- Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. Annu Rev Ecol Evol Syst 40: 481–501.
- Heliconius Genome Consortium (2012) Butterfly genome reveals promiscuous exchange of mimicry adaptation among species. Nature 487: 94–98.
- Hausdorf B (2011) Progress toward a general species concept. Evolution 65: 923–931.
- Mallet J (1995) A species definition for a modern synthesis. Trends Ecol Evol 10: 294–299
- Wu CI (2001) The genic view of the process of speciation. J Evol Biol 14: 851– 865.
- 10. Coyne FM, Orr HA (2004) Speciation. Sunderland, MA: Sinauer.
- Mallet J (2008) Hybridization, ecological races, and the nature of species: empirical evidence for the ease of speciation. Philos Trans R Soc Lond B 363: 2971–2986.
- Friberg M, Vongvanich N, Borg-Karlson AK, Kemp DJ et al. (2008) Female mate choice determines reproductive isolation between sympatric butterflies. Behav Ecol Sociobiol 62: 873–886.
- Dincă V, Lukhtanov VA, Talavera G, Vila R (2011) Unexpected layers of cryptic diversity in wood white Leptidea butterflies. Nat Commun 2: 324.
- Lorković Z (1993) Leptidea reali Reissinger, 1989 (= lorkovicii Real 1988), a new European species (Lepid., Pieridae). Nat Croat 2: 1–26.
- Dapporto L (2010) Speciation in Mediterranean refugia and post-glacial expansion of *Zerynthia polyxena* (Lepidotera, Papilionidae). J Zool Syst Evol Res 48: 229–237.
- Nazari V, Sperling FAH (2007) Mitochondrial DNA divergence and phylogeography in western Palacarctic Parnassiinae (Lepidoptera: Papilionidae): How many species are there? Insect Syst Evol 38: 121–138.
- Bookstein FL (1997) Landmark methods for forms without landmarks: localizing group differences in outline shape. Med Image Anal 1: 225–243.
- Rohlf FJ (2010) tpsDig, digitize landmarks and outlines, version 2.16. Department of Ecology and Evolution, State University of New York at Stony Brook.
- Rohlf FJ (2012) tpsUtil, file utility program. version 1.53. Department of Ecology and Evolution, State University of New York at Stony Brook.
- Rohlf FJ (2010) tpsRelw, relative warps analysis, version 1.49. Department of Ecology and Evolution, State University of New York at Stony Brook.
- 21. Borcard D, Gillet F, Legendre P (2011) Numerical Ecology with R. New York: Springer-Verlag.
- Nazari V, Zakharov EV, Sperling FA (2007) Phylogeny, historical biogeography, and taxonomic ranking of Parnassiinae (Lepidoptera, Papilionidae) based on morphology and seven genes. Mol Phylogenet Evol 42: 131–156.
- 23. Kim MI, Back JY, Kim MJ, Jeong HC, Kim KG et al. (2009) Complete nucleotide sequence and organization of the mitogenome of the red-spotted apollo butterfly, *Parnassius bremeri* (Lepidoptera: Papilionidae) and comparison with other lepidopteran insects. Mol Cells 28: 347–363.
- Aubert J, Legal L, Descimon H, Michel F (1999) Molecular phylogeny of swallowtail butterflies of the tribe Papilionini (Papilionidae, Lepidoptera). Mol Phylogenet Evol 12: 156–167.
- Hajibabaei M, deWaard JR, Ivanova NV, Ratnasingham S, Dooh RT et al. (2005) Critical factors for assembling a high volume of DNA barcodes. Philos Trans R Soc Lond B 360: 1959–1967.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M et al. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. Mol Biol Evol 28: 2731–2739.
- Bandelt HJ, Forster P, Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. Mol Biol Evol 16: 37–48.
- Polzin T, Daneschmand SV (2003) On Steiner trees and minimum spanning trees in hypergraphs. Oper Res Lett 31: 12–20.
- Dincă V, Zakharov EV, Hebert PDN, Vila R (2011c) Complete DNA barcode reference library for a country's butterfly fauna reveals high performance for temperate Europe. Proc R Soc Lond B 278: 347–355.
- Michel F, Rebourg C, Cosson E, Descimon H (2008) Molecular phylogeny of Parnassiinae butterflies (Lepidoptera: Papilionidae) based on the sequences of four mitochondrial DNA segments. Ann Soc Entomol Fr 44: 1–36.
- Posada D (2008) jModelTest: Phylogenetic model averaging. Mol Biol Evol 25: 1253–1256.

- Guindon S, Gascuel O (2003) A simple, fast and accurate algorithm to estimate large phylogenies by maximum-likelihood. Syst Biol 52: 696–704.
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. J Mol Evol 22: 160–174.
- Posada D (2003) Using Modeltest and PAUP* to select a model of nucleotide substitution. In: Baxevanis AD, Davison DB, Page RDM, Petsko GA, Stein LD, Stormo GD, editors. Current Protocols in Bioinformatics. New York: John Wiley & sons, Inc. pp. 6.5.1–6.5.14.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. J Mol Evol 16: 111–120.
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574.
- Tavaré S (1986) Some probabilistic and statistical problems in the analysis of DNA sequences. In: Miura RM, editor. Some mathematical questions in biology - DNA sequence analysis. Providence, RI: American Mathematical Society. pp. 57–86.
- Ronquist F, Deans AR (2010) Bayesian phylogenetics and its influence on insect systematics. Annu Rev Entomol 55: 189–206.
- Kudrna O, Harpke A, Lux C, Pennerstorfer J, Schweiger O et al. (2011) Distribution atlas of butterflies in Europe. Halle, Germany: Gesellschaft f
 ür Schmetterlingschultz.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. Int J Climatol 25: 1965–1978.
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modelling of species geographic distributions. Ecol Model 190: 231–259.
- 42. Austin MP (2007) Species distribution models and ecological theory: a critical assessment and some possible new approaches. Ecol Model 200: 1–19.
- 43. Elith J (2002) Quantitative methods for modeling species habitat: comparative performance and an application to Australian plants. In: Ferson S, Burgman M, editors. Quantitative methods for conservation biology. New York: Springer-Verlag. 39–58.
- Otto-Bliesner BL, Hewitt CD, Marchitto TM, Brady E, Abe-Ouchi A et al. (2007) Last Glacial Maximum ocean thermohaline circulation: PMIP2 model intercomparisons and data constraints. Geophys Res Lett 34: L12706.
- Habel JC, Rödder D, Scalercio S, Meyer M, Schmitt T (2010) Strong genetic cohesiveness between Italy and North Africa in four butterfly species. Biol J Linn Soc 99: 818–830.
- Habel JC, Husemann M, Schmitt T, Dapporto L, Rödder D et al. (2013) A forest butterfly in Sahara desert oases: isolation does not matter. J Hered 104: 234–247.
- Cianchi R, Ungaro A, Marini M, Bullini L (2003) Differential patterns of hybridization and introgression between the swallowtails *Papilio machaon* and *P. Hospiton* from Sardinia and Corsica islands (Lepidoptera, Papilionidae). Mol Ecol 12: 1461–1471.
- Wiemers M, Gottsberger B (2010) Discordant patterns of mitochondrial and nuclear differentiation in the Scarce Swallowtail *Iphiclides podalirius feisthamelii* (Duponchel, 1832) (Lepidoptera: Papilionidae). Entomol Z 120: 111–115.
- Sañudo-Restrepo CP, Dincă V, Talavera G, Vila R (2012) Biogeography and systematics of *Aricia* butterflies (Lepidoptera, Lycaenidae). Mol Phylogenet Evol (in press).
- Porter AH (2009) Ecological genetics and evolutionary ecology in hybrid zones. In: Settele J, Shreeve T, Konvička M, Van Dyck H, editors. Ecology of butterflies in Europe. Cambridge: Cambridge University Press. 296–311.
- Mallet J, Wynne IR, Thomas CD (2011) Hybridisation and climate change: brown argus butterflies in Britain (*Polyommatus* subgenus Aricia). Insect Conserv Divers 4: 192–199.
- Dincă V, Dapporto L, Vila R (2011) A combined genetic-morphometric analysis unravels the complex biogeographical history of *Polyommatus icarus* and *Polyommatus celina* Common Blue butterflies. Mol Ecol 20: 3921–3935.
- Dasmahapatra KK, Lamas G, Simpson F, Mallet J (2010) The anatomy of a 'suture zone' in Amazonian butterflies: a coalescent-based test for vicariant geographic divergence and speciation. Mol Ecol 19: 4283–4301.
- Celik T (2012) Adult demography, spatial distribution and movements of *Zerynthia polysena* (Lepidoptera: Papilionidae) in a dense network of permanent habitats. Eur J Entomol 109: 217–227.
- Bollino M, Racheli T (2012) Butterflies of the world, supplement 20, Parnassinae (partim), Parnassiini (partim), Luchdorfiini, Zerynthiini. Keltern: Goecke & Hevers.
- Balletto E, Bonelli E, Cassulo L (2007) Insecta Lepidoptera Papilionoidea. In: Ruffo S, Stoch F, editors. Checklist and distribution of the Italian fauna. 10,000 terrestrial and inland water species. Verona: Memorie del Museo Civico di Storia Naturale di Verona, Sez. Scienze della Vita, 2nd and revised edition. 17: 257–261.
- 57. Dapporto L, Habel JC, Dennis RLH, Schmitt T (2011) The biogeography of the western Mediterranean: elucidating contradictory distribution patterns of

differentiation in $Maniola\ jurtina$ (Lepidoptera, Nymphalidae). Biol J Linn Soc $103:\ 571-577.$

- Daporto L, Bruschini C, Dincă V, Vila R, Dennis RLH (2012) Identifying zones of phenetic compression in West Mediterranean butterflies (Satyrinae): refugia, invasion and hybridization. Divers Distrib 18: 1066–1076.
- Rothschild M, von Euw J, Reichstein T (1972) Aristolochic acids stored by Zeynthia polyxena (Lepidoptera). Insect Biochem 2: 334–343.
 Sime KR, Feeny PP, Haribal MH (2000) Sequestration of aristolochic acids by
- Sime KR, Feeny PP, Haribal MH (2000) Sequestration of aristolochic acids by the pipevine swallowtail, *Battus philenor* (L.): evidence and ecological implications. Chemoecology 10: 169–178.