

BIBLIOGRAFIA

- 1) Brånemark PI, Adell R, Breine U, Hansson BO, Lindström J, Ohlsson A Intra-osseous anchorage of dental prostheses. I. Experimental studies. *Scand J Plast Reconstr Surg.* 1969;3(2):81-100.
- 2) Albrektsson T, Brånemark PI, Hansson HA, Lindström J. Osseointegrated titanium implants. Requirements for ensuring a long-lasting, direct bone-to-implant anchorage in man. *Acta Orthop Scand.* 1981;52(2):155-70.
- 3) Marotti G, Ferretti M, Muglia MA, Palumbo C, Palazzini S. A quantitative evaluation of osteoblast-osteocyte relationships on growing endosteal surface of rabbit tibiae. *Bone.* 1992;13(5):363-8.
- 4) Marotti G, Ferretti M, Remaggi F, Palumbo C. Quantitative evaluation on osteocyte canalicular density in human secondary osteons. *Bone.* 1995;16(1):125-8.
- 5) Khun K. The structure of collagen. *Essays Biochem* 5:59-87,1969
- 6) Marotti G. The structure of bone tissues and the cellular control of their deposition. *Ital J Anat Embryol.* 1996;101(4):25-79.
- 7) Palumbo C. A three-dimensional ultrastructural study of osteoid-osteocytes in the tibia of chick embryos. *Cell Tissue Res.* 1986;246(1):125-31.
- 8) Price PA, Otsuka AA, Poser JW, Kristaponis J, Raman N. Characterization of a gamma-carboxyglutamic acid-containing protein from bone. *Proc Natl Acad Sci U S A.* 1976;73(5):1447-51.
- 9) Termine JD. Mineral chemistry and skeletal biology. *Clin Orthop Relat Res.* 1972;85:207-39.
- 10) Termine JD, Kleinman HK, Whitson SW, Conn KM, McGarvey ML, Martin GR. Osteonectin, a bone-specific protein linking mineral to collagen. *Cell.* 1981;26(1 Pt 1):99-105.
- 11) Rodan SB, Rodan GA, Simmons HA, Walenga RW, Feinstein MB, Raisz LG. Bone resorptive factor produced by osteosarcoma cells with osteoblastic features is PGE₂. *Biochem Biophys Res Commun.* 1981;102(4):1358-65.
- 12) Sissons HA, Kelman GJ, Ling L, Marotti G, Canè V, Muglia MA. A light and scanning electron microscopic study of osteocyte activity in calcium-deficient rats. *Calcif Tissue Int.* 1990;46(1):33-7.
- 13) Marotti G. The role of the bone cell in the regulation of phosphorus and calcium homeostasis. *G Clin Med.* 1988;69(4):243-61.
- 14) Rubinacci A, Villa I, Dondi Benelli F, Borgo E, Ferretti M, Palumbo C, Marotti G. Osteocyte-bone lining cell system at the origin of steady ionic current in damaged amphibian bone. *Calcif Tissue Int.* 1998;63(4):331-9.
- 15) Miller SC, Jee WS. The bone lining cell: a distinct phenotype? *Calcif Tissue Int.* 1987;41(1):1-5.
- 16) Volpi G, Palazzini S, Canè V, Remaggi F, Muglia MA. Morphometric analysis of osteoblast dynamics in the chick embryo tibia. *Anat Embryol (Berl).* 1981;162(4):393-401.

- 17) Lozupone E, Favia A. Density of trabecular framework and osteogenic activity in the spongiosa of long bones subjected to drastic changes in mechanical loading in the dog. *Anat Anz.* 1982;152(3):245-61.
- 18) Zallone AZ, Teti A. Animal models of bone physiology. *Curr Opin Rheumatol.* 1993;5(3):363-7.
- 19) Muglia MA, Volpi G, Remaggi F, Canè V, Palazzini S, Zaffe D, Marotti G. Activity of osteoclasts and osteocytes in compact human bone at various ages, both with and without osteoporosis. *Ital J Orthop Traumatol.* 1982;8(1):117-25.
- 20) Teti A, Marchisio PC, Zallone AZ. Clear zone in osteoclast function: role of podosomes in regulation of bone-resorbing activity. *Am J Physiol.* 1991;261(1 Pt 1):C1-7.
- 21) David JP, Neff L, Chen Y, Rincon M, Horne WC, Baron R. A new method to isolate large numbers of rabbit osteoclasts and osteoclast-like cells: application to the characterization of serum response element binding proteins during osteoclast differentiation. *J Bone Miner Res.* 1998;13(11):1730-8.
- 22) De Vernejoul MC, Horowitz M, Demignon J, Neff L, Baron R. Bone resorption by isolated chick osteoclasts in culture is stimulated by murine spleen cell supernatant fluids (osteoclast-activating factor) and inhibited by calcitonin and prostaglandin E2. *J Bone Miner Res.* 1988;3(1):69-80.
- 23) Hirano H, Urist MR. Bone-forming and bone-resorbing cell lines derived from bone marrow in tissue culture. *Clin Orthop Relat Res.* 1981;(154):234-48.
- 24) Vilmann H. Characteristics of growing bone surfaces. *Scand J Dent Res.* 1979;87(1):65-72.
- 25) Mardon HJ, Bee J, von der Mark K, Owen ME. Development of osteogenic tissue in diffusion chambers from early precursor cells in bone marrow of adult rats. *Cell Tissue Res.* 1987;250(1):157-65.
- 26) Young RW. Cell proliferation and specialization during endochondral osteogenesis in young rats. *J Cell Biol.* 1962;14:357-70.
- 27) Alhopuro S. Premature fusion of facial sutures with free periosteal grafts. An experimental study with special reference to bone formation with free periosteal grafts from the tibia, the scapula and the calvarium. *Scand J Plast Reconstr Surg Suppl.* 1978;17:1-68.
- 28) Wolff J. *Das gesetz der transformattion der knochen.* Hirschwald Berlin (Germany), 1982.
- 29) Lozupone E, Favia A. The structure of the trabeculae of cancellous bone. 2. Long bones and mastoid. *Calcif Tissue Int.* 1990;46(6):367-72.
- 30) Lipson RA, Halloran PF, Kawano H, Langer F. A microsurgical model for vascularized bone and joint transplants in rats. *Transplant Proc.* 1981 ;13(1 Pt 2):891-2.
- 31) Chen JH, Liu C, You L, Simmons CA. Boning up on Wolff's Law: mechanical regulation of the cells that make and maintain bone. *J Biomech.* 5;43(1):108-18. Epub 2009 Oct 8.
- 32) Burger EH, Klein-Nulen J. Responses of bone cells to biomechanical forces in vitro. *Adv Dent Res.* 1999;13:93-8.

- 33) Araújo MG, Wennström JL, Lindhe J. Modeling of the buccal and lingual bone walls of fresh extraction sites following implant installation. *Clin Oral Implants Res.* 2006;17(6):606-14.
- 34) Schropp L, Wenzel A, Kostopoulos L, Karring T. Bone healing and soft tissue contour changes following single-tooth extraction: a clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent.* 2003;23(4):313-23.
- 35) Botticelli D, Berglundh T, Lindhe J. Hard-tissue alterations following immediate implant placement in extraction sites. *J Clin Periodontol.* 2004;31(10):820-8.
- 36) Fickl S, Zuhr O, Wachtel H, Bolz W, Huerzeler MB. *Clin Oral Implants Res.* 2008;19(11):1111-8.
- 37) Cardaropoli G, Araujo M, Lindhe J: Dynamics of bone tissue formation in tooth extraction sites. An experimental study in dogs. *J Clin Periodontol* 2003; 30, 809–818.
- 38) Hämmerle CH, Chen ST, Wilson TG Jr. Consensus statements and recommended clinical procedures regarding the placement of implants in extraction sockets. *Int J Oral Maxillofac Implants.* 2004;19 Suppl:26-8
- 39) Tallgren A. The continuing reduction of the residual alveolar ridges in complete denture wearers: a mixed-longitudinal study covering 25 years. *J Prosthet Dent.* 1972;27(2):120-32.
- 40) Ostler MS, Kokich VG. Alveolar ridge changes in patients congenitally missing mandibular second premolars. *J Prosthet Dent.* 1994;71(2):144-9.
- 41) Atwood DA. Reduction of residual ridges: a major oral disease entity. *J Prosthet Dent.* 1971;26(3):266-79.
- 42) Cawood JI, Howell RA. A classification of the edentulous jaws. *Int J Oral Maxillofac Surg.* 1988;17(4):232-6.
- 43) Chiapasco M, Zaniboni M, Rimondini L. Dental implants placed in grafted maxillary sinuses: a retrospective analysis of clinical outcome according to the initial clinical situation and a proposal of defect classification. *Clin Oral Implant Res* 2008;19:416-28
- 44) Lekholm U, Zarb GA. Patient selection and preparation. In: Brånemark PI, Zarb GA, Albrektsson T, editors. *Tissue Integrated Prostheses: Osseointegration in Clinical Dentistry.* Chicago, Ill, USA: Quintessence; 1985. pp. 199–209
- 45) Misch CE. Bone classification, training keys to implant success. *Dent Today.* 1989 May;8(4):39-44.
- 46) Norton MR, Gamble C. Bone classification: an objective scale of bone density using the computerized tomography scan. *Clin Oral Implants Res.* 2001;12:79-84.
- 47) Turkyilmaz I, Tözüm TF, Tumer C. Bone density assessments of oral implant sites using computerized tomography. *J Oral Rehabil.* 2007;34:267-72.
- 48) Misch CE. *Contemporary implant dentistry.* St. Louis: Mosby; 1998.
- 49) Rangert B. Mechanical and biomechanical guidelines for the use of Branemark System--clinical studies. *Aust Prosthodont J* 1993; 7 Suppl 1:45-9.

- 50) Pierrisnard L, Renouard F, Renault P, Barquinis M. Influence of implant length and bicortical anchorage on implant stress distribution. *Clin Implant Dent Relat Res* 2003; 5(4):254-62.
- 51) Matsushita Y, Kitoh M, Mizuta K, Ikeda H, Suetsugu T. Two-dimensional FEM analysis of hydroxyapatite implants: diameter effects on stress distribution. *J Oral Implantol* 1990;16:6-11.
- 52) Bernard JP, Szmukler-Moncler S, Pessotto S, Vazquez L, Belser UC. The anchorage of Brånemark and ITI implants of different lengths. I. An experimental study in the canine mandible. *Clin Oral Implants Res* 2003; 14(5):593-600.
- 53) Abrahamsson I, Lindhe I et al. Early bone formation adjacent to rough and turned endosseous implant surface. An experimental study in the dog. *Clin. Oral. Impl. Res.*, 2004; 15: 381
- 54) Lossdörfer S, Schwartz Z, Wang L, Lohmann CH, Turner JD, Wieland M, Cochran DL, Boyan BD. Microrough implant surface topographies increase osteogenesis by reducing osteoclast formation and activity. *J Biomed Mater Res A*. 2004 1;70(3):361-9.
- 55) Anitua E, Orive G, Aguirre JJ, Andía I. Five-year clinical evaluation of short dental implants placed in posterior areas: a retrospective study. *J Periodontol*. 2008;79:42-8.
- 56) Das Neves FD, Fones D, Bernardes SR, Do Prado CJ, Neto AJ. Short implants-an analysis of longitudinal studies. *Int J Oral Maxillofac Implants*. 2006;21:86-93.
- 57) Fugazzotto PA. Shorter implants in clinical practice: rationale and treatment results. *Int J Oral Maxillofac Implants*. 2008;23:487-96.
- 58) Misch CE, Steingra J, Barboza E, Misch-Dietsh F, Cianciola LJ, Kazor C. Short dental implants in posterior partial edentulism: a multicenter retrospective 6-year case series study. *J Periodontol*. 2006;77:1340-7.
- 59) Romeo E, Ghisolfi M, Rozza R, Chiapasco M, Lops D. Short (8-mm) dental implants in the rehabilitation of partial and complete edentulism: a 3- to 14-year longitudinal study. *Int J Prosthodont*. 2006;19:586-92.
- 60) Ata-Ali J, Peñarrocha-Oltra D, Candel-Marti ME, Peñarrocha-Diago MA. Oral rehabilitation with tilted dental implants: A metaanalysis. *Med Oral Patol Oral Cir Bucal*. 2012 9. [Epub ahead of print]
- 61) Acocella A, Ercoli C, Geminiani A, Feng C, Billi M, Acocella G, Giannini D, Sacco R. Clinical Evaluation of Immediate Loading of Electroeroded Screw-Retained Titanium Fixed Protheses Supported by Tilted Implant: A Multicenter Retrospective Study. *Clin Implant Dent Relat Res*. 2011 Oct 20. doi: 10.1111/j.1708-8208.2011.00379.x. [Epub ahead of print]
- 62) Maló P, de Araújo Nobre M, Lopes A, Francischone C, Rigolizzo M. "All-on-4" Immediate-Function Concept for Completely Edentulous Maxillae: A Clinical Report on the Medium (3 Years) and Long-Term (5 Years) Outcomes. *Clin Implant Dent Relat Res*. 2011 Oct 18. doi: 10.1111/j.1708-8208.2011.00395.x. [Epub ahead of print]

- 63) Capelli M, Zuffetti F, Del Fabbro M, Testori T. Immediate rehabilitation of the completely edentulous jaw with fixed prostheses supported by either upright or tilted implants: a multicenter clinical study. *Int J Oral Maxillofac Implants*. 2007;22(4):639-44
- 64) Testori T, Del Fabbro M, Capelli M, Zuffetti F, Francetti L, Weinstein RL. Immediate occlusal loading and tilted implants for the rehabilitation of the atrophic edentulous maxilla: 1-year interim results of a multicenter prospective study. *Clin Oral Implants Res*. 2008;19(3):227-32
- 65) Stosić S. Mandibular reconstruction-state of the art and perspectives. *Vojnosanit Pregl*. 2008;65(5):397-403.
- 66) Sacco R, Sacco G, Acocella A, Sale S, Sacco N, Baldoni E. A systematic review of microsurgical reconstruction of the jaws using vascularized fibula flap technique in patients with bisphosphonate-related osteonecrosis. *J Appl Oral Sci*. 2011;19(4):293-300.
- 67) Urken ML, Buchbinder D, Costantino PD, Sinha U, Okay D, Lawson W, et al. Oromandibular reconstruction using microvascular composite flaps: Report of 210 cases. *Arch Otolaryngol Head Neck Surg* 1998; 124: 46–55.
- 68) Wei FC, Celik N, Chen HC, Cheng MH, Huang WC. Combined anterolateral thigh flap and vascularized fibula osteoseptocutaneous flap in reconstruction of extensive composite mandibular defects. *Plast Reconstr Surg* 2002;109: 45–52.
- 69) David DJ, Tan E, Katsaros J, Sheen R. Mandibular reconstruction with vascularized iliac crest: A 10-year experience. *Plast Reconstr Surg* 1988; 82: 792–803.
- 70) Silverstein LH, Kurtzman GM, Moskowitz E, Kurtzman D, Hahn J. Aesthetic enhancement of anterior dental implants with the use of tapered osteotomes and soft tissue manipulation. *J Oral Implantol*. 1999;25(1):18-22.
- 71) Calvo-Guirado JL, Saez-Yuguero R, Pardo-Zamora G. Compressive osteotomes for expansion and maxilla sinus floor lifting. *Med Oral Patol Oral Cir Bucal*. 2006;11(1):E52-5.
- 72) Siddiqui AA, Sosovicka M. Lateral bone condensing and expansion for placement of endosseous dental implants: a new technique. *J Oral Implantol*. 2006;32(2):87-94.
- 73) Duncan JM, Westwood M. Ridge widening for the thin maxilla. A clinical report. *Int J Oral Maxillofac Implants* 1997;12:224-227.
- 74) Koo S, Dibart S, Weber HP. Ridge splitting technique with simultaneous implant placement. *Compendium Contin Educ Dent* March 2008;29(2):106-110.
- 75) Scipioni A, Bruschi GB, Calesini G. The edentulous ridge expansion technique: A five-year study. *Int J Periodontics Restorative Dent* 1994;14:451-459.
- 76) Sethi A, Kaus T. Maxillary ridge expansion with simultaneous implant placement: 5-year results of an ongoing clinical study. *Int J Oral Maxillofac Implants* 2000;15:491-499.
- 77) Engelke WG, Diederichs CG, Jacobs HG, Deckwer I. Alveolar reconstruction with splitting osteotomy and microfixation of implants. *Int J Oral Maxillofac Implants* 1997;12:310-318.

- 78) Simion M, Baldoni M, Zaffe D. Jawbone enlargement using immediate implant placement associated with a split-crest technique and guided tissue regeneration. *Int J Periodontics Restorative Dent* 1992;12:462-473.
- 79) Ilizarov GA. The tension-stress effect on the genesis and growth of tissues. Part I. The influence of stability of fixation and soft-tissue preservation. *Clin Orthop* 1989;238:249-281.
- 80) Takahashi T, Funaki K, Shintani H, Haruoka T. Use of horizontal alveolar distraction osteogenesis for implant placement in a narrow alveolar ridge: A case report. *Int J Oral Maxillofac Implants* 2004;19:291-294.
- 81) Chiapasco M, Romeo E, Vogel G. Vertical distraction osteogenesis of edentulous ridges for improvement of oral implant positioning: A clinical report of preliminary results. *Int J Oral Maxillofac Implants* 2001;16:43-51.
- 82) Jensen OT, Cockrell R, Kuhike L, et al. Anterior maxillary alveolar distraction osteogenesis: A prospective 5-year clinical study. *Int J Oral Maxillofac Implants* 2002;17:52-68.
- 83) McAllister BS, Gaffaney TE. Distraction osteogenesis for vertical bone augmentation prior to oral implant reconstruction. *Periodontol* 2000 2003;33:54-66.
- 84) Chiapasco M, Zaniboni M, Rimondini L. Autogenous onlay bone grafts vs. alveolar distraction osteogenesis for the correction of vertically deficient edentulous ridges: a 2-4-year prospective study on humans. *Clin Oral Implants Res.* 2007;18(4):432-40.
- 85) Lopez-Cedrun JL. Implant rehabilitation of the edentulous posterior atrophic mandible: the sandwich osteotomy revisited. *Int J Oral Maxillofac Implants.* 2011;26(1):195-202.
- 86) Block MS, Haggerty CJ. Interpositional osteotomy for posterior mandible ridge augmentation. *J Oral Maxillofac Surg.* 2009;67(11 Suppl):31-9.
- 87) Bormann KH, Suarez-Cunqueiro MM, von See C, et al. Forty sandwich osteotomies in atrophic mandibles: a retrospective study. *J Oral Maxillofac Surg.* 2011;69(6):1562-70.
- 88) Chiapasco M, Brusati R, Ronchi P. Le Fort I osteotomy with interpositional bone grafts and delayed oral implants for the rehabilitation of extremely atrophied maxillae: a 1-9-year clinical follow-up study on humans. *Clin Oral Implants Res.* 2007;18(1):74-85.
- 89) Marchetti C, Pieri F, Corinaldesi G, Degidi M. A long-term retrospective study of two different implant surfaces placed after reconstruction of the severely resorbed maxilla using Le Fort I osteotomy and interpositional bone grafting. *J Oral Maxillofac Implants.* 2008;23(5):911-8.
- 90) Nyström E, Lundgren S, Gunne J, Nilson H. Interpositional bone grafting and Le Fort I osteotomy for reconstruction of the atrophic edentulous maxilla. A two-stage technique. *Int J Oral Maxillofac Surg.* 1997;26(6):423-7.
- 91) Garg AK, Morales MJ. Lateralization of the inferior alveolar nerve with simultaneous implant placement: surgical techniques. *Pract Periodontics Aesthet Dent.* 1998;10(9):1197-204

- 92) Peleg M, Mazor Z, Chaushu G, Garg AK. Lateralization of the inferior alveolar nerve with simultaneous implant placement: a modified technique. *Int J Oral Maxillofac Implants.* 2002;17(1):101-6.
- 93) Yoshimoto M, Watanabe IS, Martins MT, et al. Microstructural and ultrastructural assessment of inferior alveolar nerve damage following nerve lateralization and implant placement: an experimental study in rabbits. *Int J Oral Maxillofac Implants.* 2009;24(5):859-65.
- 94) Bidra AS, Huynh-Ba G. Implants in the pterygoid region: a systematic review of the literature. *Int J Oral Maxillofac Surg.* 2011;40(8):773-81.
- 95) Peñarrocha M, Carrillo C, Boronat A, Peñarrocha M. Retrospective study of 68 implants placed in the pterygomaxillary region using drills and osteotomes. *Int J Oral Maxillofac Implants.* 2009;24(4):720-6.
- 96) Bedrossian E, Stumpel III L, Beckely M, Indersano T. The zygomatic implant: preliminary data on treatment of severely resorbed maxillae. A clinical report. *Int J Oral Maxillofac Implants* 2002;17:861-5.
- 97) Branemark PI, Grondahl K, Ohnrell LO, Nilsson P, Petrusson B, Svensson B, et al. Zygoma fixture in the management of advanced atrophy of the maxilla: technique and long-term results. *Scand J Plast Reconstr Surg Hand Surg* 2004;38:70-85.
- 98) Malevez C, Abarca M, Durdu F, Daelemans P. Clinical outcome of 103 consecutive zygomatic implants: 6-48 months follow-up study. *Clin Oral Impl Res* 2004;15:18-22.
- 99) Mastrogiacomo M, Scaglione S, Martinetti R, Dolcini L, Beltrame F, Cancedda R, Quarto R. Role of scaffold internal structure on in vivo bone formation in macroporous calcium phosphate bioceramics. *Biomaterials.* 2006;27(17):3230-7.
- 100) Egli PS, Müller W, Schenk RK. Porous hydroxyapatite and tricalcium phosphate cylinders with two different pore size ranges implanted in the cancellous bone of rabbits. A comparative histomorphometric and histologic study of bony ingrowth and implant substitution. *Clin Orthop Relat Res.* 1988;(232):127-38.
- 101) Pallesen L, Schou S, Aaboe M, Hjørting-Hansen E, Nattestad A, Melsen F. Influence of particle size of autogenous bone grafts on the early stages of bone regeneration: a histologic and stereologic study in rabbit calvarium. *Int J Oral Maxillofac Implants.* 2002;17(4):498-506.
- 102) Stenderup K, Justesen J, Clausen C, Kassem M. Aging is associated with decreased maximal life span and accelerated senescence of bone marrow stromal cells. *Bone.* 2003;33(6):919-26.
- 103) Springer IN, Terheyden H, Geiss S, Härle F, Hedderich J, Açil Y. Particulated bone grafts--effectiveness of bone cell supply. *Clin Oral Implants Res.* 2004;15(2):205-12.
- 104) Buser D, Hoffmann B, Bernard JP, Lussi A, Mettler D, Schenk RK. Evaluation of filling materials in membrane--protected bone defects. A comparative histomorphometric study in the mandible of miniature pigs. *Clin Oral Implants Res.* 1998;9(3):137-50.

- 105) De Marco AC, Jardini MA, Lima LP. Revascularization of autogenous block grafts with or without an e-PTFE membrane. *Int J Oral Maxillofac Implants*. 2005;20(6):867-74.
- 106) Burchardt H. The biology of bone graft repair. *Clin Orthop Relat Res*. 1983;(174):28-42.
- 107) Widmark G, Andersson B, Ivanoff CJ. Mandibular bone graft in the anterior maxilla for single-tooth implants. Presentation of surgical method. *Int J Oral Maxillofac Surg*. 1997;26(2):106-9.
- 108) Tomford WW. Bone allografts: past, present and future. *Cell Tissue Bank*. 2000;1(2):105-9.
- 109) Schwartz Z, Mellonig JT, Carnes DL Jr, de la Fontaine J, Cochran DL, Dean DD, Boyan BD. Ability of commercial demineralized freeze-dried bone allograft to induce new bone formation. *J Periodontol*. 1996;67(9):918-26.
- 110) Boyan BD, Ranly DM, McMillan J, Sunwoo M, Roche K, Schwartz Z. Osteoinductive ability of human allograft formulations. *J Periodontol*. 2006;77(9):1555-63.
- 111) Kolerman R, Tal H, Moses O. Histomorphometric analysis of newly formed bone after maxillary sinus floor augmentation using ground cortical bone allograft and internal collagen membrane. *J Periodontol*. 2008;79(11):2104-11.
- 112) European Association of Musculo Skeletal Transplantation: Common standard for musculo-skeletal Tissue Banking. European Association of Musculo Skeletal Transplantation, www.eamst.org; 1997, [date last accessed 15.02.08], 1997
- 113) Committee on Biological Implants Bone graft substitutes: facts, fiction and applications. American Academy of Orthopaedic Surgeons, 70th Annual Meeting, New Orleans, Louisiana. 2003
- 114) Commissione Nazionale per il Tessuto Muscoloscheletrico Linee Guida per il prelievo, la conservazione e l'utilizzo di tessuto muscoloscheletrico approvate dal Centro Nazionale Trapianti il 17/10/2001, dal Consiglio Superiore di Sanità il 12/12/2001 e dalla Conferenza Stato - Regioni il 21/3/2002.
- 115) Legge n. 91/1999 Disposizioni in materia di prelievi e di trapianti di organi e tessuti G.U. n. 87, 15 aprile 1999
- 116) Commissione Nazionale per il Tessuto Muscoloscheletrico Linee guida, ad uso delle banche di tessuto muscoloscheletrico, per la valutazione dell'appropriatezza della richiesta di osso umano e dei suoi derivati o sostituti., Novembre 2003.
- 117) Hjorting-Hansen E, Worsaae N, Lemons JE. Histologic response after implantation of porous hydroxylapatite ceramic in humans. *Int J Oral Maxillofac Implants*. 1990 Fall;5(3):255-63.
- 118) Schwartz Z, Weesner T, van Dijk S, Cochran DL, Mellonig JT, Lohmann CH, Carnes DL, Goldstein M, Dean DD, Boyan BD. Ability of deproteinized cancellous bovine bone to induce new bone formation. *J Periodontol*. 2000;71(8):1258-69.
- 119) Wenz B, Oesch B, Horst M. Analysis of the risk of transmitting bovine spongiform encephalopathy through bone grafts derived from bovine bone. *Biomaterials*. 2001;22(12):1599-606.

- 120) Mellonig JT. Donor selection, testing, and inactivation of the HIV virus in freeze-dried bone allografts. *Pract Periodontics Aesthet Dent*. 1995;7(6):13-22; quiz 23.
- 121) Taylor JC, Cuff SE, Leger JP, Morra A, Anderson GI. In vitro osteoclast resorption of bone substitute biomaterials used for implant site augmentation: a pilot study. *Int J Oral Maxillofac Implants*. 2002;17(3):321-30.
- 122) Piattelli M, Favero GA, Scarano A, Orsini G, Piattelli A. Bone reactions to anorganic bovine bone (Bio-Oss) used in sinus augmentation procedures: a histologic long-term report of 20 cases in humans. *Int J Oral Maxillofac Implants*. 1999;14(6):835-40.
- 123) Pripatnanont P, Nuntanaranont T, Vongvatcharanon S. Proportion of deproteinized bovine bone and autogenous bone affects bone formation in the treatment of calvarial defects in rabbits. *Int J Oral Maxillofac Surg*. 2009 Apr;38(4):356-62. Epub 2009 Mar 10.
- 124) LeGeros RZ, Lin S, Rohanizadeh R, Mijares D, LeGeros JP. Biphasic calcium phosphate bioceramics: preparation, properties and applications. *J Mater Sci Mater Med*. 2003 ;14(3):201-9.
- 125) Jensen SS, Yeo A, Dard M, Hunziker E, Schenk R, Buser D. Evaluation of a novel biphasic calcium phosphate in standardized bone defects: a histologic and histomorphometric study in the mandibles of minipigs. *Clin Oral Implants Res*. 2007;18(6):752-60.
- 126) Moura J, Teixeira LN, Ravagnani C, Peitl O, Zannotto ED, Beloti MM, et al.. In vitro osteogenesis on a highly bioactive glass-ceramic (Biosilicate®). *J Biomed Mater Res A* 2007;82:545-547.
- 127) Xynos ID, Edgar AJ, Buttery LD, Hench LL, Polak JM. Ionic products of bioactive glass dissolution increase proliferation of human osteoblasts and induce insulin-like growth factor II mRNA expression and protein synthesis. *Biochem Biophys Res Commun* 2000;276:461-465.
- 128) Thomas MV, Puleo DA, Al-Sabbagh M. Bioactive glass three decades on. *J Long Term Eff Med Implants*. 2005;15(6):585-97.
- 129) Marx RE, Carlson ER, Eichstaedt RM, Schimmele SR, Strauss JE, Georgeff KR. Platelet-rich plasma: Growth factor enhancement for bone grafts. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1998;85(6):638-46.
- 130) Marx RE. Platelet-rich plasma (PRP): what is PRP and what is not PRP? *Implant Dent*. 2001;10(4):225-8.
- 131) Sánchez AR, Sheridan PJ, Kupp LI. Is platelet-rich plasma the perfect enhancement factor? A current review. *Int J Oral Maxillofac Implants*. 2003;18(1):93-103.
- 132) Park EJ, Kim ES, Weber HP, Wright RF, Mooney DJ. Improved bone healing by angiogenic factor-enriched platelet-rich plasma and its synergistic enhancement by bone morphogenetic protein-2. *Int J Oral Maxillofac Implants*. 2008;23(5):818-26.
- 133) Schuckert KH, Jopp S, Osadnik M. Modern bone regeneration instead of bone transplantation: a combination of recombinant human bone morphogenetic protein-2 and platelet-rich plasma for the

- vertical augmentation of the maxillary bone-a single case report. *Tissue Eng Part C Methods*. 2010;16(6):1335-46.
- 134) Kaya Y, Yalim M, Bahçecitapar M, Baloş K .Comparison of applying particulate demineralized bone matrix (DBM), putty DBM and open flap debridement in periodontal horizontal bone defects. A 12-month longitudinal, multi-center, triple-blind, split-mouth, randomized, controlled clinical study. Part 1 - clinical and radiographic evaluation. *J Oral Rehabil*. 2009;36(7):524-34.
- 135) Caplanis N, Lee MB, Zimmerman GJ, Selvig KA, Wikesjö UM. Effect of allogenic freeze-dried demineralized bone matrix on guided tissue regeneration in dogs. *J Periodontol*. 1998;69(8):851-6.
- 136) Fox CF, Skalak R, eds. *Tissue Engineering*. New York: Alan R. Liss, Inc., (1st Tissue Engineering Symposium Proceedings Book). 1988
- 137) Ciapetti G, Granchi D, Baldini N. The combined use of mesenchymal stromal cells and scaffolds for bone repair. *Curr Pharm Des*. 2012 1;18(13):1796-820.
- 138) Bueno EM, Glowacki J. Cell-free and cell-based approaches for bone regeneration. *Nat Rev Rheumatol*. 2009;5(12):685-97.
- 139) Mastrogiacomo M, Muraglia A, Komlev V, Peyrin F, Rustichelli F, Crovace A, Cancedda R. Tissue engineering of bone: search for a better scaffold. *Orthod Craniofac Res*. 2005;8(4):277-84.
- 140) Yilgor P, Yilmaz G, Onal MB, Solmaz I, Gundogdu S, Keskil S, Sousa RA, Reis RL, Hasirci N, Hasirci V. An in vivo study on the effect of scaffold geometry and growth factor release on the healing of bone defects. *J Tissue Eng Regen Med*. 2012 Mar 7. doi: 10.1002/term.1456.
- 141) Schofer MD, Roessler PP, Schaefer J, Theisen C, Schlimme S, Heverhagen JT, Voelker M, Dersch R, Agarwal S, Fuchs-Winkelmann S, Paletta JR. Electrospun PLLA nanofiber scaffolds and their use in combination with BMP-2 for reconstruction of bone defects. *PLoS One*. 2011;6(9):e25462.
- 142) Liu G, Sun J, Li Y, Zhou H, Cui L, Liu W, Cao Y. Evaluation of partially demineralized osteoporotic cancellous bone matrix combined with human bone marrow stromal cells for tissue engineering: an in vitro and in vivo study. *Calcif Tissue Int*. 2008;83(3):176-85.
- 143) Sabatino M, Ren J, David-Ocampo V, England L, McGann M, Tran M, Kuznetsov SA, Khuu H, Balakumaran A, Klein HG, Robey PG, Stroncek DF. The establishment of a bank of stored clinical bone marrow stromal cell products. *J Transl Med*. 2012; 6;10:23.
- 144) Kuznetsov SA, Mankani MH, Robey PG. In vivo formation of bone and haematopoietic territories by transplanted human bone marrow stromal cells generated in medium with and without osteogenic supplements. *J Tissue Eng Regen Med*. 2011 Nov 3. doi: 10.1002/term.515.
- 145) Giardino R, Nicoli Aldini N, Fini M, Tanzi MC, Faré S, Draghi L, Carpi A, Nicolini A, Giavaresi G. Bioabsorbable scaffold for in situ bone regeneration. *Biomed Pharmacother*. 2006;60(8):386-92.
- 146) Sauerbier S, Rickert D, Gutwald R, Nagursky H, Oshima T, Xavier SP, Christmann J, Kurz P, Menne D, Vissink A, Raghoobar G, Schmelzeisen R, Wagner W, Koch FP. Bone marrow

- concentrate and bovine bone mineral for sinus floor augmentation: a controlled, randomized, single-blinded clinical and histological trial--per-protocol analysis. *Tissue Eng Part A*. 2011;17(17-18):2187-97.
- 147) Schmelzeisen R, Gutwald R, Oshima T, Nagursky H, Vogeler M, Sauerbier S. Making bone II: maxillary sinus augmentation with mononuclear cells--case report with a new clinical method. *Br J Oral Maxillofac Surg*. 2011;49(6):480-2.
- 148) Brighton CT. The biology of fracture repair. *Instr Course Lect*. 1984;33:60-82
- 149) Bianchi A. *Implantologia e implanto protesi*. UTET, 1999.
- 150) Goodship AE, Kenwright J. The influence of induced micromovement upon the healing of experimental tibial fractures. *J Bone Joint Surg Br*. 1987;67(4):650-5.
- 151) Dimitriou R, Tsiridis E, Giannoudis PV. Current concepts of molecular aspects of bone healing. *Injury*. 2005;36(12):1392-404
- 152) Gerstenfeld LC, Cullinane DM, Barnes GL, Graves DT, Einhorn TA. Fracture healing as a post-natal developmental process: molecular, spatial, and temporal aspects of its regulation. *J Cell Biochem*. 2003;1;88(5):873-84. Review
- 153) Gerstenfeld LC, Cho TJ, Kon T, Aizawa T, Tsay A, Fitch J, Barnes GL, Graves DT, Einhorn TA. Impaired fracture healing in the absence of TNF-alpha signaling: the role of TNF-alpha in endochondral cartilage resorption. *J Bone Miner Res*. 2003;18(9):1584-92.
- 154) Kon T, Cho TJ, Aizawa T, Yamazaki M, Nooh N, Graves D, Gerstenfeld LC, Einhorn TA. Expression of osteoprotegerin, receptor activator of NF-kappaB ligand (osteoprotegerin ligand) and related proinflammatory cytokines during fracture healing. *J Bone Miner Res*. 2001;16(6):1004-14.
- 155) Barnes GL, Kostenuik LC, Gerstenfeld LC, Einhorn TA. Growth factor regulation of fracture repair. *J Bone Miner Res* 1999;14:1805-15
- 156) Ten Dijke P, Fu J, Schaap P, Roelen BA. Signal transduction of bone morphogenetic proteins in osteoblast differentiation. *J Bone Joint Surg* 2003;85(Suppl. 3):34-8.
- 157) Smith WC, Harland RM. Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell* 1992;70:829-40.
- 158) Massague J, Attisano L, Wrana JL. The TGF-beta family and its composite receptors. *Trends Cell Biol* 1994;4:172-8.
- 159) Cho TJ, Gerstenfeld LC, Einhorn TA. Differential temporal expression of members of the transforming growth factor beta superfamily during murine fracture healing. *J Bone Miner Res* 2002;17:513-20.
- 160) Sakou T. Bone morphogenetic proteins: from basic studies to clinical approaches. *Bone* 1998;22:591-603

- 161) Israel DI, Nove J, Kerns KM, Kaufman RJ, Rosen V, Cox KA, Wozney JM. Heterodimeric bone morphogenetic proteins show enhanced activity in vitro and in vivo. *Growth Factors*. 1996;13(3-4):291-300.
- 162) Bostrom MP. Expression of bone morphogenetic proteins in fracture healing. *Clin Orthop* 1998;355S:116-23.
- 163) Cheng H, Jiang W, Phillips FM, Haydon RC, Peng Y, Zhou L, Luu HH, An N, Breyer B, Vanichakarn P, Szatkowski JP, Park JY, He TC. Osteogenic activity of the fourteen types of human bone morphogenetic proteins (BMPs). *J Bone Joint Surg Am*. 2003;85-A(8):1544-52.
- 164) Deckers MM, van Bezooijen RL, van der Horst G, Hoogendam J, van Der Bent C, Papapoulos SE, Löwik CW. Bone morphogenetic proteins stimulate angiogenesis through osteoblast-derived vascular endothelial growth factor A. *Endocrinology*. 2002;143(4):1545-53.
- 165) Andrew JG, Hoyland JA, Freemont AJ, Marsh DR. Platelet-derived growth factor expression in normally healing human fractures. *Bone* 1995;16:455-60.
- 166) Solheim E. Growth factors in bone. *Int Orthop* 1998;22:410-6.
- 167) Canalis E. Effect of insulin-like growth factor I on DNA and protein synthesis in cultured rat calvaria. *J Clin Invest* 1980;66:709-19.
- 168) Shimasaki S, Ling N. Identification and molecular characterization of insulin-like growth factor binding proteins (IGFBP- 1, -2, -3, -4, -5 and -6). *Prog Growth Factor Res* 1991;3:243-66.
- 169) Nakamura T, Hara Y, Tagawa M, Tamura M, Yuge T, Fukuda H, Nigi H. Recombinant human basic fibroblast growth factor accelerates fracture healing by enhancing callus remodelling in experimental dog tibial fractures. *J Bone Miner Res*. 1998;13(6):942-9.
- 170) Andreassen TT, Oxlund H. Local anabolic effects of growth hormone on intact bone and healing fractures in rats. *Calcif Tissue Int* 2003;73:258-64.
- 171) Street J, Bao M, deGuzman L, Bunting S, Peale FV Jr, Ferrara N, Steinmetz H, Hoeffel J, Cleland JL, Daugherty A, van Bruggen N, Redmond HP, Carano RA, Filvaroff EH. Vascular endothelial growth factor stimulates bone repair by promoting angiogenesis and bone turnover. *Proc Natl Acad Sci U S A*. 2002;23;99(15):9656-61
- 172) Nyman S, Lindhe J, Karring T, Rylander H. New attachment following surgical treatment of human periodontal disease. *J Clin Periodontol* 1982;9:290-296.
- 173) Dahlin C, Sennerby L, Lekholm U, et al. Generation of new bone around titanium implants using a membrane technique: An experimental study in rabbits. *Int J Oral Maxillofac Implants* 1989;4:19-25.
- 174) Buser D, Dula K, Hirt HP, Schenk RK. Lateral ridge augmentation using autografts and barrier membranes: A clinical study with 40 partially edentulous. *J Oral Maxillofac Surg*. 1996 Apr;54(4):420-32.
- 175) Simion M, Trisi P, Piattelli A. Vertical ridge augmentation using a membrane technique associated with osseointegrated implants. *Int J Periodontics Restorative Dent* 1994;14:496-511.

- 176)Tinti C, Parma-Benfenati S, Polizzi G. Vertical ridge augmentation: What is the limit? *Int J Periodontics Restorative Dent* 1996;16:220-229.
- 177)Mellonig JT, Nevins M. Guided bone regeneration of bone defects associated with implants: An evidencebased outcome assessment. *Int J Periodontics Restorative Dent* 1995;15:168-185.
- 178)Buser D, Dula K, Belser UC, Hirt HP, Berthold H. Localized ridge augmentation using guided bone regeneration. II. Surgical procedure in the mandible. *Int J Periodontics Restorative Dent* 1995;15:10-29
- 179)Machtei EE. The effect of membrane exposure on the outcome of regenerative procedures in humans: A meta-analysis. *J Periodontol* 2001;72:512-516.
- 180)Simion M, Baldoni M, Rossi P, Zaffe D. A comparative study of the effectiveness of e-PTFE membranes with and without early exposure during the healing period. *Int J Periodontics Restorative Dent* 1994;14: 166-180.
- 181)Owens KW, Yukna RA. Collagen membrane resorption in dogs: A comparative study. *Implant Dent* 2001; 10:49-58.
- 182)Zhao S, Pinholt EM, Madsen JE, Donath K. Histological evaluation of different biodegradable and nonbiodegradable membranes implanted subcutaneously in rats. *J Craniomaxillofac Surg* 2000;28:116-122.
- 183)Marinucci L, Lilli C, Baroni T, Becchetti E, Belcastro S, Balducci C, Locci P. In vitro comparison of bioabsorbable and non-resorbable membranes in bone regeneration. *J Periodontol* 2001;72: 753-759.
- 184)Hurzeler MB, Quinones CR, Schupbach P. Guided bone regeneration around dental implants in the atrophic alveolar ridge using a bioresorbable barrier. An experimental study in the monkey. *Clin Oral Implants Res* 1997;8:323-331.
- 185)Wang HL, O'Neal RB, Thomas CL, et al. Evaluation of an absorbable collagen membrane in treating Class II furcation defects. *J Periodontol* 1994;65(11):1029-36
- 186)Lundgren D, Sennerby L, Falk H, et al. The use of a new bioresorbable barrier for guided bone regeneration in connection with implant installation. Case reports. *Clin Oral Implants Res* 1994;5:177-184.
- 187)Polson AM, Garrett S, Stoller NH, et al. Guided tissue regeneration in human furcation defects after using a biodegradable barrier: A multi-center feasibility study. *J Periodontol* 1995;66:377-385.
- 188)Hutmacher D, Hürzeler MB, Schliephake H. A review of material properties of biodegradable and bioresorbable polymers and devices for GTR and GBR applications. *Int J Oral Maxillofac Implants*. 1996;11(5):667-78.
- 189)Vert M, Li S, Garreau H. New insights on the degradation of bioresorbable polymeric devices based on lactic and glycolic acids. *Clin Mater*. 1992;10(1-2):3-8.

- 190) Therin M, Christel P, Li S, Garreau H, Vert M. In vivo degradation of massive poly(alpha-hydroxy acids): validation of in vitro findings. *Biomaterials*. 1992;13(9):594-600.
- 191) Postlethwaite AE, Seyer JM, Kang AH. Chemotactic attraction of human fibroblasts to type I, II, and III collagens and collagen-derived peptides. *Proc Natl Acad Sci U S A*. 1978;75(2):871-5.
- 192) Locci P, Calvitti M, Belcastro S, Pugliese M, Guerra M, Marinucci L, Staffolani N, Becchetti E. Phenotype expression of gingival fibroblasts cultured on membranes used in guided tissue regeneration. *J Periodontol*. 1997;68(9):857-63.
- 193) Schlegel AK, Möhler H, Busch F, Mehl A. Preclinical and clinical studies of a collagen membrane (Bio-Gide). *Biomaterials*. 1997;18(7):535-8.
- 194) Rothamel D, Schwarz F, Sculean A, Herten M, Scherbaum W, Becker J. Biocompatibility of various collagen membranes in cultures of human PDL fibroblasts and human osteoblast-like cells. *Clin Oral Implants Res*. 2004;15(4):443-9.
- 195) Pitaru S, Tal H, Soldinger M, Noff M. Collagen membranes prevent apical migration of epithelium and support new connective tissue attachment during periodontal wound healing in dogs. *J Periodontal Res*. 1989;24(4):247-53.
- 196) Hürzeler MB, Kohal RJ, Naghshbandi J, Mota LF, Conradt J, Hutmacher D, Caffesse RG. Evaluation of a new bioresorbable barrier to facilitate guided bone regeneration around exposed implant threads. An experimental study in the monkey. *Int J Oral Maxillofac Surg*. 1998;27(4):315-20.
- 197) Miller N, Penaud J, Foliguet B, Membre H, Ambrosini P, Plombas M. Resorption rates of 2 commercially available bioresorbable membranes. A histomorphometric study in a rabbit model. *J Clin Periodontol*. 199;23(12):1051-9.
- 198) Minabe M, Kodama T, Hori T, Watanabe Y. Effects of atelocollagen on the wound healing reaction following palatal gingivectomy in rats. *J Periodontal Res*. 1989;24(3):178-85.
- 199) Quteish D, Dolby AE. The use of irradiated-crosslinked human collagen membrane in guided tissue regeneration. *J Clin Periodontol*. 1992;19(7):476-84.
- 200) von Arx T, Brogini N, Jensen SS, Bornstein MM, Schenk RK, Buser D. Membrane durability and tissue response of different bioresorbable barrier membranes: a histologic study in the rabbit calvarium. *Int J Oral Maxillofac Implants*. 2005;20(6):843-53.
- 201) Hutmacher DW, Kirsch A, Ackermann KL, et al. A tissue engineered cell-occlusive device for hard tissue regeneration – A preliminary report. *Int J Periodontics Restorative Dent* 2001;21:49-59.
- 202) Zitzmann NU, Naef R, Scharer P. Resorbable versus nonresorbable membranes in combination with Bio-Oss for guided bone regeneration. *Int J Oral Maxillofac Implants* 1997;12:844-852.
- 203) Jovanovic SA. Protected space development for bone formation using reinforced barrier membranes In: Nevins M, Mellonig JT, eds. *Implant Therapy: Clinical Approaches and Evidence of Success*. Chicago: Quintessence Publishing; 1998:91-98.

- 204) Hurzeler MB, Strub JR. Guided bone regeneration around exposed implants: A new bioresorbable device and bioresorbable membrane pins. *Pract Periodontics Aesthet Dent* 1995;7:37-47
- 205) Simion M, Jovanovic SA, Tinti C, et al. Long-term evaluation of osseointegrated implants inserted at the time or after vertical ridge augmentation. A retrospective study on 123 implants with 1-5 year follow-up. *Clin Oral Implants Res* 2001;12:35-45.
- 206) Urist MR, Strates BS. Bone morphogenetic protein. *J Dent Res*. 1971;50(6):1392-406.
- 207) Jovanovic SA, Hunt DR, Bernard GW, Spiekermann H, Nishimura R, Wozney JM, Wikesjö UM. Long-term functional loading of dental implants in rhBMP-2 induced bone. A histologic study in the canine ridge augmentation model. *Clin Oral Implants Res*. 2003;14(6):793-803.
- 208) Jones AA, Buser D, Schenk R, Wozney J, Cochran DL. The effect of rhBMP-2 around endosseous implants with and without membranes in the canine model. *J Periodontol*. 2006;77(7):1184-93.
- 209) Hanisch O, Sorensen RG, Kinoshita A, Spiekermann H, Wozney JM, Wikesjö UM. Effect of recombinant human bone morphogenetic protein-2 in dehiscence defects with non-submerged immediate implants: an experimental study in *Cynomolgus* monkeys. *J Periodontol*. 2003;74(5):648-57.
- 210) Jung RE, Glauser R, Schärer P, Hämmerle CH, Sailer HF, Weber FE. Effect of rhBMP-2 on guided bone regeneration in humans. *Clin Oral Implants Res*. 2003;14(5):556-68.
- 211) Warnke PH, Springer IN, Wiltfang J, Acil Y, Eufinger H, Wehmöller M, Russo PA, Bolte H, Sherry E, Behrens E, Terheyden H. Growth and transplantation of a custom vascularised bone graft in a man. *Lancet*. 2004;364(9436):766-70
- 212) Gautschi OP, Frey SP, Zellweger R. Bone morphogenetic proteins in clinical applications. *ANZ J Surg*. 2007;77(8):626-31.
- 213) Meijndert L, Raghoobar GM, Schüpbach P, Meijer HJ, Vissink A. Bone quality at the implant site after reconstruction of a local defect of the maxillary anterior ridge with chin bone or deproteinised cancellous bovine bone. *Int J Oral Maxillofac Surg*. 2005;34(8):877-84.
- 214) Chiapasco M, Abati S, Romeo E, Vogel G. Clinical outcome of autogenous bone blocks or guided bone regeneration with e-PTFE membranes for the reconstruction of narrow edentulous ridges. *Clin Oral Implants Res*. 1999;10(4):278-88.
- 215) Breine U, Brånemark PI. Reconstruction of alveolar jaw bone. An experimental and clinical study of immediate and preformed autologous bone grafts in combination with osseointegrated implants. *Scand J Plast Reconstr Surg*. 1980;14(1):23-48.
- 216) Joshi A, Kostakis GC. An investigation of post-operative morbidity following iliac crest graft harvesting. *Br Dent J*. 2004 14;196(3):167-71.
- 217) Canady JW, Zeitler DP, Thompson SA, Nicholas CD. Suitability of the iliac crest as a site for harvest of autogenous bone grafts. *Cleft Palate Craniofac J*. 1993;30(6):579-81

- 218) Acocella A, Sacco R, Niardi P, Agostini T. Early implant placement in bilateral sinus floor augmentation using iliac bone block grafts in severe maxillary atrophy: a clinical, histological, and radiographic case report. *J Oral Implantol.* 2009;35(1):37-44.
- 219) Johansson B, Grepe A, Wannfors K, Hirsch JM. A clinical study of changes in the volume of bone grafts in the atrophic maxilla. *Dentomaxillofac Radiol.* 2001;30(3):157-61.
- 220) Misch CM, Misch CE, Resnik R, et al. Reconstruction of maxillary alveolar defects with mandibular symphysis grafts for dental implants: A preliminary procedural report. *Int J Oral Maxillofac Implants.* 1992; 7:360-366.
- 221) Misch CM, Misch CE. The repair of localized severe ridge defects for implant placement using mandibular bone grafts. *Implant Dent.* 1995; 4:261-267
- 222) Misch CM. Comparison of intraoral donor sites for onlay grafting prior to implant placement. *Int J Oral Maxillofac Implants.* 1997; 12:767-776.
- 223) Collins TA, Nunn W. Autogenous veneer grafting for improved esthetics with dental implants. *Compend Contin Educ Dent.* 1994; 15:370-376.
- 224) Sindet-Pedersen S, Enemark H. Reconstruction of alveolar clefts with mandibular or iliac crest bone grafts: A comparative study. *J Oral Maxillofac Surg.* 1990; 48: 554-558.
- 225) Picos, MA. Buccolingual expansion of the maxillary ridge. *Dent Implantol Update.* 1992; 3:85-87.
- 226) Picos, MA. Facilitating implant placement with chin grafts as donor sites for maxillary bone augmentation; Part I. *Dent Implantol Update.* 1995; 6:89-92.
- 227) Picos, MA. Facilitating implant placement with chin grafts as donor sites for maxillary bone augmentation: Part II. *Dent Implantol Update.* 1996; 7:1-4.
- 228) Picos, MA. Alveolar ridge augmentation with ramus buccal shelf autografts and impacted third molar removal. *Dent Implantol Update.* 1999; 4:27-31.
- 229) Misch CM. Ridge augmentation using mandibular ramus bone grafts for the placement of dental implants: Presentation of a technique. *Pract Periodontics Aesthet Dent.* 1996; 8:127-135.
- 230) Perry T. Ascending ramus offered as alternate harvest site for onlay bone grafting. *Dent Implantol Update.* 1997; 3:21-24.
- 231) Tessier P. Autogenous bone grafts taken from the calvarium for facial and cranial application. *Clin Plast Surg* 1982;9:531-8
- 232) Lenzen C, Meiss A, Bull HG. Augmentation of the extremely atrophied maxilla and mandible by autologous calvarial bone transplantation. *Mund Kiefer Gesichtschir.* 1999 ;3 Suppl 1:S40-2
- 233) Le Lorc'h-Bukiet I, Tulasne JF, Llorens A, Lesclous P. Parietal bone as graft material for maxillary sinus elevation: structure and remodelling of the donor and recipient sites. *Clin Oral Implants Res* 2005;16:244-49

- 234) Iizuka T, Smolka W, Hallermann W, Merickske-Stern R. Extensive augmentation of the alveolar ridge using autogenous calvarial split bone grafts for dental rehabilitation. *Clin Oral Implants Res* 2004;15:607-15
- 235) Smolka W, Eggensperger N, Carollo V, Ozdoba C, Iizuka T. Changes in the volume and density of calvarial split bone grafts after alveolar ridge augmentation. *Clin Oral Implants Res*. 2006;17(2):149-55
- 236) Smith JD, Abramson M. Membranous vs. endochondral bone autografts. *Arch Otolaryngol*. 1974; 99:203-210.
- 237) Zins JE, Whitaker LA. Membranous vs. endochondral bone autografts: Implications for craniofacial reconstruction. *Plast Reconstr Surg*. 1983; 72:778-786.
- 238) Hardesty RA, Marsh JL. Craniofacial onlay bone grafting: a prospective evaluation of graft morphology, orientation, and embryonic origin. *Plast Reconstr Surg*. 1990;85(1):5-14.
- 239) Yang B, Zhao M, Xiong B. Histological study and stereologic analysis of membranous versus endochondral onlay bone grafts. *Zhonghua Zheng Xing Shao Shang Wai Ke Za Zhi* 1999;15(4):280-2.
- 240) Finkelman RD, Eason AL, Rakijian DR, Tutundzhyan Y, Hardesty RA. Elevated IGF-II and TGF-beta concentrations in human calvarial bone: potential mechanism for increased graft survival and resistance to osteoporosis. *Plast Reconstr Surg*. 1994;93(4):732-8
- 241) Proussaefs P, Lozada J. The use of intraorally harvested autogenous block grafts for vertical alveolar ridge augmentation: a human study. *Int J Periodontics Restorative Dent*. 2005;25(4):351-63.
- 242) Von Arx T, Buser D. Horizontal ridge augmentation using autogenous block grafts and the guided bone regeneration technique with collagen membranes: a clinical study with 42 patients. *Clin Oral Implants Res*. 2006;17(4):359-66.
- 243) Antoun H, Sitbon JM, Martinez H, Missika P. A prospective randomized study comparing two techniques of bone augmentation: onlay graft alone or associated with a membrane. *Clin Oral Implants Res*. 2001;12(6):632-9.
- 244) Donos N, Kostopoulos L, Karring T. Augmentation of the rat jaw with autogeneic cortico-cancellous bone grafts and guided tissue regeneration. *Clin Oral Implants Res*. 2002;13(2):192-202
- 245) Verdugo F, D'Addona A, Pontón J. Clinical, tomographic, and histological assessment of periosteal guided bone regeneration with cortical perforations in advanced human critical size defects. *Clin Implant Dent Relat Res*. 2012;14(1):112-20.
- 246) Cha JK, Kim CS, Choi SH, Cho KS, Chai JK, Jung UW The influence of perforating the autogenous block bone and the recipient bed in dogs. Part II: histologic analysis. *Clin Oral Implants Res*. 2011 Aug 3. doi: 10.1111/j.1600-0501.2011.02259.x. [Epub ahead of print]

- 247) Adeyemo WL, Reuther T, Bloch W, Korkmaz Y, Fischer JH, Zöller JE, Kuebler AC. Influence of host periosteum and recipient bed perforation on the healing of onlay mandibular bone graft: an experimental pilot study in the sheep. *Oral Maxillofac Surg.* 2008;12(1):19-28
- 248) Burchardt H. The biology of bone graft repair. *Clin Orthop Relat Res* 1983;(174):28-42.
- 249) Enneking WF, Eady JL, Burchardt H. Autogenous cortical bone grafts in the reconstruction of segmental skeletal defects. *J Bone Joint Surg Am* 1980;62: 1039-1058.
- 250) Matsumoto MA, Filho HN, Francischone E, Consolaro A. Microscopic analysis of reconstructed maxillary alveolar ridges using autogenous bone grafts from the chin and iliac crest. *Int J Oral Maxillofac Implants* 2002;17:507-516.
- 251) Proussaefs P, Lozada J, Rohrer MD. A clinical and histologic evaluation of a block onlay graft in conjunction with autogenous particulate and inorganic bovine mineral (Bio-Oss): A case report. *Int J Periodontics Restorative Dent* 2002;22:567-573
- 252) Donovan MG, Dickerson NC, Hanson LJ, Gustafson RB. Maxillary and mandibular reconstruction using calvarial bone grafts and Branemark implants: a preliminary report. *J Oral Maxillofac Surg.*1994;52:588-94.
- 253) Chen NT, Glowacki J, Bucky LP, Hong HZ, Kim WK, Yaremchuk. MJ. The roles of revascularization and resorption on endurance of craniofacial onlay bone grafts in the rabbit. *Plast Reconstr Surg.* 1994;93:714-22
- 254) Iturriaga MT, Ruiz CC. Maxillary sinus reconstruction with calvarium bone grafts and endosseous implants. *J Oral Maxillofac Surg.* 2004;62:344-7.
- 255) Kline RM Jr, Wolfe SA. Complications associated with the harvesting of cranial bone grafts. *Plast Reconstr Surg.* 1995;95:5-13.
- 256) Keith JD Jr. Localized ridge augmentation with a block allograft followed by secondary implant placement: A case report. *Int J Periodontics Restorative Dent* 2004;24:11-17.
- 257) Leonetti JA, Koup R. Localized maxillary ridge augmentation with a block allograft for dental implant placement: Case reports. *Implant Dent* 2003;12:217-224.
- 258) Lyford RH, Mills MP, Knapp CI, et al. Clinical evaluation of freeze-dried block allografts for alveolar ridge augmentation: A case series. *Int J Periodontics Restorative Dent* 2003;23:417-425.
- 259) Simion M, Nevins M, Rocchietta I, Fontana F, Maschera E, Schupbach P, Kim DM. Vertical ridge augmentation using an equine block infused with recombinant human platelet-derived growth factor-BB: a histologic study in a canine model. *Int J Periodontics Restorative Dent.* 2009;29(3):245-55.
- 260) Linkow LI "Maxillary implants. A dynamic approach to oral implantology." North Haven, CT: Glarus, 1997:109-111
- 261) Tatum H, Jr. Maxillary and sinus implant reconstructions. *Dent Clin North Am* 1986;30:207-229.
- 262) Boyne PJ, James RA. Grafting of the maxillary sinus floor with autogenous marrow and bone. *J Oral Surg.* 1980;38(8):613-6.

- 263) Jensen OT, Shulman LB, Block MS, Iacono VJ. Report of the Sinus Consensus Conference of 1996. *Int J Oral Maxillofac Implants* 1998;13.
- 264) Acocella A, Sacco R, Nardi P, Agostini T: Early implant placement in bilateral sinus floor augmentation using iliac bone block grafts in severe maxillary atrophy: a clinical, histological and radiographic case report. *Journal of Oral Implantology* 2009;35(1): 37-44
- 265) Acocella A, Sacco R, Nardi P, Agostini T: Simultaneous implant placement in sinus floor augmentation using iliac bone block graft in severe maxillary atrophies: Case Report. *Implant Dent* 2008;17(4):382-388
- 266) Wallace SS, Froum SJ. Effect of maxillary sinus augmentation on the survival of endosseous dental implants. A systematic review. *Ann Periodontol.* 2003; 8: 328-343.
- 267) Aghaloo TL, Moy PK. Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement? *Int J Oral Maxillofac Implants.* 2007;22
- 268) Del Fabbro M, Testori T, Francetti L, et al. Systematic review of survival rates for implants placed in the grafted maxillary sinus. *Int J Periodontics Restorative Dent.* 2004; 24:565-577.
- 269) Wannfors K, Johansson B, Hallman M, Strandkvist T. A prospective randomized study of 1- and 2-stage sinus inlay bone grafts: 1-year follow-up. *Int J Oral Maxillofac Implants* 2000; 15:625-632.
- 270) Blomqvist JE, Alberius P, Isaksson S. Two-stage maxillary sinus reconstruction with endosseous implants: A prospective study. *Int J Oral Maxillofac Implants* 1998; 13:758-766.
- 271) Kahnberg K-E, Ekestubbe A, Grondahl K, Nilsson P, Hirsch J-M. Sinus lifting procedure. One-stage surgery with bone transplant and implants. *Clin Oral Impl Res* 2001; 12:4 79-487.
- 272) Johansson B, Wannfors K, Ekenback J, Smedberg J-1, Hirsch J. Implants and sinus inlay bone grafts in a 1-stage procedure on severely atrophied maxillae: Surgical aspects of a 3-year follow-up study. *Int J Oral Maxillofac Implants* 1999;14:811-818.
- 273) Keller EE, Tolman DE, Eckert SE. Maxillary antral-nasal autogenous bone graft reconstruction of the compromised maxilla: A 12-year retrospective study. *Int J Oral Maxillofac Implants* 1999;14:707-721.
- 274) Blomqvist JE, Alberius P, Isaksson S. retrospective analysis of one-stage maxillary sinus augmentation with endosseous implants. *Int J Oral Maxillofac Implants* 1996;11:512-521.
- 275) Froum SJ, Tarnow DP, Wallace SS, Rohrer MD, Cho S-C. Sinus floor elevation using anorganic bovine bone matrix (OsteoGraf/N) with and without autogenous bone: A clinical, histologic, radiographic, and histomorphometric analysis-Part 2 of an ongoing prospective study. *Int J Periodontics Restorative Dent* 1998; 18:529-543.
- 276) Hising P, Bolin A, Branting C. Reconstruction of severely resorbed alveolar crests with dental implants using a bovine bone mineral for augmentation. *Int J Oral Maxillofac Implants* 2001; 16:90-97.

- 277) Boyne PJ, Leslie C, Lilly BSN, et al. De novo bone induction by recombinant human bone morphogenic protein-2 (rh BMP-2) in maxillary sinus floor augmentation. *J Oral Surg* Accepted for publication 2004.
- 278) Rodriguez A, Anastassov GE, Lee H, Buchbinder D, Wettan H. Maxillary sinus augmentation with deproteinated bovine bone and platelet rich plasma with simultaneous insertion of endosseous implants. *J Oral Maxillofac Surg* 2003;61 :157-163.
- 279) Froum SJ, Wallace SS, Tarnow DP, Cho S-C. Effect of platelet-rich plasma on bone growth and osseointegration in human maxillary sinus grafts: Three bilateral case reports. *Int J Periodontics Restorative Dent* 2002;22:45-53.
- 280) Wiltfang J, Schlegel KA, Shultze-Mosgau S, Nkenke E, Zimmerman R, Kessler P. Sinus floor augmentation with beta-tricalcium phosphate (beta-TCP): Does platelet-rich plasma promote its osseous integration and degradation? *Clin Oral Imp/ Res* 2003;14:213-218.
- 281) Zuffetti F, Testori T, Fontanella W, et al. Use of platelet rich plasma in maxillary sinus lift. Preliminary histological results. *Clin Oral Impl Res* 2002;13(4):xli (Abst. 10).
- 282) Maiorana C, Sommariva L, Brivio P, Sigurta D, Santoro F. Maxillary sinus augmentation with anorganic bovine bone (Bio-Oss) and autologous platelet-rich plasma: Preliminary clinical and histologic evaluations. *Int J Periodontics Restorative Dent* 2003;23:227 -235.
- 283) Valentini P, Abensur D, Wenz B, Peetz M, Schenk R. Sinus grafting with porous bone mineral (Bio-Oss) for implant placement: A 5-year study on 15 patients. *Int J Periodontics Restorative Dent* 2000;20:245-253.
- 284) Sanchez AR, Sheridan PJ, Kupp LI. Is platelet-rich plasma the perfect enhancement factor? A current review. *Int J Oral Maxillofac Implants* 2003; 18:93-103.
- 285) Merckx MAW, Maltha JC, Stoelinga PJW. Assessment of the value of anorganic bone additives in sinus floor augmentation: A review of clinical reports. *Int J Oral Maxillofac Surg* 2003;32: 1-6.
- 286) Wallace SS, Froum SJ, Tarnow DP. Histologic evaluation of sinus elevation procedure. A clinical report. *Int J Periodontics Restorative Dent* 1996; 16:4 7-51.
- 287) Tadjoeidin ES, de Lange GL, Holzmann PJ, Kuiper L, Burger EH. Histological observations on biopsies harvested following sinus floor elevation using a bioactive glass material of narrow size range. *Clin Oral Impl Res* 2000; 11:334-344.
- 288) Valentini P, Abensur D, Densari D, Graziani JN, Hammerle CHF. Histological evaluation of Bio-Oss® in a 2-stage sinus floor elevation and implantation procedure. A human case report. *Clin Oral Impl Res* 1998;9:59-64.
- 289) Rosenlicht J, Tarnow DP. Human histologic evidence of functionally loaded hydroxyapatite-coated implants placed simultaneously with sinus augmentation: A case report 2 1/2 years post-placement. *Int J Oral Implantol* 1999;25:7-10.
- 290) Klein MO, Al-Nawas B. For which clinical indications in dental implantology is the use of bone substitute materials scientifically substantiated? *Eur J Oral Implantol*. 2011;4(5):11-29

- 291) Aguirre Zorzano LA, Rodríguez Tojo MJ, Aguirre Urizar JM. Maxillary sinus lift with intraoral autologous bone and β -tricalcium phosphate: histological and histomorphometric clinical study. *Med Oral Patol Oral Cir Bucal*. 2007;1;12(7):E532-6.
- 292) Artzi Z, Nemcovsky CE, Tal H, Dayan D. Histopathological morphometric evaluation of 2 different hydroxyapatite-bone derivatives in sinus augmentation procedures: a comparative study in humans. *J Periodontol*. 2001;72(7):911-20
- 293) Cordaro L, Bosshardt DD, Palattella P, Rao W, Serino G, Chiapasco M. Maxillary sinus grafting with Bio-Oss or Straumann Bone Ceramic: histomorphometric results from a randomized controlled multicenter clinical trial. *Clin Oral Implants Res*. 2008;19(8):796-803.
- 294) Crespi R, Mariani E, Benasciutti E, Capparè P, Cenci S, Gherlone E. Magnesium-enriched hydroxyapatite versus autologous bone in maxillary sinus grafting: combining histomorphometry with osteoblast gene expression profiles ex vivo. *J Periodontol*. 2009;80(4):586-93.
- 295) Felice P, Scarano A, Pistilli R, Checchi L, Piattelli M, Pellegrino G, Esposito M. A comparison of two techniques to augment maxillary sinuses using the lateral window approach: rigid synthetic resorbable barriers versus anorganic bovine bone. Five-month post-loading clinical and histological results of a pilot randomised controlled clinical trial. *Eur J Oral Implantol*. 2009;2(4):293-306.
- 296) Fugazzotto PA. Augmentation of the posterior maxilla: a proposed hierarchy of treatment selection. *J Periodontol*. 2003;74(11):1682-91.
- 297) Galindo-Moreno P, Avila G, Fernández-Barbero JE, Mesa F, O'Valle-Ravassa F, Wang HL. Clinical and histologic comparison of two different composite grafts for sinus augmentation: a pilot clinical trial. *Clin Oral Implants Res*. 2008;19(8):755-9.
- 298) Galindo-Moreno P, Moreno-Riestra I, Avila G, Fernández-Barbero JE, Mesa F, Aguilar M, Wang HL, O'Valle F. Histomorphometric comparison of maxillary pristine bone and composite bone graft biopsies obtained after sinus augmentation. *Clin Oral Implants Res*. 2010;21(1):122-8.
- 299) Hallman M, Sennerby L, Lundgren S. A clinical and histologic evaluation of implant integration in the posterior maxilla after sinus floor augmentation with autogenous bone, bovine hydroxyapatite, or a 20:80 mixture. *Int J Oral Maxillofac Implants*. 2002;17(5):635-43.
- 300) John HD, Wenz B. Histomorphometric analysis of natural bone mineral for maxillary sinus augmentation. *Int J Oral Maxillofac Implants*. 2004;19(2):199-207.
- 301) Lindgren C, Sennerby L, Mordenfeld A, Hallman M. Clinical histology of microimplants placed in two different biomaterials. *Int J Oral Maxillofac Implants*. 2009;24(6):1093-100.
- 302) Mangano C, Scarano A, Iezzi G, Orsini G, Perrotti V, Mangano F, Montini S, Piccirilli M, Piattelli A. Maxillary sinus augmentation using an engineered porous hydroxyapatite: a clinical, histological, and transmission electron microscopy study in man. *J Oral Implantol*. 2006;32(3):122-31.

- 303) Mangano C, Scarano A, Perrotti V, Iezzi G, Piattelli A. Maxillary sinus augmentation with a porous synthetic hydroxyapatite and bovine-derived hydroxyapatite: a comparative clinical and histologic study. *Int J Oral Maxillofac Implants*. 2007c;22(6):980-6.
- 304) Minichetti JC, D'Amore JC, Hong AY. Three-year analysis of tapered screw vent implants placed into maxillary sinuses grafted with mineralized bone allograft. *J Oral Implantol*. 2008;34(3):135-41.
- 305) Scarano A, Degidi M, Iezzi G, Pecora G, Piattelli M, Orsini G, Caputi S, Perrotti V, Mangano C, Piattelli A. Maxillary sinus augmentation with different biomaterials: a comparative histologic and histomorphometric study in man. *Implant Dent*. 2006;15(2):197-207.
- 306) Simunek A, Cierny M, Kopecka D, Kohout A, Bukac J, Vahalova D. The sinus lift with phycogenic bone substitute. A histomorphometric study. *Clin Oral Implants Res*. 2005;16(3):342-8.
- 307) Szabó G, Huys L, Coulthard P, Maiorana C, Garagiola U, Barabás J, Németh Z, Hrabák K, Suba Z. A prospective multicenter randomized clinical trial of autogenous bone versus beta-tricalcium phosphate graft alone for bilateral sinus elevation: histologic and histomorphometric evaluation. *Int J Oral Maxillofac Implants*. 2005;20(3):371-81.
- 308) Tadjoeidin ES, de Lange GL, Lyaruu DM, Kuiper L, Burger EH. High concentrations of bioactive glass material (BioGran) vs. autogenous bone for sinus floor elevation. *Clin Oral Implants Res*. 2002;13(4):428-36.
- 309) Turunen T, Peltola J, Makkonen T, Helenius H, Yli-Urpo A. Bioactive glass granules and polytetrafluoroethylene membrane in the repair of bone defects adjacent to titanium and bioactive glass implants. *J Mater Sci Mater Med*. 1998;9(7):403-7.
- 310) De Vicente JC, Hernández-Vallejo G, Braña-Abascal P, Peña I. Maxillary sinus augmentation with autologous bone harvested from the lateral maxillary wall combined with bovine-derived hydroxyapatite: clinical and histologic observations. *Clin Oral Implants Res*. 2010; 1;21(4):430-8.
- 311) Wallace SS, Froum SJ, Cho SC, Elian N, Monteiro D, Kim BS, Tarnow DP. Sinus augmentation utilizing anorganic bovine bone (Bio-Oss) with absorbable and nonabsorbable membranes placed over the lateral window: histomorphometric and clinical analyses. *Int J Periodontics Restorative Dent*. 2005;25(6):551-9.
- 312) Gonshor A, McAllister BS, Wallace SS, Prasad H. Histologic and histomorphometric evaluation of an allograft stem cell-based matrix sinus augmentation procedure. *Int J Oral Maxillofac Implants*. 2011;26(1):123-31.
- 313) Rickert D, Sauerbier S, Nagursky H, Menne D, Vissink A, Raghoobar GM. Maxillary sinus floor elevation with bovine bone mineral combined with either autogenous bone or autogenous stem cells: a prospective randomized clinical trial. *Clin Oral Implants Res*. 2011;22(3):251-8. doi: 10.1111/j.1600-0501.2010.01981.x. Epub 2010 Sep 10

- 314) Shayesteh YS, Khojasteh A, Soleimani M, Alikhasi M, Khoshzaban A, Ahmadbeigi N. Sinus augmentation using human mesenchymal stem cells loaded into a beta-tricalcium phosphate/hydroxyapatite scaffold. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008;106(2):203-9.
- 315) Sauerbier S, Stubbe K, Maglione M, Haberstroh J, Kuschnierz J, Oshima T, Xavier SP, Brunnberg L, Schmelzeisen R, Gutwald R. Mesenchymal stem cells and bovine bone mineral in sinus lift procedures--an experimental study in sheep. *Tissue Eng Part C Methods.* 2010;16(5):1033-9.
- 316) Lundgren S, Andersson S, Sennerby L. Spontaneous bone formation in the maxillary sinus after removal of a cyst: coincidental or expected reaction? *Clin Implant Dent Relat Res* 2003; 5: 78–81.
- 317) Lundgren S, Andersson S, Gualini F, Sennerby L. Bone reformation with sinus membrane elevation: a new surgical technique for maxillary sinus floor augmentation. *Clin Implant Dent Relat Res* 2004; 6: 165–173.
- 318) Palma VC, Magro-Filho O, de Oliveria JA, Lundgren S, Salata LA, Sennerby L. Bone reformation and implant integration following maxillary sinus membrane elevation: an experimental study in primates. *Clin Implant Dent Relat Res.* 2006;8(1):11-24
- 319) Winter AA, Pollack AS, Odrich RB. Sinus/alveolar crest tenting (SACT): a new technique for implant placement in atrophic maxillary ridges without bone grafts or membranes. *Int J Periodontics Restorative Dent* 2003; 23:557–565.
- 320) Thor A, Sennerby L, Hirsch JM, Rasmusson L. Bone formation at maxillary sinus floor following simultaneous elevation of the mucosa lining and implant installation without graft material: an evaluation of 20 patient treated with 44 Astra Tech Implants. *J Oral Maxillofac Surg* 2007;65:64-72
- 321) Sani E, Veltri M, Cagidiaco MC, Balleri P, Ferrari M. Sinus membrane elevation in combination with placement of blasted implants: a 3-year case report of sinus augmentation without grafting material. *Int J Oral Maxillofac Surg.* 2008 Oct;37(10):966-9.
- 322) Ellegaard B, Baelum V, Kisen-Petersen J. Non-grafted sinus implants in periodontally compromised patients: a time-to-event analysis. *Clin. Oral Impl. Res.* 2006;17: 156–164
- 323) Sohn DS, Moon JW, Lee WH, Kim SS, Kim CW, Kim KT, Moon YS. Comparison of new bone formation in the maxillary sinus with and without bone grafts: immunochemical rabbit study. *Int J Oral Maxillofac Implants.* 2011 Sep-Oct;26(5):1033-42.
- 324) Cricchio G, Palma VC, Faria PE, de Oliveira JA, Lundgren S, Sennerby L, Salata LA. Histological findings following the use of a space-making device for bone reformation and implant integration in the maxillary sinus of primates. *Clin Implant Dent Relat Res.* 2009 Oct;11 Suppl 1:e14-22. Epub 2009 Apr 16.

- 325) Hatano N, Sennerby L, Lundgren S. Maxillary sinus augmentation using sinus membrane elevation and peripheral venous blood for implant-supported rehabilitation of the atrophic posterior maxilla: case series. *Clin Implant Dent Relat Res.* 2007 Sep;9(3):150-5.
- 326) Sohn DS, Moon JW, Moon KN, Cho SC, Kang PS. New Bone Formation in the Maxillary Sinus using only absorbable gelatin sponge. *J Oral Maxillofac Surg.* 2010 Jun;68(6):1327-33.
- 327) De Boer HH. The history of bone grafts. *Clin Orthop Relat Res.* 1988;(226):292-8.
- 328) Tomford WW. Bone allografts. Past, present and future. *Cell Tissue Bank.* 2000;1(2):105-9.
- 329) Leslie HW, Bottenfield S. Donation, banking, and transplantation of allograft tissues. *Nurs Clin North Am.* 1989;24(4):891-905
- 330) Albert A, Leemrijse T, Druez V, Delloye C, Cornu O. Are bone autografts still necessary in 2006? A three-year retrospective study of bone grafting. *Acta Orthop Belg.* 2006;72(6):734-40.
- 331) Dodd CA, Fergusson CM, Freedman L, Houghton GR, Thomas D. Allograft versus autograft bone in scoliosis surgery. *J Bone Joint Surg Br.* 1988;70(3):431-4
- 332) Buck BE, Malinin TI, Brown MD. Bone transplantation and human immunodeficiency virus. An estimate of risk of acquired immunodeficiency syndrome (AIDS). *Clin Orthop Relat Res.* 1989;(240):129-36.
- 333) Simonds RJ, Holmberg SD, Hurwitz RL, et al. Transmission of human immunodeficiency virus type 1 from a seronegative organ and tissue donor. *N Engl J Med* 1992;326 726-32
- 334) Tomford, WW. Transmission of disease through transplantation of musculoskeletal allografts. *J Bone Joint Surg Am.* 1995;77 1742-54
- 335) Centers for Disease Control and Prevention. Hepatitis C virus transmission from an antibody-negative organ and tissue donor – United States, 2000-2002. *Morb Mortal Wkly Rep* 2003;52 273-6 syndrome (AIDS). *Clin Orthop Relat Res* 1989;240:129-36
- 336) Eagle MJ, Rooney P, Lomas R, Kearney JN. Validation of radiation dose received by frozen unprocessed and processed bone during terminal sterilization. *Cell Tissue Bank.* 2005;6 221-30
- 337) Grieb TA, Forng RY, Stafford RE, et al. Effective use of optimized, high dose (50kGy) gamma irradiation for pathogen inactivation of human bone allografts. *Biomaterials* 2005;26 2033-42
- 338) Smith RA, Ingels J, Lochemes JJ, et al. Gamma irradiation of HIV-1. *J Orthop Res* 2001;19 815-9
- 339) Centers for Disease Control and Prevention. Invasive *Streptococcus pyogenes* after allograft implantation – Colorado, 2003. *Morb. Mortal. Wkly. Rep.* 2003;52:1173-6
- 340) Centers for Disease Control and Prevention. Update: Allograft-associated bacterial infections – United States 2002. *Morbidity Mortal Wkly Rep* 2002;51:207-10
- 341) Lee EH, Ferguson D, Jernigan D, Greenwald M, Coté T, Bos JE, Guarner J, Zaki S, Schuchat A, Beall B, Srinivasan A. Invasive group-A streptococcal infection in an allograft recipient. A case report. *J Bone Joint Surg Am.* 2007;89(9):2044-7.

- 342) Costain DJ, Crawford RW. Fresh-frozen vs. irradiated allograft bone in orthopaedic reconstructive surgery. *Injury* 2009;40(12):1260-64.
- 343) Colwell A, Hamer A, Blumsohn A, Eastell R. To determine the effects of ultraviolet light, natural light and ionizing radiation on pyridinium cross-links in bone and urine using high-performance liquid chromatography. *Eur J Clin Invest* 1996;26:1107-14
- 344) Nguyen H, Morgan DAF, Forwood MR. Sterilization of allograft bone: Effects of gamma irradiation on allograft biology and biomechanics. *Cell Tissue Bank* 2007;8:93-105
- 345) Akkus O, Belaney RM, Das P. Free radical scavenging alleviates the biomechanical impairment of gamma radiation sterilized bone tissue. *J Orthop Res* 2005;23: 838-45
- 346) Akkus O, Belaney RM. Sterilization by gamma radiation impairs the tensile fatigue life of cortical bone by two orders of magnitude. *J Orthop Res* 2005;23 1054-8
- 347) Chen JB, Yu Y, Yang J-L, et al. BMP-7 and CBFA1 in allograft bone in vivo bone formation and the influence of γ -irradiation. *J Biomed Mat Res* 2007;80 435-43
- 348) AATB: General standards of tissue banking, technical manual for muscle-skeletal tissue banking. 1992
- 349) Gazdag AR, Lane JM, Glaser D, Forster RA. Alternative to autogenous bone graft: Efficacy and indications. *J Am Acad Orthop Surg*. 1995;3(1):1-8.
- 350) Ehrler DM, Vaccaro AR: The use of allograft bone in lumbar spine surgery. *Clin Orthop Relat Res*. 2000;(371):38-45
- 351) Scarano A, Degidi M, Iezzi G, Pecora G, Piattelli M, Orsini G, Caputi S, Perrotti V, Mangano C, Piattelli A. Maxillary sinus augmentation with different biomaterials: a comparative histologic and histomorphometric study in man. *Implant Dent*. 2006;15(2):197-207.
- 352) Cammack GV, 2nd, Nevins M, Clem DS, 3rd, Hatch JP, Mellonig JT. Histologic evaluation of mineralized and demineralized freeze-dried bone allograft for ridge and sinus augmentations. *Int J Periodontics Restorative Dent* 2005;25:231-237.
- 353) Choukroun J, Diss A, Simonpieri A, Girard MO, Schoeffler C, Dohan SL, Dohan AJ, Mouhyi J, Dohan DM. Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part V: histologic evaluations of PRF effects on bone allograft maturation in sinus lift. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006;101(3):299-303.
- 354) Marx RE: Bone and bone graft healing. *Oral Maxillofac Surg Clin North Am* 19:455, 2007
- 355) Simpson D, Kakarala G, Hampson K, Steele N, Ashton B. Viable cells survive in fresh frozen human bone allografts. *Acta Orthop* 2007;78:26-30.
- 356) Egli RJ, Sckell A, Fraitzl CR, Felix R, Ganz R, Hofstetter W, Leunig M Cryopreservation with dimethyl sulfoxide sustains partially the biological function of osteochondral tissue. *Bone*. 2003;33(3):352-61.

- 357) Egli RJ, Wingenfeld C, Hölzle M, Hempfing A, Fraitzl CR, Ganz R, Leunig M. Histopathology of cryopreserved bone allo- and isografts: pretreatment with dimethyl sulfoxide. *J Invest Surg.* 2006;19(2):87-96.
- 358) Wingenfeld C, Egli RJ, Hempfing A, Ganz R, Leunig M. Cryopreservation of osteochondral allografts: dimethyl sulfoxide promotes angiogenesis and immune tolerance in mice. *J Bone Joint Surg Am* 2002;84-A:1420-1429.
- 359) Enneking WF. Histological investigation of bone transplants in immunologically prepared animals. *J Bone Joint Surg Am.* 1957;39-A(3):597-615.
- 360) Gray JC, Elves MW. Early osteogenesis in compact bone isografts: a quantitative study of contributions of the different graft cells. *Calcif Tissue Int.* 1979;29(3):225-37.
- 361) Perrott DH, Smith RA, Kabam LB: The use of fresh frozen allogeneic bone for maxillary and mandibular reconstruction. *Int J Oral Maxillofac Surg* 1992;21(5):260-5.
- 362) Accetturi E, Germani KB, Cavalca D: Reconstruction of bone defects in the maxilla and mandibula through the use of frozen human bone. *Transplant Proc.* 2002;34(2):531-3
- 363) Rodella LF, Favero G, Boninsegna R, Borgonovo A, Rezzani R, Santoro F. TGF-beta1 and VEGF after fresh frozen bone allograft insertion in oral-maxillo-facial surgery. *Histol Histopathol.* 2010;25(4):463-71.
- 364) Acocella A, Bertolai R, Nissan J, Sacco R. Clinical, histological and histomorphometrical study of maxillary sinus augmentation using cortico-cancellous fresh frozen bone chips. *J Craniomaxillofac Surg.* 2011;39(3):192-9.
- 365) Stacchi C, Orsini G, Di Iorio D, Breschi L, Di Lenarda R. Clinical, histologic, and histomorphometric analyses of regenerated bone in maxillary sinus augmentation using fresh frozen human bone allografts. *J Periodontol.* 2008;79(9):1789-96.
- 366) Viscioni A, Dalla Rosa J, Paolin A, Franco M. Fresh-frozen bone: case series of a new grafting material for sinus lift and immediate implants. *J Ir Dent Assoc.* 2010;56(4):186-91.
- 367) Viscioni A, Franco M, Paolin A, Cogliati E, Callegari M, Zollino I, Sollazzo V, Carinci F. Effectiveness of fresh frozen and cryopreserved homologous iliac crest grafts used in sinus lifting: a comparative study. *Cell Tissue Bank.* 2011;12(4):263-71.
- 368) Acocella A, Bertolai R, Nissan J, Ellis E 3rd, Sacco R. Maxillary sinus lift using fresh frozen bone chips in presence of sinus cyst: clinical and histological report. *Cell Tissue Bank.* 2012;13(2):327-32.
- 369) Pelegrine AA, Sorgi da Costa CE, Sendyk WR, Gromatzky A. The comparative analysis of homologous fresh frozen bone and autogenous bone graft, associated or not with autogenous bone marrow, in rabbit calvaria: a clinical and histomorphometric study. *Cell Tissue Bank.* 2011;12(3):171-84.

- 370) Acocella A, Bertolai R, Sacco R, Colafranceschi M: Clinical, histological and histomorphometric evaluation of the healing of mandibular ramus bone block grafts for alveolar ridge augmentation before implant placement *J Craniomaxillofac Surg* 2010;38(3):222-30
- 371) Ham, A. W. Some histophysiological problems peculiar to calcified tissues. *J. Bone Joint Surg. Am.* 34A: 701-728, 1952.
- 372) Ellegaard, B., Karring, T., Loe, H. The fate of vital and devitalized bone grafts in the healing of interradicular lesions. *J Periodontal Res* 1975; 10: 88 97.
- 373) Chugh, S., Marks, D.S., Mangham, D.C., Thompson, A.G. Autologous bone grafting in staged scoliosis surgery. The patient as bone bank. *Spine* 1998; 23: 1793 1795.
- 374) Blomqvist JE, Alberius P, Isaksson S, Linde A, Obrant K Importance of bone graft quality for implant integration after maxillary sinus reconstruction. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1998;86(3):268-74.
- 375) Berggren, A, Weiland, A.J, Dorfman, H. The effect of prolonged ischaemia time on osteocyte and osteoblast survival in composite bone grafts revascularized by microvascular anastomoses. *Plast Reconstruct Surg* 1982;69: 290–298.
- 376) Goldberg, V.M., Stevenson, S. Natural history of autografts and allografts. *Clin Orthop* 1987; 225: 7 16.
- 377) Klein-Nulend, J., Van der Plas, A., Semeins, C.M., Ajubi, N.E., Frangos, J.A., Nijweide, P.J., Burger, E.H. Sensitivity of osteocytes to biomechanical stress in vitro. *FASEB J* 1995; 9: 441 445.
- 378) Smit, T.H., Burger, E.H. A case for strain-induced fluid-flow as regulation of bone-coupling and osteonal alignment. *J Bone Miner Res* 2002; 11: 2021 2029.
- 379) Zhao, S., Zhang, Y.K., Harris, S., Ahuja, S.S., Bonewald, L.F. MLO-Y4 osteocyte-like cells support osteoclast formation and activation. *J Bone Miner Res* 2002; 17: 2068 2079.
- 380) Bronckers, A.L.J.J., Goei, W., Luo, G., Karsenty, G., D' Souza, R.N., Lyaruu, D.M., Burger, E.H. DNA fragmentation during bone formation in neonatal rodents assessed by transferase mediated end labelling. *J Bone Miner Res* 1996; 11: 1281 1291.
- 381) Kikuyama, A., Fukuda, K., Mori, S., Okada, M., Yamaguchi, H., Hamanishi, C. Hydrogen peroxide induces apoptosis of osteocytes: involvement of calcium ion and caspase activity. *Calcif Tissue Int* 2002; 71: 243 248.
- 382) Verborgt O, Gibson GJ, Schaffler MB. Loss of osteocyte integrity in association with microdamage and bone remodeling after fatigue in vivo. *J Bone Miner Res.* 2000;15(1):60-7.
- 383) Kamijou, T., Nakajima, T., Ozawa, H. Effects of osteocytes on osteoinduction in the autogenous rib graft in the rat mandible. *Bone* 1994; 15: 629- 637.
- 384) Kingsmill VJ, Boyde A, Jones SJ. The resorption of vital and devitalized bone in vitro: significance for bone grafts. *Calcif Tissue Int.* 1999;64(3):252-6.

- 385) Zerbo IR, de Lange GL, Joldersma M, Bronckers AL, Burger EH. Fate of monocortical bone blocks grafted in the human maxilla: a histological and histomorphometric study. *Clin Oral Implants Res.* 2003;14(6):759-66.
- 386) Perrott DH, Smith RA, Kaban LB. The use of fresh frozen allogeneic bone for maxillary and mandibular reconstruction. *Int J Oral Maxillofac Surg.* 1992 Oct;21(5):260-5.
- 387) Spin Neto R, Felipe Leite C, Pereira LA, Marcantonio E, Marcantonio Jr E. Is Peripheral Blood Cell Balanced Altered by the Use of Fresh Frozen Bone Block Allografts in Lateral Maxillary Ridge Augmentation? *Clin Implant Dent Relat Res.* 2011 Aug 11. doi: 10.1111/j.1708-8208.2011.00385.x. [Epub ahead of print]
- 388) Acocella A, Bertolai R, Ellis E 3rd, Nissan J, Sacco R. Maxillary alveolar ridge reconstruction with monocortical fresh-frozen bone blocks: a clinical, histological and histomorphometric study. *J Craniomaxillofac Surg.* 2012;40(6):525-33.
- 389) Barone A, Varanini P, Orlando B, Tonelli P, Covani U. Deep-frozen allogeneic onlay bone grafts for reconstruction of atrophic maxillary alveolar ridges: a preliminary study. *J Oral Maxillofac Surg.* 2009;67(6):1300-6.
- 390) Contar CM, Sarot JR, Bordini J Jr, Galvão GH, Nicolau GV, Machado MA. Maxillary ridge augmentation with fresh-frozen bone allografts. *J Oral Maxillofac Surg.* 2009;67(6):1280-5.
- 391) Franco M, Viscioni A, Rigo L, Guidi R, Brunelli G, Carinci F. Iliac crest fresh frozen homografts used in pre-prosthetic surgery: a retrospective study. *Cell Tissue Bank.* 2009;10(3):227-33.
- 392) Viscioni A, Franco M, Rigo L, Guidi R, Spinelli G, Carinci F. Retrospective study of standard-diameter implants inserted into allografts. *J Oral Maxillofac Surg.* 2009;67(2):387-93
- 393) Franco M, Rigo L, Viscione A, De Santis B, Tropina E, Brunelli G, Guidi R, Avantaggiato A, Carinci F. CaPO₄ blasted implants inserted into iliac crest homologue frozen grafts. *J Oral Implantol.* 2009;35(4):176-80.
- 394) Carinci F, Brunelli G, Franco M, Viscioni A, Rigo L, Guidi R, Strohmenger L. A retrospective study on 287 implants installed in resorbed maxillae grafted with fresh frozen allogeneous bone. *Clin Implant Dent Relat Res.* 2010;12(2):91-8.
- 395) Franco M, Viscioni A, Rigo L, Guidi R, Zollino I, Avantaggiato A, Carinci F. Clinical outcome of narrow diameter implants inserted into allografts. *J Appl Oral Sci.* 2009;17(4):301-6.
- 396) Rigo L, Viscioni A, Franco M, Lucchese A, Zollino I, Brunelli G, Carinci F. Overdentures on implants placed in bone augmented with fresh frozen bone. *Minerva Stomatol.* 2010 Dec 15. [Epub ahead of print]
- 397) Nocini PF, Bertossi D, Albanese M, D'Agostino A, Chilosi M, Procacci P. Severe maxillary atrophy treatment with Le Fort I, allografts, and implant-supported prosthetic rehabilitation. *J Craniofac Surg.* 2011;22(6):2247-54.

- 398) Grecchi F, Zollino I, Parafioriti A, Mineo G, Pricolo A, Carinci F. One-step oral rehabilitation by means of implants' insertion, Le Fort I, grafts, and immediate loading. *J Craniofac Surg.* 2009;20(6):2205-10.
- 399) De Macedo LG, de Macedo NL, do Socorro Ferreira Monteiro A. Fresh-frozen human bone graft for repair of defect after adenomatoid odontogenic tumour removal. *Cell Tissue Bank.* 2009;10(3):221-6.
- 400) Albanese M, Procacci P, Sancassani G, Nocini PF. Fresh-frozen human bone graft to repair defect after mandibular giant follicular cyst removal: a case report. *Cell Tissue Bank.* 2012;13(2):305-13.
- 401) Albanese M, Nocini PF, Fior A, Rizzato A, Cristofaro MG, Sancassani G, Procacci. Mandibular reconstruction using fresh frozen bone allograft after conservative enucleation of a mandibular odontogenic myxoma. *J Craniofac Surg.* 2012 May;23(3):831-5. doi: 10.1097/SCS.0b013e31824dbff6.
- 402) Borgonovo AE, Tommasi F, Panigalli A, Bianchi AC, Boninsegna R, Santoro F. Use of fresh frozen bone graft in rehabilitation of maxillar atrophy. *Minerva Stomatol.* 2012;61(4):141-54.
- 403) Orsini G, Stacchi C, Visintini E, Di Iorio D, Putignano A, Breschi L, Di Lenarda R. Clinical and histologic evaluation of fresh frozen human bone grafts for horizontal reconstruction of maxillary alveolar ridges. *Int J Periodontics Restorative Dent.* 201;31(5):535-44.
- 404) Contar CM, Sarot JR, da Costa MB, Bordini J, de Lima AA, Alanis LR, Trevilatto PC, Machado MÂ. Fresh-frozen bone allografts in maxillary ridge augmentation: histologic analysis. *J Oral Implantol.* 2011 Apr;37(2):223-31.
- 405) Spin-Neto R, Landazuri Del Barrio RA, Pereira LA, Marcantonio RA, Marcantonio E, Marcantonio Jr E. Clinical Similarities and Histological Diversity Comparing Fresh Frozen Onlay Bone Blocks Allografts and Autografts in Human Maxillary Reconstruction. *Clin Implant Dent Relat Res.* 2011 Aug 11. doi: 10.1111/j.1708-8208.2011.00382.x. [Epub ahead of print]
- 406) Lumetti S, Consolo U, Galli C, Multinu A, Piersanti L, Bellini P, Manfredi E, Corinaldesi G, Zaffe D, Macaluso GM, Marchetti C. Fresh-Frozen Bone Blocks for Horizontal Ridge Augmentation in the Upper Maxilla: 6-Month Outcomes of a Randomized Controlled Trial. *Clin Implant Dent Relat Res.* 2012 Apr 24. doi: 10.1111/j.1708-8208.2012.00458.x. [Epub ahead of print]
- 407) Pimentel AC, Napimoga MH, Manzi MR, Sendyk WR. Reconstruction of the edentulous mandible with fresh frozen bone grafts and implants: a 4-year report of a prospective clinical study. *Cell Tissue Bank.* 2012 Nov 27. [Epub ahead of print]
- 408) Messori MR, Nagata MJ, Fucini SE, Pola NM, Campos N, Oliveira GC, Bosco AF, Garcia VG, Furlaneto FA. Effect of platelet-rich plasma on the healing of mandibular defects treated with bone allograft. A radiographic study in dogs. *J Oral Implantol.* 2012 Sep 26. [Epub ahead of print]

- 409) Messora MR, Nagata MJ, Pola NM, de Campos N, Fucini SE, Furlaneto FA. Effect of platelet-rich plasma on bone healing of fresh frozen bone allograft in mandibular defects: a histomorphometric study in dogs. *Clin Oral Implants Res.* 2012 Aug 28. doi: 10.1111/clr.12008. [Epub ahead of print]