



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Pancreatic fistula: A proposed percutaneous procedure

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Pancreatic fistula: A proposed percutaneous procedure / Silvia Pradella; Ernesto Mazza; Francesco Mondaini; Stefano Colagrande. - In: WORLD JOURNAL OF HEPATOLOGY. - ISSN 1948-5182. - ELETTRONICO. - 5:(2013), pp. 33-37. [10.4254/wjh.v5.i1.33]

Availability:

This version is available at: 2158/793782 since:

Published version:

DOI: 10.4254/wjh.v5.i1.33

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Pancreatic fistula: A proposed percutaneous procedure

Silvia Pradella, Ernesto Mazza, Francesco Mondaini, Stefano Colagrande

Silvia Pradella, Stefano Colagrande, Department of Experimental and Clinical Biomedical Sciences, Section of Radiodiagnosics, University of Florence, Azienda Ospedaliero-Universitaria Careggi, 50134 Florence, Italy

Ernesto Mazza, Francesco Mondaini, Department of Diagnostic Imaging, Section of Interventional Radiology, Azienda Ospedaliero-Universitaria Careggi, 50134 Florence, Italy

Author contributions: All the authors have contributed to all the phases of the article.

Correspondence to: Stefano Colagrande, Associate Professor of Radiology, Department of Experimental and Clinical Biomedical Sciences, Section of Radiodiagnosics, University of Florence, Azienda Ospedaliero-Universitaria Careggi, 50134 Florence, Italy. stefano.colagrande@unifi.it

Telephone: +39-55-7946338 Fax: +39-55-431970

Received: June 29, 2012 Revised: December 7, 2012

Accepted: December 23, 2012

Published online: January 27, 2013

Abstract

AIM: To propose a percutaneous treatment for otherwise intractable pancreatic fistula (PF).

METHODS: From 2005 to 2011, 12 patients (9 men and 3 women, mean age 59 years, median 63 years, range 33-78 years) underwent radiological treatment for high-output PF associated with peripancreatic fluid collection. The percutaneous procedures were performed after at least 4 wk of unsuccessful conservative treatments. We chose either a one or two step procedure, depending on the size and characteristics of the fistula and the fluid collection (with an arbitrary cut-off of 2 cm). Initially, 2 to 6 pigtail drainages of variable size from 8.3 (8.3-Pig Duan Cook, Bloomington, Indiana, United States) to 14 Fr (Flexima, Boston Scientific, Natick, United States) were positioned inside the collection using a transgastric approach. In a second procedure, after 7-10 d, two or more endoprosthesis (cystogastrostomic 8 Fr double-pigtail, Cook, Bloomington, Indiana, United States in 10 patients; covered Niti-S stent, TaeWoong Medical Co, Seoul, South Korea in 2 patients) were placed between the collection and the gastric lumen. In all cases the metal or plastic pros-

theses were removed within one year after positioning.

RESULTS: Four out of 12 high-output fistulas were external while 8/12 were internal. The origin of the fistulous tract was visualised by computer tomography (CT) imaging studies: in 11 patients it was at the body, and in 1 patient at the tail of the pancreas. Single or multiple drainages were positioned under CT guidance. The catheters were left in place for a varying period (0 to 40 d - median 10 and 25th-75th percentile 0-14). In one case external transgastric drainages were left in place for a prolonged time (40 d) due to the presence of vancomycin-resistant bacteria (*Staphylococcus*) and fluconazole-resistant fungi (*Candida*) in the drained fluid. In this latter case systemic and local antibiotic therapy was administered. In both single and two-step techniques, when infection was present, we carried out additional washing with antibiotics to improve the likelihood of the procedure's success. In all cases the endoprosthesis were left *in situ* for a few weeks and endoscopically removed after remission of collections, as ascertained by CT scan. Procedural success rate was 100% as the resolution of external PF was achieved in all cases. There were no peri-procedural complications in any of the patients. The minimum follow-up was 18 mo. In two cases the procedure was repeated after 1 year, due to the onset of new fluid collections and the development of pseudocysts. Indeed, this type of endoprosthesis is routinely employed for the treatment of pseudocysts. Endoscopy was adopted both for control of the positioning of the endoprosthesis in the stomach, and for its removal after resolution of the fistula and fluid collection. The resolution of the external fistula was assessed clinically and CT scan was employed to demonstrate the resolution of peripancreatic collections for both the internal and external fistulae.

CONCLUSION: The percutaneous placement of cystogastrostomic endoprosthesis can be used for the treatment of PF that cannot be treated with other procedures.

© 2013 Baishideng. All rights reserved.

Key words: Pancreas; Pancreatic fistula; Interventional radiology; Pancreatic surgery; Complications

Pradella S, Mazza E, Mondaini F, Colagrande S. Pancreatic fistula: A proposed percutaneous procedure. *World J Hepatol* 2013; 5(1): 33-37 Available from: URL: <http://www.wjgnet.com/1948-5182/full/v5/i1/33.htm> DOI: <http://dx.doi.org/10.4254/wjh.v5.i1.33>

INTRODUCTION

Postoperative pancreatic fistula (PF) is a possible complication of acute or chronic pancreatitis, partial pancreatectomy, or trauma to the pancreatic duct during upper abdominal surgery, pancreatic biopsy, or blunt abdominal trauma. PF may lead to prolonged hospitalization, increased costs, and mortality. PFs occur in about 10% to 28% of patients who undergo pancreatic resection^[1]. PFs may be internal or external, and therapeutic options for both can be conservative (medical), endoscopic, percutaneous (under radiological guidance), or surgical.

Closure of external PFs is mainly achieved by medical therapy. In selected patients who have failed with less invasive procedures, surgical treatment maybe provided, with good results, but also with significant mortality rates^[1]. Improvements in both skill levels and in medical devices, have progressively increased the success rate of endoscopic methods^[2,3]. Nonetheless, some patients must be excluded from these procedures, mainly for technical reasons. In such rare cases, for example when the pancreatic duct cannulation is difficult for anatomical reasons or because of the patient's clinical condition, the percutaneous approach is possible and may be favored.

On the basis of twelve personal cases, we propose a percutaneous radiologically guided technique for treatment of PFs, which can be used when other procedures are not indicated or would be ineffective. This method seems to be extremely helpful in achieving a complete resolution.

MATERIALS AND METHODS

Patients

From 2005 to 2011, 12 patients (9 men and 3 women, mean age 59 years, median 63 years, range 33-78 years) received radiological treatment for high-output PF associated with peripancreatic fluid collection. The percutaneous procedures were performed after at least 4 wk of inconclusive conservative treatments (total parenteral nutrition with nasojejunal tube and administration of somatostatin analogues). All patients underwent abdominal computed tomography scans in order to demonstrate the origin of the fistulous tract and to evaluate the fluid collection. In our series of patients the surgeon, in consultation with the endoscopist, considered cannulation of the pancreatic duct to be contraindicated and decided to try a percutaneous radiological approach.

Table 1 Pancreatic fistula: Characteristics of patients

No.	Sex	Age (yr)	Pathology	Drainage	Dimension (fr)	Time (d)	Endo.	Time (mo)	Proc.
1	M	61	Post-surgery PD	2	8.3	8	2	9	2
2	M	77	Post-surgery PD	2	8.3	10	2	5	2
3	M	66	Post-surgery PD	1	14	0	2	4	2
4	F	78	Post-surgery PD	2	8.3	12	2	5	2
5	M	65	Post-surgery PD	2	10	14	2	6	1
6	F	78	Post-surgery pancreatic tail resection	1	14	0	2	6	2
7	M	41	Post-traumatic	2	14	18	2	6	2
8	M	52	Post-traumatic	1	14	0	2	4	1
9	M	33	Post-traumatic	6	8.3	10	6	6	4
10	F	66	Post-pancreatitis	3	14	40	3	5	4
11	F	60	Post-pancreatitis	3	14	14	2	6	3
12	M	35	Post-pancreatitis	2	10	10	2	6	2

M: Male; F: Female; PD: Pancreatico-duodenectomy; Endo.: Endoprosthesis; Proc.: Procedure; Time (d): Period of left *in situ* of drainages; Time (mo): Period of left *in situ* of endoprosthesis.

Procedure technique

In all patients (Table 1) one or more drains (Flexima, Boston Scientific, Natick, United States) were placed under computer tomography (CT) guidance. Where necessary, the catheters were left in place for mean period of 10 d (range 0-40 d). A minimum of two prostheses (cystogastrostomic 8 Fr double-pigtail, Cook, Bloomington, Indiana, United States in 10 patients; covered Niti-S stent, TaeWoong Medical Co, Seoul, South Korea in 2 patients) were then placed under fluoroscopic guidance and endoscopic control.

All procedures were performed by an expert interventional radiologist. Depending on the dimensions of the fluid collection, we adopted two different possible procedures (Figure 1).

In cases of retrogastric pancreatic fluid collections larger than 2 cm (8 patients), a two-step procedure was adopted (Figure 2A). Initially, 2 to 6 pigtail drains of size from 8.3 (8.3-Pig Duan Cook, Bloomington, Indiana, United States) to 14 Fr (Flexima, Boston Scientific, Natick, United States) were positioned inside the collection using a transgastric approach (Figure 2B and C). In a second procedure, 7-10 d later, two or more endoprotheses (cystogastrostomic 8 Fr double-pigtail, Cook, Bloomington, Indiana, United States in 10 patients; covered Niti-S stent, TaeWoong Medical Co, Seoul, South Korea in 2 patients) were placed between the collection and the gastric lumen.

The first step of the procedure was a trans-gastric CT-guided puncture of the collection using a 18 G Chiba needle (Figure 2B). Through this needle a rigid guide (extra stiff Amplatz 0.035 inch Cook, Bloomington, Indiana, United States) was inserted, and a sufficient number of drains introduced. Correct positioning was verified by




Figure 1 Flow chart of two-step (on the left) and one-step (on the right) procedures. PF: Pancreatic fistula; CT: Computer tomography.




Figure 2 Diagnostic enhanced computer tomography scan (A), and un-enhanced computer tomography scan (B, C) performed during interventional procedure. A: Patient developed a pancreatic fistula and abdominal fluid collection after surgery for pancreatic neoplasm; B, C: Percutaneous approach, was then adopted: images show trans-gastric puncture (B) and the access and placement of 2 drains in the collection (C).




Figure 3 Prosthesis fully extended (Niti-S stent covered, TaeWoong Medical Co., Seoul, South Korea) (A) and enhanced computer tomography scans (B, C) performed after interventional procedure. Images demonstrate prosthesis *in situ* and resolution of peripancreatic collections.

CT scan. Then, the drainage was removed and the prostheses placed under fluoroscopic guidance (Figure 3).

In cases of retrogastric pancreatic fluid collections smaller than 2 cm (4 patients) with external fistulas caused by surgical or percutaneous drainage, a one step procedure was preferred (Figure 4). Through the fistula, an angiographic guide (hydrophilic-curved guidewire 0.035 inch Terumo, Tokyo, Japan) and then an extra-stiff guidewire (Amplatz extra stiff inch 0.035 Boston Scientific, Natick, United States) were inserted. Through this latter guide a TIPSS cannula (transjugular intrahepatic Access set Cook, Bloomington, Indiana, United States) was positioned, the guide removed, and a mandrel inserted.

Then, under fluoroscopic guidance, the cannula was pushed into the gastric lumen, the mandrel removed, and a new guide/s inserted, as described in the literature^[4]. Thus, using a prosthesis, a by-pass is created between the




Figure 4 The external fistula opens on the skin surface (A), a catheter has been positioned through the fistula (B) and two prostheses have then been placed through these two guides (C).

origin of the fistula and the lumen of the stomach. The procedure (performed under local anesthesia associated with a mild analgo-sedation Remifentanyl and Midazolam based) was carried out under endoscopic surveillance. Prosthesis positioning follow-up was guided by CT (Figure 3B and C). In all cases, the metal or plastic prostheses were removed within one year after positioning. Recently a new device (Niti-S stent covered, TaeWoong Medical Co., Seoul, South Korea) has been introduced into clinical practice consisting of a self-expandable, coated, removable Nitinol metal stent, spool-shaped, 2 cm long, narrower in the central portion (12 mm) and wider at the ends (29 mm) (Figure 3A). This device, positioned between the collection and the gastric lumen, does not clog simply because it is large. We employed this tool in the last two patients. It will probably replace the plastic prostheses in the treatment of pancreatic fistulas.

RESULTS

The patients' characteristics are shown in Table 1. In all patients, one or more drains were positioned under CT guidance (Table 1). Four of the 12 fistulas were external while 8/12 were internal, all were high-output fistulas. External fistulas occurred in 2 patients with pancreatitis, and in 2 other patients post-surgical drainage. In all patients the CT examination showed a discontinuity of the main pancreatic duct, allowing us to localize the origin site of the fistula: this origin was at the body in 11 patients and at the tail of the pancreas in 1 patient. The catheters were left in place for 0 to 40 d (median 10 and 25th-75th percentile 0-14). The longest period was in a case showing the presence of vancomycin-resistant bacteria (*Staphylococcus*) and fluconazole-resistant fungi (*Candida*) in the drained fluid. In this case systemic and local (via catheter) antibiotic therapy was administered. The endoprotheses were left *in situ* for few weeks and endoscopically removed after remission of collections, demonstrated by CT scan.

The resolution of external PF was achieved in all cases, and there were no peri-procedural complications. In two cases the procedure was repeated after 1 year us-

ing new external drains, following the onset of pseudocyst and new fluid collections.

DISCUSSION

The percutaneous method we have described, seems to be extremely promising for achieving a complete resolution of PF under radiologic guidance.

Even though PF is a frequent complication of pancreatic resection, its management is still not standardized^[3]. PFs may be internal or external and both result from leakage of pancreatic juice from a disrupted pancreatic duct. External PFs extend from the pancreas (or peripancreatic fluid collection) to the skin surface, and a pancreatic fluid collection is frequently associated with a fistula. On the basis of the amount of secretions, PFs are classified into high-output (> 200 mL per 24 h), and low-output (< 200 mL per 24 h) type^[1].

The choice of treatment depends on the characteristics the patient and the fistula. The overall success rate of the medical approach is high, 68%-100% of cases^[1], and in failed cases, endoscopic treatment has been suggested as a reasonable next step to achieve healing a PF within a few weeks^[4]. Surgical treatment is indicated only for patients who have failed the less invasive attempts and has a success rate of about 90%-92%, but with a significant mortality rate of 6%-9%^[1]. Endoscopic procedures include various drainage techniques, comprehensive cannulation of the injured pancreatic duct and stent positioning. The case studies reported in the literature, however, involve few patients, and success is dependent on the experience of physicians^[2,5].

Some patients are not eligible for the above mentioned treatments, mainly due to technical limitations of endoscopic retrograde cholangio pancreatography, related to duodenal and pancreatic duct stricture or post-surgical anatomy, that make duct cannulation complex or impossible^[6].

In these patients radiological treatment can be considered although not all cases are eligible, as with other minimally invasive methods. The interventional radiologist evaluates each case, and when the access is ad-

equate, may decide to use the method described above, a procedure already well established in cases of pancreatic pseudocyst and characterized by low complication and mortality rates. The prostheses are left in position for a few weeks to allow the normal flow or egress of pancreatic juices into the gut, facilitating the closure of the fistula. In our experience, this technique allows the resolution of the fistula without significant complications. Obviously, the cases treated were few because this procedure is rarely necessary. However, we have described this technique because we believe that it can be a feasible option or may be used in addition to other treatments.

COMMENTS

Background

The pancreatic fistula (PF) is a possible complication, especially in pancreatic surgery, still associated with a significant mortality. In almost all cases, medical therapy leads to healing but problems exist in a few cases. These patients have different treatment options. The procedure described here is an additional alternative that does not exclude other therapeutic options.

Research frontiers

Modern surgery of the pancreas is still burdened with complications, such as fistulas, and the challenge is to reduce resulting mortality. The minimally invasive procedures proposed in the literature are all based on only a few cases. The procedure described here fits into the puzzle of customized treatments.

Innovations and breakthroughs

In selected cases, described in the paper, the authors adopted a treatment previously used in pancreatic pseudocysts for the treatment of patients with PFs. They observed a remission of fistula associated with the drainage of pancreatic fluid within the stomach. So, the procedure described is not "new" but is an innovative employment of a method generally used for pancreatic pseudocyst treatment.

Applications

The technique represents a further option within the framework of minimally invasive procedures of intractable fistulas in selected patients. This procedure does not preclude surgery or other treatments.

Peer review

The manuscript proposes a new technique to solve persistent high-output PF.

REFERENCES

- 1 **Butturini G**, Daskalaki D, Molinari E, Scopelliti F, Casarotto A, Bassi C. Pancreatic fistula: definition and current problems. *J Hepatobiliary Pancreat Surg* 2008; **15**: 247-251 [PMID: 18535760 DOI: 10.1007/s00534-007-1301-y]
- 2 **Hirota M**, Kanemitsu K, Takamori H, Chikamoto A, Hayashi N, Horino K, Baba H. Percutaneous trans fistulous pancreatic duct drainage and interventional pancreatojejunostomy as a treatment option for intractable pancreatic fistula. *Am J Surg* 2008; **196**: 280-284 [PMID: 18639662 DOI: 10.1016/j.amjsurg.2007.05.055]
- 3 **Cicek B**, Parlak E, Oguz D, Disibeyaz S, Koksak AS, Sahin B. Endoscopic treatment of pancreatic fistulas. *Surg Endosc* 2006; **20**: 1706-1712 [PMID: 16960673 DOI: 10.1007/s00464-005-0764-7]
- 4 **Seewald S**, Ang TL, Kida M, Teng KY, Soehendra N. EUS 2008 Working Group document: evaluation of EUS-guided drainage of pancreatic-fluid collections (with video). *Gastrointest Endosc* 2009; **69**: S13-S21 [PMID: 19179137 DOI: 10.1016/j.gie.2008.12.035]
- 5 **Saeed ZA**, Ramirez FC, Hepps KS. Endoscopic stent placement for internal and external pancreatic fistulas. *Gastroenterology* 1993; **105**: 1213-1217 [PMID: 8405869]
- 6 **Barkay O**, Sherman S, McHenry L, Yoo BM, Fogel EL, Watkins JL, DeWitt J, Al-Haddad MA, Lehman GA. Therapeutic EUS-assisted endoscopic retrograde pancreatography after failed pancreatic duct cannulation at ERCP. *Gastrointest Endosc* 2010; **71**: 1166-1173 [PMID: 20303489 DOI: 10.1016/j.gie.2009.10.048]

P- Reviewer Vigano L S- Editor Cheng JX
L- Editor Hughes D E- Editor Li JY

