



REGIONE MARCHE
AZIENDA SANITARIA
UNICA REGIONALE
ZT 12




OSPEDALE MADONNA
DEL SOCCORSO
UNITÀ OPERATIVA
DI NEUROLOGIA

NEUROSONOLOGIA
STROKE-UNIT
DIRETTORE
L. CURATOLA

S. SANGUIGNI
T. CARBONI
R. GOBBATO
C. PACI
G. D'ANDREAMATTEO

NEUROSONOLOGY AND BRAIN PATHOLOGY

Training courses
in transcranial colour-coded
duplex sonography - 2009



The neurosonology of the intracerebral
vessels and parenchyma

Edited by Sandro Sanguigni

ABSTRACT BOOK

The role of ultrasound examination in disease of the thoracic aorta

Sergio Castellani

Clinical and Experimental Angiology Unit, University of Florence, Italy
E-mail: scaste.sergio@gmail.com

Introduction

Stroke and transient ischaemic attacks are often the consequence of a disease of the aortic arch, especially in elderly patients. The thoracic aorta is a common site of embolising atheromas (1-5); in addition, the dissecting complication of aortic aneurysms can extend to the origin of the cerebral feeding arteries (6,7). A recent extensive metanalysis of a large number of studies clearly showed that when the transverse aortic diameter exceeds 3.4 cm, it is positively and strongly related to the incidence of cervical artery dissections (8). Given the possible link between aortic disease and sudden neurological symptoms, during the past three decades ultrasound techniques have been extensively used to achieve an anatomical characterisation of the aortic arch, to define the presence of atherosclerotic debris, and to diagnose aneurysmal disease and/or its dissecting complications.

Aortic aneurysms

Ultrasound examination of the ascending aorta is feasible both by transthoracic and transoesophageal echocardiography, but the yield of the two techniques is quite different. The first reports on applications of transthoracic echocardiography (TTE) for the diagnosis of aortic disease date back to the early 1970s (9). Subsequently, the diagnostic accuracy of transoesophageal echocardiography (TEE) proved superior to that of TTE in detecting aortic aneurysmal disease and its complications; sensitivity and specificity ranged from 94 to 100% (10-12). Indeed, TTE was extensively shown to be less reliable than TEE examination due to the poor acoustic thoracic windows both from parasternal and jugular approaches. Aortic dissection is an acute medical emergency with a high mortality; early recognition and appropriate treatment is crucial to improving survival. Ultrasound examination of the aorta is recommended by American College of Emergency Physicians guidelines as a first level test (13) and given the superiority of the technique, TEE has become the preferred US modality for urgent evaluation of a patient with a suspected acute complication of an aneurysm of the thoracic aorta. Although desirable, TEE is not always readily available as the initial test, particularly in unstable patients with suspected aortic dissection in an acute setting soon after hospitalisation in the emergency room. On the other hand, data collected in limited case series (14,15) indicate that even TTE performed by emergency room physicians can allow reliable diagnosis of aortic dissections and can be a valuable resource especially in the presence of a haemodynamically unstable patient and/or in the presence of an acute cerebral ischaemia, when the findings of the cardiac examination can prompt rapid and often life-saving therapeutic decisions.

Aortic debris

Atheroma of the aortic arch is an important and poorly recognised factor in the aetiology of stroke (16). The risk of stroke is correlated with the presence and thickness of the atherosclerotic lesion and with the coexistence of an atherosclerotic plaque (17). The progressive enlargement of the elderly population is associated with an increasing prevalence of systemic and aortic atherosclerotic disease. In addition, during the last two decades the astonishing increase in the number of coronary angiographies

nipulation or catheter movements through the aortic arch (18). Detecting the presence of aortic debris and characterising its severity, particularly in patients older than 60 years, makes it possible to draw a more accurate profile of individual risk of embolic events. Aortic plaques are most often seen in elderly patients with hypertension, hypercholesterolaemia and a history of smoking (19). Screening for atherosclerotic lesions of the aorta in the cardiac surgical population may become even more important than in the general elderly population because most cardiac surgical patients are at greater risk of clinically significant aortic atherosclerosis (20). It has been reported, for example, that in patients undergoing surgery for aortic stenosis, the stroke rate triples (reaching 16%) in those showing complex arch atheroma (i.e., a thicker plaque with protrusion of mobile debris or ulceration or both) (20,21). Thorough knowledge of aortic anatomy thus can help us to choose the most appropriate management strategy, such as axillary cannulation, less traumatic cannula tips, balloon occlusion, soft cross clamps, circulatory arrest with aorta grafting, variations of "no touch" techniques and the recently introduced transapical valve replacement with off-pump grafting (20). Transoesophageal echocardiography has been reported to allow accurate assessment of the degree of atherosclerosis in the thoracic aorta (10). The presence of complex arch atheromas on TEE appears to be associated with an increased risk of perioperative stroke (20). However the mid and distal segments of the ascending aorta are often obscured by interposition of the tracheal tree. Unfortunately this "blind spot" on the aorta is the portion most often manipulated during surgery. Intraoperative epi-aortic ultrasound has been shown to be more sensitive than either palpation or TEE for detecting aortic plaques (22), but it is not routinely used because most surgical teams consider the traditional method of palpation equally reliable (23). Despite technical limitations and caveats aortic atheroma and stroke risk can be reasonably screened for preoperatively with the standard TEE and the routine use of ultrasound guidance in cardiac surgery may improve postoperative outcomes in this patient population. According to the guidelines of the American Echocardiography Society (24), the indication of TEE for evaluating the cardiovascular source of an embolic event in a patient who has a normal TTE and normal ECG and/or a history of atrial fibrillation/flutter is uncertain.

Different investigators have found a positive and significant correlation between atherosclerotic plaque dimension and embolic activity measured by transcranial Doppler monitoring (18,25,26). While all studies in subjects undergoing coronary angiographies (18,25,27) failed to demonstrate any correlation between emboli counts and clinical events, a positive and significant correlation has been found between transcranial microembolic signals and the risk of cerebral ischaemic complications among patients undergoing cardiac surgery (28). Therefore, in these subjects, an integrated evaluation (i.e., TEE plus TCD monitoring) may provide a more useful and accurate predictive index of embolic risk.

References

1. Atherosclerotic disease of the aortic arch as a risk factor for recurrent ischemic stroke. The French Study of Aortic Plaques in Stroke Group. *N Engl J Med* 1996;334:1216-1221
2. Kronzon I, Tunick PA. Aortic atherosclerotic disease and stroke. *Circulation* 2006;114:63-75
3. Ueno Y, Kimura K, Iguchi Y et al. Mobile aortic plaques are a cause of multiple brain infarcts seen on diffusion-weighted imaging. *Stroke* 2007;38:2470-2476
4. Di Tullio MR, Homma S, Jin Z, Sacco RL. Aortic atherosclerosis, hypercoagulability, and stroke: The APRIS (Aortic Plaque and Risk of Ischemic Stroke) study. *J Am Coll Cardiol* 2008;52:855-861
5. Mazighi M, Labreuche J, Gongora-Rivera F, Duyckaerts C, Hauw JJ, Amarenco P. Autopsy prevalence of proximal extracranial atherosclerosis in patients with fatal stroke. *Stroke* 2009;40:713-718
6. Uchino K, Estrera A, Calleja S, Alexandrov AV, Garami Z. Aortic dissection presenting as an acute ischemic stroke for thrombolysis. *J Neuroimaging* 2005;15:281-283
7. Grupper M, Eran A, Shifrin A. Ischemic stroke, aortic dissection, and thrombolytic therapy – the importance of basic clinical skills. *J Gen Intern Med* 2007;22:1370-1372
8. Khandheria BK, Tajik AJ, Taylor CL et al. Aortic dissection: review of value and limitations of two-dimensional echocardiography in a six-year experience. *J Am Soc Echocardiogr* 1989;2:17-2
9. Victor MF, Mintz GS, Kolter MN, Wilson AR, Segal BL. Two-dimensional echocardiographic diagnosis of aortic dissection. *Am J Cardiol* 1981;48:1155-1159
10. Nihoyannopoulos P, Joshi J, Athansopoulos G, Oakley CM. Detection of atherosclerotic lesions in the aorta by transesophageal echocardiography. *Am J Cardiol* 1993;71:1208-1212
11. Keren A, Kim CB, Hu BS et al. Accuracy of biplane and multiplane transesophageal echocardiography in diagnosis of typical acute aortic dissection and intramural hematoma. *J Am Coll Cardiol* 1996;28:627-636
12. Laissy JP, Blanc F, Soyer P et al. Thoracic aortic dissection: diagnosis with transesophageal echocardiography versus MR imaging. *Radiology* 1995;194:221-226

14. Fojtik JP, Costantino TG, Dean AJ. The diagnosis of aortic dissection by emergency medicine ultrasound. *J Emerg Med* 2007;32:191-196
15. Blaivas M, Sierzenski PR. Dissection of the proximal thoracic aorta: a new ultrasonographic sign in the subxiphoid view. *Am J Emerg Med* 2002;20:344-348
16. Macleod MR, Amarenco P, Davis SM, Donnan GA. Atheroma of the aortic arch: an important and poorly recognised factor in the aetiology of stroke. *Lancet Neurol* 2004;3:408-414
17. Fujimoto S, Yasaka M, Otsubo R, Oe H, Nagatsuka K, Minematsu K. Aortic arch atherosclerotic lesions and the recurrence of ischemic stroke. *Stroke* 2004;35:1426-1429
18. Bladin CF, Bingham L, Grigg L, Yapanis AG, Gerraty R, Davis SM. Transcranial Doppler detection of microemboli during percutaneous transluminal coronary angioplasty. *Stroke* 1998;29:2367-2370
19. Kronzon I, Tunick PA. Aortic atherosclerotic disease and stroke. *Circulation* 2006;114:63-75
20. Goland S, Trento A, Czer LS et al. Thoracic aortic arteriosclerosis in patients with degenerative aortic stenosis with and without coexisting artery disease. *Ann Thorac Surg* 2008;85:113-119
21. Culliford A. Invited Commentary to: Goland S, Trento A, Czer LS et al. Thoracic aortic arteriosclerosis in patients with degenerative aortic stenosis with and without coexisting artery disease. *Ann Thorac Surg* 2008;85:113-119
22. Suvarna S, Smith A, Stygall J et al. Intraoperative assessment of the ascending aorta: a comparison of digital palpation, transesophageal echocardiography, and epiaortic ultrasonography. *J Cardiothorac Vasc Anesth* 2007;21:805-809
23. Whitley WS, Glas KE. An argument for routine ultrasound screening of the thoracic aorta in the cardiac surgery population. *Semin Cardiothorac Vasc Anesth* 2008;12:290-297
24. Douglas P, Khandheria B, Stainback RF et al. American College of Cardiology Foundation Quality Strategic Directions Committee Appropriateness Criteria Working Group; American Society of Echocardiography; American College of Emergency Physicians; American Society of Nuclear Cardiology; Society for Cardiovascular Angiography and Interventions; Society of Cardiovascular Computed Tomography; Society for Cardiovascular Magnetic Resonance; American College of Chest Physicians; Society of Critical Care Medicine. ACCF/ASE/ACEP/ASNC/SCAI/SCCT/SCMR 2007 appropriateness criteria for transthoracic and transesophageal echocardiography: a report of the American College of Cardiology Foundation Quality Strategic Directions Committee Appropriateness Criteria Working Group, American Society of Echocardiography, American College of Emergency Physicians, American Society of Nuclear Cardiology, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and the Society for Cardiovascular Magnetic Resonance endorsed by the American College of Chest Physicians and the Society of Critical Care Medicine. *J Am Coll Cardiol* 2007;50:187-204
25. Castellani S, Bagnoli M, Polidori C, Giuglioli C, Margheri M, Gensini GF. Embolic signals from the middle cerebral arteries in patients undergoing coronary angiographies. *Coronary Artery Disease. From Prevention to Intervention*. Prague 2001:1-4
26. Rundek T, Di Tullio MR, Sciacca RR et al. Association between large aortic arch atheromas and high-intensity transient signals in elderly stroke patients. *Stroke* 1999;30:2683-2686
27. Hamon M, Gomes S, Clergeau MR, Fradin S, Morello R, Hamon M. Risk of acute brain injury related to cerebral microembolism during cardiac catheterization performed by right upper limb arterial access. *Stroke* 2007;38:2176-2179
28. Dittrich R, Ringelstein EB. Occurrence and clinical impact of microembolic signals during or after cardiosurgical procedures. *Stroke* 2008;39:503-511

