



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Ovarian response to Gonadotrophin stimulation in patients with previous endometriotic cystectomy

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Ovarian response to Gonadotrophin stimulation in patients with previous endometriotic cystectomy / ME Coccia; F Cammilli; L Ginocchini; F Borruto; F Rizzello.. - ELETTRONICO. - (2006), pp. 95-99. (Intervento presentato al convegno SEGI annual congress tenutosi a Milano nel May 10-13, 2006).

Availability:

The webpage <https://hdl.handle.net/2158/782007> of the repository was last updated on

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Ovarian Response to Gonadotrophin Stimulation in Patients with Previous Endometriotic Cystectomy

M.E. Coccia, F. Cammilli, L. Ginocchini, F. Borruto* and F. Rizzello

*Dept Gynaecology Perinatology and Human Reproduction,
University of Florence, Florence, Italy*

**Dept of Obstetrics and Gynaecology, University of Verona, Italy*

Summary

Aim of this study is to investigate the follicular response of ovaries after laparoscopic ovarian cystectomy for endometriotic cyst. Data from 298 infertile patients with endometriosis referred for IVF-ET/ICSI were reviewed. Forty five women who underwent laparoscopic excision of a monolateral endometriotic ovarian cyst before IVF-ET/ICSI were selected. Follicular responses of postcystectomy ovary and normal ovaries were compared.

The mean (\pm DS) number of follicles > 15 mm was 1.5 ± 1.6 in postcystectomy ovaries and 3.6 ± 2.8 in the control ovaries ($P < 0.05$). For women ≥ 35 years of age, the mean follicular response of normal ovaries and mean number of follicle of ≥ 15 mm were reduced significantly when compared to control ovaries in women < 35 years of age (4.1 ± 3.5 versus 8.7 ± 4.4 ; 1.9 ± 1.8 versus 5.1 ± 2.7);, while post-surgery ovaries showed a similar reduced response in both age groups.

Postcystectomy ovaries showed reduced follicular response particularly in women ≥ 35 years of age. Ovarian surgery in reproductive age, when proposed, should be performed with very cautious technique to preserve normal residual ovarian tissue.

Introduction

Laparoscopic surgery has become the gold standard for the treatment of ovarian endometriomas (*Canis M et al., 2001*). Endometriotic cysts represent one of the more common ovarian cysts treated by laparoscopic cystectomy also in women who suffered of the concomitant problem of infertility.

It has been proven that cystectomy may induce loss of follicular reserve during IVF, and ovarian cystectomy could provoke loss of normal ovarian tissue either by removing ovarian stroma with oocytes together with the capsule or by thermal damage provoked by coagulation. Studies have shown conflicting results, a detrimental effect or no adverse effect on ovarian response during ovarian stimulation for IVF after laparoscopic cystectomy (*Wong BC et al., 2004*). Loh et al demonstrated that postcystectomy ovaries have reduced follicular response in natural and stimulated cycles (*Loh FH et al., 1999*). Donnez, on the other hand, showed that internal vaporization of endometrioma does not impair IVF outcome (*Donnez J et al., 2001*).

Aim of our study was to evaluate the ovarian response to controlled ovarian hyperstimulation (COH) for ART in patients with a history of endometriomas treated by laparoscopic surgery.

Material and Methods

Data from 298 infertile patients with endometriosis referred at Department of Gynaecology Perinatology and Human Reproduction, University of Florence were reviewed. Forty-five women who underwent laparoscopic excision of a monolateral endometriotic ovarian cyst before IVF-ET/ICSI were included.

Patients

The average age of these 45 patients was 33.5 years (range, 24-41 years). 15 (33. %) had moderate disease, whereas 30 (66.7%) had severe disease, patient presented mild disease. All patients had unilateral endometriotic cyst. The mean size of the endometriotic cysts excised was 5.13 ± 2.9 cm.

All patients had transvaginal ultrasound evaluation of the ovaries pre-surgery. Contra lateral normal ovaries of patients with unilateral endometriotic cysts, based on ultrasound findings, were used as controls.

The ovarian response of 45 postcystectomy ovaries and 43 normal ovaries (controls) during monitored cycles was analyzed.

Ovarian Stimulation and Follicular Monitoring

The IVF-ET protocol used for ovarian stimulation was the long protocol with GnRH analogue plus gonadotropins. GnRHa (Decapeptyl® 0, 1 mg, IPSEN Italy) was injected subcutaneously starting on day 21 of the cycle. After down-regulation (adequate ovarian suppression at pelvic ultrasonography and circulating E_2 below 35 pg/ml), purified urinary FSH (Gonal -F®, 75 IU, sc Serono, Italy), 3 to 4 ampoules (225-300 IU), was administered. Human chorionic gonadotropins (hCG) 10.000 IU (Gonasi® 5000 IU, two ampoules; AMSA, Italy) was injected when at least two or more follicles reached a diameter 17-18 mm.

Ovulation was monitored with transvaginal ultrasound scans. Ovarian follicles were measured in two dimensions and the average diameter was calcu-

lated. The number of follicles with an average diameter ≥ 15 mm developed in each ovary just before HCG injection was noted. Follicular responses of postcystectomy ovary and normal ovaries were compared.

Statistical Analysis

All analyses were performed by using Statistical Package for the Social Science (SPSS for windows, Microsoft, version 13). Data are expressed in percentages or means \pm SD. Statistical analysis was performed with the *t* test for parametric data or χ^2 test for categorical data.

Statistical significance was set at $P < 0.05$.

Results

The overall results showed that the mean follicular response of post-cystectomy ovaries was reduced significantly compared with normal ovaries. Specifically, the mean number of follicles was 3.7 ± 3 in the previously operated ovary and 6.7 ± 5 in the control ovary ($P < 0.001$). The number of follicles ≥ 15 mm was significantly diminished in the previously operated ovary when compared with control ovary, too (1.5 ± 1.6 vs 3.6 ± 2.8).

Because age is an important determinant in ovarian response and decline in fecundity begin to accelerate after the age of 35 years (Hull MG *et al.*, 1996), the ovarian responses were stratified according to the age groups of < 35 or ≥ 35 years (Table 1).

Considering the group of younger women < 35 years of age the mean follicular response, the mean follicular response of postcystectomy ovaries was still reduced significantly compared with normal ovaries. For the older patients ≥ 35 years of age, even the normal ovaries showed poor response, and there was no statistically significant difference between the ovarian responses of postcystectomy and normal ovaries.

For women ≥ 35 years of age, the mean follicular response of normal ovaries and mean number of follicle of ≥ 15 mm were reduced significantly when compared to control ovaries in women < 35 years of age (4.1 ± 3.5 versus 8.7 ± 4.4 ; 1.9 ± 1.8 versus 5.1 ± 2.7), while post-surgery ovaries showed a similar reduced response in both age groups.

Table 1: Mean number of follicles produced by postcystectomy ovaries and normal ovaries, stratified according to age group of patients

	Age of < 35 years			Age of ≥ 35 years		
	Normal ovary	Postcystectomy ovary	<i>P</i>	Normal ovary	Postcystectomy ovary	<i>P</i>
Mean number of follicles \pm SD	8.7 \pm 4.4	4.3 \pm 3.4	<0.05	4.1 \pm 3.5	2.8 \pm 2.1	NS
Mean number of follicles ≥ 15 mm \pm SD	5.1 \pm 2.7	1.7 \pm 1.9	<0.05	1.9 \pm 1.8	1.3 \pm 1	NS

NS = not significant

Conclusions

Data literature shows controversy with respect to the surgical intervention for endometriotic cyst of the ovary before IVF-ET.

Excision of the cyst wall of the endometrioma may be removing part of the ovarian tissue and primordial follicles along with it. Ho et al (*Ho HY et al., 2002*) compared ovarian response in women previously treated by surgery for unilateral endometrioma. Diseased ovaries showed a significantly lower number of dominant follicles than normal ovaries.

Loh et al. noted that postcystectomy ovaries showed a reduced follicular response in natural and clomiphene-stimulated cycles, while they observed a response comparable to normal ovaries when stimulated with gonadotropins (*Loh FH et al., 1999*). Canis, in a retrospective cohort study, showed that number of oocytes was not significantly decreased by laparoscopic cystectomy (*Canis M et al., 2001*). In the same way, Marconi confirmed no negative effect of cystectomy (*Marconi G et al., 2002*). In a more recent study Garcia-Velasco demonstrates that laparoscopic cystectomy for endometriomas before commencing an IVF cycles does not compromise the number and quality of oocytes obtained (*Garcia-Velasco JA et al., 2004*), but he asked why perform surgery at all?

In our data, postcystectomy ovaries showed reduced follicular response in younger women. The reduction in response in women e" 35 years of age seems strictly connected with the age and no statistical significant difference in number of follicles was observed by comparing postcystectomy ovaries in both age groups.

There is still a debate over whether the presence of endometriomas may affect IVF adversely. In symptomatic patients, there is a clear benefit to surgery. On the other hand, the benefit of surgery in asymptomatic patient before controlled ovarian hyperstimulation is still matter of debate.

It is important not to generalize the data to all patients with endometriomas. Ovarian surgery should be proposed in selected cases (based on the size, symptoms) and a very cautious technique is recommended to preserve normal residual ovarian tissue, particularly in younger women.

References

- Canis M, Pouly JL, Tamburro S, et al. Ovarian response during IVF-embryo transfer cycles after laparoscopic ovarian cystectomy for endometriotic cysts of >3 cm in diameter. *Hum Reprod*,16:2583–2586 (2001)
- Donnez J, Wyns C, Nisolle M. Does ovarian surgery for endometriomas impair the ovarian response to gonadotropin? *Fertil Steril*,76(4):662-5 (2001).
- Garcia-Velasco JA, Mahutte NG, Corona J, Zuniga V, Giles J, Arici A, Pellicer A. Removal of endometriomas before in vitro fertilization does not improve fertility outcomes: a matched, case-control study. *Fertil Steril*.,81(5):1194-7 (2004).
- Ho HY, Lee RK, Hwu YM, Lin MH, Su JT, Tsai YC. Poor response of ovaries with endometrioma previously treated with cystectomy to controlled ovarian hyperstimulation. *J Assist Reprod Genet.*,19(11):507-11 (2002).
- Hull MG, Fleming CF, Hughes AO, McDermott A. The age-related decline in female fecundity: a quantitative controlled study of implanting capacity and survival of

- individual embryos after in vitro fertilization. *Fertil Steril.*,65(4):783-90 (1996).
- Loh FH, Tan AT, Kumar J, Ng SC. Ovarian response after laparoscopic ovarian cystectomy for endometriotic cysts in 132 monitored cycles.*Fertil Steril.*,72(2):316-21 (1999).
- Marconi G, Vilela M, Quintana R, Sueldo C. Laparoscopic ovarian cystectomy of endometriomas does not affect the ovarian response to gonadotropin stimulation. *Fertil Steril.*,78(4):876-8 (2002).
- Wong BC, Gillman NC, Oehninger S, et al. Results of in vitro fertilization in patients with endometriomas: is surgical removal beneficial? *Am J Obstet Gynecol*,191(2):597-606; discussion 606-7 (2004).