

# Lifestyle interventions in preventing new type 2 diabetes in Asian populations

Pietro Amedeo Modesti<sup>1</sup> · Giorgio Galanti<sup>2</sup> · Piergiuseppe Cala<sup>3</sup> · Maria Calabrese<sup>4</sup>

Received: 10 June 2015 / Accepted: 27 August 2015  
© SIMI 2015

**Abstract** The aim of this study was to review current evidence on interventional studies aimed at the prevention of type 2 diabetes in Asian population with lifestyle interventions. Prevalence of type 2 diabetes sharply increased in most Asian countries during the last decades. This issue has now also relevant implication for Europe where different surveys are also consistently revealing an higher prevalence of type 2 diabetes and other and major CVD risk factors among subjects originating from Asian Countries than in the native population. Nutrition and lifestyle transition seem to play a role in disclosing the predisposition for the development of type 2 diabetes and great interest is now shown toward the possibility to intervene with lifestyle intervention on at risk populations. A meta-analysis of Randomized Controlled Trials showed that lifestyle interventions are highly effective also in the Asian population. All studies were, however, conducted with an individual approach based on the identification of high-risk individuals. When ethnic minority groups have to be addressed, an approach directed to the community rather than to the individual might, however, be

more effective. This review reinforces the importance for policy-makers to consider the involvement of the whole community of minority immigrant groups with lifestyle intervention programs.

**Keywords** Type 2 diabetes prevention · Ethnicity · Ethnic minority groups · IDF geographical regions · Meta-analysis

## Diabetes in Asia; which implications for Europe?

Asia is the epicentre of the global epidemic of type 2 diabetes mellitus (T2DM) [1]. Over 138 million people with diabetes live in the Western Pacific area (98.4 million in China) and 72 million in the South-East Asia area (65.1 million in India). Most Asian countries experienced a relevant nutrition and lifestyle transition in the past few decades that played a role in disclosing the predisposition of Asian populations to the development of T2DM. China presented an innovative approach showing the importance of lifestyle intervention in the prevention of T2DM [2]. Other studies in the USA, Finland, Japan and India reached similar rates of success [3–6]. In the same line, the importance of physical exercise is now recognized in prevention, and the discipline of sports medicine has begun to widen its interests from athletes to cardiovascular and metabolic diseases in populations [7–9]. The current T2DM epidemic might thus be limited, and sustainable programs have been launched at national levels [10–16].

Changes occurring in Asian countries have now implications also for Europe where a high prevalence of T2DM is observed among Asian migrants originating from India, Pakistan, Bangladesh, Sri Lanka, Nepal, Bhutan and the Maldives [17–20]. For the majority of European countries,

---

**Electronic supplementary material** The online version of this article (doi:10.1007/s11739-015-1325-2) contains supplementary material, which is available to authorized users.

---

✉ Pietro Amedeo Modesti  
pamodesti@unifi.it

<sup>1</sup> Department of Medicina Sperimentale e Clinica, University of Florence, Largo Brambilla 3, 50134 Florence, Italy

<sup>2</sup> Sports Medicine Center, University of Florence, Florence, Italy

<sup>3</sup> Direzione generale Diritti di cittadinanza e Coesione Sociale, Regione Toscana, Florence, Italy

<sup>4</sup> U.O. Diabetologia, ASL 4 Prato, Ospedale Misericordia e Dolce, Prato, Italy

where immigration is a recent phenomenon, a common assumption is that health promotion interventions effective in the general population are also effective in ethnic minorities. Therefore, only a few countries specifically consider minority groups in their national health plans for metabolic and cardiovascular prevention. Cultural factors may, however, limit the ability of interventions addressed at the native population to reach minority groups [21]. The current relevant wave of immigration requires implementation of prevention strategies specifically addressed to minority ethnic groups [22].

### Ethnic origin and the risk of type 2 diabetes

Overweight and obesity are the main risk factors for the development of new T2DM, and the importance of body weight control is well recognized in prevention [21, 23]. Modest weight loss (5–10 % of body weight) and modest physical activity (30 min daily) indeed offer a variety of benefits in addition to the possibility of preventing or delaying T2DM [21]. Rendering targeted advice to at-risk individuals is thus especially imperative for physicians. The main question is: Who is to be considered at high risk? Subjects originating from South Asia, China, and Africa develop T2DM at a higher rate, at an earlier age, and at lower ranges of BMI than their European counterparts [24]. A clustering of different genetic defects or polymorphisms, the “thrifty gene” hypothesis, a genetic susceptibility to insulin resistance and higher central adiposity at similar BMI levels, or epigenetic changes occurring during gestation, might play a role in influencing metabolic phenotypes of Asian populations [25–27]. For these reasons, the WHO consultation group recommends a lower cutoff of BMI for Asians with respect to native European populations [28]. The identified diagnostic cutoff for being overweight is 23 kg/m<sup>2</sup> in India, and 24 kg/m<sup>2</sup> in China [29, 30]. Proper consideration of this issue by European physicians when facing these new patients in their offices might increase opportunities for education, intervention, and behaviour and lifestyle changes [31]. However, the main scientific societies in Europe still give little attention to the different cutoffs for overweight of ethnic minorities [21].

Furthermore, although ethnic origin has a value “per se” in the estimation of the risk for T2DM, doctors often base their prediction on blood glucose measured at opportunistic screening [32–36]. The risk of developing T2DM is over 5 times higher in subjects with impaired glucose tolerance (IGT) being 12 times higher in those with both IGT and impaired fasting glucose (IFG) compared to normoglycemic individuals [37]. However, blood glucose has limitations because (1) it is an invasive, costly, and time

consuming procedure; (2) it has a large random variation; (3) when taken in the aggregate, age, family history of T2DM, ethnicity, waist-to-hip ratio, BMI, blood pressure, and lipid levels, combined with plasma glucose levels are more predictive of future T2DM than glucose levels by themselves; (4) most importantly, the opportunities for an immigrant to have a blood test may be low, because of cultural barriers (varying rates of literacy, limited motivation linked to the lack of education on healthy lifestyle) and socioeconomic status [38–40]. Finally, primary prevention should be addressed at high-risk subjects when they are still in a normoglycemic state, and interventions should prevent their transition from normoglycemia to IFG and IGT. For these reasons, inexpensive, easily administered, cost-effective, and validated non-invasive risk scores (based on non-laboratory clinical variables) have been made available [41, 42]. These non-invasive scores identify a high risk of T2DM (*C* statistics  $\geq 0.8$ ) with acceptable to good discriminatory power across diverse settings in Europe [42–47]. Five non-invasive scores, the ARIC 2005, ARIC 2009, AUSDRISK, DPoRT, and QD Score, include the ethnic origin in the model [48–52]. To screen subjects at risk for T2DM, the American Diabetes Association (ADA) currently recommends a non-invasive tool that includes ethnicity (available at the URL: <http://www.ndep.nih.gov/am-i-at-risk/diabetes-risk-test.aspx>). A study performed in the USA compared three risk scores in a multi-ethnic population living in the same country [53]. In Europe, non-invasive tools were validated in different settings, although none of the diverse setting included ethnic minority groups [42].

The possibility of preventing T2DM was demonstrated in different studies, all including subjects on the basis of blood glucose values [54]. Some randomized controlled clinical trials were specifically conducted in Asian populations.

### Lifestyle interventions and prevention of type 2 diabetes in Asian populations

PUBMED and EMBASE were systematically searched for randomized controlled clinical trials investigating the effect of lifestyle interventions in preventing new T2DM in Asian populations. Eligible studies were controlled randomized clinical trials performed in Asian populations, reporting incidence of new cases of T2DM, whatever the duration of the study, published between 1994 and December 2014. Randomized clinical trials including patients with diabetes and duplicate publications or sub-studies of included trials were excluded. Two investigators independently abstracted the data from the eligible studies. Disagreements were resolved via consensus. Search was

conducted using terms, including type 2 diabetes mellitus, non-insulin-dependent diabetes, prevention, pre-diabetes, impaired fasting glucose (IFG), IGT, obesity, and lifestyle modification, limiting search to randomized controlled clinical trials and human studies. A manual search was also performed on reference lists from articles, reviews, and editorials. The measure of the effect of treatment was the difference in the incidence of new cases of T2DM in the intervention group and in the control group. For each study included in the meta-analysis, the incidences of T2DM in the control group and intervention group were used to process the forest plot. Intervention effect (new cases of T2DM) was expressed as odds ratio (OR), with 95 % confidence intervals (CIs); meta-analysis was performed by a random-effects model. Heterogeneity was examined with the  $I^2$ , where  $I^2$  values of 75 % or more indicated a high level of heterogeneity. The meta-analysis followed the PRISMA Checklist (Supplemental information) and was conducted using Rev Man 5.0 software (Fig. 1).


Eight randomized controlled clinical trials studies evaluated diet and physical activity interventions in Asian adults [2, 5, 6, 55–59]. The main characteristics of the included studies are reported in Table 1. The eight prospective randomized clinical studies include a total population of 2721 subjects at high risk for T2DM. All T2DM prevention trials performed in Asian populations have been based on a high-risk status defined by blood tests. In all studies, the inclusion criteria indeed required

IGT, assessed with oral glucose tolerance test (OGTT), or IFG [2, 5, 6, 55–59]. All eight RCTs were suitable for inclusion in the meta-analysis showing a 45 % reduction in the incidence of new T2DM in Asian subjects assigned to the intervention arm (OR 0.55; 95 % CI 0.44–0.70) (Fig. 2). Most importantly, a low level of heterogeneity was found among studies performed in different settings ( $I^2$  8 %).

The earliest Chinese study, published in 1997, started activities in 1986 when the prevalence of T2DM in China was still low (1–2 %) [2, 60, 61]. Pan et al. screened 577 men and women with IGT from a sample of 110,660 subjects. Subjects were then randomized to a programme of diet, exercise, or both [2]. Dietary intervention was focused on increased amounts of vegetables and reduced consumption of alcohol and simple sugars; overweight individuals (those with a BMI >25 kg/m<sup>2</sup>) were encouraged to lose weight. The exercise group was instructed to increase their daily activity by an equivalent of 20 min of moderate activity (brisk walking), and the diet-plus-exercise group was asked to do both exercise and dietary modification. After 6 years of follow-up, all three interventions were similarly effective, with risk reductions of 31–46 % compared to the control group. A recent long-term follow-up shows that the reduction in the 20-year cumulative incidence in the intervention groups (80 vs 93 % in the control group) [62] is paralleled by a reduction in the long-term cumulative incidence of CVD mortality (11.9 % in the intervention group vs 19.6 % in the control group, HR 0.59; 95 % CI 0.36–0.96;  $p = 0.033$ ) and all-cause mortality (28.1 vs 38.4, HR 0.71; 95 % CI 0.51–0.99;  $p = 0.049$ ) [62]. These data are the first available showing the impact on mortality of lifestyle intervention and are of great value in a country where 40 % of subjects aged 19–29 years were recently found to have a blood glucose level in the range of pre-diabetes [63].

In the study by Kosaka et al., 458 Japanese men with IFG and IGT were assigned to lifestyle ( $n = 102$ ) or standard intervention ( $n = 356$ ) [5]. OGTT was repeated every 6 months, and at the 4-year follow-up the cumulative incidence of T2DM was 9.3 and 3.0 % in the control and in the intervention groups, respectively. The reduction in the risk of T2DM is associated with larger body weight reduction (0.39 and 2.18 kg, respectively).

In the Indian Diabetes Prevention Programme (IDPP-1), 10,839 native Asian Indians underwent the screening and IGT was detected in 1027 subjects [6]. 1025 responded when called for a confirmatory OGTT, and a total of 531 subjects diagnosed with IGT on both tests (persistent IGT) were randomized into four groups (standard advice; lifestyle intervention; metformin; lifestyle plus metformin). Lifestyle intervention proved to be effective in this population with a very high 3-year cumulative incidence of




**Fig. 1** Selection of studies for the meta-analysis

**Table 1** Randomized controlled clinical trials (RCT) performed in Asian populations

Author	Year	Inclusion criteria	Ethnicity (Country)	Duration (months)	Arms	Outcome
Pan et al. [2]	1997	IGT	Chinese (China)	72	Control: general written instructions on diet and physical activity Intervention: personalized instruction and counselling for diet and physical exercise	New type 2 diabetes
Kosaka et al. [5]	2005	Men; IFG and IGT	Japanese (Japan)	48	Control: for BMI 24 kg/m <sup>2</sup> , take 5–10 % smaller meals than they had been taking and increase physical activity (lose weight); for BMI <24 kg/m <sup>2</sup> avoid gaining weight by diet and exercise Intervention: for BMI ≥22 kg/m <sup>2</sup> subjects were informed of their desirable body weight individually (suggestion to reduce weight at a rate of 0.5–1.0 kg/month); for BMI <22 kg/m <sup>2</sup> maintain weight and not gain weight	New type 2 diabetes
Ramachandran et al. [6]	2006	IGT on two occasions (persistent IGT)	Indians (India)	30	Control: subjects involved in physical labour, who had to walk or cycle for >30 min/day, or performing exercises regularly, were asked to continue their routine activities. Sedentary subjects were advised and regularly motivated to walk briskly for at least 30 min each day Intervention: subjects received advice on both healthy diet and regular physical activity. Scores were given based on the subjects' adherence. Diet modification including reduction in total calories, refined carbohydrates and fats, avoidance of sugar, and inclusion of fibre-rich foods	New type 2 diabetes
Sakane et al. [56]	2011	IFG and IGT	Japanese (Japan)	36	Control: one group session on a healthy lifestyle and prevention of diabetes. No individual guidance was given during the study period Intervention: specific advice on both healthy diet and regular physical activity. Goals of intervention: reduce body weight by 5 % in overweight and obese subjects, and to increase energy expenditure due to leisure time physical activity by 700 kcal per week. Advise: take the proper amount of calories; decrease the mean percent of energy derived from dietary fat (<25 %); restrict daily alcohol consumption (<160 kcal); eat three meals a day and avoid eating late at night; achieve the exercise goal, aerobic exercise such as walking is recommended	New type 2 diabetes
Saito et al. [55]	2011	Overweight with IFG and IGT	Japanese (Japan)	36	Control: reduce total energy intake and increase physical activity (5 % reduction in body weight), through the help of nurses, dieticians, physical therapists, and physicians. Instructions: four times at 12-month intervals Intervention: reduce total energy intake and increase physical activity (5 % reduction in body weight), through the help of nurses, dieticians, physical therapists, and physicians. Individual instructions and follow-up support from the medical staff at least nine times. Follow-up visits and use of self-monitoring sheets for recording body weight, pedometer counts, and how close they came to attaining their goals. Dietary intervention: reduce total energy intake mainly by restricting excess intake of fat or carbohydrates (fat intake at 20–25 % of total energy intake; carbohydrate intake at 55–60 % of total energy intake)	New type 2 diabetes

**Table 1** continued

Author	Year	Inclusion criteria	Ethnicity (Country)	Duration (months)	Arms	Outcome
Iqbal Hydrie et al. [57]	2012	Age >30 years; IGT	Pakistani (Pakistan)	18	Control: subjects received general diet and exercise information at baseline and at subsequent visits; no intensive individual specific counselling was given Intervention: training sessions and counselling were given to subjects in the intervention group about intervention goals (body weight loss $\geq 5\%$ via diet control and physical exercise, total fat intake <30% of energy consumed, fibre intake of 15 g/1000 kcal, and moderate exercise >30 min/day). Sessions with dietitians and physical trainers at each visit and individual counselling to increase physical activity (endurance exercises such as walking, jogging, and cycling)	New type 2 diabetes
Ramachandran et al. [58]	2013	Men; owning a mobile phone; familial history of type 2 diabetes; BMI >23 kg/m <sup>2</sup> ; IGT	Indians (India)	20	Control: personalized education and motivation about health lifestyle principles, and written information about diet and physical activity Intervention: in addition, a manager website delivered cyclically phone message reminders (2–4 message per week, not the same message in a 6-month period)	New type 2 diabetes
Bhopal et al. [59]	2014	IFG; IGT; waist: >90 cm M or >80 cm in W	South Asians (UK)	36	Control: standardized written and verbal advice Intervention: consultation with a dietitian (both participants and family volunteers were part of this intervention) with 15 visits over 3 years (baseline, monthly for the first 3 months, then every 3 months) on achieving weight loss through a calorie-deficit diet and physical activity (>30 min daily brisk walking), using culturally adapted and translated resources (information on shopping and cooking; 3-day food diaries; dietary patterns questionnaire). Annual group sessions, including a food shopping tour and brisk walking. Pedometers were given to provide step counts for motivation through self-monitoring and for the dietitians to assess progress. Bodyweight and waist circumference data were used as motivational devices by dietitians	Weight change at 3 years

**Fig. 2** Odds ratio for the reduction of type 2 diabetes in randomized clinical trials which evaluated the effects of non-pharmacologic lifestyle interventions (physical exercise and diet) in Asian adults

T2DM (55, 39, 40 and 39 % in groups 1–4, respectively) [6].

Sakane et al. performed a lifestyle intervention program in a primary health-care setting on 304 middle-aged Japanese subjects with IGT [56]. The 3-year cumulative incidence tended to be lower in the intervention group (14.8 vs 8.2 %).

The incidence of T2DM was high in the 641 Japanese overweight subjects (aged 30–60 years) with IFG and IGT enrolled by Saito et al. [55]. At 36 months follow-up, the estimated cumulative incidences of T2DM were 12.2 % in the frequent intervention group ( $n = 311$ ) and 16.6 % in the control group ( $n = 330$ ).

In the study performed in Pakistan by Iqbal Hydrie et al., 5000 people attended screening camps where 2300 persons filled the questionnaire and 1825 subjects were identified to be at high risk [57]. Of the 1739 subjects who took the oral glucose tolerance test, 317 were identified as IGT and randomized into three groups (control; intense lifestyle modification advice; intense lifestyle modification advice and metformin 500 mg twice daily). The overall incidence was 4 cases per 1000 person-months, with an incidence of 8.6 cases in the control group, 2.5 cases in the lifestyle modification advice group, and 2.3 cases in the lifestyle modification advice + drug group. The overall compliance rate was 86 %.

An innovative technological approach based on mobile phone messaging in the prevention of T2DM was explored by Ramachandran et al. who screened male Indian subjects with positive familial history of T2DM or BMI >23 kg/m<sup>2</sup> before performing the IGT test [45, 58]. In this Indian study, the 537 working Indian men with IGT were randomly allocated to receive mobile phone messages ( $n = 271$ ) or standard care ( $n = 266$ ). All participants received diet and exercise advice at baseline. The cumulative incidence of T2DM was significantly reduced in the intervention group compared with the standard-care group [50 (18 %) of 271 participants vs 73 (27 %) of 266; hazard ratio 0.64, 95 % CI 0.45–0.92;  $p = 0.015$ ]. The trial is the first to report a reduction in the onset of T2DM with a mobile messaging intervention. The investigators recorded a high response rate of 96 % ( $n = 517$ ) with a duration of follow-up of 2 years. The nature of the intervention prevented masking of the participants and field staff; however, the laboratory staff and the principal and co-investigators were masked [58]. If the results of this trial are replicated in other settings, mobile phone messaging might be a practical and affordable way to deliver lifestyle advice to delay or prevent the onset of T2DM.

The first study investigating the effects of lifestyle modifications on progression to T2DM in South Asians (Indian or Pakistani) living in the UK was recently performed by Bhopal et al. [59]. Subjects aged 35 years or

older, with waist circumference 90 cm or greater in men or 80 cm or greater in women ( $n = 1319$ ), were screened by oral glucose tolerance test; 196 (15 %) had IGT or IFG and 171 entered the trial and were randomized to intervention (78 families with 85 participants) or control (78 families with 86 participants). Interestingly, the protocol considered the supporting role of the family as important, because participants in the same family were not randomized separately. At 3 years, progression to T2DM (either doctor diagnosed or by oral glucose tolerance test at 3 years) was observed less frequently in the intervention group (15 %) than the control group (21 %), although the difference was not statistically significant (95 % CI 0.27–1.67;  $p = 0.3705$ ).

This meta-analysis is associated with several important limitations. First, this review is limited to published randomized controlled trials, while observational studies and additional unpublished literature are excluded. Although randomized controlled trials generally provide maximum validity and causal inference, non-randomized studies performed at the community level may have been lost. Second, only one study included in the meta-analysis enrolled Asian subjects living in Europe. Further research is therefore needed to better identify effective interventions for the prevention of T2DM in Asian minority groups living in Europe, an issue now crucial for Europe [64]. Third, we did not include subjects from America or Africa, which may limit the generalizability of our results to African or other populations.

Notwithstanding these limitations, current randomized clinical trials seem to indicate that T2DM can be prevented in Asian individuals who are at high risk. However, when the intervention is targeted at Asian communities living in Europe, some aspects have to be considered.

### **A new perspective for lifestyle interventions in minority groups: from individual to collectivist goals**

The main limitations in the high-risk approach to minority groups are represented by costs of the screening procedure, low compliance at follow-up, and the problematic contact with undocumented migrants.

A comparison of potential screening strategies and subsequent interventions for the prevention and treatment of T2DM was performed by Gillies et al., who considered a hypothetical population of the UK, aged 45 years, at the time of screening [65]. In their estimation, lifestyle intervention strategy shows a small clinical potential benefit in terms of average years spent without T2DM and cases of T2DM prevented, because discounted QALYs gained compared with no screening were 0.09 (0.03–0.17) for

screening and lifestyle interventions. The hypothetical population had 17 % of either IGT or undiagnosed T2DM at the time of screening. However, as noted by Gillies et al., when an increased prevalence of IGT and T2DM are considered, the QALYs decrease, and most importantly the total costs of the screening strategy increase [65]. When the model was run for a South Asian cohort, with high prevalence of T2DM, the results for QALYs were reduced and costs of screening increased. Notwithstanding these limitations, Gillies et al. consider lifestyle interventions to be cost-effective compared with no screening in an “at-risk” population [65].

A recent trial performed in the Netherlands studied the effectiveness of an intensive, culturally targeted lifestyle intervention in general practice for weight status and metabolic profile of South Asians at risk of T2DM [66]. Although T2DM incidence is not included among outcomes, the trial is of great interest because limitations connected with the high-risk approach in minority groups are clearly shown. Low initial response rate and laborious recruitment, high dropout rate, and the lack of effect of the lifestyle intervention on weight change and other metabolic parameters indeed led the authors to raise the question whether the high-risk strategy is the optimal approach to prevent T2DM in minority groups [66]. Admiraal et al. conclude that health gain might be better achieved by focusing on prevention strategies that tackle the high risk of T2DM among South Asians at an earlier stage [67].

The large number of subjects not included in population statistics (hidden population of undocumented migrants) is also often not considered in trials and cost-effectiveness analysis. Approximately, 1.7 million Chinese people are now estimated to live in Europe, mostly in the UK, France, Italy, Germany, and Spain, and migration from China to Europe has now been mainly concentrated in countries of Southern Europe, e.g. Italy and Spain [68]. In China, a prevalence of T2DM of 11.6 %, with over 40 % of young adults (aged 18-to-29 years) at risk was found in 2010 [63, 69]. Should the prevalence of prediabetes and T2DM in the Chinese living in Europe be as high as in China, the prevention of T2DM could be a need for our health system [63, 70]. Many minority patients have difficulty communicating with their health-care providers and other cultural barriers may exist [71]. A qualitative study performed in the UK identifies a main barrier for the Bangladeshi community in the complex value hierarchy of what is accepted to be healthy (small portion size, limited rich and fatty food, regular activity) and what is important for the social norms of hospitality, the religious requirement for modesty, and the cultural rejection of a “sporting” identity or dress (especially for women, older people, and senior members of the society) [72]. Contemporary health

promotion, usually based on assumptions of a self-investment, should thus leave the approach to individuals when the aim is to involve societies with a collectivist history [73]. This paradigm change is now required by an ever-changing society.

On this basis, a new strategic plan aimed at the prevention of new T2DM in the Chinese community settled in Prato has been approved recently in Italy by the Regione Toscana. The project aim is to encourage weight control and physical activity with interventions on the Chinese community. The local Chinese school will be the setting to contact families. The project will include: distribution of the population of Chinese training materials on T2DM and the correct rules on nutrition and lifestyles; training courses on the importance of nutrition in the prevention of T2DM directed at teachers of the Chinese School and to canteen managers of Chinese factories; promotion of physical activity through the creation of sports groups aggregation of Chinese entities for the realization of projects of sport activity (formation of teams with structured training courses, realization of tournaments and amateur competitions). Objectives for action include the reduction of 5 % loss of body weight through the control diet (total intake of fat less than 30 % of energy consumed, fibre intake per day of 15 g/1000 kcal) and the promotion of moderate exercise (30 min/day with resistance exercises such as walking, jogging, and cycling).

## Conclusion

The review identifies three issues to be considered in primary care: (1) ethnic origin is an important element for risk assessment of future T2DM, and specific diagnostic cutoffs for overweight have to be considered in general practice; (2) lifestyle interventions are particularly important to reduce the future incidence of T2DM in the Asian population living in Europe; (3) the approach to individual high-risk subjects followed in randomized controlled trials is effective, although limitations exist when pursued in minority groups and a new perspective for an approach specifically involving the whole communities is now needed for Europe.

It is now important to launch public health promotion programs of primary prevention, specifically addressed to minority groups, and the cooperation of ethnic communities is essential. Educational prevention programs are important to limit the development of new T2DM in Europe.

## Compliance with ethical standards

**Conflict of interest** None.

**Statement of human and animal rights** This article does not contain any studies with human participants or animals performed by any of the authors.

**Informed consent** For this type of study formal consent is not required.

## References

- Goldberg R, Temprosa M, Otvos J, Brunzell J, Marcovina S, Mather K, Arakaki R, Watson K, Horton E, Barrett-Connor E (2013) Lifestyle and metformin treatment favorably influence lipoprotein subfraction distribution in the Diabetes Prevention Program. *J Clin Endocrinol Metab* 98(10):3989–3998. doi:10.1210/jc.2013-1452
- Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV (1997) Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 20(4):537–544. doi:10.2337/diacare.20.4.537
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM, Diabetes Prevention Program Research Group (2002) Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346(6):393–403. doi:10.1056/NEJMoa012512
- Lindstrom J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, Hamalainen H, Harkonen P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Mannelin M, Paturi M, Sundvall J, Valle TT, Uusitupa M, Tuomilehto J (2006) Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet* 368(9548):1673–1679. doi:10.1016/S0140-6736(06)69701-8
- Kosaka K, Noda M, Kuzuya T (2005) Prevention of type 2 diabetes by lifestyle intervention: a Japanese trial in IGT males. *Diabetes Res Clin Pract* 67(2):152–162. doi:10.1016/j.diabres.2004.06.010
- Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V (2006) The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia* 49(2):289–297. doi:10.1007/s00125-005-0097-z
- Bushman B (2014) Promoting exercise as medicine for prediabetes and prehypertension. *Curr Sports Med Rep* 13(4):233–239. doi:10.1249/jsr.0000000000000066
- Lambert EV, Bull F (2014) Public health recommendations for physical activity in the prevention of type 2 diabetes mellitus. *Med Sport Sci* 60:130–140. doi:10.1159/000357343
- Fernandes RA, Coelho ESMJ, Spiguel Lima MC, Cayres SU, Codogno JS, Lira FS (2015) Possible underestimation by sports medicine of the effects of early physical exercise practice on the prevention of diseases in adulthood. *Curr Diabetes Rev* 11(3):201–205
- Hopper I, Billah B, Skiba M, Krum H (2011) Prevention of diabetes and reduction in major cardiovascular events in studies of subjects with prediabetes: meta-analysis of randomised controlled clinical trials. *Eur J Cardiovasc Prev Rehabil* 18(6):813–823. doi:10.1177/1741826711421687
- Schellenberg ES, Dryden DM, Vandermeer B, Ha C, Korownyk C (2013) Lifestyle interventions for patients with and at risk for type 2 diabetes: a systematic review and meta-analysis. *Ann Intern Med* 159(8):543–551. doi:10.7326/0003-4819-159-8-201310150-00007
- Hamilton MT, Hamilton DG, Zderic TW (2014) Sedentary behavior as a mediator of type 2 diabetes. *Med Sport Sci* 60:11–26. doi:10.1159/000357332
- Saaristo T, Moilanen L, Korpi-Hyovalti E, Vanhala M, Saltevo J, Niskanen L, Jokelainen J, Peltonen M, Oksa H, Tuomilehto J, Uusitupa M, Keinanen-Kiukkaanniemi S (2010) Lifestyle intervention for prevention of type 2 diabetes in primary health care one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D). *Diabetes Care* 33(10):2146–2151. doi:10.2337/Dc10-0410
- Tuomilehto J, Schwarz P, Lindstrom J (2011) Long-term benefits from lifestyle interventions for type 2 diabetes prevention time to expand the efforts. *Diabetes Care* 34:S210–S214. doi:10.2337/Dc11-S222
- Assoc AD (2013) Economic costs of diabetes in the US in 2012. *Diabetes Care* 36(4):1033–1046. doi:10.2337/Dc12-2625
- Hermansen K, Baekdal TA, Daring M, Pietraszek A, Mortensen LS, Jorgensen H, Flint A (2013) Liraglutide suppresses postprandial triglyceride and apolipoprotein B48 elevations after a fat-rich meal in patients with type 2 diabetes: a randomized, double-blind, placebo-controlled, cross-over trial. *Diabetes Obes Metab* 15(11):1040–1048. doi:10.1111/dom.12133
- McKeigue PM, Marmot MG, Syndercombe Court YD, Cottier DE, Rahman S, Riemersma RA (1988) Diabetes, hyperinsulinaemia, and coronary risk factors in Bangladeshis in east London. *Br Heart J* 60(5):390–396
- Shai I, Jiang R, Manson JE, Stampfer MJ, Willett WC, Colditz GA, Hu FB (2006) Ethnicity, obesity, and risk of type 2 diabetes in women—a 20-year follow-up study. *Diabetes Care* 29(7):1585–1590. doi:10.2337/Dc06-0057
- Khoo CM, Sairazi S, Taslim S, Gardner D, Wu Y, Lee J, van Dam RM, Tai ES (2011) Ethnicity modifies the relationships of insulin resistance, inflammation, and adiponectin with obesity in a multiethnic Asian population. *Diabetes Care* 34(5):1120–1126. doi:10.2337/Dc10-2097
- Wang C, Li J, Xue H, Li Y, Huang J, Mai J, Chen J, Cao J, Wu X, Guo D, Yu L, Gu D (2014) Type 2 diabetes mellitus incidence in Chinese: contributions of overweight and obesity. *Diabetes Res Clin Pract*. doi:10.1016/j.diabres.2014.09.059
- Ryden L, Grant PJ, Anker SD, Berne C, Cosentino F, Danchin N, Deaton C, Escaned J, Hammes HP, Huikuri H, Marre M, Marx N, Mellbin L, Ostergren J, Patrono C, Seferovic P, Uva MS, Taskinen MR, Tendera M, Tuomilehto J, Valensi P, Zamorano JL (2013) ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 34(39):3035–3087. doi:10.1093/eurheartj/eh108
- Rechel B, Mladovsky P, Ingleby D, Mackenbach JP, McKee M (2013) Migration and health in an increasingly diverse Europe. *Lancet* 381(9873):1235–1245. doi:10.1016/S0140-6736(12)62086-8
- Barba C, Cavalli-Sforza T, Cutter J, Darnton-Hill I, Deurenberg P, Deurenberg-Yap M, Gill T, James P, Ko G, Miu AH, Kosulwat V, Kumanyika S, Kurpad A, Mascie-Taylor N, Moon HK, Nishida C, Noor MI, Reddy KS, Rush E, Schultz JT, Seidell J, Stevens J, Swinburn B, Tan K, Weisell R, Wu ZS, Yajnik CS, Yoshiike N, Zimmet P, Consultation WE (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363(9403):157–163
- Chiu M, Austin PC, Manuel DG, Shah BR, Tu JV (2011) Deriving ethnic-specific BMI cutoff points for assessing diabetes risk. *Diabetes Care* 34(8):1741–1748. doi:10.2337/dc10-2300
- Carulli L, Rondinella S, Lombardini S, Canedi I, Loria P, Carulli N (2005) Review article: diabetes, genetics and ethnicity. *Aliment Pharmacol Ther* 22:16–19. doi:10.1111/j.1365-2036.2005.02588.x



26. Yajnik CS, Lubree HG, Rege SS, Naik SS, Deshpande JA, Deshpande SS, Joglekar CV, Yudkin JS (2002) Adiposity and hyperinsulinemia in Indians are present at birth. *J Clin Endocr Metab* 87(12):5575–5580. doi:[10.1210/jc.2002-020434](https://doi.org/10.1210/jc.2002-020434)
27. McKeigue PM, Pierpoint T, Ferrie JE, Marmot MG (1992) Relationship of glucose-intolerance and hyperinsulinemia to body-fat pattern in South Asians and Europeans. *Diabetologia* 35(8):785–791
28. WHO (2004) Appropriate body-mass index in Asian populations and its implications for policy and intervention strategies. *Lancet* 363(9412):902
29. Misra A, Chowbey P, Makkar BM, Vikram NK, Wasir JS, Chadha D, Joshi SR, Sadikot S, Gupta R, Gulati S, Munjal YP, Concensus G (2009) Consensus statement for diagnosis of obesity, abdominal obesity and the metabolic syndrome for Asian Indians and recommendations for physical activity, medical and surgical management. *J Assoc Phys India* 57:163–170
30. Bei-Fan Z, Working CMAG (2002) Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults: study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Asia Pac J Clin Nutr* 11:S685–S693. doi:[10.1046/j.1440-6047.11.s8.9.x](https://doi.org/10.1046/j.1440-6047.11.s8.9.x)
31. Hsu WC, Araneta MRG, Kanaya AM, Chiang JL, Fujimoto W (2015) BMI cut points to identify at-risk Asian Americans for type 2 diabetes screening. *Diabetes Care* 38(1):150–158. doi:[10.2337/Dc14-2391](https://doi.org/10.2337/Dc14-2391)
32. Edelstein SL, Knowler WC, Bain RP, Andres R, BarrettConnor EL, Dowse GK, Haffner SM, Pettitt DJ, Sorkin JD, Muller DC, Collins VR, Hamman RF (1997) Predictors of progression from impaired glucose tolerance to NIDDM—an analysis of six prospective studies. *Diabetes* 46(4):701–710. doi:[10.2337/diabetes.46.4.701](https://doi.org/10.2337/diabetes.46.4.701)
33. Shaw JA, Zimmet PZ, de Courten M, Dowse GK, Chitson P, Gareeboo H, Hemraj F, Fareed D, Tuomilehto J, Alberti KGMM (1999) Impaired fasting glucose or impaired glucose tolerance—what best predicts future diabetes in Mauritius? *Diabetes Care* 22(3):399–402. doi:[10.2337/diacare.22.3.399](https://doi.org/10.2337/diacare.22.3.399)
34. Shaw JE, de Courten M, Boyko EJ, Zimmet PZ (1999) Impact of new diagnostic criteria for diabetes on different populations. *Diabetes Care* 22(5):762–766. doi:[10.2337/diacare.22.5.762](https://doi.org/10.2337/diacare.22.5.762)
35. Gabir MM, Hanson RL, Dabelea D, Imperatore G, Roumain J, Bennett PH, Knowler WC (2000) The 1997 American Diabetes Association and 1999 World Health Organization criteria for hyperglycemia in the diagnosis and prediction of diabetes. *Diabetes Care* 23(8):1108–1112. doi:[10.2337/diacare.23.8.1108](https://doi.org/10.2337/diacare.23.8.1108)
36. de Vegt F, Dekker JM, Jager A, Hienkens E, Kostense PJ, Stehouwer CDA, Nijpels G, Bouter LM, Heine RJ (2001) Relation of impaired fasting and postload glucose with incident type 2 diabetes in a Dutch population—the Hoorn study. *JAMA* 285(16):2109–2113. doi:[10.1001/jama.285.16.2109](https://doi.org/10.1001/jama.285.16.2109)
37. Gerstein HC, Santaguida P, Raina P, Morrison KM, Balion C, Hunt D, Yazdi H, Booker L (2007) Annual incidence and relative risk of diabetes in people with various categories of dysglycemia: a systematic overview and meta-analysis of prospective studies. *Diabetes Res Clin Pract* 78(3):305–312. doi:[10.1016/j.diabres.2007.05.004](https://doi.org/10.1016/j.diabres.2007.05.004)
38. Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer HM, Byrd-Holt DD (1998) Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in US adults—the Third National Health and Nutrition Examination Survey, 1988–1994. *Diabetes Care* 21(4):518–524. doi:[10.2337/diacare.21.4.518](https://doi.org/10.2337/diacare.21.4.518)
39. Stern MP, Williams K, Haffner SA (2002) Identification of persons at high risk for type 2 diabetes mellitus: do we need the oral glucose tolerance test? *Ann Intern Med* 136(8):575–581
40. Modesti PA, Bianchi S, Borghi C, Cameli M, Capasso G, Ceriello A, Ciccone MM, Germano G, Maiello M, Muiesan ML, Novo S, Padeletti L, Palmiero P, Pillon S, Rotella CM, Saba PS, Scicchitano P, Trimarco B, Volpe M, Pedrinelli R, Di Biase M (2014) Cardiovascular health in migrants: current status and issues for prevention. A collaborative multidisciplinary task force report. *J Cardiovasc Med* 15(9):683–692. doi:[10.2459/Jcm.0000000000000069](https://doi.org/10.2459/Jcm.0000000000000069)
41. American Diabetes A (2013) Standards of medical care in diabetes—2013. *Diabetes Care* 36(Suppl 1):S11–S66. doi:[10.2337/dc13-S011](https://doi.org/10.2337/dc13-S011)
42. Kengne AP, Beulens JWJ, Peelen LM, Moons KGM, van der Schouw YT, Schulze MB, Spijkerman AMW, Griffin SJ, Grobbee DE, Palla L, Tormo MJ, Arriola L, Barengo NC, Barricarte A, Boeing H, Bonet C, Clavel-Chapelon F, Dartois L, Fagherazzi G, Franks PW, Huerta JM, Kaaks R, Key TJ, Khaw KT, Li KR, Muhlenbruch K, Nilsson PM, Overvad K, Overvad TF, Palli D, Panico S, Quiros JR, Rolandsson O, Roswall N, Sacerdote C, Sanchez MJ, Slimani N, Tagliabue G, Tjonneland A, Tumino R, van der A DL, Forouhi NG, Sharp SJ, Langenberg C, Riboli E, Wareham NJ (2014) Non-invasive risk scores for prediction of type 2 diabetes (EPIC-InterAct): a validation of existing models. *Lancet Diabetes Endocrinol* 2(1):19–29. doi:[10.1016/S2213-8587\(13\)70103-7](https://doi.org/10.1016/S2213-8587(13)70103-7)
43. Lindstrom J, Tuomilehto J (2003) The diabetes risk score—a practical tool to predict type 2 diabetes risk. *Diabetes Care* 26(3):725–731. doi:[10.2337/diacare.26.3.725](https://doi.org/10.2337/diacare.26.3.725)
44. Buijse B, Simmons RK, Griffin SJ, Schulze MB (2011) Risk assessment tools for identifying individuals at risk of developing type 2 diabetes. *Epidemiol Rev* 33(1):46–62. doi:[10.1093/epirev/mxq019](https://doi.org/10.1093/epirev/mxq019)
45. Collins GS, Mallett S, Omar O, Yu LM (2011) Developing risk prediction models for type 2 diabetes: a systematic review of methodology and reporting. *BMC Med*. doi:[10.1186/1741-7015-9-103](https://doi.org/10.1186/1741-7015-9-103)
46. Noble D, Mathur R, Dent T, Meads C, Greenhalgh T (2011) Risk models and scores for type 2 diabetes: systematic review. *BMJ* 343:d7163. doi:[10.1136/bmj.d7163](https://doi.org/10.1136/bmj.d7163)
47. Abbasi A, Peelen LM, Corpeleijn E, van der Schouw YT, Stolk RP, Spijkerman AMW, van der A DL, Moons KGM, Navis G, Bakker SJL, Beulens JWJ (2012) Prediction models for risk of developing type 2 diabetes: systematic literature search and independent external validation study. *BMJ*. doi:[10.1136/bmj.e5900](https://doi.org/10.1136/bmj.e5900)
48. Schmidt MI, Duncan BB, Bang H, Pankow JS, Ballantyne CM, Golden SH, Folsom AR, Chambless LE (2005) Identifying individuals at high risk for diabetes—the Atherosclerosis Risk in Communities study. *Diabetes Care* 28(8):2013–2018. doi:[10.2337/diacare.28.8.2013](https://doi.org/10.2337/diacare.28.8.2013)
49. Kahn HS, Cheng YJ, Thompson TJ, Imperatore G, Gregg EW (2009) Two risk-scoring systems for predicting incident diabetes mellitus in US adults age 45–64 years. *Ann Intern Med* 150(11):741–751
50. Chen L, Magliano DJ, Balkau B, Colagiuri S, Zimmet PZ, Tonkin AM, Mitchell P, Phillips PJ, Shaw JE (2010) AUSDRISK: an Australian type 2 diabetes risk assessment tool based on demographic, lifestyle and simple anthropometric measures. *Med J Aust* 192(4):197–202
51. Rosella LC, Manuel DG, Burchill C, Stukel TA, team ftP-D (2011) A population-based risk algorithm for the development of diabetes: development and validation of the Diabetes Population Risk Tool (DPoRT). *J Epidemiol Commun Health* 65(7):613–620. doi:[10.1136/jech.2009.102244](https://doi.org/10.1136/jech.2009.102244)
52. Hippisley-Cox J, Coupland C, Robson J, Sheikh A, Brindle P (2009) Predicting risk of type 2 diabetes in England and Wales:

- prospective derivation and validation of QDScore. *BMJ*. doi:10.1136/bmj.b880
53. Mann DM, Bertoni AG, Shimbo D, Carnethon MR, Chen HY, Jenny NS, Muntner P (2010) Comparative validity of 3 diabetes mellitus risk prediction scoring models in a multiethnic US cohort. *Am J Epidemiol* 171(9):980–988. doi:10.1093/Aje/Kwq030
  54. Merlotti C, Morabito A, Pontiroli AE (2014) Prevention of type 2 diabetes; a systematic review and meta-analysis of different intervention strategies. *Diabetes Obes Metab* 16(8):719–727. doi:10.1111/dom.12270
  55. Saito T, Watanabe M, Nishida J, Izumi T, Omura M, Takagi T, Fukunaga R, Bandai Y, Tajima N, Nakamura Y, Ito M (2011) Lifestyle modification and prevention of type 2 diabetes in overweight Japanese with impaired fasting glucose levels A Randomized Controlled Trial. *Arch Intern Med* 171(15):1352–1360
  56. Sakane N, Sato J, Tsushita K, Tsujii S, Kotani K, Tsuzaki K, Tominaga M, Kawazu S, Sato Y, Usui T, Kamae I, Yoshida T, Kiyohara Y, Sato S, Kuzuya H (2011) Prevention of type 2 diabetes in a primary healthcare setting: three-year results of lifestyle intervention in Japanese subjects with impaired glucose tolerance. *BMC Public Health* 11(1):40. doi:10.1186/1471-2458-11-40
  57. Iqbal Hydrie MZ, Basit A, Shera AS, Hussain A (2012) Effect of intervention in subjects with high risk of diabetes mellitus in pakistan. *J Nutr Metab* 2012:867604. doi:10.1155/2012/867604
  58. Ramachandran A, Snehalatha C, Ram J, Selvam S, Simon M, Nanditha A, Shetty AS, Godsland IF, Chaturvedi N, Majeed A, Oliver N, Toumazou C, Alberti KG, Johnston DG (2013) Effectiveness of mobile phone messaging in prevention of type 2 diabetes by lifestyle modification in men in India: a prospective, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol* 1(3):191–198. doi:10.1016/S2213-8587(13)70067-6
  59. Bhopal RS, Douglas A, Wallia S, Forbes JF, Lean ME, Gill JM, McKnight JA, Sattar N, Sheikh A, Wild SH, Tuomilehto J, Sharma A, Bhopal R, Smith JB, Butcher I, Murray GD (2014) Effect of a lifestyle intervention on weight change in south Asian individuals in the UK at high risk of type 2 diabetes: a family-cluster randomised controlled trial. *Lancet Diabetes Endocrinol* 2(3):218–227. doi:10.1016/s2213-8587(13)70204-3
  60. National\_Diabetes\_Research\_Group (1981) A mass survey of diabetes mellitus in a population of 300,000 in 14 provinces and municipalities in China (author's transl). *Zhonghua Nei Ke Za Zhi* 20(11):678–683
  61. Pan XR, Yang WY, Li GW, Liu J (1997) Prevalence of diabetes and its risk factors in China, 1994. *Diabetes Care* 20(11):1664–1669. doi:10.2337/diacare.20.11.1664
  62. Li GW, Zhang P, Wang JP, Gregg EW, Yang WY, Gong QH, Li H, Li HL, Jiang YY, An YL, Shuai Y, Zhang B, Zhang JL, Thompson TJ, Gerzoff RB, Roglic G, Hu YH, Bennett PH (2008) The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 371(9626):1783–1789. doi:10.1016/S0140-6736(08)60766-7
  63. Xu Y, Wang L, He J, Bi Y, Li M, Wang T, Wang L, Jiang Y, Dai M, Lu J, Xu M, Li Y, Hu N, Li J, Mi S, Chen CS, Li G, Mu Y, Zhao J, Kong L, Chen J, Lai S, Wang W, Zhao W, Ning G (2013) Prevalence and control of diabetes in Chinese adults. *JAMA* 310(9):948–959. doi:10.1001/jama.2013.168118
  64. Modesti PA (2015) Ethnicity: new challenge for cardiovascular prevention in Europe. *Hypertension* 66(3):464–465. doi:10.1161/HYPERTENSIONAHA.115.05777
  65. Gillies CL, Lambert PC, Abrams KR, Sutton AJ, Cooper NJ, Hsu RT, Davies MJ, Khunti K (2008) Different strategies for screening and prevention of type 2 diabetes in adults: cost effectiveness analysis. *BMJ* 336(7654):1180. doi:10.1136/bmj.39545.585289.25
  66. Admiraal WM, Vlaar EM, Nierkens V, Holleman F, Middelkoop BJ, Stronks K, van Valkengoed IG (2013) Intensive lifestyle intervention in general practice to prevent type 2 diabetes among 18–60-year-old South Asians: 1-year effects on the weight status and metabolic profile of participants in a randomized controlled trial. *PLoS One* 8(7):e68605. doi:10.1371/journal.pone.0068605
  67. Rose G (1981) Strategy of prevention: lessons from cardiovascular disease. *Br Med J (Clin Res Ed)* 282(6279):1847–1851
  68. Latham K, Wu B (2013) Chinese immigration into the EU: new trends, dynamics and implications. 1–65. [http://www.chathamhouse.org/sites/files/chathamhouse/public/Research/Asia/0313ecran\\_lathamwu.pdf](http://www.chathamhouse.org/sites/files/chathamhouse/public/Research/Asia/0313ecran_lathamwu.pdf)
  69. Zhou M, Astell-Burt T, Bi Y, Feng X, Jiang Y, Li Y, Page A, Wang L, Xu Y, Wang L, Zhao W, Ning G (2015) Geographical variation in diabetes prevalence and detection in china: multilevel spatial analysis of 98,058 adults. *Diabetes Care* 38(1):72–81. doi:10.2337/dc14-1100
  70. Modesti PA, Han Y, Jing Y, Xiaoling W, Mengyue Z, Zihua Y, Jia G, Perruolo E, Bini L, Camera M, Biggeri A, Rapi S, Hongsheng C, Li Z, Zengli W, Jianbin H, Xiaoyue Z, Zhao D (2014) Design and arrangement of the CHIP (CHinese In Prato) study. *Epidemiol Prev* 38(6):357–363
  71. Stuart-Shor EM, Berra KA, Kamau MW, Kumanyika SK (2012) Behavioral strategies for cardiovascular risk reduction in diverse and underserved racial/ethnic groups. *Circulation* 125(1):171–184. doi:10.1161/circulationaha.110.968495
  72. Grace C, Begum R, Subhani S, Kopelman P, Greenhalgh T (2008) Prevention of type 2 diabetes in British Bangladeshis: qualitative study of community, religious, and professional perspectives. *BMJ* 337:a1931. doi:10.1136/bmj.a1931
  73. Artinian NT, Fletcher GF, Mozaffarian D, Kris-Etherton P, Van Horn L, Lichtenstein AH, Kumanyika S, Kraus WE, Fleg JL, Redeker NS, Meininger JC, Banks J, Stuart-Shor EM, Fletcher BJ, Miller TD, Hughes S, Braun LT, Kopin LA, Berra K, Hayman LL, Ewing LJ, Ades PA, Durstine JL, Houston-Miller N, Burke LE, American Heart Association Prevention Committee of the Council on Cardiovascular N (2010) Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association. *Circulation* 122(4):406–441. doi:10.1161/CIR.0b013e3181e8edf1