


Effect-based activities on air pollution:

What is the state of the natural and anthropogenic Italian ecosystems?

Edited by Ilaria D'Elia, Alessandra De Marco, Giovanni Vialetto



Effect-based activities on air pollution: what is the state of the natural and anthropogenic Italian ecosystems?

Edited by Ilaria D'Elia, Alessandra De Marco, Giovanni Vialetto

Chapter 1

Chapter 2

Authors: Giovanni Vialetto, Alessandra De Marco, Tiziano Pignatelli, Antonio Ballarin-Denti

Chapter 3

Authors: Ilaria D'Elia, Luisa Ciancarella, Antonio Piersanti, Giovanni Vialetto

Chapter 4

Authors: Marco Ferretti, Giada Bertini, Filippo Bussotti, Laura Canini, Mario Cammarano, Roberto Canullo, Stefano Carnicelli, Guia Cecchini, Gianfranco Fabbio, Angela Farina, Daniele Giorgini, Matteo Feducci, Aldo Marchetto, Maurizio Marchi, Giorgio Matteucci, Khawla Zouglami

Chapter 5

Authors: Elisabetta Salvatori, Alessandra Campanella, Maria Chiesa, Lorenzo Cotrozzi, Angelo Finco, Lina Fusaro, Giacomo Gerosa, Giacomo Lorenzini, Riccardo Marzuoli, Cristina Nali, Romina Papini, Elisa Pellegrini, Mariagrazia Tonelli, Fausto Manes

Chapter 6

Authors: Maria Francesca Fornasier, Patrizia Bonanni, Marcello Vitale, Michele De Sanctis, Giuliano Fanelli, Fabio Attorre, Alessandra De Marco, Silvano Fares, Luca Salvati

Chapter 7

Authors: Michela Rogora, Aldo Marchetto, Rosario Mosello

Chapter 8

Authors: Pasquale Spezzano, Giovanni Vialetto

Chapter 9

Authors: Elena Paoletti, Elisabetta Salvatori, Fausto Manes,

Chapter 10

Authors: Antonio Piersanti, Pierluigi Altavista, Carla Ancona, Giovanna Berti, Ennio Cadum, Luisella Ciancarella, Ilaria D'Elia, Francesco Forastiere, Marina Mastrantonio, Francesca Pacchierotti, Gaia Righini, Raffaella Uccelli

Chapter 11

Authors: Ilaria D'Elia, Giovanni Vialetto, Luisella Ciancarella, Alessandra De Marco

2016 ENEA
National Agency for New Technologies, Energy and
Sustainable Economic Development

ISBN 978-88-8286-344-9

Copy editing: Giuliano Ghisu

Cover design: Flavio Miglietta

Printing: Laboratorio Tecnografico ENEA – Frascati

Photos on the cover and on page 81 taken by Eleonora Mambrini.

PREFACE

The Division Models and Technologies for Risks Reduction of the ENEA Sustainability Department was established in July 2015 with the specific aim of assemble all the resources, both human and instrumental, that could propose reflections, analyzes and assessments about the interactions among energy strategies, emissions of pollutants and greenhouse gases, air quality, climate change and resulting impacts on health and ecosystems.

The use of numerical models, at different spatial scales, to describe the physical and chemical processes affecting the atmosphere and the ocean is at the hearth of the history and activities of the research groups involved. Nevertheless, great effort and commitment was also dedicated to the development of integrated models to explore links and feedback between the different aspects of the same system. These unique platforms called IAM (Integrated Assessment Modelling) help and support policy-makers to develop and promote sustainable policies.

With this vision we have been developing the national model MINNI (Integrated National Model in support to the International Negotiation on Air Pollution) since 2003. It includes the Integrated Assessment Model GAINS-Italia, which helped our Country: to negotiate revisions of European and International Directives and Protocols with technical data and national scenario simulations; to support technically the actions needed to deal with European infringement procedures; to help the Regional Authorities to fully understand and use the measurements until they have become totally autonomous; to support the early season of the Air Quality Regional Plans together with the assessment of the measures identified in them.

A parallel and equally far-sighted vision has been offered by our colleagues of the actual Division who have been dealing for years with the “pollution effects”. They hold significant roles within the International Convention on Long Range Transboundary Air Pollution (CLRTAP) of the United Nations Economic Commission for Europe (UNECE) both as national representatives in the Executive Body and National Focal Point or Experts within technical Task Forces.

To date a lot of work has been done but we need more and more of each other’s expertise to move forward in the complexity. We cannot have disciplinary “exclusivity” because atmospheric pollution and its effects are no longer (and have never been) a single discipline theme. At the same time we must avoid that integration could generate only simplification.

The main intent of this publication and of this day is to take stock of the situation in Italy, sharing what has been done both on the atmospheric pollution and effects side as well as the many and important experiences of comparison and integration. We would like to start from here in order to deal with what Europe strongly advocates: that the definition of policies to face air pollution considers as a priority the reduction of its effects.

Rome, March 2017.

Gabriele Zanini

(Head of Division Models and Technologies for the Risks Reduction)

CONTENTS

PREFACE	3
CHAPTER 1 – Introduction.....	7
CHAPTER 2 - Effect Oriented Activities in the LRTAP Convention	10
CHAPTER 3 - Sources and Emissions of air pollutants.....	16
CHAPTER 4 - Actual and potential impact of air pollution on Italian forests: results from the long-term national forest monitoring networks under the ICP Forests	22
4.1 Introduction.....	22
4.2 Pressures	23
4.2.1 Status.....	23
4.2.2 Trend.....	25
4.3 Impacts.....	26
4.3.1 Forest health.....	26
4.3.2. Forest growth	27
4.3.3 Forest biodiversity.....	29
4.3.4. Forest nutrition.....	32
4.4. Risk assessment.....	34
4.4.1 Risk for forest health.....	34
4.4.2. Risk for forest growth	35
4.4.3. Risk for forest biodiversity	35
4.4.4. Risk for forest nutrition.....	36
4.5 Conclusions.....	37
CHAPTER 5 - Effects of air pollution on crops and semi-natural vegetation.....	40
5.1 Introduction.....	40
5.2 Pressures	40
5.2.1 Ozone	40
5.2.2 Nitrogen	42
5.3 Impacts.....	42
5.3.1 Ozone	42
5.3.2 Nitrogen	46
5.4 Risk assessment.....	48
5.4.1 Ozone	48
5.4.2 Nitrogen	52
CHAPTER 6 - Biodiversity as an important indicator of soil acidity and eutrophication: the role of the modelling in preserving it	54
6.1 Introduction.....	54
6.2 Measurements of climate variable for modelling simulation.....	55
6.3 Impacts.....	59

6.3.1 Biodiversity analysis with dynamic models.....	60
6.3.2 Biodiversity analysis with statistical models.	62
6.4 Risk assessment.....	66
6.5 Conclusions.....	67
CHAPTER 7 - The contribution of Italy to the ICP WATERS Programme	70
7.1 Introduction.....	70
7.2 Atmospheric pollution pressures on surface waters.....	72
7.3 Main impacts of atmospheric pollution on rivers and lakes	75
7.4 Deposition scenarios and risk assessment.....	78
7.5 Conclusive remarks.....	80
CHAPTER 8 - Are technical materials and cultural heritage exposed to air pollution risk? The contribution of Italy to ICP Materials	82
8.1 INTRODUCTION	82
8.2 Effects on materials.....	83
8.3 Effects on historic and cultural monuments and buildings.	85
8.4 Concluding remarks	87
8.5 Acknowledgments.....	87
CHAPTER 9 - Vegetation and urban air quality: recent findings	90
CHAPTER 10 – Effects of Air Pollution on Health	96
10.1 Introduction.....	96
10.2 Results.....	97
CHAPTER 11 - What remains to be done to reduce air pollution?	104
REFERENCES.....	109

CHAPTER 1 – INTRODUCTION

In the last twenty years, a large effort has been carried out in the Italian community working on the impacts of atmospheric pollution on different ecosystems, both natural or anthropogenic. For long time the work was done without the necessary interconnection between the different sub-networks belonging to the Working Group on Effects. In order to bridge the gaps between scientists and policy makers, the communication between the work done in the different sub-groups has been recently improved and the realization of this monographic text is one of its first tangible result.

The intent of the text is to collate information from the important fields of measurements and modeling to obtain dose-response relationships able to estimate the impacts of air pollutant on the different ecosystems.

Italy is the richest country of the world in terms of biodiversity, agricultural products and number of cultural heritage sites exposed to negative impacts of air pollution and climate change. The Italian country is formed by different peculiar ecosystems, from coastal area to forests, to cities included in the UNESCO list for cultural heritage. It is thus very important to protect our environmental and cultural heritage from environmental pressure and from anthropogenic footprint. A wide choice of measures to control atmospheric pollution and decrease the negative impacts on health and ecosystems is available and possible alternative methodologies, like for example the change in the human behaviours, could have a high importance in a future sustainable society.

More in general, the analysis of the Italian monitoring network shows that the health of the main natural ecosystems has improved, even if there is still some to be done at local level and for specific pollutants. For instance, due to the peculiar conditions of sun exposure and climate, the situation for ozone is still very critical in a Mediterranean environment.

In some cases, the efficacy of the strategies of controlling air pollution can be impaired by climate change, while in some others, strategies to reduce air pollution can be considered as win-win policies, because are beneficial from several points of view.


The revision of the National Emission Ceilings Directive (NECD) has established emission reduction targets focusing on the reduction of the impacts of air pollution on ecosystems and human health. This important change captured more attention to the monitoring activities of the impacts of air pollution on natural and anthropogenic ecosystems. This is an emerging issue to promote and to give the due importance to the activities of the Working Group on Effects. In this context, the Italian work has been developed to provide policymakers with a clear picture of what has already been achieved and what can still be done to protect our environment.

The key findings emerged from this work and from the different communities and networks are summarized in the following text.

- **The implementation of the Convention on Long-range Transboundary Air Pollution and its protocols has allowed reaching a significant success in reducing emissions, especially for sulphur, concentrations and deposition trends. Despite this success, exceedances of critical loads and level, of the air quality limit values and high population exposure still exist.**

- **Forest condition has improved in Italy since the 1990s, with limited evidence of an impact due to tropospheric ozone. On the other side, there was a distinct effect of nitrogen deposition on forest nutrition, growth and carbon sequestration.**
- **Biodiversity indices and habitat suitability index in forest sites are recovering consequently to decreasing in pollutant deposition.**
- **Increasing background O₃ levels in Italy still affect health, productivity and quality of crops and (semi-) natural vegetation. The risk assessment should be based on the effective O₃ dose absorbed through stomata, taking into account the different species/cultivar sensitivity, and the concurrent effect of other stress factors, such as the reactive nitrogen deposition and climatic conditions typical of the Mediterranean area.**
- **The interactions between air quality and urban vegetation are potentially of great interest, although understanding is still imperfect. In Italy, a few studies have been carried out to estimate potential uptake of pollutant by trees, in particular in two cities: Rome and Florence, in order to estimate and map this Regulating Ecosystem Service.**
- **Surface water showed a widespread response to decreasing deposition of acidity, sulphate and nitrogen compounds, but the recovery was somewhat delayed, due to the interacting effect of several factors, such as nitrogen saturation of soils in the catchments and climate change.**
- **Nitrogen deposition will continue to have a prominent role in the acidification processes and in the nitrogen status of surface water. The recovery patterns in the next future will be more and more influenced by climatic factors.**
- **Atmospheric pollution is a key factor in the deterioration of sensitive materials and materials used in historical and modern cultural heritage are the most vulnerable to air pollution. Even if the decrease of concentrations of air pollutants has led to the decrease of deterioration rates (mainly due to the decrease of SO₂), current corrosion rates and soiling are still unacceptably high. The cost associated with the damages is enormous but difficult to estimate.**
- **Health impact assessment of air pollution at national scale in Italy has been carried out in several projects, based on measured data and modelling techniques. Results agreed on showing that PM₁₀ and NO₂ induce several thousands of premature deaths and hospital admissions per year, due to cardiovascular and respiratory diseases. Confirming conclusions of previous European-scale assessments, air pollution is proved to be a major risk for human health in Italy.**
- **Many solutions are available and only an integrated approach takes into account the co-benefits of linking air pollution and climate change. The integration of all the different tools, from measures to models, and the coordination among science sectors and different research teams could help in identifying effective environmental policy.**

Effect Oriented Activities in the LRTAP Convention



Chapter coordinator: Giovanni Vialetto

Contributors: Giovanni Vialetto¹, Alessandra De Marco¹, Tiziano Pignatelli², Antonio Ballarin-Denti³

¹ENEA, Laboratory of Atmospheric Pollution, ²ENEA, Strategic Technical Support,

³University of Brescia "La Cattolica"

CHAPTER 2 - EFFECT ORIENTED ACTIVITIES IN THE LRTAP CONVENTION

The Convention on Long-range Transboundary Air Pollution (CLRTAP – see web site <http://www.unece.org/env/lrtap/welcome.html>) has achieved successful results in reducing the emissions of a wide range of atmospheric pollutants and consequently in decreasing acidification and eutrophication processes, in soothing ozone peak levels of ozone and photochemical smog, and in ensuring improvements in nitrogen atmospheric levels and deposition rates. The Convention has also proved to be flexible and dynamic in responding to new challenges and problems raised by the transboundary air pollution. The Convention has been also a powerful driver to promote a sound science-for-policy approach to relevant environmental problems not just transferring valuable knowledge and datasets from the scientific community to the decision-makers and involved stakeholders, but also supporting and fostering the whole policy-making process.

However, despite the progress achieved under the Convention, the air quality in the UNECE region is still matter of concern, given its persistent impacts on human health and ecosystems while new environmental problems are emerging (ECE, 2010). A solid and widely recognized scientific basis has been a key success factor for the Convention's outcomes in air pollution abatement. This was enabled by the development of a common and shared knowledge-providers network based on a large system of scientific facilities s managed by authoritative research organizations and aimed at monitoring and modelling a wide set of environmental parameters and ensuring an intense and interdisciplinary exchange of scientists across all the UNECE countries. In addition, the Convention has provided an open-innovation platform for scientists and policymakers to exchange information which has led to a growing and fruitful international cooperation, creating mutual trust, higher policy-harmonization, and the dissemination of good practices in environmental monitoring and assessment. In order to organize its work, the Convention has created an appropriate functional structure, based on expert groups, task forces and research centres. The updated structure of the Convention is showed in Fig. 1.1.

Acidification of soil and water was the main environmental problem in the seventies, caused by the high levels of sulphur and nitrogen atmospheric emissions. Therefore, the reduction of these pollutants was considered as the most urgent need. In the following years, new harmful atmospheric pollutants drew the attention of both the scientific and political community and were thus included into the Convention's protocols. At the same time, new pollutants were being considered in the upcoming policies, such as Non-Methane Volatile Organic Compounds (NMVOC), Heavy Metals (HMs), Persistent Organic Pollutants (POPs), Ozone (O₃), Particulate Matter (PM) and, more recently Black Carbon (BC).

While the first protocols developed under the Convention focused on technologies capable to reduce emissions, protocols negotiated since the 1990s adopted an effect-oriented approach, aiming at the best cost-effective way to reach the set reduction targets. At the same, it was also recognized that various air pollutants may interact in the atmosphere, thus leading to combined or even synergistic impacts and often depending on the same emission sources. This made a substance-by-substance approach less efficient and was the reason why the so-called multi-pollutant-multi-effect approach was developed. The first protocol based on this new approach was the 1999 Gothenburg Protocol to Abate Acidification, Eutrophication and Ground-level Ozone (EB, 1999).




Figure 1.1 – Updated structure of the Convention on Long Range Transboundary Air Pollution.

With the revision of the Gothenburg Protocol, approved in 2012 (ECE, 2012a,b), a further step forward was made, and the particulate matter (PM_{2.5} fraction) was also introduced into the Protocol. The focus of the Protocols is targeted to the health protection, which becomes the basis for the optimized emission reduction allocation and commitments for each Party and in doing so, the Convention has achieved considerable success in solving environmental and health problems. These successes are largely due to the scientific grounds of the Convention and the unique way in which science affects any further policy development (look for example Maas et al., 2016). Another key success factor is represented by the Convention’s wide geographical coverage. In fact, the Convention embraces most of the northern hemisphere from the West Coast of North America to the Pacific Coast of the Russian Federation (see fig. 1.2).



Figure 1.2 – Map showing the Convention on Long-Range Transboundary Air Pollution signatories.

Italy has a long tradition of activities in the UN-ECE Convention on Long Range Transboundary Air Pollution (UN-ECE LRTAP), since its signature on 14 November 1979, in Geneva and the ratification by the Parliament, occurred on 15 July 1982. The Convention was adopted on 13 November 1979 and left open for signature until 16 November 1979 at the United Nations Office in Geneva. Nowadays, the Convention comprises 32 signatories and 51 Parties.

Table 1.1 – Status of signature and ratification by Italy of the Protocols of the LRTAP Convention.

Treaty	Signature	Ratification, Acceptance, Approval, Accession, Succession
LRTAP Convention	14-Nov-79	15 Jul 1982
The 1984 Geneva Protocol on Long-term Financing of the Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP)	28-Sep-84	12 Jan 1989
The 1985 Helsinki Protocol on the Reduction of Sulphur Emissions or their Transboundary Fluxes by at least 30 per cent	09-Jul-85	5 Feb 1990
The 1988 Sofia Protocol concerning the Control of Emissions of Nitrogen Oxides or their Transboundary Fluxes	01-Nov-88	19 May 1992
The 1991 Geneva Protocol concerning the Control of Emissions of Volatile Organic Compounds or their Transboundary Fluxes	19-Nov-91	30 Jun 1995
The 1994 Oslo Protocol on Further Reduction of Sulphur Emissions	14-Jun-94	14 Sep 1998
The 1998 Aarhus Protocol on Heavy Metals	24-Jun-98	Not yet ratified
The 1998 Aarhus Protocol on Persistent Organic Pollutants (POPs)	24-Jun-98	20 Jun 2006
1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone to the Convention on Long-range Transboundary Air Pollution (Gothenburg Protocol)	01-Dec-99	Not yet ratified

In the following decades 8 Protocols were developed, adopted, signed and ratified by a number of Parties. The table 1.1 summarizes the status of signature and ratification by Italy.

The Protocols on Heavy Metals, on POPs and the Protocol to Abate Acidification, Eutrophication and Ground-level Ozone, have been amended and adopted, respectively on 13 December 2012 (Heavy Metals), on 18 December 2009 (POPs), on 4 May 2012 (Gothenburg Protocol). The ratification process of these last three Protocols is still in progress for Italy, although the European Union has accepted both the amended Protocol on Heavy Metals and the amended Protocol on POPs. The last three Protocols, in the amended version mentioned above, are not yet entered into force because the minimum number of ratification by the Parties has not yet been achieved.

The Italian experts participate actively in the works of many technical bodies of the LRTAP Convention. Notably, within the Working Group on Effects, the International Cooperative Programme on Effects of Air Pollution on Materials, including Historic and Cultural Monuments sees the Co-Chair of Italy, since 2005, currently in the person of Pasquale Spezzano. As well, the Task Force on Techno-Economic Issues has the Italian Co-Chair Tiziano Pignatelli, involved since 2006. Moreover, several scientific studies are being conducted under the leadership and /or responsibility of the Italian experts.

Moreover, during the very first discussions on the Convention, it was immediately recognized that a good understanding of the harmful effects of air pollution was a prerequisite for reaching agreement on effective pollution control. To develop the necessary international cooperation in the research on and the monitoring of pollutant effects, the Working Group on Effects (WGE) was established under the Convention in 1980.

The Working Group on Effects provides the Convention’s Executive Body with sound and shared scientific information on the degree and geographic extension of the impact of major air pollutants on the human health and some relevant natural and human-heritage targets like water, forests, vegetation and materials.

Such information is the result of monitoring activities on the effects caused by atmospheric pollution across Europe and North America, and are based upon scientific research on dose-response relationships, critical loads and damage evaluation.

WGE is composed by six International Cooperative Programmes (ICPs) and the Task-force on Health. Each ICP includes a Task-Force and a Coordination Centre with the aim of collecting data from monitoring networks. For each ICP, a National Focal Centre (NFC) is established, with the aim to coordinate the national activities, participate in the works of the task-forces, and submit data to the Coordination Centres.

Italy, as Party of the Convention, is therefore committed to appoint the NFCs, to establish and control the monitoring networks, to submit the required data and to participate in the activities of the task-forces. The head of the Italian Delegation to WGE is Alessandra De Marco (ENEA), and the other delegation members are Antonio Ballarin-Denti (Catholic University of Brescia) and Sergio Cinnirella (CNR). All the national Focal Points are reported in table 1.2.

Table 1.2 – All the National Focal Points to the different International Cooperative Programme under the LRTAP Convention.

WGE Task Force	Description	National Focal Point	Institution
ICP Forests	ICP on assessment and monitoring of air pollution effects on forests	Angela Farina	State Forestry Authority
		Laura Canini	State Forestry Authority
ICP Integrated Monitoring	ICP on integrated monitoring of air pollution effects on ecosystems	Angela Farina	State Forestry Authority
		Laura Canini	State Forestry Authority
ICP Modelling & Mapping	ICP on modelling and mapping of critical levels and loads and air pollution effects, risks and trends	Patrizia Bonanni	Isprambiente
		Francesca Fornasier	Isprambiente
		Alessandra De Marco	ENEA
		Marcello Vitale	“Sapienza” University of Rome
ICP Materials	ICP on effects of air pollution on materials, including historic and cultural monuments	Pasquale Spezzano (co-chair ICP and NFP)	ENEA
ICP Vegetation	ICP on effects of air pollution on natural vegetation and crops	Fausto Manes	“Sapienza” University of Rome
		Elisabetta Salvatori	“Sapienza” University of Rome
		Alessandra De Marco	ENEA
ICP Waters	ICP on assessment and monitoring of the effects of air pollution on rivers and lakes	Michela Rogora	CNR

Sources and Emissions of air pollutants



Chapter coordinator: Ilaria D'Elia

Contributors: Ilaria D'Elia¹, Luisa Ciancarella¹, Antonio Piersanti¹, Giovanni Vialetto¹

¹ENEA, Laboratory of Atmospheric Pollution


CHAPTER 3 - SOURCES AND EMISSIONS OF AIR POLLUTANTS

Air pollution is a very important environmental and social issue that poses multiple challenges in terms of management and mitigation (EEA, 2015). Air pollution represents the single largest environmental health risk in Europe and particulate matter has become a major concern for public health whose carcinogenicity has been recognized by the International Agency for Research on Cancer (IARC, 2015). Many measures and policies have been adopted in the past decades at European, national, regional and even local level but a large portion of European population and ecosystems is still exposed to concentrations that exceed the European Union (EC, 2008) and the World Health Organization (WHO, 2005) air quality standards. In this view, the Gothenburg protocol was amended in 2012 and, in 2013, the European Commission adopted the Clean Air Policy Package (COM, 2013) with the aim to further reduce the impacts of harmful emissions on human health and the environment.

Air pollutants are emitted from anthropogenic and natural sources; they may be either emitted directly or formed in the atmosphere and they have a number of impacts on health, ecosystems, the built environment and the climate.

Figure 3.1 shows the trend of the Italian emissions of sulphur oxides (SO_x), nitrogen oxides (NO_x), ammonia (NH₃), primary PM with a diameter of 10 µm or less (PM₁₀) and PM with a diameter of 2.5 µm or less (PM_{2.5}), non-methane volatile organic compound (NMVOC), carbon monoxide (CO) and methane (CH₄) between 1990 and 2014. Similarly, figure 3.1 shows the trend in emissions of the toxic metals arsenic (As), cadmium (Cd), nickel (Ni), lead (Pb), mercury (Hg), Polycyclic Aromatic Hydrocarbon (PAH) and dioxins (Diox). In the period 1990-2014, in Italy almost all the primary and precursor emissions have decreased (IIR, 2016). Reductions are especially relevant for the main pollutants, SO_x (-93%); NO_x (-61%); CO (-69%); NMVOC (-57%) and Pb (-94%), while NH₃ shows the smallest reduction (-17%).

Effective action to reduce the impacts of air pollution requires a good understanding of its causes, how pollutants are transported and transformed in the atmosphere, and how they impact on humans, ecosystems and the climate.




Source: Based on IIR, 2016.

Figure 3.1 – Trend in Italian emissions of SO₂, NO_x, NH₃, PM₁₀, PM_{2.5}, NMVOC, CO and CH₄ (top) and of As, Cd, Cr, Hg, Ni, Pb, PAH, Diox (bottom), 1990-2014 (% of 1990 levels).

The major drivers for the trend are reductions in the industrial and road transport sectors, due to the implementation of various European Directives which introduced new technologies, plant emission limits, the limitation of sulphur content in liquid fuels and the shift to cleaner fuels (fig. 3.2). Emissions have also decreased for the improvement of energy efficiency as well as the promotion of renewable energy.

In the following plots, an analysis of the Italian emission trend by sector in the period 1990-2014 is shown. The main source sectors contributing to emissions of air pollutants in Italy are transport, energy, industry, the commercial, institutional and households sector, agriculture and waste.

The transport sector has considerably reduced its emissions of air pollutants in Italy since 1990, as figures 3.2 and 3.3 show, except for PAH emissions, which have increased by 26% from 1990 to 2014. The highest emission reductions from transport between 1990 and 2014 were registered for SO_x (90%) and for Pb (100%).




Source: Based on IIR, 2016.

Figure 3.2 – Trend in Italian emissions from main sources of SO_x, NO_x, NH₃, PM₁₀, PM_{2.5}, NMVOC, CO and CH₄, 1990-2014 (% of 1990 levels).

With the exception of SO_x emissions, the commercial, institutional and household sector has significantly increased its emissions, whose trend follows the fuelwood consumption variation. This emission increase is due to the use of household wood and other biomass combustion for heating, owing to government incentives/subsidies, rising costs of other energy sources, or an increased public perception that it is a ‘green’ option (EEA, 2015). Biomass is being promoted as a renewable fuel that can assist with climate change mitigation and contribute to energy security.


Energy production and industry considerably reduced their air pollutant emissions between 1990 and 2014, with the exception of As emissions from the industrial sector.

Agriculture is the main sector in which emissions of air pollutants have least decreased while waste sector shows a great emission reduction in Diox emissions.



Source: Based on IIR, 2016.

Figure 3.3 – Trend in Italian emissions from main sources of As, Cd, Ni, Pb, Hg, PAH and Diox, 1990-2014 (% of 1990 levels).



Source: Based on IIR, 2016.

Figure 3.4 – Main emitting sectors for the main pollutants in the year 2014.

The main emitting sectors in the year 2014 according to the last national emission inventory (IIR, 2016), are shown in fig. 3.4.

Industry is the largest source of SO_x, As, Cd, Ni, Pb, Hg and Diox, whose emissions respectively represent the 49%, 90%, 53%, 41%, 65%, 60% and 49% of total national emissions.

The transport sector is the biggest contributor to NO_x emissions, accounting for 69% of total Italian emissions in 2014. However, NO_x emissions from road transport have not been reduced as much as expected with the introduction of the Euro standards, since emissions in real-life driving conditions for diesel vehicles are often higher than those measured during the approval test.

The commercial, institutional and household sector is by far the most important sector of primary PM₁₀, PM_{2.5} and PAH emissions contributing respectively to 57%, 65% and 71% of total PM₁₀, PM_{2.5} and PAH total emissions. Moreover, it is the second most significant emitter of Diox, CO, Cd, Pb and Hg. This sector surely represents a key sector in reducing air pollutant emissions.

The agricultural sector is the greatest emitter of NH₃ and was responsible for 96% of total NH₃ emissions in Italy in 2014. NH₃ emissions have decreased by only 17% from 1990 to 2014 because European policies have cut PM precursor gas emissions, with the exception of NH₃ from agriculture (EEA, 2015).

Emissions have decreased in many sectors except for the commercial, institutional and households sector, whose emissions showed an increase for almost all pollutants and agriculture which showed the lowest reduction. As a consequence, the commercial, institutional and households and agricultural sectors still have a high potentiality to reduce emissions and improve air quality.

In order to reduce the harmful effects of atmospheric pollutants, the National Emission Ceilings Directive (NECD) (EC, 2001) sets EU Member States individual air pollutant emission limits, or ‘ceilings’, restricting emissions for four important air pollutants: nitrogen oxides (NO_x), non-methane volatile organic compounds (NMVOC), sulphur dioxide (SO₂) and ammonia (NH₃).

Table 3.1 – Italian progress in meeting ceilings set out in NECD Annexes (EC, 2001).

Pollutant	NEC Ceilings 2010 (kt)	Emission inventory (kt)				
		2010	2011	2012	2013	2014
SO ₂	475	217	195	176	145	131
NO _x	990	978	950	867	816	790
NM VOC	1159	1046	954	942	909	849
NH ₃	419	389	402	415	402	393

As of 2010, all Member States are required to meet their emission ceilings. Italy respects its emissions ceilings for all the four pollutants during the period 2010 to 2014 as shown in table 3.1.

Many measures and policies have been adopted in the past decades at European, national, regional and even local level but a large portion of European population and ecosystems is still exposed to concentrations that exceed the World Health Organization (WHO, 2005) and the European Union air quality standards (EC, 2008). In this view, in 2013, the European Commission adopted the Clean Air Policy Package (COM, 2013), with the aim to further reduce the impacts of harmful emissions on human health and the environment, where the proposal for a revised National Emission Ceilings Directive (NECD) plays a major role.

Actual and potential impact of air pollution on Italian forests: results from the long-term national forest monitoring networks under the ICP Forests



National Focal Point: Angela Farina and Laura Canini (State Forestry Authority)

Chapter coordinator: Marco Ferretti

Contributors: Marco Ferretti^{1,2}, Giada Bertini³, Filippo Bussotti⁴, Laura Canini⁵, Mario Cammarano⁶, Roberto Canullo⁷, Stefano Carnicelli⁸, Guia Cecchini⁸, Gianfranco Fabbio³, Angela Farina⁵, Daniele Giorgini⁷, Matteo Feducci⁴, Aldo Marchetto⁹, Maurizio Marchi³, Giorgio Matteucci¹⁰, Khawla Zougami⁷

¹Swiss Federal Research Institute WSL; ²formerly at TerraData environmetrics; ³CREA-Centro di ricerca per la selvicoltura; ⁴Università di Firenze - Dipartimento di Biotecnologie Agrarie, Sezione di Botanica Ambientale ed Applicata; ⁵State Forestry Authority; ⁶CNR-Istituto di Biologia Agroambientale e Forestale; ⁷Università di Camerino - School of Biosciences and Veterinary Medicine, Plant Diversity and Ecosystems Management unit;

⁸Università di Firenze- Dipartimento di Scienze della Terra; ⁹CNR-Istituto per lo Studio degli Ecosistemi;

¹⁰CNR- Istituto per i Sistemi Agricoli e Forestali del Mediterraneo.

CHAPTER 4 - ACTUAL AND POTENTIAL IMPACT OF AIR POLLUTION ON ITALIAN FORESTS: RESULTS FROM THE LONG-TERM NATIONAL FOREST MONITORING NETWORKS UNDER THE ICP FORESTS

This Chapter was prepared for the most part on the basis of evaluations carried out within the LIFE project “Sustainable Monitoring And Reporting To Inform Forest and Environmental Awareness and Protection – SMART4Action” [LIFE13 ENV/IT/000813].

4.1 Introduction

Forests in Italy cover ca. 8.76×10^6 , stock 1,242 Mt carbon and offer a series of invaluable products and services to the entire society (Gasparini and Tabacchi, 2011; Gasparini et al., 2013). Preserving forest integrity is therefore essential. Although effects are in general less obvious and frequent in comparison with other biotic and abiotic stressors, air pollution may have an impact on several compartments of forest ecosystem, and may endanger forest health, productivity and diversity (Bobbink et al., 2010; Karnosky et al., 2003). In this chapter, we will present status, trends of pressure, impacts and risks placed to forests by the most important air pollution issues in Italy as emerged from data collected by national forest monitoring networks since the mid of the 1990s. To do this, we will concentrate on data originated from harmonized measurements carried out at the national Level I and Level II networks that are parts of the UNECE ICP Forests (Figure 4.1).




Figure 4.1 – Level I (small black dots) and Level II (large colored symbols) networks in Italy. Colors refer to main tree species in Level II plot (source: LIFE SMART4Action). Sample plots (Phase 3+) of the National Forest Inventory are also reported.

4.2 Pressures

Deposition of oxydized (NO_x) and reduced nitrogen (NH_y) and tropospheric ozone (O_3) are the most important air pollution forest-related issues in Italy and elsewhere. Both N deposition and ozone levels have been measured at the sites of the Italian ICP Forests Level II network since the 1990s. Atmospheric deposition is collected in the open field and under forest canopy (throughfall) by different set of funnels (Figure 4.2A). Stemflow collectors (Figure 4.2B) are used in beech plots to collect precipitation running along stems. Ozone concentration is measured using passive samplers (Figure 4.2C) since 1996. A further set of ozone measurements have been carried out over the period 2007 - 2011 on 15 Level I plots in Trentino (Northern Italy) (Gottardini et al., 2010).




Figure 4.2 – Collectors used for open field and throughfall (A), stemflow (B), and ozone passive samplers (C).

4.2.1 Status

Sampling and analysis of atmospheric deposition and of ozone was carried out on a variable number of plots, depending on funding. In particular, in 2009-2011 this activity was carried out and validated on 21 plots. Throughfall deposition represents a good estimate of the deposition reaching the forest soil. The geographic distribution of average throughfall deposition for 2009-2011 of some selected ions and of the pH values obtained from the volume weighted average of hydrogen ion deposition is shown in Figure 4.3.

All the plots show pH values higher than > 5 , mostly around the equilibrium value of 5.6. In the case of sulphate deposition (corrected for the amount deriving by marine aerosol - (CLRTAP, 2004), a geographic pattern is evident, with lower values on the Alps, and higher values in the Southernmost plots. Due to the episodic deposition of Saharan dust, this is leading to episodes with high sulphate and calcium concentration. In the case of nitrate, ammonium and total N (which includes N in organic compounds), high values were measured in the Po plain (where most of industries and farming are located) and in the surrounding hills, while lower values were detected in high elevation Alpine sites.




Figure 4.3 – Annual deposition of selected ions and average ozone concentration during the growing season at Level II sites, 2009-2011. Site codes in Figure 4.1. (Source: LIFE SMART4Action).

As for ozone, average concentration during the growing season was in general > 40 ppb in all sites, with slightly higher values in the Southern regions and in the Adriatic part of the peninsula (Figure 4.3).




Figure 4.4 – Mean 2000-2003 AOT40 at 26 Italian Level II sites (dots) and 41 background conventional monitoring stations over the period 2000-2003. After Ferretti et al., 2007.

Table 4.1. Ozone flux estimates available for Level II sites in Italy.

Site	Year	Reporting unit	Flux estimate	Reference
CAL1	2000-2002	seasonal AFst, (mmol O ₃ m ⁻²)	8-10	Schaub et al., 2007
TOS1	2005	seasonal AFst1.6, (mmol O ₃ m ⁻²)	53	Bussotti and Ferretti, 2007
TOS2	2005	seasonal AFst1.6, (mmol O ₃ m ⁻²)	49	Bussotti and Ferretti, 2007
LAZ2	2005	seasonal AFst1.6, (mmol O ₃ m ⁻²)	47	Bussotti and Ferretti, 2007
TRE1	1999-2011	seasonal AFst1.6, (mmol O ₃ m ⁻²)	22-35	Gottardini et al., 2012

Accumulated ozone above threshold 40 ppb (AOT40) was estimated to exceed up to 10 times the Critical Level set to protect vegetation (Gerosa et al., 2003, 2007; Ferretti et al., 2007) (Figure 4.4). In terms of ozone flux, estimates were obtained for some Level II sites at individual years (Table 4.1) (Bussotti and Ferretti, 2007).

4.2.2 Trend

Figure 4.5 shows the temporal trend of sulphate, nitrate, and ammonium deposition and ozone concentration in nine forest sites with complete 1997-2013 dataserries. A marked decrease in sulphate deposition is evident in all sites at the end of the 1990s, while the reduction is less evident in the more recent period. Sulphate reduction resulted statistically significant (Seasonal Kendall test on monthly data, $p < 0.01$) in eight out of nine sites. In the case of nitrate and ammonium, a slight decrease was evident only in the last years and it was significant ($p < 0.05$) only in four and one site, respectively.




Figure 4.5 – Temporal trend in deposition of selected ions in nine Level II forest plots in Italy. (Source: LIFE SMART4Action).

Ozone concentration increased between 1997 and 2005, and decreased between 2005 and 2011 (Figure 4.5). The decreasing pattern is confirmed over the most recent years. As for ozone flux trend, information is available only for one Level II plot, namely TRE1 (Gottardini et al., 2012). At this site, ozone flux in terms of accumulated flux above 1.6 (AFst1.6) follows the same time pattern as AOT40, augmenting from 25 to 35 mmol m² PLA between 1999 and 2005, and decreasing to ca. 27 mmol m² PLA between 2006 and 2009 (Gottardini et al., 2012).

4.3 Impacts

N deposition may affect soil chemistry, nutrient balance, alter species-host interactions, biomass allocation, sensitivity to climate and to ozone. Ozone may reduce photosynthesis, cause premature senescence and direct injury to foliage. All together, this may have direct and indirect impact on forest health, growth, plant species diversity, and nutrition.

4.3.1 Forest health

In Italy, forests health has been measured under quality assured conditions since 1997. The most reliable variable adopted was (and is) defoliation, assessed both on Level I (ca. 250 plots) and Level II (31 plots) networks (see Figure 4.1). Direct injury to foliage due to ozone has been also assessed on some sites.

On Level I, mean defoliation for the 1997-2014 period was higher in broadleaves (39.7% of trees with defoliation higher than 25%) than in conifers (24.1%). Defoliation shows lower level in the South (30.7%) with respect to the North (35.5%) and Central (36.0%) Italy. Broadleaves were more defoliated (43.6%) in Northern Italy and less defoliated (35.8%) in Southern Italy. In conifers, defoliation was higher in Central Italy (36.9%) than in Southern Italy (12.3%).




Figure 4.6 – A) Percentage of trees with defoliation higher than 25% in the Level I plots, year -2014. B) 1997-2014 trend of the percentage of trees with defoliation >25%. Plots with at least 10 years of observations were analyzed with Mann-Kendall test. Significant increases (positive, red) and decreases (negative, green) of defoliation are indicated. (Source: LIFE SMART4Action).

The distribution of the defoliated trees (>25%) at each Level I plot (Figure 4.6A) reveals higher frequency in the North-Western regions of Italy (Piedmont, Lombardy, Liguria and partly the Northern areas of Tuscany). The analysis of the 1997-2014 trends (Figure 4.6B) evidenced that plots with significant increase of defoliated trees were concentrated especially in the Northern regions and in the Alpine area. All in all, however, there is a significant ($P < 0.01$) decreasing trend for defoliation, mostly driven by broadleaves.

Level II results confirm substantially the pattern described for the level I. Defoliation is increasing in Northern Italy (with the exception of VAL1) and decreasing in Central and Southern Italy (with the exception of LAZ1).

Foliar symptoms attributable to ozone injury were assessed at 12 Level II sites over the period 2003-2009 (Bussotti and Ferretti, 2007, 2009; Gottardini et al., 2012) and on a subset of Level I plots in Trentino in 2008-2009 (Gottardini et al., 2012). While no symptom was reported for Level I plots in Trentino, 46 *taxa* were found symptomatic at the investigated Level II sites. Off-plot investigation carried out nearby Level I plots in Trentino revealed symptoms on additional species, noticeably on *Viburnum lantana* (Gottardini et al., 2012).

4.3.2. Forest growth

Forest growth has been monitored since 1997 on Level II plots, and 17 plots have full coverage of the monitoring period 1997-2015 (i.e. five subsequent inventories). Current volume increment (IcV , the difference in volume between two subsequent years) and mean volume increment (ImV , the ratio between the volume and tree's age) were calculated on the basis by the volume functions provided

by the Italian National Forest Inventory (Tabacchi et al., 2011). Concerning absolute values and expected trends, data are summarized in Table 4.2.

Table 4.2 – Estimated volume and increments for the 17 CONECOFOR plots (hf=high forests; sc=stored coppices; tc=transitory crops). Standard deviation is reported in italic. (Source: LIFE SMART4Action).

Plot	Age	Type	Volume (m ³ ha ⁻¹)					Current volume increment (m ³ ha ⁻¹ yr ⁻¹)				Mean volume increment (m ³ ha ⁻¹ yr ⁻¹)				
			1997	2000	2005	2010	2015	2000	2005	2010	2015	1997	2000	2005	2010	2015
ABR1	130	hf	430.3	476.9	524.9	596.0	629.6	15.5	9.6	14.2	6.7	3.8	4.1	4.4	4.8	4.8
CAL1	130	hf	539.2	596.5	682.7	743.9	795.4	19.1	17.2	12.2	10.3	4.8	5.2	5.7	6.0	6.1
CAM1	120	hf	711.3	777.5	835.2	882.4	948.1	22.1	11.5	9.5	13.1	7.0	7.4	7.6	7.7	7.9
FRI2	120	hf	740.3	808.0	892.3	962.6	935.9	22.6	16.9	14.1	-5.3	7.3	7.7	8.1	8.4	7.8
LOM1	100	hf	369.3	416.4	478.9	561.3	676.8	15.7	12.5	16.5	23.1	4.5	4.9	5.3	5.9	6.8
PUG1	95	hf	563.0	605.7	658.5	689.9	754.8	14.2	10.6	6.3	13.0	7.3	7.6	7.7	7.7	7.9
TRE1	210	hf	667.8	742.9	762.6	813.4	866.2	25.0	3.9	10.2	10.6	3.5	3.8	3.8	4.0	4.1
VAL1	160	hf	481.8	503.9	549.5	598.7	648.6	7.4	9.1	9.8	10.0	3.4	3.5	3.7	3.9	4.1
VEN1	140	hf	455.2	502.3	552.2	608.1	658.5	15.7	10.0	11.2	10.1	3.7	4.0	4.2	4.5	4.7
EMI1	65	sc	199.6	209.1	225.2	239.1	238.2	3.2	3.2	2.8	-0.2	4.2	4.2	4.1	4.0	3.7
EMI2	65	sc	215.4	241.4	289.2	320.8	355.6	8.7	9.6	6.3	7.0	4.6	4.8	5.3	5.3	5.5
LAZ1	55	sc	152.6	177.4	190.3	211.7	233.1	8.3	2.6	4.3	4.3	4.1	4.4	4.2	4.2	4.2
MAR1	55	sc	235.0	248.8	266.1	294.2	332.3	4.6	3.5	5.6	7.6	6.4	6.2	5.9	5.9	6.0
SAR1	70	sc	246.9	261.9	261.7	281.9	311.6	5.0	0.0	4.0	5.9	4.7	4.8	4.4	4.3	4.5
TOS1	70	sc	198.6	215.5	238.5	257.7	291.5	5.6	4.6	3.8	6.8	3.8	3.9	4.0	4.0	4.2
PIE1	80	tc	253.0	279.9	307.3	341.1	372.9	9.0	5.5	6.8	6.4	4.1	4.3	4.4	4.5	4.7
SIC1	70	tc	196.3	216.6	211.8	226.9	241.9	6.8	-1.0	3.0	3.0	3.8	3.9	3.5	3.5	3.5
High forests			550.9	603.3	659.6	717.4	768.2	17.5	11.3	11.5	10.2	5.0	5.4	5.6	5.9	6.0
			<i>±130.9</i>	<i>±142.6</i>	<i>±146</i>	<i>±142.8</i>	<i>±125</i>	<i>±5.4</i>	<i>±4.1</i>	<i>±3.1</i>	<i>±7.4</i>	<i>±1.7</i>	<i>±1.7</i>	<i>±1.8</i>	<i>±1.7</i>	<i>±1.6</i>
Stored coppices			208.0	225.7	245.2	267.6	293.7	5.9	3.9	4.5	5.2	4.6	4.7	4.6	4.6	4.7
			<i>±33.2</i>	<i>±31</i>	<i>±34.9</i>	<i>±39.4</i>	<i>±49.8</i>	<i>±2.2</i>	<i>±3.2</i>	<i>±1.3</i>	<i>±2.9</i>	<i>±0.9</i>	<i>±0.8</i>	<i>±0.8</i>	<i>±0.8</i>	<i>±0.9</i>
Transitory crops			224.7	248.2	259.5	284.0	307.4	7.9	2.3	4.9	4.7	3.9	4.1	4.0	4.0	4.1
			<i>±40.1</i>	<i>±44.8</i>	<i>±67.5</i>	<i>±80.8</i>	<i>±92.6</i>	<i>±1.6</i>	<i>±4.6</i>	<i>±2.6</i>	<i>±2.4</i>	<i>±0.2</i>	<i>±0.3</i>	<i>±0.6</i>	<i>±0.7</i>	<i>±0.9</i>

High forests include stands from 95 to 210 years old and are characterized by an average volume of 550.9 m³ ha⁻¹ in 1997 and 768.2 m³ ha⁻¹ in 2015. Stored coppices include stands from 55 to 70 years with an average volume of 208.0 m³ ha⁻¹ in 1997 and 293.7 m³ ha⁻¹ in 2015. Transitory crops included just two plots ranging from 70 to 80 years and with an average volume of 224.7 m³ ha⁻¹ in 1997 and 307.4 m³ ha⁻¹ in 2015.

A positive trend in standing volume has been observed over the concerned period for all the forest types considered (Figure 4.7A). Between 1997 and 2015 high forests had an average increase of 11.84 m³ ha⁻¹ per year, more than double if compared to stored coppices and transitory crops (4.62 m³ ha⁻¹ per year and 4.34 m³ ha⁻¹ per year respectively), which are characterized also by similar trends.




Figure 4.7 – Trends in volume (A), current increment (B) and mean increment (C) at Level II plots in Italy. (Source: LIFE SMART4Action).

The analysis of increments (Figures 4.7 B, C) revealed that high forests are characterized by a decreasing IcV ($10.2 \text{ m}^3 \text{ ha}^{-1}$ per year in 2010-2014), but still higher and distant from ImV ($6.0 \text{ m}^3 \text{ ha}^{-1}$ per year in 2010-2014). For stored coppices and transitory crops, IcV is decreasing ($5.2 \text{ m}^3 \text{ ha}^{-1}$ per year and $4.7 \text{ m}^3 \text{ ha}^{-1}$ per year in 2010-2014, respectively) and very close to ImV ($4.7 \text{ m}^3 \text{ ha}^{-1}$ per year and $4.1 \text{ m}^3 \text{ ha}^{-1}$ per year in 2010-2014), meaning that the latter is very close to the culmination. The strongest variation of IcV among subsequent inventory periods (Figure 4.7B) has been recorded for the 2000-2005 as a possible effect of the well-known 2003 heat wave, as reported by various Authors across Europe (Ciais et al., 2005; Leuzinger et al., 2005; Bertini et al., 2011). This variation was much more evident in high forests and less obvious in coppices and transitory crops.

4.3.3 Forest biodiversity

Forest biodiversity has been assessed mostly as diversity of vascular plant species in the Italian forest monitoring networks, Level I and Level II.

Level I was surveyed in 2007 by means of four 10x10 m Sampling Units (SU) on 201 selected plots, according to a harmonised manual (Canullo et al., 2007). It is considered as the representative baseline for the assessment of vascular plant diversity in Italian forests. As grouping factors, the Biogeographical regions and the EFCTs Forest types (EEA 2007, 2012) have been assumed. The spatial distribution of α -diversity across Level I plots in Italy is given in Figure 4.8, with vascular species density (no. of species $\times 100 \text{ m}^{-2}$) ranging from 2.8 to 54.




Figure 4.8 – Species density (no. of species $\times 100 \text{ m}^{-2}$) as result of mean species richness between 4 SU on each site. (Source: LIFE SMART4Action).

Non Metric Multidimensional Scaling (NMDS) analysis ordinated the sites in terms of similarity of species composition: the most clustered groups by EFTC Forest types are well separated and distributed according both to Latitude and Biogeographical regions (Alpine coniferous forests, Mountainous beech forest, Thermophilous deciduous forests, Broadleaved evergreen forests; Figure 4.9).




Figure 4.9 – Non Metric Multidimensional Scaling plot shows the relation between species composition (Jaccard dissimilarity index) and the EFTC. $R = 76.8\%$; stress $\approx 22\%$. 3 - Alpine coniferous forests; 5 - Oak-hornbeam forests; 6 - Beech forests; 7 - Mountainous beech forest; 8 - Thermophilous deciduous forests; 9 - Broadleaved evergreen forests; 10 - Coniferous forests of the Mediterranean region; 13 - Native plantations; 14 - Exotic plantations and woodlands. (Source: LIFE SMART4Action).

The sampling design in the Level II sites includes 12 10x10 m SUs (Canullo et al., 2013) and proved to capture at least 80% of the expected number of terricolous plant species (including Vascular, Lichens and Bryophytes). For vascular plants, species density was proved to be a good proxy for plot richness (Ferretti et al., 2006). Validated, continuous and comparable datasets cover the period 1999-2012 with 11 plots (Figure 4.10).




Figure 4.10 –Time pattern of total number of species at individual Level II plots. (Source: LIFE SMART4Action).

Linear model for plant species density (no. of species * 100 m⁻²) was tested for the selected plots. (within-subject contrasts RMANOVA. Table 4.3). Significant trends (P<0.05) were detected at CAM1 (Campania, South Italy, beech) where the beech forest shows there is an increasing species density, and at LOM1 (Lombardy, North Italy, Norway spruce), with a noticeable reduction in α -diversity of the transitional Spruce community (R²=0.79).

Table 4.3 – Linear model for species density (no. of species * 100 m⁻²) at the selected Level II plots over the period 1999-2012. Italics: *p*<0.10; bold: *p*<0.05. (Source: LIFE SMART4Action).

Level II site	R ²	<i>p</i>	SE _{regr}	<i>b</i>
ABR1	0,323	<i>0,061</i>	4,815	-0,095
CAL1	0,028	<i>0,099</i>	4,671	-0,111
CAM1	0,118	0,023	9,608	0,171
EMI1	0,046	<i>0,053</i>	3,616	0,067
FRI1	0,055	0,141	4,526	0,111
LAZ1	0,008	0,626	1,562	0,042
LOM1	0,789	0,029	12,153	-0,259
MAR1	0,081	0,149	5,221	-0,110
TOS1	0,104	0,469	2,482	-0,055
VAL1	0,024	0,380	2,204	-0,053
VEN1	0,006	0,371	1,947	0,035

Trends in the species density appear a combination of previous management impact, present changes due to stand dynamics, and external factors. Even if not significant in term of linear trends, is worth to note the visible effect of a strong moth attack (*Lymantria dispar*) in 2002-2003 combined with the 2003 and 2007 heat waves at LAZ1 site (Turkey oak). Beech forest in VEN1 recovered quickly from a severe hailstorm of 1988, leading to still visible effects in 1999. The spruce community in VAL1 has no directional trends, but the annual variability is progressively reducing. The absolute minimum at CAL1 is not apparently linked to known factors.

4.3.4. Forest nutrition

Forest nutrition has been examined in terms of soil and foliar nutritional status.

(i) Soil solid phase. Soil chemistry is measured at Level I and Level II plots. Main results are reported in Figure 4.11 for Level I. In particular:

- carbon content is most frequent between 15-30 g kg⁻¹ and C:N ratio between 10-15;
- base saturation (BS) is most frequently >75%;
- cation exchange capacity (CEC) is most frequently between 10 and 20 mmolc⁺·kg⁻¹, and ca. 50% of the observed plots have a CEC > 20 mmolc⁺·kg⁻¹;
- although ca. 50% of plots have a pH >5.5, there is a considerable amount of plots with pH <4.5.




Figure 4.11 – Soil condition on Level I plots in Italy (n: 239). From top left, clock-wise: frequency of plots in different class of carbon (C), C:N, Cation Exchange Capacity (CEC), Base Saturation (BS) and pH.
(Source: LIFE SMART4Action).

(ii) Soil solution. Soil solution is measured only at Level II sites. It provides important information about changes in the chemistry of soil water that may reflect changes in the atmospheric input and/or changes in ecosystem structure and processes. As such, careful interpretation is always necessary. Main results obtained after evaluating data collected over different time windows (3-15 yrs) and processed by means of Seasonal Mann-Kendall test (SMK) are summarized in Table 4.4.

Table 4.4 – Variable considered, layer examined, no. of plots, range of values among plots and occurrence of significant/non significant trends in soil solution data. (Source: LIFE SMART4Action).

Variable	Layer	No of plot	Range	Significant increasing trend	Significant decreasing trend	Non significant trend
pH	forest floor	6	5.09-6.09	0	3	3
	topsoil	8	4.24-6.59	2	0	6
	subsoil	8	5.04-7.42	4	0	4
NO ₃ -N	forest floor	6	0.65-4.03	2	2	2
	topsoil	8	0.09-8.85	1	1	6
	subsoil	8	0.15-2.89	0	2	6
NH ₄ -N	forest floor	6	0.08-1.34	0	2	4
	topsoil	3	0.03	1	-	-
	subsoil	8	-	-	-	-
SO ₄ -S	forest floor	6	0.5-1.63	0	2	4
	topsoil	8	0.31-2.3	0	5	3
	subsoil	8	0.44-4.71	2	4	2
Base cations	forest floor	6	6.01-15.09	1	3	2
	topsoil	8	0.72-13.88	3	3	2
	subsoil	8	0.85-16.44	2	3	3

(iii) Foliar nutrients. Bi-annual collection and chemical analysis of leaves and needles have been carried out at the Level II plots over the period 1995-2013. Major nutrients have been analyzed: N, S, P, Ca, Mg, K. Main results for the most important tree species are as follows (Figure 4.12A, B):

- in terms of nutritional status, nutrient concentrations are for the most part of species and plots within the acceptable range. Possible exceptions are: high N values reported for some beech sites (Figure 4.12A), and low values reported for holm oak plots; possible low S values at Norway spruce sites; possible low K values at holm oak plots; generalized high values for Ca;
- in terms of time trends, a generalized significant decreasing trend for N (beech, Norway spruce, Turkey oak), S (Norway spruce, Turkey oak). For other nutrients, the time trend is less generalized and more based on species-plot combination (Figure 4.12B).




Figure 4.12 – Foliar nutrients. A) mean N values at six beech plots; B) summary of time trends, all sites. Vertical axis shows the % of significant positive trends (above zero) and the % of negative trends (below zero). The % of non significant trend is not explicitly showed. Numbers above bars indicate the number of analyzed sites for the given nutrient. (Source: LIFE SMART4Action).

4.4. Risk assessment

Risk is evaluated here in terms of possible effects of ozone and N deposition on the various impact indicators reported above.

4.4.1 Risk for forest health

No specific study has been undertaken to investigate the effect of N deposition on forest health in Italy. A European-scale study conducted on Level II sites (including Italy), however, revealed that N deposition and related N variables in soil and foliage improve tree defoliation models. The estimated effect is different according to the species being considered (Ferretti et al., 2015). As for ozone, a slight effect has been found for defoliation on beech (Ferretti et al., 2003; Ferretti et al., 2007; Bussotti and Ferretti, 2009). Results from a subset of Level I plots investigated in Trentino (N. Italy) over the period 2007-2011 show that ozone concentration has no significant effect on tree health of assorted species, with Norway spruce and larch being the most frequent ones (Gottardini et al., 2012).

As for direct effect on foliage, symptoms attributed to ozone were assessed at Level II sites and a subset of Level I plots in Trentino. Although several species were found symptomatic (Bussotti and Ferretti, 2007, 2009; Ferretti et al., 2003; Gottardini et al., 2012), statistical relationship with ozone exposure was always weak (Bussotti and Ferretti, 2009). Recent results obtained with single-species approach and *Viburnum lantana* as *in situ* biomonitor were promising, at least at local level (Gottardini et al., 2014; Gottardini et al., submitted).

4.4.2. Risk for forest growth

Distinct effect of N deposition has been detected on growth and C sequestration (Figure 4.13) (Ferretti et al., 2014). In relative terms, the maximal annual response of basal area increment (BAI) was estimated at 0.074–0.085% for every additional kgN. This corresponds to an annual maximal relative increase of 0.13–0.14% of carbon sequestered in the above-ground woody biomass for every additional kgN, i.e. a median value of 159 kgC per kgN ha⁻¹ per year (range: 50–504 kgC per kgN, depending on the site). The importance of N related variables was further confirmed by a study on 15 Level I plots in Trentino, where N:Mg was the most important predictor of BAI.

No significant effect of ozone has been detected, neither on Level II plots (Ferretti et al., 2003; Ferretti et al., 2014), nor on a subset of Level I plots in Trentino (Gottardini et al., 2012).

All in all, variables related to stand, N deposition, nutrition, soil and climate are by far the most important and significant predictors of growth (Ferretti et al., 2003; 2014).




Figure 4.13 – Annual BAI_{rel} (A) and annual estimated C sequestered (B) in 2000–2009 plotted against annual N deposition over the same time window at Level II plots in Italy. Solid line: all species; dashed line: broadleaves only. FS: *Fagus sylvatica*; PA: *Picea abies*; QD: deciduous oak (only *Quercus cerris* in this diagram); QI: *Quercus ilex*. (after Ferretti et al., 2014).

4.4.3. Risk for forest biodiversity

Possible air pollution effect on species diversity at Level II plots has been examined in two respects: the expected sensitivity of species composition in relation to ozone and according to the list of sensitive species by the ICP Forests (Ferretti et al., 2003) and the expected effects of stand, soil, meteorology and deposition data on species density (Ferretti et al., 2006).

As for the frequency of ozone sensitive species, an ozone vulnerability index (OVI) that takes into account trees, shrubs, and herbs was calculated. According to this index, beech forests in north and central Italy were potentially the most sensitive to ozone.

As for the set of factors affecting vascular species diversity, multivariate (Generalized Linear Models; Ordinary Least Square Regression) and univariate (Spearman rank order correlation) statistical methods were tested on the set of data collected over the period 1999-2003. All in all, results of the multivariate approach revealed that soil (C, N, C/N, C/P, K, P) and stand (tree species, tree species in the upper layer, leaf litter) are the most important variables: species diversity was found to decrease at increasing level of soil C, N, and C/N.




Figure 4.14 – Species density at beech Level II sites vs. exceedance of critical loads (CL) for N for the same sites. CL and exceedances were derived from measurements at the very site. Drawn after Ferretti et al., 2006.

Univariate analysis carried out for beech plots revealed that species diversity decreases as N deposition and exceedance of Critical Load for N increases ($0.001 < P < 0.05$) (Figure 4.14).

4.4.4. Risk for forest nutrition

Distinct effect of N deposition on soil pH and Basic Cation Exchangeable (BCE) was documented, especially for broadleaved forests (Ferretti et al., 2014) (Figure 4.15). Significant effect on foliar nutrient ratios was also reported (Ferretti et al., 2014) (Figure 4.16). With respect to ozone, no study has been undertaken so far to evaluate possible effects on soil biota and foliar nutrients.




Figure 4.15 – pH (a), C:N (b) and BCE (c) of the mineral topsoil plotted against actual N deposition. Soil data are those obtained after the 1995-1996 survey. Deposition data are mean annual values 2000-2009. Regressions represent always the best fit for the given dataset. Continuous line: all species; dashed line: broadleaves only. FS: *Fagus sylvatica*; PA: *Picea abies*; QD: deciduous oaks; QI: *Quercus ilex*. After Ferretti et al., 2014.




Figure 4.16 – Foliar N:P (a), N:Ca (b), N:Mg (c) and N:K (d) plotted against measured throughfall deposition. Foliar data are mean values after sampling carried out at years 2001, 2003, 2005, 2007, 2009. Deposition data are mean annual values 2000-2009. Continuous line: all species; dashed line: broadleaves only. FS: *Fagus sylvatica*; PA: *Picea abies*; QD: deciduous oak; QI: *Quercus ilex*. After Ferretti et al., 2014.

4.5 Conclusions

When using measured data for both pressure and impact, evidence from the Italian forest monitoring networks (UNECE ICP Forests Level I and II) can be summarized as follows:

(i) trends in air pollution issues. There is a significant reduction of sulphate; for nitrate and ammonium the decrease is much slighter, although significant at some sites. After an increase between 1996-2005, mean summer ozone concentration is now decreasing. A decreasing 2000-2013 trend was also reported at the European scale (Sanders et al., 2016).

(ii) Measurable impact. Forest health improved over the 1997-2014 period, with a significant ($P < 0.01$) decreasing trend for defoliation, mostly driven by broadleaves. Forests at Level II sites show positive trends in terms of average increment, and the standing volume is increasing. This latter point is not surprising, as sites are not actively managed and accumulate biomass. Forest biodiversity shows huge variability (Level I) but no clear trend (Level II) over the period 1999-2012. Forest nutrition revealed that acidic soils occur at ca. 50% of Level I plots. In soil solution data collected at level II sites, a decreasing trend for sulphate is evident. There is also significant

evidence of reduced basic cation leaching, which are important in relation to the impacts of acidifying deposition on soil chemistry. Foliar nutrients revealed a composite picture, with mostly decreasing trend for N and S, increasing for K, and much less obvious pattern for P, Ca, and Mg.

(iii) Evidence for risk. Although in terms of reported levels, ozone is potentially a serious risk for Italian forests, measurable effects are quite limited on both Level I and II. Therefore, evidence for actual risk is weak, and limited effect on health and growth were reported by the various study undertaken since early 2000s. On the other end, evidence for N deposition effect on forest nutrition and growth is outstanding. Current level of N deposition was proven to impact soil and foliar chemistry, tree growth and carbon sequestration. Effects on plant diversity were also reported, mainly for beech forests.

Effects of air pollution on crops and semi-natural vegetation



National Focal Point: Fausto Manes and Elisabetta Salvatori (UNIROMA1)

Chapter coordinator: Fausto Manes

Contributors: Elisabetta Salvatori¹, Alessandra Campanella², Maria Chiesa³, Lorenzo Cotrozzi², Angelo Finco³, Lina Fusaro¹, Giacomo Gerosa³, Giacomo Lorenzini², Riccardo Marzuoli³, Cristina Nali², Romina Papini², Elisa Pellegrini², Mariagrazia Tonelli², Fausto Manes¹

¹Department of Environmental Biology, Sapienza University of Rome; ²Department of Agriculture, Food and Environment, University of Pisa; ³Department of Mathematics and Physics, Catholic University of Brescia

CHAPTER 5 - EFFECTS OF AIR POLLUTION ON CROPS AND SEMI-NATURAL VEGETATION

5.1 Introduction

The International Cooperative Programme on Effects of Air Pollution on Natural Vegetation and Crops (ICP Vegetation, formerly ICP Crops) was established in 1987 as a subsidiary body of the Working Group on effects of the UNECE Convention on long-range transboundary air pollution (LTRAP). ICP Vegetation is an international research programme investigating the impacts of air pollutants (particularly tropospheric ozone (O₃), heavy metals and nitrogen) on crops and semi-natural vegetation.

Participants meet each year at the Task Force Meeting (TFM) to discuss recent results and the future development of the programme. Recently, Italy hosted the 25th and the 28th of the TFM, respectively organized by the Math and Physics Department of the Catholic University of Brescia in 2012, and by the National Focal Centre, Department of Environmental Biology, Sapienza University of Rome, in 2014. Some of the works that were presented during the Rome meeting are published in *Annali di Botanica*, Vol. 5, 2015.

5.2 Pressures

Tropospheric ozone (O₃), nitrogen deposition and heavy metals are currently identified by the ICP Vegetation as the main sources of pressure for crops and (semi)-natural vegetation.

5.2.1 Ozone

O₃ is considered as the main oxidizing agents in the near-surface atmosphere of the Mediterranean region (Cristofanelli and Bonasoni, 2009), also acting as a major greenhouse gas (Cooper et al., 2014). Despite the increasing efforts to regulate the precursor emissions this secondary air pollutant (Directive 2008/50/EC, EC, 2008), background O₃ levels are continuously increasing, particularly in Southern Europe (Coll et al., 2009; Cooper et al., 2014). Although the reduction of emissions has decreased the magnitude of O₃ peaks over the last decade, high daily summer O₃ concentrations still occur in the Mediterranean area, favored by high values of irradiance and temperature (Fernández-Fernández et al., 2011; Sicard et al., 2013). Furthermore, nocturnal O₃ pollution episodes are also frequent in the Mediterranean area, particularly along coastal sites located downwind of large urban conurbations, thus posing a concrete risk to vegetation (Fares et al., 2009; Mereu et al., 2009).

Fig. 5.1 describes the temporal trend of the target value for the protection of vegetation, indicated by the EU Directive (EC, 2008) as the AOT40 calculated over a given period using only the one-hour values measured between 08:00 to 20:00 CET in selected monitoring stations with different characteristics. The 5-y average of 9 ppm h (target value for vegetation protection) is systematically exceeded in most of them (see fig. 5.1).




Figure 5.1 – AOT40 (expressed in ppm h) 5-y running averages as observed in 8 Tuscan monitoring stations (Data source: Tuscan Regional Environmental Agency, ARPAT, Florence). The dashed horizontal line shows the target value for the protection of vegetation. GR-Maremma is a rural site; AR-Casa Stabbi is rural/background; AR-Acropoli, FI-Settignano, PT-Montale, LU-Carignano, PI-Passi, PI-Montecerboli are suburban.

It is also interesting to estimate the biological additive effects of O₃ pollution and summer heat waves. This because the meteorological conditions typical of the heat waves (i.e. high temperature, intense solar radiation, dryness) are also responsible of the formation and build-up of photochemically generated ozone (Lorenzini et al., 2014) (Fig. 5.2).




Figure 5.2 – Diurnal variation of ozone hourly concentration in the rural/background monitoring station of Casa Stabbi (Arezzo Province) (1 July-15 August) in 2012 (two years with dramatic summer heat waves), in comparison with an historical series (1999-2002 + 2004-2011) (After Lorenzini et al., 2014).

5.2.2 Nitrogen

Among the main concerns for the health of Mediterranean vegetation, increasing attention is given to anthropogenic perturbation of the nitrogen (N) cycle (Phoenix et al., 2006; Ochoa-Hueso et al., 2011). Nitrogen is often a limiting factor in many terrestrial ecosystems, but its cycle at global scale has been altered by post-industrial human activities, mainly because of combustion of fossil fuels and fertilization practices (Haber-Bosch reaction) aimed at increasing agricultural production, and resulting in increased N deposition (mainly nitrates and ammonium) (Galloway et al., 2008). NO_x and NH₃ emitted to the air are the main nitrogen pollutants contributing to the amount of reactive nitrogen rising levels of atmospheric deposition on soils, vegetation surfaces and waters (The European Nitrogen Assessment, 2011).

5.3 Impacts

5.3.1 Ozone

O₃ is known to affect plant photosynthetic function in different ways, and the effects of the O₃ stress can be observed at biochemical, microscopic and macroscopic level. The first interaction between O₃ and plants occurs at the level of the guard cells of stomata, in which O₃ triggers the production of Reactive Oxygen Species (ROS), thus activating a signalling cascade inducing an overall efflux of anions and K⁺ (Vahisalu et al., 2010; Vainonen and Kangasjärvi, 2015). The consequent loss of guard cells turgor, leading to stomatal closure, reduces the gas uptake through stomata; this mechanism restricts the O₃ flux into the leaves (“avoidance mechanism”, Castagna and Ranieri, 2009) but, at the same time, also limits the photosynthetic CO₂ assimilation (Astorino et al., 1995). The O₃ molecules that enter the sub-stomatal chamber can directly peroxidate the membrane lipids of the mesophyll cells (Vitale et al., 2008) or, in the apoplastic space, can generate toxic ROS (like OH[•], O₂^{-•}, H₂O₂), which can then move inside the cells (Vainonen and Kangasjärvi, 2015). Furthermore, O₃ triggers the ROS production from different endogenous, enzymatic sources, a process known as “oxidative burst” (Manes et al., 1990a; Scalet et al., 1995). Despite acting as signalling molecules, that leads to the activation of several plant defence responses (Vainonen and Kangasjärvi, 2015), ROS are also responsible of direct oxidative damages to different molecules involved in the photosynthetic process, such as chlorophylls a and b, and Rubisco, whose activity and content have been reported to decline under O₃ stress (Goumenaki et al., 2010). Moreover, detrimental effects of O₃ on the photosynthetic electron transport chain, and in particular on Photosystem II (PSII) function, have been demonstrated (Guidi et al., 2002; Pellegrini, 2014). A reversible, photoprotective down regulation of PSII photochemistry is, however, the most commonly observed PSII response to O₃, being a consequence of the reduced demand of NADPH and ATP from the Calvin cycle; the latter can be caused by both stomatal closure and biochemical limitations (Bussotti et al., 2011; Mereu et al., 2011; Salvatori et al., 2013).

Also, the role of Photosystem I (PSI) in plant photosynthetic response to O₃ has received increasing attention in the last years. In fact, the measurement of prompt chlorophyll “a” fluorescence (PF) and the JIP test analysis (Strasser et al., 2004; 2010) have shown that the I-P part of the PF transient, which correlates to PSI content and activity (Ceppi et al., 2012; Schansker et al., 2005), is particularly sensitive to O₃ and other oxidative stress factors (Oukkarroum et al., 2009; Bussotti et al., 2011; Pollastrini et al., 2014; Bernardini et al., 2016).

In particular, the amplitude of the I-P phase of the PF (ΔV_{I-P}) was reduced by O₃ stress in many studies, indicating a negative effect of this pollutant on the efficiency of electron transport through PSI, to reduce the end acceptors beyond PSI (Bussotti et al., 2011; Mereu et al., 2011; Pollastrini et al., 2014). Recent studies carried out with the innovative technique of multi signal fluorescence measurement (Fusaro et al., 2016a; Salvatori et al., 2015) have highlighted that, in both crop and natural species, it is possible to distinguish an early O₃ response of the photochemical apparatus, involving PSII only, and a late response, occurring when O₃ cumulative stress becomes more severe, in which PSI activity and content are also modulated.

Ultrastructural alterations and visible leaf injury are reported in sensitive species after ozone stress. The appearance and diffusion of visible leaf injuries due to ozone exposure were reported in several experiments on bean (Gerosa et al., 2009) and durum wheat (Gerosa et al., 2014; Monga et al., 2015). Results showed that microscopical leaf symptoms, assessed as cell death and H₂O₂ accumulation, preceded by three-four days the appearance of visible symptoms.

Iriti et al. (2006) obtained similar results on currant tomato, suggesting the potential use of this species for ozone biomonitoring protocols, since there was a linear relationship between the intensity and diffusion of visible leaf injuries and the hourly ozone concentration mean.

Basile et al. (2010) have applied Transmission Electron Microscope (TEM) analysis on O₃-resistant (NC-R) and O₃-sensitive (NC-S) clones of *Trifolium repens* L. cv. Regal, exposed to ambient O₃ conditions in the Botanical Garden of the Sapienza University of Rome, Italy, from June to October 2005, following the ICP Vegetation Experimental Protocol. This work has highlighted that ultrastructural injuries appeared to be widespread in NC-S leaves, showing dead cells and heavily damaged living cells, while the NC-R clones showed much less damage: the chloroplast maintained an almost intact organization of thylakoids (granal and intergranal regions), but the chloroplast side facing the apoplastic space appeared with no thylakoids. Interestingly, Hsp70 levels, an important stress marker, were only slightly increased in the O₃-sensitive clone with respect to the O₃-resistant clone; in contrast, a strong decrease (–60%) in phosphoenolpyruvate carboxylase (PEPCase) protein levels was measured in the ozone-resistant clone.

Pellegrini et al. (2015a) conducted a comparative study on functional leaf traits and the diurnal dynamics of photosynthetic processes on plants of two grape (*Vitis vinifera*) varieties (Aleatico, ALE, and Trebbiano giallo, TRE), exposed under controlled conditions to realistic concentrations of O₃ (80 ppb for 5 h day⁻¹, 8:00-13:00 h, + 40 ppb for 5 h day⁻¹, 13:00-18:00 h). At the constitutive level, morphological functional traits of TRE improved leaf resistance to gas exchange, suggesting that TRE is characterized by a potential high degree of tolerance to ozone. At the end of the treatment, both varieties showed typical visible injuries on fully expanded leaves and a marked alteration in the diurnal pattern of photosynthetic activity. This was mainly due to a decrease in stomatal conductance (-27 and -29% in ALE and TRE, respectively, in terms of daily values in comparison to controls) and mesophyll functioning (+33 and +16% of the intercellular carbon dioxide concentration). Although the genotypic variability of grape regulates the response to oxidative stress, similar detoxification processes were activated, such as an increased content of total carotenoids (+64 and +30%, in ALE and TRE), enhanced efficiency of thermal energy dissipation within photosystem II (+32 and +20%) closely correlated with the increased de-epoxidation index (+26 and +22%) and variations in content of some osmolytes.

In summary, it can be concluded that: the daily photosynthetic performance of grapevine leaves was affected by a realistic exposure to ozone. In addition, the gas exchange and chlorophyll *a* fluorescence measurements revealed a different quali-quantitative response in the two varieties. The genotypic variability of *V. vinifera* and the functional leaf traits seem to regulate the acclimation response to oxidative stress and the degree of tolerance to ozone. Similar photoprotective mechanisms were activated in the two varieties, though to a different extent.

Valletta et al. (2016) have investigated the physiological and metabolic effects of O₃ on two wine cultivars of *V. vinifera*, a red grape (San Giuseppe) and a white grape (Maturano) (Fig. 5.3), both autochthonous cultivars of the Latium region and having an economic importance at local scale. The results showed that the white grape cultivar appeared more sensitive to O₃ stress. Moreover, differently from what expected O₃ did not activate stilbene production but, in both cultivars, influenced the content of chlorogenic acid (CGA), an important molecule in the biosynthetic pathway of polyphenols and an antioxidant itself, which decreased in the fumigated samples and recovered after 8 days. The decrease in CGA not necessarily indicates a decrease in its biosynthesis, since it is highly probable that CGA is consumed or as antioxidant, or as precursor to other phenolic compounds.



Figure 5.3 – “Walk-in” Chamber facility of the Department of Environmental Biology, Sapienza University of Rome, with *V. vinifera* plants fumigated with O₃.

As a result of the above described invisible and visible alterations, and of the increase in metabolic cost for detoxification and repair, a reduction of total biomass, as well as a decreased carbon allocation to heterotrophic tissues (i.e. roots and fruits) (Andersen, 2003), is frequently reported for different species (Salvatori et al., 2013). In crops, such effects translate in a reduction of yield, as well as of food quality, with consequent economic losses to the agricultural sector that are mostly reported for Italy and Spain. In particular, Nali et al. (2002), on the basis of O₃ concentrations recorded by ten monitoring stations in Tuscany, have estimated yield losses due to ozone, which varied in Florence from 8% for corn and alfalfa to 27% for soybean, in Pisa from 5% for corn to 24% for soybean, in Lucca from 3% for corn to 17% for soybean. A preliminary economic estimate for corn, wheat, barley, soybean, tomato and alfalfa, calculated annual damage to be 4.6 MEuro in Florence, 0.5 MEuro in Lucca and MEuro in Pisa.



Figure 5.4 – Open-Top chambers facility of C.R.IN.ES. at Curno (Bergamo).

Several experiments on horticultural yield losses due to ozone were carried out in the Po plain climatic conditions at the CRINES research site (Centro di Ricerca Inquinamento atmosferico ed Ecosistemi, Curno, Bergamo) by the Environmental Physics and Ecophysiology research group of the Catholic University of Brescia (Fig. 5.4).

Results showed mean yield reductions of 40% on bean (cv. “borlotto nano lingua di fuoco” Gerosa et al., 2009), 36% on tomato (cv. “Oxheart”, Gerosa et al., 2008), 18% on lettuce cv. “Romana” and 14% on lettuce cv. “Canasta” (Marzuoli et al., 2016a). In the same experimental site Monga et al. (2015) performed an experiment on durum wheat, highlighting yield losses up to 16% (in cv. “Sculptur”) in plants that were exposed to O₃ enriched air (+50% of the ambient ozone).

Dry biomass reductions up to 25% due to O₃ were observed on alfalfa in the Po plain, and significant differences of forage quality parameters were also found for the same species. However, in all of the above mentioned experiments, a strong intraspecific variability of the response to ozone was observed. For example, ozone caused a slight increase of fruit yield in the tomato cv “San Marzano” (Gerosa et al., 2008), Monga et al. (2015) in a varietal screening experiment found that only 2 of the 5 studied cultivars of durum wheat were ozone sensitive.

The dataset of the experiment on tomato contributed to the definition of a dose-response relationship and critical levels based on ozone flux and fruit yield loss (Gonzalez-Fernandez et al., 2014) that was included in current version of the “Manual on Methodologies and Criteria for Modelling and Mapping Critical Loads and Levels and Air Pollution Effects, Risks and Trends” (CLRTAP, 2010). Other experimental datasets have been required for the definition of analogous dose-response relationships for leafy crops species (lettuce, Marzuoli et al., 2016a), bean (Gerosa et al., 2009) and durum wheat (Monga, 2015; Gerosa et al., *in preparation*) in order to define their specific critical levels.

In addition to controlled conditions experiments (with Open-Top Chambers), some open field experiments were conducted in order to characterize the ozone uptake at agricultural ecosystem level, and to quantify the partition of the total ozone flux between the stomatal and non-stomatal components of the deposition pathway. These studies are strictly necessary to properly calculate the

ozone dose absorbed by vegetation for the regional level risk assessment, and for the application of the critical levels.

In this context, it is important to mention the field experiments performed on winter wheat (Gerosa et al., 2003) and soybean (Gerosa, 2002) that highlighted the presence of a significant non-stomatal component of the ozone total flux that could cause serious overestimation of the effects in case it would not be taken into account (Bassin et al., 2004; Tuovinen et al., 2007). These results contributed in the fine calibration of the EMEP model (which contain the DO₃SE module embedded) for the estimation of the non-stomatal component (Tuovinen et al., 2004).

Finally, a dataset of measurements on onion fields (Gerosa et al., 2007) was used to further improve the ozone flux partition process, identifying the agrometeorological parameters that mostly can influence the quality of the flux estimations.

The ecophysiological and biochemical effects of realistic O₃ exposure under controlled conditions upon medicinal species have been also investigated.

In particular, *Salvia* (Pellegrini et al., 2015b) and the perennial herbaceous lamiacea *Melissa officinalis* (known as lemon balm) have been considered. Lemon balm has been selected as a test species to elucidate the variations on biochemical composition of an officinal plant under short- and long-term exposure to O₃. Whole plants and cell cultures were investigated and biotechnological practical implications envisaged. A consistent fraction of this project (the one focused on rosmarinic acid biosynthesis at the molecular level - Döring et al., 2014a, b) has been performed jointly with a German team. Signaling molecules, membrane integrity, secondary metabolism, programmed cell death, photosynthetic performances are amongst the other key issues taken into account (Pellegrini et al., 2011, 2013b; Tonelli et al., 2015; D'Angiolillo et al., 2015).

The lichen response to O₃ exposure has been evaluated (Pellegrini et al., 2014a), thorough description of the biochemical and physiological mechanisms that are at the basis of the O₃-tolerance of lichens. Chlorophyll *a* fluorescence emission, histochemical ROS localization in the lichen thallus, and biochemical markers [enzymes and antioxidants involved in the ascorbate/glutathione(AsA/GSH) cycle; H₂O₂ and O₂ were used to characterize the response of the epiphytic lichen *Flavoparmelia caperata* exposed to O₃ (250 ppb, 5 h d⁻¹, 2 weeks) at different watering regimes and air relative humidity in a fumigation chamber. After a two-week exposure Chl*a*F was affected by the watering regime but not by O₃. The watering regime influenced also the superoxide dismutase activity and the production of ROS. By contrast O₃ strongly influenced the AsA/GSH biochemical pathway, decreasing the AsA content and increasing the enzymatic activity of ascorbate peroxidase, dehydroascorbate reductase and glutathione reductase independently from the watering regime and the relative humidity applied.

5.3.2 Nitrogen

Increasing in N deposition is a growing threat to semi-natural grasslands that are listed as a priority habitat for biodiversity conservation in European Union Habitats Directive (92/43/CEE). A study carried out on grassland in Central Italy highlighted that N deposition markedly increases aboveground biomass, maintaining low species diversity (Bonanomi et al., 2006). The strong growth response the authors found out using a relatively low level of nitrogen enrichment, clearly indicated a nitrogen limited community.

Nitrogen eutrophication could exacerbate the deleterious effects of land abandonment on grassland biodiversity (Bonanomi et al., 2006) through enhanced above-ground competition and litter accumulation.

Moreover, since stressors like O₃ or drought can have an impact on roots biomass (Fig. 5.5), puzzling interaction effects might occur. In order to understand the effects of ongoing global environmental change on Mediterranean ecosystems, it is necessary take into account how the multiple stresses affect vegetation since a comprehensive picture is far from being given (Tattini and Loreto, 2014). A recent work by Fusaro et al. (2016b) has investigated how functional and structural traits of two Mediterranean species with different leaf habits (*Fraxinus ornus* and *Quercus ilex*) shift because of nitrogen (N) addition (30 kg ha⁻¹ y⁻¹), also exploring the effect that nitrogen has on the water stress response. Their results have shown that the early successional, deciduous *F. ornus* tends to invest more nitrogen at leaf level enhancing photosynthetic machinery, whereas late successional evergreen *Q. ilex* invests resources on non-photosynthetic biomass, keeping constant N content at leaf level. These differences between species can modify their competitive relations, through influence temperate forest community structure and dynamics. This study also highlights that N can have role in water stress response in both species, but in different way. In *F. ornus* N has ameliorative effect on water stress, determining high gas exchange rates and photosystems functionality. On the contrary, in *Q. ilex* the N addition seems to increase the susceptibility to water stress, possibly because of changes in biomass partitioning due to N. Moreover, studies relative interaction between N deposition and O₃, highlighted an adverse synergistic effect, since exposure to ambient O₃ concentrations was shown to reduce the Nitrogen Use Efficiency. On the other hand, the growth of plants in response to N (i.e. root development), mitigates impact on biomass and physiology due to O₃ (Marzuoli et al., 2016b). Mechanism of action of N on tree species can contribute to provide a more reliable risk assessment, and accordingly it should be implemented for natural and semi-natural vegetation.




Figure 5.5 – Root biomass of NC/R and NC/S clover clones, harvested after 1, 2, 3, 4 and 5 months of exposure to ambient O₃ in the Botanical Garden of Sapienza University of Rome.

5.4 Risk assessment

The methodologies developed within UN/ECE to assess the impacts of air pollution on crops are described in the Chapter 3 of the “Manual on Methodologies and Criteria for Modelling and Mapping Critical Loads and Levels and Air Pollution Effects, Risks and Trends.” (UN/ECE, 2010).


5.4.1 Ozone

Chapter 3 of the mapping manual considers two main approaches to set critical levels to protect vegetation from tropospheric ozone: the concentration-based critical level (O_3 accumulated over a threshold of x ppb, AOT x) and the uptake-based critical level (accumulated O_3 dose above a threshold of Y or phytotoxic ozone dose, PODY). The critical levels are defined as the dose that causes a reduction by 5% of the yield at 95% confidence level (UN/ECE, 2010). The two approaches differ by their data requirements (UN/ECE, 2015a), because PODY is linked to stomatal uptake that is limited by environmental condition such as temperature, light, soil moisture, vapor pressure deficit, phenology and wind (Mills et al., 2011). Since their first establishment in 1996, the O_3 critical levels have been intensively debated among scientists. In particular, many studies carried out in Italy have shown that, despite O_3 exposure at Italian background sites exceeds the AOT40-based critical level, the occurrence of adverse effects of O_3 on forests and crops appears controversial (Manes et al., 2005; Ferretti et al., 2007). At this regard, the use of the stomatal uptake concept (PODY) is highly recommended since, besides pollutant concentration, it takes into account vegetation characteristics and environmental constraints that affect the dose of O_3 effectively absorbed by the plant, an aspect that is of particularly importance under Mediterranean climatic conditions (Ferretti et al., 2007; Manes et al., 2005; Anav et al., 2016; De Marco et al., 2015).

The most common methodology to define the risk assessment is biomonitoring. Biological monitoring can be defined as the measurement of the response of living organisms to changes in the air quality of their environment (Nali et al., 2006), and may provide integrated information on air quality impacts. As plants are more sensitive in terms of physiological reaction to the most prevalent air pollutants than humans and animals (Nali et al., 2006), they are more suitable to be used for biomonitoring. Biomonitoring can be performed through the analysis on the vegetation already present in a given study area (so-called passive biomonitoring) or carried out with selected test plants introduced at the study site (active biomonitoring) (Nali and Lorenzini, 2007). The selection of plants with different sensitivity/resistance to O_3 has been a challenge since 1950's (Karlsson et al., 2003), with the purpose of biomonitoring phytotoxic effects of this pollutant under conditions of ambient exposure. From 1987, these efforts have been coordinated under the ICP Vegetation (Harmens et al., 2015a). Many plant species have been tested as active O_3 biomonitors on pan-European scale, such as radish (*Raphanus sativus* L. cv. Cherry Belle, Manes et al., 1990b; Allegrini et al., 1994), *Phaseolus vulgaris* cv Lit (Astorino et al., 1995), and *Trifolium subterraneum* cv Geraldton. From 1996 to 2004, the ICP Vegetation international biomonitoring programme has involved exposure of ozone sensitive (NC-S) and ozone resistant (NC-R) biotypes of white clover (*T. repens* L. cv. Regal) (Heagle et al., 1994) (Figg. 5.6, 5.7). The clover system has proven to be useful to detect detrimental effects of ambient O_3 (Manes et al., 2003; Nali et al., 2009), however visible injury and biomass reductions proved to be more related to stomatal O_3 uptake than to air O_3 concentration, expressed as AOT40 (Mills et al., 2011).



Figure 5.6 – “Biomonitoring station” with NC-S and NC-R clover clones, in the Botanical Garden of Sapienza University of Rome.



(a) (b)
Figure 5.7 – Healthy (a) and O₃-injured (b) leaves of white clover, NC-S clone.

Moreover, studies carried out in Mediterranean climatic conditions, have shown the importance to consider the effect of environmental variables, such as high air temperatures and Vapour Pressure Deficit (VPD), which can act as confounding factors on the O₃ dose-response relationship of the clover system, even if plants are grown following standard protocols under non limiting water availability (Ferretti et al., 2007; Manes et al., 2005).

Ozone-hypersensitive tobacco (*Nicotiana tabacum* L.) Bel-W3 has been also used worldwide as a reliable and sensitive bioindicator of ozone, showing a characteristic and specific foliar response. The typical foliar lesions induced by ozone under realistic exposure to ambient air are bi-facial greyish necrotic spots, scattered over the lamina (Manes et al., 1990b; Lorenzini et al., 2000). The tobacco system has been extensively investigated, also from the quantitative point of view. Biomonitoring campaigns have been successfully performed all over the world. Ozone-resistant tobacco plants (cv. Bel-B) are routinely inserted in the plots; their sensitivity threshold, in terms of visible injury, for 2 h exposures is 220 ppb vs. 100 ppb of Bel-W3.

Therefore, the appearance of injury on Bel-W3 but not on Bel-B provides further confirmation that such injury is due to ozone. Conventional biomonitoring is performed with adult tobacco plants (about 2 months old, 60 cm in height). However, some years ago, an innovative miniaturized kit for ozone biomonitoring was developed and patented at the University of Pisa, employing plates with tobacco germlings (typically (Lorenzini, 1994; Lorenzini et al., 1995 – Fig. 5.8). These plates are exposed to ambient air for 7 days in shaded conditions, and the intensity of the injury of cotyledons and the first leaf is visually assessed and related to the dose of ozone to which the seedlings have been exposed (Nali et al., 2004; 2007). More precisely, at the University of Pisa two are the applicative fields based on the tobacco mini-kit, i.e. (i) educational projects of environmental education with young pupils (and their teachers and families) (e.g. Nali and Lorenzini, 2007; Pellegrini et al., 2014b), and (ii) regional wide season-long campaigns to fulfill commitments of local/national environmental authorities to allow permitting of industrial plants such as thermal power stations. Under the guidance of their teachers, the pupils had several opportunities to practice with many basic and applied study areas and disciplines and were initiated into the scientific method in a simple and absorbing manner. Though primarily an educational exercise, the survey provided sound research elements and the picture of pollution that emerged has increased the knowledge of air quality in the investigated area.



Figure 5.8 – The miniaturized kit for biomonitoring of O_3 with tobacco Bel-W3 germlings. The external dimensions are 12.5x8.5 cm. Please note necrotic lesions induced by the exposure to ambient air in the presence of O_3 .

Recently, ozone sensitive (S156) and ozone resistant (R123) genotypes of snap bean (*Phaseolus vulgaris* L.), selected through genetic crosses, have been proposed as a new biomonitoring system (Burkey et al., 2005) (Fig. 5.9). Several studies have tested the new system under semi-controlled environmental conditions, showing that pod yield was similar in S156 and R123 under O_3 concentrations lower than 30 ppb (S156/R123 pod yield ratio equal to 1), and consistently reduced by increased O_3 levels in the sensitive genotype only (S156/R123 pod yield ratio < 1) (Burkey et al., 2005, 2012; Flowers et al., 2007). Given these promising results, the ICP Vegetation O_3 programme has conducted field trials using the snap bean system, following a standardised experimental protocol at pan-European scale since 2008 (Fig. 5.10), the advantage being also the cost-effectiveness of the seed-propagated bean plants, in respect to the vegetative-propagated clover clones (Burkey et al., 2005).

The results of the field trials, however, have shown that the extent of leaf injury on S156 variety, as well as the S156/R123 pod yield ratio, were often not directly related to the AOT40 at the experimental site. This was particularly true for Mediterranean sites, such as the Castelporziano Estate (Rome, Italy), in which the response of the snap bean system during three consecutive summer periods (years 2008-2010), characterized by different meteorological conditions and O₃ levels, was not related to seasonal ozone concentration (Fusaro et al., 2015). Recently, Salvatori et al. (2013) pointed out that the O₃ response of S156 and R123 bean lines varied within plant growth stage, with the different O₃ sensitivity of the two genotypes being more apparent during vegetative growth and flowering. Such differences can be due to different stomatal conductance of the sensitive and resistant plants, and to different effect of O₃ on the I-P phase of the fluorescence transient (Salvatori et al., 2015). A large-scale application of the snap bean biomonitoring system, under different climatic conditions and O₃ levels found at pan-European scale, appears therefore unsuitable.



Figure 5.9 – Ozone sensitive (S156) and ozone resistant (R123) genotypes of snap bean (*Phaseolus vulgaris* L.), during a fumigation experiment carried out in the “walk-in” chambers of the Department of Plant Biology, Sapienza University of Rome.



Figure 5.10 – O₃-resistant (R123) and O₃-sensitive (S156) snap bean plants, after three months of ambient air exposure in the Castelporziano Presidential Estate, following the ICP Vegetation biomonitoring protocol (Fusaro et al., 2015).

5.4.2 Nitrogen

The risk assessment of nitrogen pollution in European Mediterranean natural and semi-natural ecosystems is difficult since only few experiments were carried out in the whole region. In Chapter 5 of the Mapping Manual, a list of critical loads ($\text{kg N ha}^{-1} \text{y}^{-1}$) is reported for natural and semi-natural ecosystems classified according to the European Nature Information System (EUNIS). For example, over a range of 15 and 25 $\text{kg N ha}^{-1} \text{y}^{-1}$ Mediterranean xeric grasslands are expected to increase production and dominance of graminoids. Between 20 and 30 $\text{kg N ha}^{-1} \text{y}^{-1}$, Mediterranean shrub ecosystem changes species richness and composition; however, critical loads have not yet been set.

Biodiversity as an important indicator of soil acidity and eutrophication: the role of the modelling in preserving it



National Focal Point: Patrizia Bonanni, Francesca Fornasier, Marcello Vitale, Alessandra De Marco

Chapter coordinator: Maria Francesca Fornasier

*Contributors: Maria Francesca Fornasier¹, Patrizia Bonanni¹, Marcello Vitale², Michele De Sanctis²,
Giuliano Fanelli², Fabio Attorre², Alessandra De Marco³, Silvano Fares⁴, Luca Salvati⁴*

*¹ISPRA, Italian National Institute for Environmental Protection and Research; ²Sapienza University of Rome;
³ENEA, Laboratory of Atmospheric Pollution; ⁴CREA, Council for Agricultural Research and Economics*

CHAPTER 6 - BIODIVERSITY AS AN IMPORTANT INDICATOR OF SOIL ACIDITY AND EUTROPHICATION: THE ROLE OF THE MODELLING IN PRESERVING IT

6.1 Introduction

The activities linked to biodiversity protection in Italy (and Europe) are the main issue of the ICP Modelling and Mapping Task Force. The objective to reach no net loss of biodiversity is the target for 2020, and modelling is the most suitable methodology at regional and national scale in order to investigate biodiversity losses.

Biodiversity provides ecosystem services crucial for human well-being. However, rapid population growth, coupled with socio-cultural changes, climate change and the unpredictable nature of economic change and its transition patterns, create challenges for decision-makers. Air pollution is a serious threat to the diversity of life. In general, it can be said that the effects of air pollutants on biological diversity usually affect more lower life forms than higher forms, and, as a result of air pollution, the most affected species decline, while some increase. In general, sensitivity varies from species to species within each group of organisms, but also due, for example, to the pollution load, the stage of life at which the individual is exposed, and the way competition is altered within a particular ecosystem. The effects found on plants affect mostly lichens, bryophytes, fungi, herbaceous flowering species, and trees.

Since 1970s, some evidences concerning negative effects of acid deposition on natural ecosystems have occurred. At the beginning of 1980s, extensive forest damage in central Europe linked to high levels of acid deposition led European Economic Commission to adopt the “Convention on Long-Range Transboundary Air Pollution” to mitigate the impact of air pollution on human health and natural ecosystems through research efforts and the adoption of air pollution abatement policies. At the same time, Critical Loads were derived as indicators of excessive nitrogen and sulphur atmospheric deposition.

Some studies have suggested that in Europe most of the increase in forest growth can be accounted for N deposition (Sutton et al., 2008) and very little by elevated CO₂ concentrations, but this does not seem to apply in all regions. Rehfuss et al. (1999) reported that the combination of CO₂ rise and elevated N deposition accounted for a 15-20% increase in forest net primary productivity. However, the N deposition effect on growth is expected to saturate or even decline in ecosystems with high N inputs (Brumme and Khanna, 2008). Nitrogen emissions and deposition of nitrogen compounds have decreased since 1990 but relatively little compared to sulphur emissions. Agriculture and transport are the main sources of nitrogen pollution (EEA, 2007). In addition, nitrogen components can lead to eutrophication of ecosystems. When these pollutants exceed certain levels (‘critical load’), there is a negative effect on biodiversity. A critical load is defined as “a quantitative estimate of an exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge” (Nilsson and Grennfelt, 1988). Exceedances of critical loads by current or future nitrogen loads indicate risks for adverse effects on biodiversity. The currently used methodologies to derived critical loads are all described in Mapping Manual (www.icpmapping.org).

Furthermore, excess nitrogen is one of the major threats to biodiversity. Excessive levels of reactive forms of nitrogen in the biosphere and atmosphere constitute a major threat to biodiversity in

terrestrial, aquatic and coastal ecosystems. On land, it causes loss of sensitive species and hence biodiversity by favouring a few nitrogen tolerant species over less tolerant ones.

6.2 Measurements of climate variable for modelling simulation

The monitoring activities of meteorological parameters carried out within the national program for integrated control of forest ecosystems (CONECOFOR) represent the Italian application of ICP-FOREST monitoring activities. Meteorological variables have been collected by CREA (Council for Agricultural Research and Economics) in collaboration with the State Forestry Corps through ground monitoring systems of a wide network of automatic stations (in function since 1997 in numerous forest areas of the monitoring network).

The data collection, integrated with other variables surveyed in the CONECOFOR Programme, support integrated studies on forest ecology and plant physiology and, more generally, contribute to define the health status of the Italian forests.

6.2.1 Activities description

Permanent monitoring areas for acquisition of meteorological data, inside the forest (“in the plot”) and in “open field” within the radius of 2 km from the first, have been established in accordance with EU Regulations n. 1091/94 and n. 690/95.

This monitoring activity started in 1997 and involved initially 8 permanent areas, for a total of 13 stations. The stages of the implementation of the survey included (i) automatic stations purchase and installation, (ii) periodical collection of data related to the selected variables, as well as (iii) data integration, control, validation and processing. Currently, the geographical structure of the monitoring network has undergone numerous improvements, totalling 26 working stations, of which 23 “open field” and 16 “in the plot”: the current distribution of the stations is shown in figure 6.1.



Figure 6.1 – Location of the meteorological network on the Italian territory.

The stations are installed and managed directly by CREA-RPS. The data analysis is performed in the laboratory of biometeorology at CREA-under the supervision of Dr Silvano Fares and Dr Luca Salvati. Some technical solutions for reducing sampling and data elaboration costs (such as the

upgrade with remote controls, see figure 6.2) are being tested in a small number of forest plots. Until now, 4 areas (ABR1, VEN1, LAZ1 and EMI1) have been implemented with ftp data transmission (datalogger Campbell CR-1000 and modem GPRS Telit GT863-PY).

The main variables registered with a hourly time resolution are:

Open field (Datalogger CR10x, CR1000, CR200x)

1. Wind speed (10 and 2 m) and direction (10 m);
2. Solar radiation (2 m);
3. Temperature and Relative humidity (air 10, 2 and 0.1 m);
4. Soil Temperature (20 cm);
5. Precipitation;
6. Snow depth;
7. Soil volumetric water content and Temperature profiles (10, 30 and 60 cm).

In the plot (Datalogger CR10x, CR1000)

1. Soil volumetric water content and Temperature profiles (10, 30 and 60 cm);
2. Temperature and Relative humidity (air 2 and 0.1 m);
3. Soil Temperature (20 cm);
4. Precipitation;
5. Snow depth;
6. Soil moisture (10, 25 and 60 cm mod. Delta T).




Figure 6.2 – “Open field” and “in the plot” automatic meteorological stations upgraded.

The received meteorological data are processed into a database in two steps: first, descriptive metadata of the permanent site (geographic and physical characteristics), metadata of the stations (acquisition mode, instrumentation installed), metadata of the sensors (specifications), and metadata of the measured parameters and of their processing are registered.

Successively, meteorological data from each weather station are logged and undergo a series of controls and automatic processing for data quality.

The information on the “control” field allow to determine both the type of sensor reporting abnormal data, and the impact on measurement of the error duration. These metadata are useful to assess the completeness of meteorological data in terms of percentage of working time.

Annual activities of the equipment ordinary maintenance are provided in order to ensure the normal operation of the acquisition system, as well as extraordinary interventions for the resolution of specific technical problems.

Finally, the collected data are validated and made available on the website of the ICP forest.

The meteorological data of the 13 selected monitoring sites with the longest time series are reported in table 6.1 as a set of descriptive statistics calculated for five main meteorological variables.

Figures 6.3 and 6.4 show precipitation and temperature trends. The data time lapse is 1st January 1998 – 31st December 2013, with 1-day sampling interval.




Figure 6.3 – Time trend of average rainfall for the 13 main sites with the longest time series.




Figure 6.4 – Time trend of average temperature at 2 m for the 13 main sites with the longest time series.

Table 6.1 – Summary statistics, including the mean and median, standard deviation and coefficient of variation values, calculated for each variable, relative to 13 selected sites during the entire analyzed time period (1998-2013).

Note that the variation coefficient (CV) is a dimensionless quantity.

Stations	Statistics	AT01 (°C)	AT2 (°C)	RH01 (%)	RH2 (%)	PR (mm)
01-ABR1	Mean	6.63	7.21	79.62	83.12	2.06
	Median	7.1	7.6	82.1	86.4	0
	Std Dev	6.66	6.74	16.49	15.87	6.61
	CV	100.5	93.48	20.71	19.1	320.65
03-CAL1	Mean	10.83	11	89.17	86.4	3.91
	Median	10.8	11	95.5	93	0
	Std Dev	6.23	6.24	13.87	15.64	10.22
	CV	57.53	56.71	15.56	18.1	261.12
05-EMI1	Mean	12.22	12.02	81.16	80.64	1.92
	Median	12.5	12.3	84.75	85.2	0
	Std Dev	8.17	8.06	17.28	18.37	6.1
	CV	66.89	67.03	21.29	22.78	318.06
09-LAZ1	Mean	11.25	11	79.63	75.62	2.51
	Median	11.1	10.6	82	78.2	0
	Std Dev	7.3	6.92	15.18	17.51	6.99
	CV	64.86	62.93	19.06	23.16	279.01
12-PIE1	Mean	6.09	7.31	85.94	79.27	4.54
	Median	6.1	7.3	93.9	84.9	0
	Std Dev	6.96	6.82	17.23	20.1	15.52
	CV	114.35	93.19	20.05	25.35	341.81
20-VEN1	Media	4.5	5.72	91.34	85.75	4.72
	Median	4.8	6	93.4	90.1	0
	Dev std	7.28	7.2	8.98	14.2	13.85
	CV	161.67	125.89	9.83	16.56	293.21
06-EMI2	Mean	9.8	9.54	82.07	80.06	3.64
	Median	10	9.3	85.9	84.2	0
	Std Dev	7	7.03	16.36	17.36	9.95
	CV	71.45	73.67	19.94	21.69	273.66
08-FRI2	Mean	6.16	5.91	92.63	91.36	3.62
	Median	6.6	6.3	96.3	95.5	0
	Std Dev	7.56	7.64	8.79	10.89	11.05
	CV	122.7	129.15	9.49	11.92	305.05
10-LOM1	Mean	-	7.55	-	71.1	2.96
	Median	-	8.2	-	73	0
	Std Dev	-	6.71	-	17.34	8.13
	CV	-	88.82	-	24.38	274.13
13-PUG1	Mean	-	11.91	-	78.7	2.09
	Median	-	11.7	-	83	0
	Std Dev	-	6.87	-	13.71	6.99
	CV	-	57.71	-	17.41	334.7
17-TRE1	Mean	1.8	2.21	94.8	86.39	2.19
	Median	1.1	2	99.7	91.2	0
	Std Dev	7.24	7.29	9	14.44	5.89
	CV	401.24	330.18	9.49	16.71	268.88
22-LAZ2	Mean	-	15.72	-	77.42	1.3
	Median	-	15.3	-	79	0
	Std Dev	-	5.96	-	12.34	4.43
	CV	-	37.9	-	15.94	340.25
27-BOL1	Mean	-	4.45	-	72.75	3.01
	Median	-	4.7	-	75	0
	Std Dev	-	6.91	-	18.04	8.91
	CV	-	155.35	-	24.79	295.97

6.3 Impacts

Adverse effects of excessive deposition, like vegetation changes or forest dieback, does therefore not necessarily lead to immediate damages to ecosystems. To reconstruct and/or predict the temporal development of a soil and vegetation system, dynamic models, that include relevant time-dependent processes, are required. In the last three decades, different dynamic models have been developed, such as FORSAFE (Belyazid et al., 2006), MAGIC (Cosby et al., 1985, 2001), SMART/SMARTml (Bonten et al., 2011) and VSD/VSD+ (Posch and Reinds 2009; Reinds et al., 2001). The last one is used at site level in Italy.

More precisely, a suite of four models has been set to estimate the time required for a new (steady) state to be achieved when chemical and biological parameters response to a change in deposition. Figure 6.5 summarizes the suite of models used to estimate critical loads and biodiversity indices per single site.




Figure 6.5 – Schematic representation of two methods to estimate Critical load: steady-state and dynamic. The second approach is well represented by the combination of 4 models, able to estimate biodiversity indices.

The **MetHyd model** is a meteo-hydrological pre-processor. It is working elaborating Temperature, Photosynthetic Active Radiation and Precipitation surplus to estimate reduction factor of nitrification rates and of mineralization rates.

The **GrowthUp model** is a pre-processor able to compute N and base cation uptake from user specified tree-growth inputs.

The **Very Simple Dynamic (VSD) model** is the (minimal) extension of the Simple Mass Balance (SMB) steady-state model into a dynamic soil (acidification) model. The **VSD+ model** is an extension of the VSD model with detailed C and N dynamics. The model requires only a minimum set of inputs (compared to more detailed models) and execution time is minimised by reducing the set of model equations to a single non-linear equation. To facilitate the exploration of model behaviour at individual sites, the model is linked to a graphical user interface (GUI). This GUI allows easy (Bayesian) calibration, forward simulation (scenario analyses) and can also be used to compute target loads and delayed times between deposition reductions and ecosystem recovery.

PROPS module is the last step of the suite devoted to compute each potential species occurrence probability. The potential biotic is defined using EUNIS class corresponding to Corine land cover class.

6.3.1 Biodiversity analysis with dynamic models.

The described dynamic modelling suite has been applied in 5 forest Italian sites. The application of the modelling suite to one Italian site is described in Figure 6.6. The site chosen in this figure is IT10/LOM1 (Val Masino) from the ICP-Integrated Monitoring and/or ICP-Forests networks respectively. This site is characterized by high naturalistic value (protected by Habitat and Bird Directives), high sensitivity (low critical loads) and high pollutant exposure (critical loads were exceeded in the year 2000). The Val Masino site is a secondary *Picea abies* dominated forest with *Abies alba* and *Vaccinium myrtillus*. It belongs to CONECOFOR Programme since 1995 and is included in the ICP Forest European network. This site is in the Central Alps at 900-1190 m a.s.l., and its forest type is classified as EUNIS class G4.6. The VSD+ suite model was applied to simulate soil solution chemistry behaviour in response to net input from atmosphere, net element uptake by vegetation, net nitrogen immobilisation and element weathering.




Figure 6.6 – VSD+ run results. Parameter in soil solution and atmospheric deposition are shown.

The outputs obtained from VSD+ were used as inputs for running the PROPS vegetation model. Vegetation types (32 typical species) have been hypothesized from the EUNIS class G4.6. PROPS module estimated the Habitat Suitability Index (HSI), that is defined as the arithmetic mean of the normalized occurrence probabilities of the typical ecosystem species (Figure 6.7A). The PROPS model also computed isolines of normalized occurrence probability as a function of pH and N concentration (Figure 6.7B).

The HSI trend modelled by PROPS in 5 Italian forest sites, characterised by different climatic and environmental conditions is shown in Fig 6.8. The analysis shows an increasing trend for all the sites, even if with different slope, demonstrating a biodiversity recovery after the year 2000, when a decrease in Nitrogen deposition is found all over Italy. The higher rate of increase for the HSI is in the site called IT05, that is in Central Italy, where the dominant tree species is European beech.




Figure 6.7 – Site IT10 VSD+ results. Habitat Suitability Index during time (A) and occurrence probability isolines as pH and N concentration dependent (B).




Figure 6.8 – HSI in five Italian sites from year 1880 to 2100.

In the same way, it is possible to derive N and S range deposition to allow 80% of typical plant species for a specific ecosystem to survive. Those values are N and S biodiversity-based critical loads.

Figure 6.9 shows how derive N and S critical values to ensure the surviving of 80% typical species in three example sites.




Figure 6.9 – Habitat suitability Index as N and S concentration dependent in three different Italian forest sites. Biodiversity CL (CLNmin, CLSmax, CLNmax, CLSmin) indicates the range of values for S and N indispensable for the maintenance of at least 80% of the typical species for each biocenosis.

6.3.2 Biodiversity analysis with statistical models.

The impacts of air pollution on biodiversity are affected by the presence of other factors which themselves have an influence on ecosystems, such as meteorological conditions, forest management practices, soil types, and introduced plant diseases. Sorting out the most important factors is often difficult and sometimes impossible. Some of these factors are interrelated; trees may for instance be so weakened by air pollution as to be especially susceptible to disease. High altitude environments will be among the first to show the effects of acidification.

Although pollution often decreases with altitude, deposition can remain high because precipitation increase. Severe climatic conditions also make plants unable to absorb additional atmospheric nitrogen, which instead leaks by run-off.

We have chosen three differently test sites belonging to the Italian forestry monitoring network (see table 6.2 show sites information):

Table 6.2 – Test sites main information.

Site	IT05/ABR1	IT09/LAZ1	IT10/LOM1
Name	Selvapiana	Monte Rufeno	Val Masino
Latitude	41.8475	42.8306	46.2378
Longitude	13.5975	11.9139	93.5211
Altitude (m slm)	1500	690	1190
No. Of species	24	60	60
Forest type	beech forest	oak forest	spruce (and fir) forest
Age	123	48	93
EMEP50 cod	84036	80035	71039
Protection	Birds and Habitat directives applied	0	Birds and Habitat directives applied
EUNIS	G1.6	G1.7	G4.6

The Pielou's evenness index

Because the values of the diversity indices are not always comparable among them and depending on the extent to which they can actually vary, we used the evenness as a measure of diversity normalized on a fixed scale (e.g. from 0 to 1), allowing to carry out these comparisons among the three test sites.

$$H' = - \sum p_i \ln(p_i)$$

where H' is the Shannon-Weaver Diversity Index (Shannon and Weaver 1949), p_i is the relative abundance of each group of organisms.

$$H'_{max} = \ln S$$

where S is the species number. The Pielou's evenness (Pielou, 1966) is derived from the Shannon-Weaver index as follows:

$$J = \frac{H'}{H'_{max}}$$

J is constrained between 0 and 1; the less variation in communities among species implies higher values of J .

In our study, in order to measure the overall plant species diversity of a given transect, a modified version of the Shannon index (H'), named $H_{dunestd}$ (Grunewald & Schubert 2007), was used:

$$H_{dunestd} = - \frac{1}{\ln(k)} \sum p_i \times \ln(p_i)$$

where p_i = % cover of the i^{th} species and k is number of sampled species.

$H_{dunestd}$ has proved to be more useful than the Shannon diversity index in limiting habitats such as the coastal dunes (De Luca et al., 2011), where natural stressful conditions determine the presence of a few species with high dominance (Martinez et al., 2004).

$H_{dunestd}$, in fact, uses the abundance of species (as cover percentage) in relation to a constant sampling area, and hence, unlike H , is able to detect changes both in species diversity and total cover.

Characterisation of the ecological niche for plant species: the Ellenberg's indicators.

Ellenberg defined a set of indicator values for the vascular plants of central Europe (Ellenberg 1979, 1988; Ellenberg et al., 1991). The latest edition of Ellenberg's indicator values applies a 9-point scale for each of six gradients: soil acidity, soil productivity or fertility, soil humidity, soil salinity, climatic continentality and light availability. These have been widely used, both in central Europe and in adjacent parts of western Europe. The basis of indicator values is the realised ecological niche. Plants have a certain range of tolerance of temperature, light, soil pH, and so on. If we wish to make inferences about the ecological conditions pertaining at a site, much useful information can be obtained from the flora. These values are not i.e. mean pH values, but are on an arbitrary scale reflecting soil pH though not directly based on measurements. However, an advantage of indicator values is that they may be more sensitive to the requirements of plants than to a selected physical variable.




Figure 6.10 – Four Ellenberger's indices (Nutrient availability, pH, Continentality and Temperature) calculated for the three Italian study sites. They well define the sites' characteristics either soil properties or climate.

When the Ellenberg's indices were applied to the three test sites, they showed the ecological ranges that defined peculiar characteristics for each forest site (Figure 6.10). Significant differences of indices were observed among sites (ANOVA test, $p < 0.05$).

The application of two biodiversity indices J and $H_{dunestd}$ showed completely different trends among sites (Fig. 6.11). It is useful to remember that $H_{dunestd}$ is a modified Shannon-Weaver index that was focused on the cover percentage of each species (Attorre et al., 2013). The Pielou's evenness did not shown any difference among test sites (all sites have high equitability i.e. individuals are highly distributed among species), whereas $H_{dunestd}$ exhibited different biodiversity values, pointing out different patterns of plant species distribution related to the peculiar habitats of forest sites. In fact, a preliminary analysis of species richness at community level revealed that the lowest values occurred in beech forests (ABR1) and the highest in Turkey oak forests (LAZ1), whereas spruce forests (LOM1) were intermediate.

These features highlighted different vegetation dynamics characterised by fluctuations as the commonest on-going process (i.e. LOM1). The regeneration dynamic is also widespread due to the recent abandonment of wood exploitation and coppice management (i.e. ABR1) and low values of $H_{dunestd}$, whereas the regression dynamic was predominant in Turkey oak forests as in LAZ1 (and high values of $H_{dunestd}$).




Figure 6.11 – Biodiversity indices (*J* and *H_{dune}*, light and grey boxes, respectively) calculated for the three study sites. Both indices have been calculated by using the same phytosociological surveys, showing however, different values.

Correlation among variables (biodiversity indices, Ellenberg’s indices and exceedances above (BOF N) and under canopy (BSC N) (Both with trend in Figure 6.12) calculated for the three test sites did not highlight a clear aspect for the three test sites. Correlation analysis showed that diversity indices did not correlate with the Ellenberg’s indicators except *J* in the LAZ1 site (-0.72 Ellen. Temp; -0.88 Ellen. Cont.; 0.75 Ellen. Soil Moist.), but only *H_{dunestd}* correlated with BOF N in LOM1.




Figure 6.12 – Exceedances above (BOF N) and under canopy (BSC N) for the three study sites.

These simple analyses highlighted that different diversity indices (one based on the species proportion and the other based on proportional coverage area) could suggest different trends, which were not in correlation with climatic data and N depositions for three herbaceous communities growing under different ecological niches. It seems that one decade of data it is not sufficient for assessing a change in herbaceous community's composition. Inferences made on the air pollutant-induced effects in affecting plant community's composition should be carried out with extreme caution.

Our results are highlighting that the application of different biodiversity indices and ecological indicators should be applied in extensive way to other European herbaceous plant communities, to assess if air pollutants, and/or climate, and or anthropogenic activities are causal effects for a changing plant communities. Furthermore, it is important to have a long historical series of Ellenberg indices to assess the spatio-temporal dynamics for the European and Italian forest sites. Some trends are noticeable such as an increase of the Ellenberg Soil pH for ABR1 site or a reduction of Ellenberg Soil N for LOM1 site, although significances are questionable.

Cause-effect relationships between pollutants and primary production can be difficult to find in the field because a multitude of interactions concurrently could act on the response variable. For this reason, non-linear statistical techniques are becoming of increasing interest. The synergic or antagonistic roles of some limiting factors (such as high temperature, high ozone concentration) in affecting physiological processes (gas exchange, stomatal conductance) are very evident. Further, local variations of ox-N and red-N affected NPP in a complementary manner, acting as positive or negative drivers.

The impacts of air pollution and climate change on crown defoliation were different for each tree species, suggesting species-dependent effects on forests health and vitality. The vulnerability of forest tree species not only depends on exposure to climate change and air pollution but also on adaptive ability of the tree species (Lindner et al., 2010). Changes in climate will be associated to biotic (frequency and consequence of pest and disease outbreak) and abiotic disturbances (changes in fire occurrence and wind storm), causing strong implications for forest ecosystem (Lindner et al., 2010). In this frame nitrogen deposition, affects tree physiology, carbon allocation and plant interactions, resulting in complex relations with other environmental limiting factors such as drought (Matyssek et al., 2006).

6.4 Risk assessment

The risk assessment for critical loads for nutrient nitrogen in Italy for the year 2015 is shown in Figure 6.13.

The deposition over Italy are obtained by GAIN-Italy model with a grid resolution of 20 km and the exceedance map is obtained by difference between nitrogen deposition and critical loads exceedances. Exceedances are present almost at all the latitude in Italy, even if higher levels of exceedances are in the Alps in the Northern Region (> 10 kg/ha/y), due to higher emission levels of nitrogen compounds and consequently higher nitrogen deposition in this area.




Figure 6.13 – Nitrogen nutrient exceedance in Italy in the year 2015, based on GAINS-Italy model total nitrogen deposition exceedance.

6.5 Conclusions

To estimate the impacts of nitrogen pollution on biodiversity two kind of approaches are suitable, the first one based on dynamic modeling and the second one based on statistical models. Dynamic models estimate the biodiversity indices on a large time frame from 1980 till 2100, while the statistical ones analyse the non-linear relationship between environmental parameters and biodiversity. All the results show that there is a decreasing trend in nitrogen deposition, due to the policies developed to control nitrogen emissions. The decrease is coupled with a recovery trend in biodiversity indices. Despite such recovery trend in Italy, especially in Northern region, there are still nitrogen nutrient exceedance areas, where the ecosystem is exposed to the pressure of nitrogen pollution over the critical loads limits. More effort is needed to further reduce nitrogen pollution in Northern area, and this effort should be addressed to the agricultural sector that is the main constraint to completely recover biodiversity.

The contribution of Italy to the ICP WATERS Programme



National Focal point: Michela Rogora

Contributors: Michela Rogora¹, Aldo Marchetto¹, Rosario Mosello¹

¹CNR Institute of Ecosystem Study, Verbania Pallanza

CHAPTER 7 - THE CONTRIBUTION OF ITALY TO THE ICP WATERS PROGRAMME

7.1 Introduction

The international cooperative programme on assessment and monitoring of air pollution on rivers and lakes (ICP WATERS) was established under the Executive Body of the UNECE Convention on Long-range Transboundary Air Pollution (LRTAP) in July 1985. The ICP WATERS Programme Centre is hosted by the Norwegian Institute for Water Research (NIVA), while the Norwegian Climate and Pollution Agency leads the programme.

The main aim of the ICP Waters Programme is to assess, on a regional basis, the degree and geographical extent of the impact of atmospheric pollution, in particular acidification, on surface waters. More than 20 countries in Europe and North America participate in the programme and provide data on a regular basis.

ICP Waters is based on existing surface water monitoring programmes in the participating countries, implemented by voluntary contributions. The ICP site network is geographically extensive and includes long-term data series (more than 20 years) for many sites. At present, the network includes about 200 sites in Europe and North America (Fig. 7.1). The programme yearly conducts chemical and biological intercalibrations (e.g. Escudero, 2015; Fjellheim et al., 2015).




Figure 7.1 – The ICP WATERS sites in Europe (left) and the Italian network in the area of Lake Maggiore watershed, Northern Italy (right). Blue circles: lake sites; green circles: river sites; red triangles: atmospheric deposition sampling stations (including the Swiss network run by “Ufficio aria, clima e energie rinnovabili, SPAAS, DT Canton Ticino”). Site codes as in Tab. 7.1.

Since its establishment, ICP Waters has been an important contributor to document the effects of implementing the Protocols under the Convention. Numerous assessments, workshops, reports and publications covering the effects of long-range transported air pollution on surface water have been published over the years. Reports, scientific papers, Task Force proceedings and other publications are made available through the ICP WATERS website: <http://www.icp-waters.no>. Beside periodic reports describing the long-term chemical trends at the ICP WATERS sites, reports or papers on specific issues have also been produced: for instance, in depth analysis of biological data providing

evidence of recovery from acidification, or investigation on the so-called “confounding factors”, i.e. factors, other than deposition, which may affect long-term change in water chemistry and biology (Wright et al., 2005). As an example, several studies were devoted to the role of climate change on the reversibility of water acidification (e.g. Wright et al., 2006; Wright and Jenkins, 2001). A further confounding factor which has been addressed using ICP WATERS data was nitrate leaching, which proved to be important in delaying acidification recovery at sites subject to high N deposition (de Wit and Lindholm, 2010).

The Institute of Ecosystem Study of the National Research Council of Italy (CNR ISE; formerly “Istituto Italiano di Idrobiologia”) in Verbania Pallanza has been the National Focal Point for the ICP WATERS since 1995, under the direction and coordination of the Italian Ministry for the Environment, Land and Sea. The regular participation of Italy to the ICP WATERS has led to:

- the development of an Italian network of sites, consisting of three subalpine rivers, one subalpine lake and two high altitude alpine lakes in the area of Lake Maggiore watershed, Piedmont region (Fig. 7.1, Tab. 7.1), to follow the long-term evolution of acidification and recovery, and more generally the response of surface waters in sensitive areas to changing deposition;
- the harmonisation and standardisation of monitoring practices, both for sampling and analysis, following the provision of the Programme Centre (NIVA); this was also allowed by the regular participation of the CNR ISE hydrochemical laboratory to chemical intercomparisons;
- specific research performed on selected sites or areas, to better understand site-specific processes and dynamics at the catchment scale (e.g. nitrogen leaching and response to changing nitrogen deposition; acidification recovery under climate change scenarios).

Table 7.1 – Main characteristics of the Italian sites included in the ICP WATERS network.

Site name	Code	Altitude (m a.s.l.)	Catchment area (km ²)	Lake area (km ²)	Yearly average flow (m ³ s ⁻¹)	Data since	Data frequency
Lake Paione Inferiore	IT01	2002	1.26	0.0068		1984	1-2 per year
Lake Mergozzo	IT02	194	10.43	1.83		1978	2-4 per year
Lake Paione Superiore	IT03	2269	0.50	0.0086		1984	1-2 per year
River Cannobino	IT04	193	110.4	-	5.04	1978	Monthly
River Pellino	IT05	290	17.5	-	0.92	1984	Monthly
River Pellesino	IT06	290	3.4	-	0.19	1986	Monthly

The Italian network and the monitoring activities are described in details in Mosello et al. (2000). The CNR ISE has been regularly sending data to the Programme Centre since the establishment of the network. Collected data included base chemical variables (pH, alkalinity, conductivity), nutrients (phosphorus and nitrogen compounds), major ions (Ca, Mg, Na, K, SO₄, Cl) and selected trace metals (Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn). Besides monitoring surface water bodies, the CNR ISE also runs a network for the assessment of long-term change in atmospheric deposition (Rogora et al., 2016). Analytical methods and QA/QC procedure adopted in the laboratory are described in details at the website <http://www.idrolab.ise.cnr.it>

Results have been presented at national and international conferences and published both in ICP WATERS reports and in scientific papers (e.g. Stoddard et al., 1999; Skjelkvåle et al., 2005; Garmo et al., 2014). In particular, a fruitful cooperation exists with the Swiss colleagues of the “Ufficio aria, clima e energie rinnovabili, SPAAS, DT Canton Ticino” and of the University of Applied Sciences and Arts of Southern Switzerland, who act as ICP WATERS National Focal Point for Switzerland (Steingruber 2015).

7.2 Atmospheric pollution pressures on surface waters

Within the ICP Waters aims the main pressures identified as important at the Italian sites of the network are acidification and nitrogen deposition. Furthermore, especially in recent times, attention has been paid to the effects of climate change, in interaction with the other drivers.

Acidification of surface waters due to rain acidity has been a problem in the 1970s and 1980s for some high altitude lakes in the Central Alps, characterised by a low alkalinity pool and a limited buffering capacity due to the geological composition of the catchments (Marchetto et al., 1994). Beside the two lakes included in the ICP WATERS network (Lake Paione Superiore and Inferiore; Tab. 7.1), the CNR ISE has been regularly monitoring from the chemical point of view about 30 high altitude lakes since the early 1980s. Most of these lakes, which underwent acidification in the 1980s, partially recovered from the mid-1990s as a response to the decreasing deposition of acidifying compounds, showing an increase of alkalinity and pH (mainly as sulphate (SO₄)) (Rogora et al., 2001; 2013). However, few sites remain acidic or still show a high sensitivity to acidification. This is particularly evident at the snowmelt, when alkalinity may be fully depleted by the incoming waters rich in acidifying compounds. At present, nitrate is the dominant acidifying agent in the high altitude lakes, due to the high input of nitrogen compounds from atmospheric deposition (Rogora et al., 2013).

One of the main risk related to water acidification is the dissolution of **trace metals**, which may be harmful for the aquatic biota (especially Al, Cd, Pb, Cu, Ni). For this reason, in addition to standard chemical variables also trace metals have been regularly analysed at the ICP WATERS sites in Italy. Levels proved to be low at all the sites, including high altitude lakes (Tornimbeni and Rogora, 2012). Within the ICP WATERS network, Italian sites proved to be among the most affected by nitrogen (N) inputs from the atmosphere. They were indeed threatened by **N enrichment** due to the N saturation of soils in the catchment and following nitrate (NO₃) leaching to surface waters (Rogora and Mosello, 2007; Rogora, 2007). This is mainly due to the high levels of N deposition which affected the study sites, located north of the Po Plain, one of the most densely inhabited and most industrialised and urbanised areas of Europe. Depositions in the alpine and subalpine areas are particularly high as a combined effect of high pollutant concentrations and high precipitation amounts for orographic effects (Rogora et al., 2006).

Long-term studies at the ICP WATERS sites in Italy also revealed the important role that **climate** drivers may have on the response of surface waters to changing deposition (Rogora and Mosello, 2007; Rogora et al., 2003a). Climatic factors interact with atmospheric deposition affecting the long-term changes in lake water chemistry and biology, with an overall effect which may both favour or contrast recovery patterns according to the specific processes at each site. Studies at the Italian sites also demonstrated that temporal variations of N compounds in surface water may be affected by climatic factors: both increasing temperature and change in precipitation regime proved to be important in the NO₃ long- and short- term dynamics in rivers and lakes (Rogora, 2007;

Rogora et al., 2013). The assessment of the atmospheric inputs of acidity and S and N compounds in the area of Lake Maggiore watershed have been performed since the beginning of the monitoring through a cooperation between Italy and Switzerland. This area is indeed shared almost equally between the two countries, and ICP WATERS sites, both Italian and Swiss sites, are located here (Fig. 7.1). An extensive network for the study of atmospheric deposition chemistry exists in this area, consisting of 14 stations in total (Rogora et al., 2016). In Italy, emissions of sulphur dioxide (SO_2) reached its maximum between 1965 and 1980, while nitrogen oxides (NO_x) peaked around 1985. Successively, emissions decreased by 90%, 58% and 14%, respectively for SO_2 , NO_x and NH_3 compared to the values in 1990 (Romano et al., 2014). This resulted in a sharp decrease in the deposition of sulphate, acidity and, at a lesser extent, of N compounds.

The long-term trends of reduced (NH_4) and oxidised (NO_3) N deposition at sites in the Italian part of Lake Maggiore showed a slight tendency to decrease starting in 2006 (Fig. 7.2). The decrease of nitrogen loads was widespread and affected both the northern alpine stations and the southern ones (Rogora et al., 2016). The decrease was more evident for oxidized N than for reduced N. Therefore, deposition of ammonium acquired an increasing importance in time, especially at the southern, more polluted sites: the relative contribution of reduced N to wet N deposition passed from about 50% in the early 1990s to 56-57% in recent years (Rogora et al., 2016).




Figure 7.2 – Trends of NO_3 and NH_4 annual deposition at the atmospheric deposition sampling sites in Lake Maggiore watershed, North-Western Italy. For site location, see Fig. 7.1.

To assess both temporal change and spatial gradients in the **atmospheric deposition of S and N compounds** in the area of Lake Maggiore, maps of average deposition for 5-year periods were produced (Fig. 7.3). A spatial gradient in the deposition of sulphate and nitrogen compounds was evident both in the 1990s and in recent times (2008-2012), with highest values in the south-eastern part of the area, close to the major emission sources, and decreasing values towards the Alps. However, gradients became less and less evident in time: because of the declining concentrations of acidifying compounds observed in the last two decades, which affected in particular the more polluted southern sites. Despite these recent changes, deposition of inorganic N in North Western Italy, as the sum of ammonium and nitrate, is still between 110 and 140 meq m⁻² y⁻¹ (15-20 kg N ha⁻¹ y⁻¹) and it has been estimated that total N deposition, including the contribution of dry and organic N deposition, may be 30-40% higher (Rogora et al., 2016).




Figure 7.3 – Maps of SO₄ and NO₃ deposition over the area of Lake Maggiore watershed in the 1990s compared with the recent period (2008-12). For the location of the sites see Fig. 7.1.

7.3 Main impacts of atmospheric pollution on rivers and lakes

The effects of the main pressures on water quality and biota have been assessed at the Italian ICP WATERS sites both through the analysis of long-term data and in-depth studies at specific sites. In the first period since the establishment of the network (1995-2000), the focus has been on acidification and related problems (pH and alkalinity drop, heavy metals dissolution; Mosello et al., 2000; Rogora et al., 2001). Extensive studies were performed by the Programme centre, in cooperation with the National Focal Points, to assess the regional extension of the acidification problem and its relevance at European scale (Stoddard et al., 1999). In Italy, major effects of acid rain were detected at a limited number of sites in the Central Alps, showing significant pH decrease (below 5.6), full depletion of the alkalinity pool and increase of aluminium concentration. Although only few lakes were effectively acidified, most of the lakes in this part of the Alps proved to be sensitive to acidic inputs and potentially threatened, if deposition had remained at the same level (Marchetto et al., 1994). Effects of acidification were evident on both the lake flora and fauna: benthic diatoms assemblage was shifted towards acidophilus species, and zooplankton lost the dominant species, *Arctodiaptomus alpinus*. Palaeolimnological studies outlined that lake acidification paralleled the increasing input of long-range transported industrial pollutants, traced by spherical carbonaceous particles (Guilizzoni et al., 1996; Marchetto et al., 2004).

The Convention on Long Range Transboundary Air Pollution (CLRTAP) entered into force in 1983 and was subsequently extended by eight specific protocols. The Gothenburg Protocol, for instance, for the abatement of acidification, eutrophication and ground-level ozone, was adopted in 1999 and sets emission ceilings for 2010. As an effect of emission reduction, in particular of SO₂, rain acidity substantially decreased and a chemical recovery of alpine lakes immediately started: for instance, the ICP WATERS site Lakes Paione Inferiore and Superiore showed positive trends of pH and alkalinity since the mid-1990s (Fig. 7.2; Mosello et al., 2000; Marchetto et al., 2004). First signs of biological recovery were identified, such as change in diatom flora and appearance of sensitive species among benthic insects (Marchetto et al., 2004).

Despite this positive response to changing deposition, several critical issues still affect high altitude lakes and sensitive sites in general: a recent study performed on about 40 lakes in Italy (Piedmont) and Switzerland (Canton Ticino) showed that some lakes are still acidic or show a high sensitivity to acidification. This sensitivity is particularly evident at the snowmelt, when alkalinity is still fully depleted in some lakes (Rogora et al., 2013). The study also highlighted the prominent role of N deposition in this area: now, nitrate is the dominant acidifying agent in the studied lakes, due to the high input of nitrogen compounds from atmospheric deposition. As an example, Fig. 7.4 shows the SO₄ to NO₃ ratio in Lake Paione Superiore: as an effect of the decreasing concentration of SO₄ in lake water, NO₃ has become more and more important in time and presently contribute for 40-45% of the sum of acid anions in this lake. Wet deposition of inorganic N (sum of ammonium and nitrate) affecting the ICP WATERS sites has remained fairly constant over a 30-year period and close to 25 kg N ha⁻¹ y⁻¹. These levels of N deposition proved to be among the highest in Europe (Evans et al., 2001), because of the location downwind of the major emission sources (Rogora et al., 2006a). This huge flux of N from the atmosphere caused N saturation of terrestrial catchments and increasing levels of NO₃ in rivers and lakes (Rogora and Mosello 2007; Rogora, 2007). According to pan-European studies, among the few sites in Europe, the Italian sites located in the Lake Maggiore area were showing a significant increase of NO₃ concentrations in the 1980s and 1990s (Skjelkvåle et al., 2005).




Figure 7.4 - Trends of pH and alkalinity (left panel) and of the SO_4 to NO_3 ratio (right panel) in the ICP WATERS site Lake Paione Superiore (IT03).

Furthermore, detailed N budgets performed for the single river catchments showed that the atmospheric input of N cannot be fully retained by soil and vegetation, with N retention varying from 60-70% of the input in the northern part of the area, to 20-30% or even lower in the southern catchments (Rogora et al., 2006b). In a previous analysis of long-term trends in stream water chemistry, the increasing NO_3 levels and the limited seasonal pattern of NO_3 were identified as signals of an aggrading level of N saturation in time; according to monthly NO_3 data, most of the river sites could be classed as being at a medium or high stage of N saturation (Rogora, 2007).

Recently, monitoring data for both rivers and lakes showed a reversal in NO_3 trends (Rogora et al., 2012), as a response to the decreasing N inputs from the atmosphere observed since 2005-2006. A study performed considering both the ICP WATERS sites and other monitoring sites in the same area, showed how this change was widespread, affecting both high-altitude lakes in the Alps and subalpine lakes and rivers, and occurred at almost the same time at all sites (Fig. 7.5). High altitude lakes show a wide range of concentrations; however, values in the 1980s reached 30-40 $\mu\text{eq L}^{-1}$ in some lakes, while they are mostly below 20 $\mu\text{eq L}^{-1}$ in recent years (Fig. 7.5, right panel). The data for sites subject to a continuous monitoring highlight a coherent pattern, with a dominant positive trend till 2000-2005, followed by a decreasing one (Fig. 7.5, left panel).

The decrease in most recent period was particularly evident for stream sites: for instance, at the ICP WATERS sites Rivers Pellino and Pellesino NO_3 decreased from 120-140 $\mu\text{eq L}^{-1}$ around 2000 to the present values of about 90 $\mu\text{eq L}^{-1}$ (Fig. 7.3)

Among ICP WATERS lake sites, the watershed of Lake Mergozzo, located in the subalpine part of the area (Fig. 7.1), was probably affected by a state of N saturation: NO_3 sharply increased in the 1980s and 1990s (Fig. 7.5), then stabilized at around 50 $\mu\text{eq L}^{-1}$ since 2000. The N enrichment of the lake water was ascribed entirely to atmospheric inputs (Rogora et al., 2012). Due to their remote location, lakes Paione Inferiore and Superiore are subject to lower levels of N deposition, and are characterised by distinctly lower NO_3 concentrations compared to the other sites (Fig. 7.5). Nevertheless, the mean level of NO_3 in the water of these lakes (20-25 $\mu\text{eq L}^{-1}$) is higher than is recorded at other mountain sites in remote areas of the world (Rogora et al., 2008a), so that the Paione lakes and the other mountain lakes in the same area can be considered as possibly impacted by atmospheric N transported from downwind emission sources (Rogora et al., 2012).




Figure 7.5 – Long -term trends of yearly NO_3 concentrations (top) at the surface water monitoring sites in the area of Lake Maggiore watershed, including the ICP WATERS sites, and at high altitude lakes in the Central Alps ($n=30$) measured in different periods (bottom). The thick line shows the median value for each survey.

Mer: Lake Mergozzo (IT02); LPS: Lake Paione Superiore (IT03); LPI: Lake Paione Inferiore (IT01); Can: River Cannobino (IT04); Pel: River Pellino (IT05); Pls: River Pellesino (IT06); Sbe: River S. Bernardino; Sgi: River S. Giovanni; Mag: River Maggia; Gio: River Giona.

Besides the similarity of trends, a further feature common to most of the data series was the presence of a high interannual variability in the years between 1999 and 2004; 2002 and 2003 in particular were characterized by peak values of NO₃ at several sites (Fig. 7.5). To better investigate the possible drivers of this short-term variability in water chemistry, climate factors were considered: in particular, the role of temperature and precipitation regime and trends in changing NO₃ concentrations in surface waters was assessed (Rogora, 2007; Rogora and Mosello, 2007). Temperature, especially extreme values, proved to be important in NO₃ export from river catchments: high air temperature and prolonged dry conditions, such as those of the summer of 2003, affected NO₃ dynamics in some of the subalpine rivers in the Lake Maggiore area (Rogora, 2007). These results were explained by the effect of warm periods on temperature-dependent processes such as mineralization and nitrification (Rogora et al., 2008b).

Also, the analysis of other chemical trends (e.g. sulphate, base cations) demonstrated that climatic factors interact with atmospheric deposition affecting the long-term changes in lake water. Some high altitude lakes, for instance, showed an increasing trend of sulphate concentrations, despite the huge decrease of sulphate input from the atmosphere (Rogora et al., 2013). Besides sulphate, also base cations increased in these lakes, with an overall effect on conductivity. An analysis of climate data for the study area puts in evidence how the last two decades (1990-2010) have been characterised by the highest temperature rise and this change mainly affected spring and summer months. Data collected also confirmed a decrease in snowfall as well in the length of the snow cover period (Meteo Svizzera, 2012; Rogora et al., 2003a). A climate related effect can be suggested to explain trends in water chemistry contrasting with those in atmospheric deposition. First of all the presence of less and less snow on the ground and a more exposed portion of the catchments lead to increasing export of weathering products to lake water (Rogora et al., 2003a). In addition, as suggested by some recent studies, the cryosphere, and particularly permafrost degradation processes, may drive lake chemical changes (Thies et al., 2007). For this reason, besides the discrete monitoring of lake water, high frequency measurements of some basic parameters (lake water temperature, conductivity) have been started in the last few years at selected study sites.

7.4 Deposition scenarios and risk assessment

To assess the impacts of different deposition scenarios on lake and river water chemistry, dynamic modelling has been performed at Italian ICP WATERS sites (Rogora, 2004; Rogora et al., 2003b). Some cooperative studies were also performed at a European level to assess and predict the extent of acidification and recovery, also in relation to confounding factors (Wright et al., 2006; Helliwell et al., 2014). The model used, named MAGIC (Model of Acidification of Groundwater In Catchments), is a process-oriented, intermediate-complexity dynamic model which has been in use for more than 20 years and extensively tested on catchments in Europe and North America for the long term reconstruction and future prediction of soil and surface water acidification at the catchment scale (Cosby et al., 2001). The outputs clearly demonstrated the benefits of achieving the emission reductions in both S and N compounds agreed under the Gothenburg Protocol. It was also clear that, besides the substantial reduction of SO₄ deposition from the peak levels of the 80s, N deposition too had to be reduced to protect freshwaters from further acidification (Rogora et al., 2003b). The results also showed how including other factors specific to the Mediterranean area,

such as dust deposition and climate change, may improve the fit of experimental data and the reliability of the model forecast.

For this reason, a further modelling exercise was performed, in cooperation with researchers from several European countries, adopting a common protocol at 14 intensively-studied sites in Europe and Eastern North America, including Italian sites (Wright et al., 2006). The results suggested that modelling of recovery from acidification should take into account possible concurrent climate changes and, especially in the case of the Italian sites, on the climate-induced changes in nitrogen retention and solute export from the catchments.

A further assessment of the efficacy of deposition reduction on water quality at sensitive sites was performed within the ICP WATERS in the so-called “ex-post analysis” (Wright et al., 2011). The already available MAGIC calibration was used to simulate the evolution of lake chemistry at the site Lake Paione Superiore in response to the deposition scenarios issued by the Coordination Centre for Effects (CCE) in 2010. In particular, projections were made under the scenario of national emission estimates (NAT) and emissions with maximum feasible reductions (MFR) (Fig. 7.6). Under both scenarios, a sharp increase of pH and of the acid neutralising capacity (ANC) of lake water was predicted and an evident improvement of lake water chemical status should be expected. This can be mainly ascribed to the decreasing concentrations of SO_4 , which will probably decrease further in the next few years and then stabilise (Fig. 7.6).




Figure 7.6 – Resulting projections for ANC (top) and SO_4 (bottom) in Lake Paione Superiore (ICP WATERS site IT03) using the ex-post scenarios provided by CCE. The observed values measured in lake water (autumn samplings) are also shown. NAT: national emission estimates (NAT); MFR: maximum feasible reductions.

However, reduction of N emission in source regions, located in the lowlands, will be crucial in the next few years for the recovery of these types of lakes from acidification and for their nitrogen status as a whole.

Alkalinity, or alternatively ANC (Acid Neutralising Capacity), has been widely used as an index of the acidification status, and of acid sensitivity of surface waters. In Europe, the value of $20 \mu\text{eq L}^{-1}$ has been identified as the minimum level required for ecosystem protection under UNECE protocols. However, proposed critical limits vary depending on the target group of organisms. Furthermore, even when considering the same target (e.g. macroinvertebrates), critical limits may vary, depending on the typical fauna of sensitive species and their adaptations to native water chemistry. In the high Alps for instance an ANC limit of $30 \mu\text{eq L}^{-1}$ has been suggested (Raddum and Skjelkvåle, 2001). Some of the monitored lakes in the Alps still show values of ANC below this limit, especially at snowmelt (Rogora et al., 2013). However, it must be also pointed out that some of these lakes are characterised by a very limited “natural” alkalinity pool, due to the lithological composition of their catchments. As a consequence, the alkalinity or ANC critical limits identified at European and international level will be hardly achieved in these sensitive lakes, even under the most optimistic deposition scenarios. Furthermore, when thinking to biological recovery, it should be considered that benthic species in high altitude lakes, which are frequently used as biological indicators, in the future will be probably more affected by changing physical conditions, due to the effects of meteorology and climate, than by the chemistry of lake waters.

In the framework of the UNECE CLRTAP, empirical critical loads of nitrogen (CLN) have been defined for natural and semi-natural ecosystems (Bobbink and Hettelingh, 2011). For instance, for permanent oligotrophic lakes, ponds and pools, a CLN of $3\text{-}10 \text{ kg N ha}^{-1} \text{ y}^{-1}$ has been identified. At present, although deposition of acidity at ICP WATERS sites in Italy mostly falls below the critical limits, the atmospheric deposition of nitrogen remains still too high with respect to critical levels and further reductions are needed to prevent nitrogen saturation of terrestrial and aquatic ecosystems.

7.5 Conclusive remarks

Overall the results of long-term studies at ICP WATERS sites in Italy emphasise the benefits of achieving emission reduction targets. Deposition clearly responded to emission decrease, which has been particularly evident for SO_2 . Surface water response to changing deposition was widespread but somewhat delayed, due to the interacting effect of several factors, such as N saturation of soils in the catchments and climate change.

Despite the current tendency toward recovery, atmospheric deposition and other global changes will probably keep affecting freshwater quality in the future, especially in the alpine area. The recovery patterns, both from acidification and from N saturation, will be more and more influenced by climatic factors, such as temperature and precipitation, also through indirect effects (snow cover change, retreating glaciers, permafrost degradation).

Nitrogen deposition, both as oxidised and reduced nitrogen, will continue to have a prominent role in the acidification processes and in the nitrogen status of surface water. From this perspective, further reductions in the emissions of N compounds should be the target of national and international policy.

Are technical materials and cultural heritage exposed to air pollution risk? The contribution of Italy to ICP Materials



Co-Chair ICP Materials and National Focal Point: Pasquale Spezzano

Chapter coordinator: Pasquale Spezzano

Contributors: Giovanni Vialetto¹, Pasquale Spezzano¹

¹ENEA, Laboratory of Atmospheric Pollution

CHAPTER 8 - ARE TECHNICAL MATERIALS AND CULTURAL HERITAGE EXPOSED TO AIR POLLUTION RISK? THE CONTRIBUTION OF ITALY TO ICP MATERIALS

8.1 Introduction

Air pollutants in combination with climatic parameters are key factors in the corrosion and deterioration of several metallic and non-metallic materials. This reduces the operating life of technical materials and threatens objects of cultural heritage, an important component of our individual and collective identity. The impact of pollutants emitted into the atmosphere on materials is cumulative and irreversible because, unlike natural ecosystems, materials have no possibility of self-regeneration. This impact causes massive economic losses for protective measures, substitution of degraded materials, cleaning, maintenance, and restoration work on buildings and historical and cultural monuments exposed outdoors.

The damage to the materials exposed to the atmosphere is the result of complex interactions between chemical, physical and biological parameters. The main forms of degradation caused by atmospheric pollutants on materials and in particular on built cultural heritage are corrosion (loss of material due to chemical attack) and soiling (due to the deposition of particulates on the surfaces). Since materials degradation also occurs in the absence of pollutants, it is important to quantify how much air pollution due to human activities influences and accelerates the background (largely unaffected by human activities) degradation of the materials.

Sulphur dioxide (SO_2) is the main pollutant responsible for the corrosion of materials, but also nitrogen oxides (NO_x), ozone (O_3), particulate matter, carbon dioxide (CO_2), and sea salt from sea spray play an important role. Wet and dry deposition both contribute to the deterioration of materials. Generally, sulphur and nitrogen oxides are oxidised in the atmosphere to sulphuric acid (H_2SO_4) and nitric acid (HNO_3) that exert their corrosive action on materials by lowering the pH of the precipitation (acid rain). SO_2 is also the main cause of the sulphation process of stone and bronze surfaces. The deposition of airborne particles on surfaces of buildings and historic monuments causes soiling and accelerates chemical degradation of the materials, thus impacting on both the aesthetic appeal and the decay of such structures. O_3 has an indirect role as it oxidise sulphur oxides and nitrogen oxides to H_2SO_4 and HNO_3 and exerts a direct role in the degradation of metals such as copper and copper alloys and in the oxidation of polymeric materials. Chloride content in sea salt is an effective corrosive agent.

CO_2 is generally not considered a pollutant itself, but a climate-altering gas. However, CO_2 from the air dissolves in rainwater making it slightly acidic (carbonic acid, H_2CO_3). The rainwater may react with materials that are largely made from calcium carbonates (CaCO_3) – limestone being a common example - transforming the calcium carbonate, slightly soluble, into the more soluble calcium bicarbonate ($\text{Ca}(\text{HCO}_3)_2$). This is washed away causing the material to be weathered (karst effect).

Calcareous stones such as limestone and marble, which are used in most heritage buildings, are the most vulnerable to degradation. Natural degradation factors such as changes in climate and micro-climate around the surfaces, freeze-thaw phenomena, salt crystallization, etc., add to human factors, mainly represented by air pollution, leading to several forms of deterioration, including the build-up of the so-called “black crusts”, heterogeneous deposits consisting of gypsum (calcium sulphate, CaSO_4), calcite resulting from dissolution and subsequent re-precipitation of calcium carbonate, carbonaceous particles and other components.

In addition, air pollutants deposited on materials could enrich it with nutrients, thus favouring the biological colonization (bacteria, fungi, algae, lichens and plants), a further damage factor of the surfaces of architectural works.

8.2 Effects on materials

The effects of air pollutants as well as climate parameters on the atmospheric corrosion and soiling of various materials, including materials used in objects of cultural heritage, are investigated by the International Co-operative Programme on Effects on Materials, including Historic and Cultural Monuments (ICP Materials <http://www.corr-institute.se/icp-materials>).

ICP Materials is lead by Sweden, which provides the programme with the Main Research Centre, Swerea KIMAB AB. Since 2005 the chairmanship is shared by Sweden and Italy (ENEA), which together are responsible for the co-ordination and organisation of the programme. A Task Force consisting of representatives from all countries participating in ICP Materials is responsible for the implementation of the programme.

Italy through ENEA is responsible for the sub-centre for stock of materials at risk and cultural heritage which includes use of results including mapping, stock at risk and economic evaluations aimed especially at objects of cultural heritage. Czech Republic, France, Spain, Sweden, Switzerland, and United Kingdom are providing the programme with materials sub-centres. Each sub-centre is responsible for a material or group of materials and prepare, distribute and evaluate corrosion effects after exposure on samples of materials, regardless of where they were exposed. Norway is providing the programme with the environmental sub-centre, which maintains the environmental database and evaluates trends of the environmental data.

The exposure of materials is performed in a network of test sites, located in countries that are Parties of the Convention, covering different climatic conditions and different levels of air pollution. The measurements include a wide range of pollutants, precipitation and climate parameters. A wide range of materials has been selected and exposed that are representative both for technical materials and materials used in objects of cultural heritage. Currently, the ICP Materials network consists of 26 monitoring and exposure stations located in 18 different countries (Figure 8.1).




Figure 8.1 – Map of test sites of the monitoring programme of ICP Materials (left) and test site in Venice with material samples on the rack (right).

Italy participates in the programme since its inception with four sites: Rome (Istituto Superiore di Sanità), Rome-Casaccia (ENEA), Milan (ARPA Lombardia), and Venice (ARPA Veneto).

Figure 8.2 shows the change in the yearly corrosion rates of carbon steel, weathering steel, traditional zinc, blasted zinc, copper, and limestone and the change in soiling of modern glass samples observed at the four Italian stations. Overall, the effects observed in Italy resemble the trend observed by the ICP Materials network (Tidblad et al, 2014; Tidblad et al., 2016) and reflect the decline in emissions of SO₂ and NO_x. Although there are considerable fluctuations between the measurement campaigns, also attributable to fluctuations in the climatic parameters, corrosion rates of carbon steel declined considerably over the investigated period. Based on only two exposures (1987 and 2011) also the corrosion rate of weathering steel (a low-alloyed steel with increased resistance to atmospheric corrosion) shows similar results after one year of exposure. Corrosion rate of traditionally grinded zinc also declined significantly until the mid-90s, but the trend has diminished in the following years (up to 2000). Glass blasted zinc, investigated from 1997 onwards, shows no clear downward trend (the corrosion of blasted zinc is generally higher because of the rougher surface given by this treatment). Copper samples were exposed during five exposure periods. In general, a significant decrease in the corrosion rate between 1987 and 1997 can be observed while in the following years the improvement is less evident. For limestone, declining trends could be observed initially, particularly in urban areas where the reduction of SO₂ emissions was greater, after that no recognisable downward trend could be observed. Evaluation of soiling of modern glass has been investigated by ICP Materials since 2005. In general, and in analogy with all the sites of the ICP Materials network, haze does not show a clear trend.

Although the degradation of materials in Europe is today significantly reduced, mainly due to reduction of sulphur pollution, the current rates of corrosion and soiling of materials are overall still unacceptably high. Air pollution is still a problem, but more complicated by the superimposition of the effects of a multitude of harmful air pollutants with those of climate parameters such as temperature, relative humidity and amount of rainfall. Effective strategies for the reduction of the effects of air pollutants on materials need to go hand in hand with strategies to cope with climate change.

The research programmes undertaken by ICP Materials lead to the derivation of statistically reliable dose-response functions linking the corrosion or deterioration rate of several materials to the levels of pollutants in combination with climatic parameters. Two sets of dose-response functions have been derived: functions for the SO₂ dominating scenario (Tidblad et al., 1998) and functions for the multi-pollutant scenario (UNECE, 2015). These dose-response functions can be used to calculate the degree of material corrosion or soiling starting from ambient data, and then identify those areas where the risk of damage is greater.

Threshold values of degradation rates, below which impacts are “acceptable” (for materials used in technical constructions) or “tolerable” (for materials used in objects of cultural heritage), have been defined, to which correspond acceptable/tolerable levels, i.e. those concentrations or loads of pollutants that do not lead to unacceptable/intolerable increase in the rate of corrosion or deterioration. Acceptable/tolerable corrosion rates are expressed as a multiple of the background corrosion rate. Such calculations can be used to produce maps showing increased risk of corrosion and soiling at many different scales, from a continent to individual cities.




Figure 8.2 – Change in the yearly corrosion rates of carbon steel, weathering steel, traditional zinc, blasted zinc, copper, and limestone and the change in soiling of modern glass samples observed at the four Italian stations.

8.3 Effects on historic and cultural monuments and buildings.

The ultimate goal of the ICP Materials program is the calculation of cost of damage caused by deterioration of materials and attributable to atmospheric pollution. It is possible to estimate the difference in cost between two alternative scenarios, one representing the current (or future) air pollution situation and another that quantifies the damage in a “background” scenario. The difference in the deterioration rate can be then used to estimate the cost associated with the air pollution. Performing inventories of stock of materials at risk is one important pieces of information that is required for a cost calculation.

The main results from the Italian sub-centre in this area are the case studies in Italy (Doytchinov et al., 2009; Doytchinov et al., 2010) and the participation in a review of available data on stock of materials at risk (Tidblad et al., 2010a) and economic evaluations (Tidblad et al., 2010b). The Italian sub-centre has also co-authored the chapter on stock at risk studies (Watt et al., 2009) in a book on effects of air pollution on cultural heritage, which was the final product of the European CULT-STRAT project and included significant parts from the ICP Materials programme.

In recent years, the Italian sub-centre has conducted a “Pilot study on inventory and conditions of stock of materials at risk at five UNESCO cultural heritage sites”. The study is presented in four individual ICP Materials reports during the period 2011–2015: Part I: Methodology (Doytchinov et al., 2011); Part II: Determination of stock of materials at risk for individual monuments (Doytchinov et al., 2012); Part III: Economic evaluation (Doytchinov et al., 2014); and Part IV: The relationship between the environment and the artefact (Spezzano et al., 2015).

Five important UNESCO World Cultural Heritage Site in Europe were studied: Greece, Athens, Acropolis, (The Parthenon); France, Paris, The Facades in the Centre of City; Czech Republic, Prague, The National Library; Germany, Berlin, The New Museum; and UK, Bath, Royal Crescent. The study included the evaluation of the dimensions of the monuments and the nature and amount of any material used for its realization by means of field inspection and examination of images, photos and other documents available in literature and on the internet. As the dominating material of the studied monuments is limestone/marble, the multi-pollutant dose-response function for limestone was applied to determine the corrosion and soiling of the materials used in the construction of the monuments.

The main conclusions from the study are as follows: being located in the heart of European capitals, the studied UNESCO sites are impacted by air pollution, mainly due to HNO₃ (a product of NO₂ oxidation) and PM₁₀, two pollutants that currently seem to play a prominent role in determining damage of limestone. The improvement of air quality between 2000 and 2010, mainly attributable to a significant reduction of air concentration of SO₂, produced a small decrease in the recession rate for limestone, first year exposure, which for the studied sites was quantified in about 5-8 per cent. Calculated recession rates after one year of exposure are above the background corrosion rate (3.2 μm year⁻¹) and generally close to the target for the year 2050 (6.4 μm year⁻¹) or even at one case close to the target for 2020 (8.0 μm year⁻¹). Corrosion due to air pollution would result in material deterioration costs ranging from €9.2 per square metre per year (m⁻² year⁻¹) to €43.8 m⁻² year⁻¹, depending on the pollution level and the climatic conditions. These costs add to the cost in background areas, estimated from €14 m⁻² year⁻¹ to €28 m⁻² year⁻¹.

The predicted loss of reflectance after five years of exposure to the surrounding environment was still unacceptably high for the studied monuments. Predicted soiling rate of limestone indicate that a “tolerable soiling before action”, which represents the threshold triggering significant adverse public reaction of what constitutes acceptable soiling and generally set at 35%, will be reached within 4-7 years after any restoration work. For cultural heritage objects a period of 10-15 years is considered to be appropriate.

In continuation of the pilot study, ICP Materials has launched a Call for Data on “Inventory and condition of stock of materials at UNESCO World Cultural Heritage Site”. The official letter of the Call for Data, a template for submission of data, an explanatory note with instructions on the use of the reporting template, and a brochure exemplifying the step by step approach for the previously assessed UNESCO sites were provided by the Call. These documents have been also made available for downloading on the ICP Materials website.

Main objective of the Call for Data is to invite Parties to participate in studies evaluating material deterioration due to air pollution at UNESCO World Cultural Heritage Site. The ultimate objective is to provide policy makers the evidence of the effects of air pollution not on a generic material or a generic artefact but on easily recognizable symbols of our culture and history.

This Call for Data requires qualitative and quantitative data on both the historic/cultural monument and on the environment. In view of the complexity of the call, the deadline for submission of the data is set to July 2017. Six Parties to the Convention have announced their intention to participate in the Call: Croatia, Germany, Italy, Norway, Sweden, and Switzerland.

8.4 Concluding remarks

Air pollution is a major environmental problem and has an impact not only on human health but also on the surface of materials and particularly on the surfaces of buildings and cultural monuments exposed outdoors. Interactions between air pollutants and materials lead to an early deterioration and soiling of buildings and monuments. This prompts more frequent maintenance and restoration activities, which is a cost, and which adversely affect the artefact. Most of the cultural heritage in Europe, and Italy in particular, are located in the heart of cities, where higher levels of pollutants are usually found.

Although decreasing trends in corrosion were observed for all materials in connection to the decrease of concentrations of air pollutants (mainly SO₂), current corrosion rates and soiling are still unacceptably high. The effects observed in Italy, which resemble the trend observed by the ICP Materials network, show that for some material (i.e. carbon steel and copper) a decreasing trend is still evident while for zinc and limestone no decrease can be detected in recent years. A trend can hardly be identified for the soiling of modern glass.

Currently, HNO₃ (a product of NO₂ oxidation) and PM₁₀ seem to play a prominent role in determining damage of limestone. In developing future policies on air quality, it is important to specifically address impacts on cultural heritage and the built environment. Air pollutants and climate act together in determining corrosion and degradation of materials. Further reduction of air pollutants is one way of compensating for increased risk of corrosion due to climate change.

In general, the damage caused by air pollution to historic and cultural monuments and buildings should be better studied and understood, including the assessment of the cost associated with the damage and the interaction of pollution and global climate change.

8.5 Acknowledgments

We would like to express our appreciation to Elena Dell'Andrea (Agenzia Regionale per la Prevenzione e Protezione Ambientale del Veneto), Marcello Ferdinandi (Istituto Superiore di Sanità) and Matteo Lazzarini (Agenzia Regionale per la Protezione dell'Ambiente della Lombardia) for their continuous support and assistance in the management of test sites for exposure of materials and measurement of environmental parameters.

We also wish to acknowledge present and former members of the ICP Materials Task Force, especially those responsible for materials sub-centres (Johan Tidblad, Katerina Kreislova, Markus Faller, Daniel de la Fuente, Tim Yates and Aurélie Verney-Carron).

Vegetation and urban air quality: recent findings



Chapter coordinator: Elena Paoletti

Contributors: Elena Paoletti¹, Elisabetta Salvatori², Fausto Manes²,

¹CNR, Italy; ²Department of Environmental Biology, Sapienza University of Rome

CHAPTER 9 - VEGETATION AND URBAN AIR QUALITY: RECENT FINDINGS

As urbanization increases, also the significance of urban forests (UFs; including individual trees, parks and forests) for improving the environmental quality of life in the cities is increasing (Roy et al., 2012). Among the many benefits that UF provide to people, a top relevance is up to air quality improvement, expected to improve human health by removing gaseous air pollutants and particulate matter (PM) from the air (Weber, 2013). Urban air pollution is a major threat to citizens' health (Pascal et al., 2013). A plethora of primary pollutants, e.g. nitrogen oxides (NO_x), sulphur dioxide (SO₂), PM, are directly emitted by combustion in industrial processes and vehicles. Secondary pollutants, e.g. ozone (O₃) and secondary organic aerosol (SOA), are formed by reactions in the atmosphere among precursors like NO_x and VOC. Among the major air pollutants, PM and O₃ have the largest impact on human health at present (Pascal et al., 2013). UFs act as sink for pollutants and have been argued to phytoremediate the air (Nowak, 2006; Manes et al., 2012). Areas with high urban forest density, in fact, have lower PM than other sites (Irga et al., 2015). Such filtering capacity usually translates into little percent improvements of air quality, but such little values translate into significant savings in terms of human health (Nowak et al., 2013; 2015).

Pollution removal by plants occurs through a combination of two pathways, including deposition to plant surfaces and/or stomatal uptake (Grote et al., 2016). Dry and wet deposition includes scavenging of pollutants by the foliage or bark and, in the cases of reactive air pollutants such as ozone, also in the gas-phase due to emitted reactive substances. Removal depends on air pollution concentrations, meteorological conditions, and resistance in the crown space, through the boundary layer adjacent to surfaces, and uptake of gases through stomata, i.e. the tiny pores on leaf surfaces. These resistances are controlled by vegetation properties at different scales: community (e.g. single trees, green corridors, parks, and forests), canopy (e.g. crown size, shape and density) and foliage (e.g. leaf shape, surface properties and physiology). Larger tree crowns have a higher potential of ameliorating air quality by maximising pollutant deposition (Paoletti et al., 2004), and thus the characteristics of the tree cover - in particular density and continuity of crowns - and of individual tree crowns - size, architecture, and the leaf area per unit ground surface area (*LAI*) - are important drivers for air quality improvement. Also leaf surface characteristics, e.g. cuticular morphology, leaf wettability and hairiness, affect particulate deposition (Kardel et al., 2012). Another important leaf trait affecting the air quality is the persistence of foliage throughout the year (evergreen species) or only during the growing season (deciduous species). As particulate pollution is typically higher in winter (Sieghardt et al., 2005), evergreen species should be recommended for maximizing the deposition of particles (Manes et al., 2014; Fares et al., 2016). In contrast, gaseous pollutants and in particular O₃ are higher during the growing season (Paoletti, 2006; 2009), thus deciduous species are better suited for filtering gaseous pollution.

The capacity to remove pollutants may largely diverge under stress conditions. In a study carried out in the city of Rome, Fusaro et al. (2015) have shown that urban and peri-urban forests were affected by different environmental stressors and forest management practices. Such factors exert a strong effect on the functionality of the two forest sites, thus affecting their ozone removal capacity and the resulting air quality improvement. Understanding these effects is therefore an essential step for a reliable quantification of the air quality amelioration provided by vegetation in metropolitan areas, and for a better management of the Green Infrastructure (sensu Tzoulas et al., 2007) of the

city. At this regard, also species selection for air pollution mitigation should consider the ability of tree species to adapt to local conditions.

Non-stomatal removal processes also include chemical deposition resulting from gas-phase reactions between pollutants (mostly O₃ and NO_x) and biogenic VOC (BVOC) emitted from the ecosystem (e.g. plants or soil) (Fares et al., 2010). Deciduous plants are typically high BVOC emitters, with highest emissions occurring during spring and summer at midday (Holzinger et al., 2006). BVOCs contribute to the formation of important air pollutants e.g. ozone, secondary organic aerosols and PM (Calfapietra et al., 2013). Adopting low BVOC-emitting species in UF is crucial for controlling air quality (Benjamin and Winer, 1998; Ren et al., 2014). In addition to the species-specific BVOC emission factor, however, also the amount of emitting leaves affects the total production of BVOC by trees. Therefore, larger crowns constitute a negative indicator of air quality in BVOC-emitting species. The main BVOCs are isoprene and monoterpenes.

Evidence is emerging that the presence of roadside trees in street-canyons reduces the upwards transport of air pollutant emissions, increases their storage in the canyon and reduces the penetration of clean air from aloft. As a result, higher pollutant concentrations can be observed at pedestrian level within vegetated canyons (Amorim et al., 2013; Salmond et al., 2013; Vos et al., 2013; Gromke and Blocken, 2015). Further investigation of the complex inter-relations between plant characteristics, microclimate, street configuration and pollutant emission, however, is still needed.

Importantly, trees may also affect air quality by emitting primary particles (pollen) and BVOCs (Churkina et al., 2015). Pollens may act as allergens and are possibly more potent in combination with other urban pollutants (Beck et al., 2013). In Europe, 113 million citizens suffer from allergic rhinitis and 68 million from allergic asthma (EFA, 2011), a number that will likely increase due to climate change (Forsberg et al., 2012). Pollen affects human health by triggering those allergic reactions (Bartra et al., 2007; Traidl-Hoffman et al., 2003). Pollen deposition on leaf surfaces, however, helps abating pollen concentration in the air (Dzierzanowski et al., 2011; Terzaghi et al., 2013), likely by mechanisms similar to those regulating particle deposition.

Based on physiological as well as anatomical species-specific traits, Nowak and Crane (1998) developed the iTree model (Formerly, UFORE) to calculate deposition rates of SO₂, NO_x, CO, O₃ and PM per leaf area and per tree from climatic and air pollution boundary conditions. This model has been used for several European case studies (e.g. Paoletti, 2009; Paoletti et al., 2011).

By focusing on the species-specific tree properties (i.e. traits) that determine canopy and foliage interaction with major air pollutants, Grote et al. (2016) classified the major tree species of European cities according to their potential of ameliorating air quality (Figure 9.1).

In addition to allergenicity, calculated as pollination duration x intensity x toxicity (Cariñanos et al., 2016), and BVOC emission potential, calculated as lumped isoprene and monoterpene emission potentials under standard conditions (based Karl et al., 2009), Grote et al. (2016) focused also on PM removal efficiency (based on Yang et al., 2015), water use efficiency (WUE, from Wang et al., 2013), and shading capacity calculated as leaf area index x relative leaf abundance throughout the year (based on Tiwary et al., 2016) x crown width/tree height (based on <https://www.horticopia.com/hortpip/index.shtml>). The efficiency of water use makes a plant well adapted to face water stress, which is a common issue for UFs.

Shading is highly appreciated as UF environmental service, because it reduces the urban heat island, i.e. the typical increase of air and surface temperatures that occur in the cities.

The suitability of a tree species for a particular combination of demands is highly case specific. The relative trade-off or synergistic benefit of different traits also depends on the trees immediate surrounding and the importance of the respective ecosystem service. *Aesculus*, for example might be favored for its shading ability but abundance is restricted to sites with good water supply because its WUE is low. *Pinus* species can be favored in Southern Europe, because they not only efficiently remove pollution but are also relatively stress tolerant. In particular, they are drought resistant, a trait less relevant in Northern regions. Stress tolerance may be the first selection criteria in polluted areas even if the gain in ecosystem services is small.




Figure 9.1 – Classification of major tree species in European cities according to their water use efficiency, shading capacity, particulate removal efficiency, isoprene and monoterpene emission potential and allergenicity, following Grote et al. (2016).

In Italy, several studies have recently investigated the ability of urban tree species to remove O₃ and PM from the air. Removal in the city of Florence was estimated per forest type and mapped by Paoletti et al. (2011) and Bottalico et al. (2016).

In the Municipality of Rome, Manes et al. (2012) have quantified the effects of tree diversity on the removal of tropospheric ozone (O₃) by means of a spatial analysis, integrating system dynamic modeling and GIS. Two years were considered (2003 and 2004), differing in climatic conditions and ozone levels. The results showed that different tree functional groups (evergreen broadleaves, deciduous broadleaves and conifers), have complementary O₃ uptake patterns, related to tree physiology and phenology, thus maintaining a stable community function across different climatic conditions. Based on published unitary costs of externalities and of mortality associated with O₃, the ecosystem service of O₃ removal of the Rome urban forest has been valued to roughly US\$2 and \$3 million/year, respectively. The removal of PM₁₀ has been also quantified in the same area and for the same years (Manes et al., 2014), and the Ecosystem Service of PM₁₀ removal by the three functional groups in the five Sanitary Districts of the Municipality has been mapped. Given the spatial uniformity of PM₁₀ levels in the urban area, the highest amount of PM₁₀ deposition rates, during the whole period, were those of the Sanitary District with the largest vegetation cover.

The highest PM₁₀ depositions for the three functional groups were estimated for the 2004 summer period, in concurrence with the highest mean values of Leaf Area Index. Large urban parks can also significantly contribute to improve air quality locally, i.e. at neighborhood level. At this regard, Silli et al. (2015) have estimated the potential PM₁₀ deposition to vegetation in Villa Ada, a historical park located in the Rome downtown, surrounded by densely built areas and by high-traffic density roads. The results showed that trees may effectively abate suspended particles, with evergreen broadleaved trees being most effective during summer, reducing the average air concentration of PM₁₀. During the year 2012, the woody vegetation of Villa Ada removed in total 4417.2 kg of PM₁₀.

In a recent study (Marando et al., 2016), the quantification of seasonal PM₁₀ removal capacity of urban and peri-urban forests has been carried out for the year 2015, considering the whole Metropolitan City (MC) of Rome. This is an administrative unit, which corresponds to the former Province of Rome, and covers an area of 5352 km². Around 22% of this surface is covered by forests, and the overall monetary value of the PM₁₀ removal service can be estimated to 161.78 million Euros for the year 2015.

At national scale, the PM₁₀ and O₃ removal from urban and periurban forests has been estimated in ten Metropolitan Cities (Turin, Venice, Milan, Genoa, Bologna, Florence, Rome, Naples, Bari and Reggio Calabria), taking into account the main Physiognomic-Structural Vegetation Categories. The findings remark the importance of the Leaf Area Index in PM₁₀ removal, and of functional diversity, which is related to stomatal conductance, in the O₃ sequestration process. The overall monetary value of this Ecosystem Service was estimated to be equal to 47 and 297 million USD for O₃ and PM₁₀, respectively, in the year 2003 (Manes et al., 2016). This represents, however, a gross estimate, which do not take into account the cost of management and maintenance of the GI.

In a recent study performed in the periurban forest of Castelporziano (Rome), Fares et al. (2016) have measured fluxes of PM₁, PM_{2.5} and PM₁₀ with fast optical sensors and eddy covariance technique, observing that PM₁ is mainly deposited during the central hours of the day, while fluxes of PM_{2.5} and PM₁₀ were negligible. Furthermore, a Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT v4) was applied to simulate the PM emission from traffic in the city of Rome, showing that a significant portion of PM is removed by vegetation in the days when the plume trajectory meets the peri-urban forest.

Guidolotti et al. (2016) implemented the EMEP MSC-W model to scale-down to tree-level O₃, NO₂ and PM₁₀ removal in a urban area in Northern Italy, compared the results with the outputs of UFORE (nowadays i-Tree) and found a good agreement with the estimates by this new methodology. In the case of O₃, pros and cons of laboratory, field, and modeling approaches have been recently discussed and a combination of the three levels of investigation has been recommended as essential for estimating O₃ removal by urban trees (Calfapietra et al., 2016).

Effects of air pollution on health



Chapter coordinator: Antonio Piersanti¹

Contributors: Antonio Piersanti¹, Pierluigi Altavista², Carla Ancona³, Giovanna Berti⁴, Ennio Cadum⁴, Luisella Ciancarella¹, Ilaria D'Elia¹, Francesco Forastiere³, Marina Mastrantonio², Francesca Pacchierotti², Gaia Righini¹, Raffaella Uccelli²

¹ENEA, Laboratory of Atmospheric Pollution, ²ENEA, Laboratory of Biosafety and Risk assessment,

³ Department Of Epidemiology, Lazio Regional Health Service, ⁴Piedmont Regional Environmental Protection Agency

CHAPTER 10 – EFFECTS OF AIR POLLUTION ON HEALTH

Part of this Chapter was presented at the 8th International Congress on Environmental Modelling and Software of the International Environmental Modelling and Software Society (iEMSs) – 10-14 July, 2016, in Toulouse (France).

10.1 Introduction

According to the World Health Organization (WHO, 2015), air pollution is the largest environmental risk for human health, with estimates of 600000 premature deaths and 1.6 billion US\$ of economic cost due to mortality and diseases in Europe in 2010 (WHO Regional Office for Europe, 2015). The assessment of the impacts of air quality on health is the endpoint of European and national cost-effective policies, which can be translated into emission reductions and control strategies.

Modelling tools connecting atmospheric dynamics, human response to air pollution and policy options, allow to optimize the use of dedicated resources and data, such as high performance computing and epidemiological cohort studies. In Italy, two recent projects (VIAS and EU LIFE+ MED HISS) estimated the health effects of air pollution on the national scale, following the Health Impact Assessment (HIA) scheme (Figure 10.1), based on existing estimates of the relative risk from previous epidemiological studies, which are applied to the specific exposure of the studied population. The use of a reference national air quality model (MINNI – National Integrated Model to support the international negotiations on atmospheric pollution), combined with monitoring data, allowed the coverage of the whole territory and thus the use of national population data for a comprehensive calculation of exposure and health outcomes.

VIAS (Integrated Assessment of the Impact of Air Pollution on the Environment and Health, 2013-2015; Ancona, 2015) project was funded by the Centre for Disease Control of the Italian Ministry of Health and coordinated by the Department of Epidemiology of the Lazio Region Health Service. VIAS estimated the mortality (from respiratory disease, cardiovascular disease, lung cancer and total) and the months of life lost due to exposure to air pollution. Estimations were carried out both on Italy as a whole and on each of the 20 Italian Administrative Regions. Health effects of PM_{2.5}, NO₂ and O₃ were quantified.




Figure 10.1. Air quality Health Impact Assessment scheme.

10.2 Results

To quantify the health gains of Italian population following changes in targets and policies, VIIAS carried out a baseline assessment on years 2005 and 2010. Using model projections on different scenarios (current legislation scenario - CLE, compliance with EU and Italian air quality standards – CLE + Target 1, CLE concentrations reduced by 20% - CLE + Target 2) an estimate of health population for 2020 was produced for future evaluation.

Both the national concentration maps of the analyzed pollutants and the future emission scenarios were provided by the MINNI model system (www.minni.org), funded by the Italian Ministry of Environment, which runs an atmospheric modeling system (AMS, Mircea et al., 2014) and an integrated assessment model (GAINS-Italy, D'Elia et al., 2009). The AMS performances have been assessed against measured concentrations in Italy for several simulation years (Mircea et al., 2014, 2016; Ciancarella et al., 2016).

The population data (people aged 30 and over) used for the exposure assessment are related to the year 2005 and were obtained by interpolation of the official national censuses of 2001 and 2011. The concentration of pollutants was provided by the model on a 4x4 km² grid, while population data were available on sub-municipal census zones. Therefore, for the calculation of exposure, population data were aggregated on the model grid.

The population exposure was associated with health outcomes by using concentration-response functions (CRFs), which put in relation air pollutants concentrations and health damages. A CRF provides an estimate of the relative risk, i.e. the increment of a health effect associated with a single-unit increment of the ambient concentration of the pollutant. CRFs are site specific, being obtained from observational epidemiological studies. In VIIAS, the WHO guidelines have been applied, using CRFs for mortality and coronary events taken from the WHO HRAPIE review conducted in 2013 (WHO, 2013). Pollutant-specific CRFs on single health outcomes were used. Thresholds of 10 µg/m³ for PM_{2.5}, 20 µg/m³ for NO₂ and 70 µg/m³ for O₃ were applied in the assessment, following WHO recommendations on the HIA procedure. The adult population (age 30 years and more) was taken into account following HRAPIE recommendations, as most of the evidence on PM_{2.5} long term effects a mortality comes from studies that focused on populations around 30 years of age and above.

VIIAS provided both national aggregated figures and results on 3 geographic areas (North, Center, South and Islands), 20 Member Regions, 2 macro areas (urban, non-urban). A summary of the results at the national level is presented in the following table 10.1.

Total figures in 2005 evidence the relevant impact of PM_{2.5} and NO₂ on mortality, ranging from 23000 to 34000 attributable deaths. The average exposure of the population to PM_{2.5} is 20.1 µg/m³, very close to the EU limit value of 25 µg/m³ (EC, 2008). Remarkable differences were found among the 20 Member Regions, with Northern Italy showing a higher number of deaths from PM_{2.5} (22485) and NO₂ (14008), and between urban and non-urban areas, with deaths attributable to NO₂ being more than doubled (16736 versus 6651) and PM_{2.5} deaths only slightly higher in urban areas (19358 versus 15194), confirming the different dispersion pattern and main sources (road traffic and residential heating for NO₂, secondary aerosol formation for PM_{2.5}). The different 2020 scenarios provide a forecast of prevented deaths according to different targets. As an example, attaining the EU limits could save 11000 lives on PM_{2.5} and 14000 on NO₂. The health gains are dependent on the geographical area, as emission reductions are different between Member Regions.

Table 10.1. Average population exposure, attributable deaths for PM_{2.5}, NO₂ and O₃, and months of life lost for PM_{2.5}, in Italy. Results of the VIAS project.

			2005	2010	2020 CLE	2020 CLE - Target1	2020 CLE - Target2
PM_{2.5}	general mortality	population exposure (µg/m ³)	20.1	15.8	18.1	16.2	14.5
		attributable deaths (95% confidence interval)	34552 (20608-43215)	21524	28595	23170	18511
		months of life lost	9.7	5.5	7.7	5.9	4.2
NO₂	general mortality	population exposure (µg/m ³)	24.7	17.9	16.6	16.1	13.3
		attributable deaths (95% confidence interval)	23387 (21514-50283)	11993	10117	9021	5247

It is worth noting that the 2020 CLE scenario shows higher exposure and effects for PM_{2.5} in 2020 than in 2010, due to an increase in PM_{2.5} emissions. This growth is caused by the large spread of wood combustion for residential heating, due both to climate change mitigation policies (encouraging carbon-neutral fuels such as biomasses) and to persistent effects of the economic crisis (leading people to use self-provided wood for heating). This growth overcomes scenario mitigation measures, resulting in an increase of PM_{2.5} concentrations and population exposure.

Results were obtained on the MINNI geographical reference, providing national maps of health outcomes, e.g. death rates per 100000 inhabitants from PM_{2.5} (Figure 10.2). Large zones in Northern Italy, Tuscany, Rome and Naples show significant death rate values in 2005, with peaks above 250. A decreasing trend is observed between 2005 and 2020, which can be attributed to the growing efficacy of policy mitigation options. Although compliance with the current EU legislation would have a large impact on the health of Italian residents, there is a large scope for further improvement.

Other uncertainties are present in the health outcomes results (synergies and antagonisms between pollutants were not quantified), in the concentration-response functions (conclusions of epidemiological studies conducted in different conditions and populations are extrapolated to Italy) and in the model outputs (coarse resolution for NO₂, scenario assumptions are realistic but subjected to periodical revisions).




Figure 10.2. Death rates attributable to $PM_{2.5}$ each 100000 inhabitants in the 5 case studies. Results of the VIAS project.

MED HISS (Mediterranean Health Interview Surveys Studies: long term exposure to air pollution and health surveillance) is a recently completed EU LIFE+ Pilot project (2013-2016) involving four countries (Italy, France, Slovenia and Spain). MED HISS is aimed to set up a surveillance system of long term effects of air pollution, based on the common availability of routine air quality and health data. Three main information sources were used: 1) the National Health Interview Surveys (NHIS, Hupkens et al., 1999), available in all countries, representative of the total population and covering both urban and rural areas; 2) mortality and hospital admissions registries; 3) air pollution models.

The use of NHISs data already available, which include individual information on main potential confounders (smoking, BMI, occupation, education, etc.), allows the recruiting of retrospective cohorts, with a remarkable saving of resources, that are normally allocated on building the information on the observed population. Furthermore, NHISs data on entire national populations try to overcome the existing limitations on the health effect of air pollution, historically based on cohort studies from outside EU, or conducted in Europe, but with restrictions on age or location.

Cohorts are being followed-up for mortality and morbidity, and each subject will be assigned a level of exposure to air pollution (PM_{10} , $PM_{2.5}$, NO_2 , O_3), derived from national-scale dispersion models. As population and health data are available at municipality level, a dedicated work package up-scaled the model gridded concentrations on municipalities, using a weighted block averaging procedure (Ignaccolo, 2012) that accounts for built-up surface percentages collected from CORINE LAND COVER data (<http://www.eea.europa.eu/data-and-maps/data/corine-land-cover-2006-raster-3>). MED HISS cohorts are representative of all populations and areas of residence (urban, rural, metropolitan) and long term effect will be evaluated for a wide range of diseases.

Long term effects of air pollution on mortality and hospital admissions (the latter rarely assessed in epidemiological studies) were calculated for Italy with the 1999-2000 NHIS, with different follow-up periods and number of cohort members using the Cox proportional hazard model (Cox, 1972), with pollutants and age time-varying variables, adjusting for other variables (age, gender, educational level, activity status, living alone, BMI, smoking, physical activity).

Hazard ratios related to $10 \mu\text{g}/\text{m}^3$ of $\text{PM}_{2.5}$ increase are in line with other cohort studies for mortality for all causes (1.04, confidence interval 95% 1.02-1.06) and for circulatory system diseases (1.03, 1.00-1.06) and lung cancer (1.12, 1.04-1.21). Risks of first-ever hospital admissions related to $\text{PM}_{2.5}$ were found to be significant for circulatory system diseases (1.04, 1.01-1.07), lung cancer (1.18, 1.08-1.29), kidney cancer (1.18, 1.02-1.36) and myocardial infarction (1.15, 1.10-1.21).

The relative risks allowed to assess the health outcomes in Italy for 2010 related to $\text{PM}_{2.5}$: 33533 (20429-41368) attributable deaths, 14 years of life lost on average for each death, 9.2 months of reduction of life expectancy for each individual. Northern Italy and urban areas, being more polluted, show higher mortality.

As MED HISS proposed a new methodology, several practical issues emerged during the project.

Data availability is different in the 4 countries, both generated by air quality models (in France-CHIMERE, in Italy-MINNI, in Slovenia-ARSO and in Spain-CALIOPE) and for health data. Model datasets are inhomogeneous because available on different years, due to differences in the national practices and non-mandatory use of models for regulatory purposes (e.g., MINNI model data on Italy required assimilation of measured data, to be comparable to the other countries).

Inhomogeneous anonymization and linkage procedures and schemes of NHIS questionnaires are in force across Europe. The individual linkage between NHIS sampled subjects and mortality/morbidity data was not possible in Spain and Slovenia for preserving privacy and was substituted with an ecological approach based on aggregated data, which allowed to calculate relative risk rates for mortality from $\text{PM}_{2.5}$.

National-scale epidemiological studies can be as well based on measured data of pollutant concentrations. A recent epidemiological study has been carried out through a voluntary collaboration between researchers from ENEA, Italian National Institute for Environmental Protection and Research (ISPRA) and Istituto Superiore di Sanità (ISS) (Uccelli et al., 2016). It concerns female lung cancer mortality and long term exposure to particulate matter (PM) in Italy starting from data detected by the official monitoring stations. It is the first Italian epidemiological study on female lung cancer mortality on all the municipalities of province capital cities with available measured mean annual concentrations of PM_{10} and $\text{PM}_{2.5}$. PM is one of the main responsible of the impact of air pollution on human health and it has recently been classified in Group 1 by IARC, besides outdoor air pollution (IARC, 2015). Even though in Italy a reduction of the emissions has been observed in the last 25 years, PM concentrations are still high in comparison with both the European targets and the WHO guidelines of air quality (WHO, 2006). The study considered female population only, in order both to reduce the confounding effect of occupational exposures and to focus on an association previously investigated mainly in men. The dose-response relationship between female lung cancer mortality in the 2000-2011 period and PM_{10} and/or $\text{PM}_{2.5}$ available mean annual outdoor concentrations (respectively in 64 and 32 municipalities) and the burden of death due to such exposures were computed.

Standardized mortality rates (SMRate) were calculated by the ENEA's epidemiological database, which includes the Italian mortality data (both general and cause specific) up to 2013, the 3rd International Classifications of Diseases (ICD VIII, IX and X), and the Italian decennial census populations from 1961 and their annual interpolations. Mortality data, codified and recorded by the National Institute of Statistics (ISTAT), are the only health data immediately available in Italy for all municipalities. Multiple regression analysis of SMRates, as a function of PM concentrations, considering percentage of smokers and deprivation indexes as additional explanatory variables, was performed for PM₁₀ only due to the relatively low number of PM_{2.5} monitoring stations' measures. An SMRate increase of 0.325 for 1 µg/m³ increment of PM₁₀ concentration was calculated.

On the basis of such an increase and of the attributable risk evaluated from the overall difference of SMRates between the 2 subgroups of municipalities equal/below and above the WHO guideline of 20 µg/m³, a proportion ranging between 13-16% female lung cancer mortality could be attributed to PM₁₀ levels exceeding the WHO guideline. Therefore, about 300 female lung cancer deaths could be prevented each year in the investigated municipalities with an overall annual population of 8, 146, 520.

What remains to be done to reduce air pollution?



Chapter coordinator: Ilenia D'Elia

Contributors: Ilenia D'Elia¹, Giovanni Vialetto¹, Luisella Ciancarella¹, Alessandra De Marco¹

¹ENEA, Laboratory of Atmospheric Pollution

CHAPTER 11 - WHAT REMAINS TO BE DONE TO REDUCE AIR POLLUTION?

The previous chapters have shown how air pollution and its environmental effects have considerably improved over the past 20 years. This important result has been achieved also as a consequence of the implementation of the different protocols under the Convention on Long Range Transboundary Air pollution (CLRTAP), and in particular of the national emission ceilings under the Gothenburg Protocol of 1999 and the NEC Directive (EC, 2001). However, critical issues remain, such as the outstanding effect of N deposition on forest nutrition and growth and on water quality. Current levels of N deposition are still too high to prevent impact on soil, foliar and water chemistry, tree growth and carbon sequestration.

The implementation of the current protocols and directives contributes to reduce, but not eliminate, the negative impacts on human health, materials and environment and further measures are needed in order to eliminate as much as possible negative effects. The impact assessment of the Clean Air Policy Package (COM, 2013) shows, for example, that by 2030 the clean air policy package, compared to the business as usual scenario, is estimated to avoid 58000 premature deaths, save 123000 km² of ecosystems from nitrogen pollution, save 56000 km² protected Natura 2000 areas from nitrogen pollution and save 19000 km² of forest ecosystems from acidification. Moreover, health benefits alone will save €40-140 billion in external costs and provide about €3 billion in direct benefits (Amann et al., 2014).

The last two most important negotiation processes, the Gothenburg Protocol at the international level and the NEC Directive at the European level, have introduced for the first time a cost-effectiveness and effect-based principle as the rationale to derive quantitative and differentiated national reduction obligations based on the reduction of health impact (Amann et al., 2011; COM, 2013; Amann et al., 2015). This new approach highlighted how the reductions of the effects on human health and ecosystems are leading all the air pollution policies.

To support the methodological aspects of the policy design through this new approach and make it applicable at a regional level within a country, the Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA) has developed MINNI (*National Integrated Model to support the International Negotiation on atmospheric pollution*) (www.minni.org), funded by the Italian Ministry of Environment, Land and Sea (see box 11.1).

MINNI is an Integrated Modelling System that links atmospheric science with the economics of emission abatement measures and policy analysis and consists of several interdependent and interconnected components: the national AMS (*Atmospheric Modeling System*, Mircea et al., 2014) and the national GAINS-Italy (Ciucci et al., 2016; D'Elia et al., 2009). They interact in a feedback system through ATMs (*Atmospheric Transfer Matrices*) and RAIL (*RAINS-Atmospheric Inventory link*).

BOX 11.1

MINNI (National Integrated Model to support the International Negotiation on atmospheric pollution)

The MINNI project started as an agreement between the Italian Ministry of Environment and ENEA, in collaboration with AriaNet and IIASA to support the Italian emission reduction policies. MINNI has then evolved into a scientific project, coordinated by ENEA, to investigate thematic concerns concerning the air pollution field, including the development and maintaining of a state-of-the-art National Integrated Model and its validation using experimental data.

The project focuses on natural and anthropogenic causes of atmospheric pollution, both from national and trans-boundary origin, and on its impact on human health, ecosystems and country's cultural heritage. In particular air concentrations of ozone, heavy metals, Persistent Organic Pollutants (POPs), primary and secondary particulate matter are studied, along with acid depositions and depositions leading to eutrophication.


MINNI consists of two main modeling systems:

- **Atmospheric Model System (AMS)**
- **GAINS-Italy (Greenhouse Gas - Air Pollution Interactions and Synergies)**

The Atmospheric Model System (AMS) belongs to the family of chemical transport models (CTM), which describe physico-chemical processes in the atmosphere. The outputs of such models are four dimensions pollutant concentrations (considering the time), starting from known meteorological and emissions conditions with a spatial resolution of 4 km and temporal of one hour. Such models allow to link emissions to concentrations, a crucial point for management of recovery actions.

The integrated assessment model system GAINS-Italy allows evaluation of impacts and costs. Starting from information on emission abatement technologies and economic scenarios of the energy and productive sectors, GAINS-Italy produces alternative and/or future emission scenarios and provides support for the evaluation of the cost-efficiency of abatement options. The efficiency is expressed in terms of concentration reductions.

The two components are linked in a feedback system, by ATM Atmospheric Transfer Matrices (ATM) and RAINS-Atmospheric Inventory Link (RAIL).



In accordance with the most updated emission scenario elaborated by ENEA and ISPRA (D'Elia and Peschi, 2016), the baseline scenario, which assumes full implementation of current legislation, both at a European and national level, will make it possible for Italy to comply with the emission reduction commitments of the Gothenburg Protocol for all pollutants, while the new NEC targets from the year 2030 will require additional measure for all pollutants (see fig. 11.1).




Figure 11.1 – Projected emission reduction in the last national emission scenario and comparison with the NEC targets.

Different and cost-efficient measures are available to reduce emissions, for example the Ecodesign Directive in the residential sector could reduce PM_{2.5} emissions in a range between 35% and 55% in the year 2030 with respect to the base year 2005 depending on the implementation hypotheses; while measures on urea-based fertilizers and of the low N feeding for livestock could lead to an ammonia reduction of 22% (D’Elia and Peschi, 2016). Scenarios are not forecasts and they merely describe future developments based on major driving forces and on their impacts. In order to define the additional reduction needed and the most efficient measures that should be put in place to reach the emission reduction commitments, sensitivity analyses are unavoidable, especially on Euro 6 standards on diesel cars whose complete failure would let the NOx target unreachable.

The amended Gothenburg protocol (ECE, 2012a; 2012b) and the other last new protocols and the new NEC Directive (EC, 2016) also address newly recognized challenges. Emission reduction commitments have been introduced for a long list of new pollutants, from fine particulate matter (PM_{2.5}) to some heavy metals (lead, cadmium and mercury) and a list of persistent organic pollutants (including inter alia dioxins, PAHs, DDT, PCB etc.). In particular, measures for fine particulate matter also include black carbon, that not only may have harmful health effects, but it is also a powerful greenhouse gas. It is very important to consider climatic policies in combination with air quality policies to have a win-win policy.

Ozone, mercury and POPs also show the importance of the global transport of atmospheric pollutants. From this point of view, cooperation between CLRTAP and other northern hemispheric and worldwide organizations is becoming increasingly important and is highly encouraged.

Another possibility to reduce the air pollution is linked to the so called unconventional measures. One of the most representative examples of unconventional measure is the reduction of meat consumption, strictly linked to human diet and personal choices. An example of the change in pollutants emissions linked to change in the human consumption was reported by Westhoek et al. (2014) that demonstrated a reduction in the emission of nitrogen compounds around 40% halving meat consumption all over Europe.

Changes toward a more plant-based diet could help substantially in mitigating emissions of GHGs, because the 24% of GHGs emissions (an average for all the selected countries) is due to food consumption. Unfortunately, this is a largely unexplored area of climate policy. Few authors have proposed changes that lower meat consumption.

Indeed, anthropogenic emissions of greenhouse gases (GHGs) related to food production accounts for about 15% at a world level. Carlsson-Kanyama (1998 and 2003) have shown that food choices and diet can influence the energy requirements for the provision of human nutrition and the associated GHG emissions. Meals similar in caloric content may differ by a factor lasting from 2 to 9 in GHG emissions (Engstrom et al., 2007). An analysis of the energy inputs showed that meals with similar nutritional value had a difference in GHG emissions of up to a factor of 4, depending on the items chosen (Carlsson-Kanyama et al., 2003). All of these studies identified certain foods as more resource demanding/polluting, including animal products and certain vegetable-intensive ways produced.

Many improvements have been reached through the Convention on Long-range Transboundary Air Pollution and its protocols but long-term risks due to ozone, nitrogen, heavy metals and persistent organic pollutant continue to exist and a significant proportion of the Italian population is exposed to high concentrations of fine particles and ozone. Lots solutions are available and an integrated approach to air pollution and climate change could lead to significant co-benefits. Effective environmental policy could only be developed implementing monitoring programmes, developing models to predict concentrations and deposition levels, and assessing their effects and measures. The integration of all the different tools, from measures to models, and the coordination among science sectors and different research teams could help in identifying cost-effective solutions.

REFERENCES

CHAPTER 2

EB, 1999. Protocol to the 1979 Convention on Long-Range Transboundary Air Pollution to abate Acidification, Eutrophication and Ground-level Ozone, EB.AIR/1999/1.

<http://www.unece.org/fileadmin/DAM/env/lrtap/full%20text/1999%20Multi.E.Amended.2005.pdf>

ECE, 2010. Decision 2010/18. Long-term strategy for the Convention on Long-range Transboundary Air Pollution and Action Plan for Its Implementation. ECE/EB.AIR/106/Add.1.

http://www.unece.org/fileadmin/DAM/env/lrtap/ExecutiveBody/Decision_2010.18.pdf

ECE, 2012a. Decision 2012/1. Amendment of annex I to the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone. ECE/EB.AIR/111/Add.1.

http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE_EB.AIR_111_Add.1_ENG_DECISION_1.pdf

ECE, 2012b. Decision 2012/2. Amendment of the text of and annexes II to IX to the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone and the addition of new annexes X and XI.

ECE/EB.AIR/111/Add.1

http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE_EB.AIR_111_Add.1_ENG_DECISION_2.pdf

Maas, R., et al., 2016. Towards Cleaner Air. Scientific Assessment Report 2016. May 2016.

http://www.unece.org/fileadmin/DAM/env/lrtap/ExecutiveBody/35th_session/CLRTAP_Scientific_Assessment_Report_-_Final_20-5-2016.pdf

CHAPTER 3 - SOURCES AND EMISSIONS OF AIR POLLUTANTS

COM, 2013. COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT, THE COUNCIL, THE EUROPEAN ECONOMIC AND SOCIAL COMMITTEE AND THE COMMITTEE OF THE REGIONS. A Clean Air Programme for Europe. Brussels, 18.12.2013, COM(2013) 918 final.

http://ec.europa.eu/environment/air/clean_air_policy.htm

EC, 2001. Directive 2001/81/EC of the European Parliament and of the Council of 23 October 2001 on national emission ceilings for certain atmospheric pollutants. Official Journal of the European Union, L. 309 of 27/11/2001.

EC, 2008. Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and a cleaner air for Europe. EC Official Journal L. 152 of 11.06.2008.

EEA, 2015. Air Quality in Europe – 2015 report. Report n. 5/2015. <http://www.eea.europa.eu/publications/air-quality-in-europe-2015>.

IARC, 2015. Outdoor Air Pollution. IARC monographs on the evaluation of carcinogenic risk to humans. Volume 109. Lyon, France: IARC; 2015. <http://monographs.iarc.fr/ENG/Monographs/vol109/index.php>

IIR, 2016. Italian Emission Inventory 1990-2014. Informative Inventory Report 2016. ISPRA Technical Report, n. 240/2016. ISBN 978-88-448-0765-8.

WHO, 2005. WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide. Global update 2005. Summary of risk assessment. WHO/SDE/PHE/OEH/06.02.

http://apps.who.int/iris/bitstream/10665/69477/1/WHO_SDE_PHE_OEH_06.02_eng.pdf

CHAPTER 4 - FOREST

Bertini, G., Amoriello, T., Fabbio, G., Piovosi, M., 2011. Forest growth and climate change. Evidences from the ICP-Forests intensive monitoring in Italy. iForest 4: 262-267.

Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L., De Vries, W.,

2010. Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Applicat*, 20: 30-59.
- Bussotti, F., Ferretti, M., 2007. OzoneFlux. Measure and modelling of Ozone flux in Evergreen Mediterranean stands of the EU Intensive Monitoring of Forest Ecosystems (Level II) – An approach at different intensity levels. Final report – Italy. Jointly prepared by Corpo Forestale dello Stato, Italia; Ministero de Medio Ambiente, Direccion para la Biodiversidad, España: 161 ps.
- Bussotti, F., Ferretti, M., 2009. Visible injury, crown condition, and growth responses of selected Italian forests to ozone. Evidence from the National Monitoring Programme. *Environmental Pollution*, 157: 1427–1437.
- Canullo, R., Campetella, G., Petriccione, B., 2007. Manuale per le operazioni di campionamento della vegetazione. Progetto BioSoil – biodiversity: Valutazione della biodiversità forestale sulla Rete sistematica di Livello I. Università di Camerino – MIPAAF, 20 pp.
- Canullo, R., Allegrini, M.C., Campetella, G., 2013. Manuale nazionale di riferimento per la raccolta dei dati di vegetazione nella rete italiana CONECOFOR LII (Programma Nazionale per il Controllo degli Ecosistemi Forestali – UN/ECE, ICP Forests). *Braun-Blanquetia* 48: 1-65. [<http://www.scienzadellavegetazione.it/sisv/libreria/braun-blanquetia/BRBL48.pdf>] (accessed May 2016).
- Ciais, P., Reichstein, M., Viovy, N., et al., 2005. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437:529–533. doi: 10.1038/nature03972.
- CLRTAP, 2004. Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution effects, risks and trends. Convention on Long-Range Transboundary Air Pollution: 251 ps.
- EEA, 2007. European forest types – Categories and types for sustainable forest management reporting and policy. 2nd edition, EEA Technical report 9/07, European Environment Agency, Copenhagen: 111 ps.
- EEA, 2012. Biogeographic regions in Europe. [<http://www.eea.europa.eu/data-and-maps/figures/biogeographical-regions-in-europe-1>]. (accessed May 2016).
- Ferretti, M., Calderisi, M., Marchetto, A., et al., 2015. Variables related to nitrogen deposition improve defoliation models for European forests. *Annals of Forest Science*, 72, 897-906. DOI 10.1007/s13595-014-0445-6.
- Ferretti, M., Marchetto, A., Arisci, S., et al., 2014. On the tracks of Nitrogen deposition effects on temperate forests at their southern European range - an observational study from Italy. *Global Change Biol*, 20: 3423-3438.
- Ferretti, M., Fagnano, M., Amoriello, T., et al., 2007. Measuring, modelling and testing ozone exposure, flux and effects on vegetation in southern European conditions - what does not work. A Review from Italy. *Environmental Pollution*, 146: 648-658.
- Ferretti, M., Fabbio, G., Bussotti, F., Petriccione, B., 2006. Aspects of biodiversity in selected forest ecosystems in Italy. *Annali Istituto Sperimentale per la Selvicoltura*, 30, Suppl. 2.
- Ferretti, M., Bussotti, F., Fabbio, G., Petriccione, B., 2003. Ozone and Forest Ecosystems in Italy. Second report of the Task Force on Integrated and Combined (I&C) evaluation of the CONECOFOR programme. *Annali Istituto Sperimentale per la Selvicoltura, Special Issue, Arezzo Anno 1999 Vol. 30, Suppl. 1.*
- Gasparini, P., Di Cosmo, L., Pompei, E., 2013. Il contenuto di carbonio delle foreste italiane. *Inventario Nazionale delle Foreste e dei serbatoi forestali di Carbonio INFC2005. Metodi e risultati dell'indagine integrativa.* Ministero delle Politiche Agricole, Alimentari e Forestali, Corpo Forestale dello Stato; Consiglio per la Ricerca e la Sperimentazione in Agricoltura, Unità di ricerca per il Monitoraggio e la Pianificazione Forestale. Trento, 284 pp.
- Gasparini, P., Tabacchi, G., 2011. L'Inventario Nazionale delle Foreste e dei Serbatoi Forestali di Carbonio INFC 2005. Secondo inventario forestale nazionale italiano. Metodi e risultati. Ministero delle Politiche Agricole, Alimentari e Forestali, Corpo Forestale dello Stato; Consiglio per la Ricerca in Agricoltura, Unità di Ricerca per il Monitoraggio e la Pianificazione Forestale. Edagricole, Milano: pp. 653.
- Gerosa, G., Ferretti, M., Bussotti, F., Rocchini, D., 2007. Estimates of ozone AOT40 from passive sampling in forest sites in South-Western Europe. *Environmental Pollution*, 145: 629-635.

Gerosa, G., Ferretti, M., Buffoni, A., Spinazzi F., 2003. Vegetation exposure to ozone at the Permanent Monitoring Plots of the CONECOFOR Programme in Italy: estimating AOT40 by means of passive samplers. In: Ferretti, M., Bussotti, F., Fabbio, G., Petriccione, B., (Eds.), *Ozone and Forest Ecosystems in Italy*. Second report of the Task Force on Integrated and Combined (I&C) evaluation of the CONECOFOR programme. *Annali Istituto Sperimentale per la Selvicoltura, Special Issue, Arezzo Anno 1999 Vol. 30, Suppl. 1 2003: 53-62.*

Gottardini, E., Cristofolini, F., Cristofori, A. et al., 2012. Ozono e foreste in Trentino – Risultati del progetto ozono EFFORT 2007-2011, Fondazione Edmund Mach, San Michele all’Adige (TN), 144 ps.
www.fmach.it/content/download/18642/181979/.../9130_Report_Ozone_EFFORT.pdf

Gottardini, E., Cristofolini, F., Cristofori, A., Ferretti, M., 2014. Ozone risk and foliar injury on *Viburnum lantana* L.: a meso-scale epidemiological study. *Science of Total Environment*, 493: 954–960.

Gottardini, E., Cristofolini, F., Ferretti, M., submitted. Foliar symptoms on *Viburnum lantana* reflects annual changes in ozone concentration in Trentino (northern Italy). *Ecological Indicators*.

Gottardini E., Cristofori A., Cristofolini F., Ferretti M., 2010. Variability of ozone concentration in a montane environment, northern Italy. *Atmospheric Environment*, 44: 147-152

Karnosky, D.F., Percy, K.E., Chappelka, A.H., et al., 2003. Air pollution, global change and forests in the New Millennium. *Elsevier DES*, 3: 469 ps.

Leuzinger, S., Zotz, G., Asshoff, R., Körner, C., 2005. Responses of deciduous forest trees to severe drought in Central Europe. *Tree Physiol* 25:641–650. doi: 10.1093/treephys/25.6.641.

Sanders, T.G.M., Michel, A.K., Ferretti M., 2016. 30 years of monitoring the effects of long-range transboundary air pollution on forests in Europe and beyond. *UNECE/ICP Forests, Eberswalde*, 67 p.

Schaub, M., Emberson, L., Bueker, P., Kräuchi, N., 2007. Preliminary results of modeled ozone uptake for *Fagus sylvatica* L. trees at selected EU/UN-ECE intensive monitoring plots. *Environmental Pollution* 145 (3): 636-643.

Tabacchi, G., Di Cosmo, L., Gasparini, P., Morelli, S., 2011. Stima del volume e della fitomassa delle principali specie forestali italiane, Equazioni di previsione, tavole del volume e tavole della fitomassa arborea epigea.

CHAPTER 5 - EFFECTS OF AIR POLLUTION ON CROPS AND SEMI-NATURAL VEGETATION

Allegrini, I., Cortiello, M., Manes, F., Tripodo, P., 1994. Physicochemical and biological monitoring as integrated tools in evaluating tropospheric ozone in urban and semirural areas. *Science of the Total Environment* 141, 75-85.

Anav, A., De Marco, A., et al., 2016. Comparing concentration-based (AOT40) and stomatal uptake (PODY) metrics for ozone risk assessment to European forests. *Global Change Biology* 22, 1608–1627.

Andersen, C.P., 2003. Source–sink balance and carbon allocation below ground in plants exposed to ozone. *New Phytologist* 157, 213–228.

Astorino, G., Marganti, I., Tripodo, P., Manes, F., 1995. The response of *Phaseolus vulgaris* L. cv. Lit. to different dosages of the anti-ozonant Ethylenediurea (EDU) in relation to chronic treatment with ozone. *Plant Science* 111, 237-248.

Basile, A., Esposito, S., Sorbo, S., Napolitano, E., Conte, B., Salvatori, E., Manes F., 2010. Ultrastructural alterations induced by tropospheric ozone: comparison between resistant and sensitive clones of *Trifolium repens* L. CV. Regal. *International Journal of Environment and Health* 4(2-3), 260-277.

Bassin, S., Calanca, P., Weidinger, T., Gerosa, G., Fuhrer, J., 2004. Modeling seasonal ozone fluxes to grassland and wheat: model improvement, testing and application. *Atmospheric Environment* 38, (15) 2349–2359

Bernardini, A., Salvatori, E., Guerrini, V., Fusaro, L., Canepari, S., Manes, F., 2016. Effects of high Zn and Pb concentrations on *Phragmites australis* (Cav.) Trin. Ex. Steudel: photosynthetic performance and metal accumulation capacity under controlled conditions. *International Journal of Phytoremediation* 18 (1), 16-24.

Bonanomi, G., Caporaso, S., Allegranza, M., 2006. Short-term effects of nitrogen enrichment, litter removal and cutting on a Mediterranean grassland. *Acta Oecologica* 30(3), 419-425.

- Burkey, K.O., Booker, F.I., Ainsworth, E.A., Nelson, R.L., 2012. Field assessment of a snap bean ozone bioindicator system under elevated ozone and carbon dioxide in a free air system. *Environmental Pollution* 166, 167-171
- Burkey, K.O., Miller, J.E., Fiscus, E.L., 2005. Assessment of ambient ozone effects on vegetation using snap bean as a bioindicator species. *Journal of Environmental Quality* 34, 1081–1086.
- Bussotti, F., Desotgiu, R., Cascio, C., Pollastrini, M., Gravano, E., Gerosa, G., Marzuoli, R., Nali, C., Lorenzini, G., Salvatori, E., Manes, F., Schaub, M., Strasser, R.J., 2011. Ozone stress in woody plants assessed with chlorophyll a fluorescence. A critical reassessment of existing data. *Environmental and Experimental Botany* 73, 19-30.
- Castagna, A., Ranieri, A., 2009. Detoxification and repair process of ozone injury: from O₃ uptake to gene expression adjustment. *Environmental Pollution* 157, 1461–1469.
- Cepi, M.G., Oukkaroum, A., Çiçeka, N., Strasser, R.J., Schansker, G., 2012. The IP amplitude of the fluorescence rise OJIP is sensitive to changes in the Photosystem I content of leaves: a study on plants exposed to magnesium and sulfate deficiencies, drought stress and salt stress. *Physiologia Plantarum* 144, 277–288.
- Coll, I., Lasry, F., Fayet, S., Armengaud, A., Vautard, R., 2009. Simulation and evaluation of 2010 emission control scenarios in a Mediterranean area. *Atmospheric Environment* 43, 4194–4204.
- Convention on Long-Range Transboundary Air Pollution (CLRTAP), 2010. Manual in Methodologies and Criteria for Modelling and Mapping Critical Loads and Levels and Air Pollution Effects, Risks and Trends. ICP-Vegetation Co-ordination Centre, UK. Available at: <http://www.icpmapping.org>.
- Cooper, O.R., Parrish, D.D., Ziemke, J., Balashov, N.V., Cupeiro, M., Galbally, I.E., Gilge, S., Horowitz, L., Jensen, N.R., Lamarque, J.-F., Naik, V., Oltmans, S.J., Schwab, J., Shindeli, D.T., Thompson, A.M., Thouret, V., Wang, Y., Zbinden, R.M., 2014. Global distribution and trends of tropospheric ozone: An observation-based review. *Elementa: Science of the Anthropocene* 2: 000029, doi: 10.12952/journal.elementa.000029.
- Cristofanelli, P., Bonasoni, P., 2009. Background ozone in the southern Europe and Mediterranean area: Influence of the transport processes. *Environmental Pollution* 157, 1399–1406.
- D'Angiolillo, F., Tonelli, M., Pellegrini, E., Nali, C., Lorenzini, G., Pistelli, L., Pistelli, L., 2015. Can ozone alter the terpenoid composition and membrane integrity of in vitro *Melissa officinalis* shoots? *Natural Product Commun.*, 13: 1055-1058.
- De Marco, A., Sicard, P., Vitale, M., Carriero, G., Renou, C., Paoletti E., 2015. Metrics of ozone risk assessment for Southern European forests: testing the potential of canopy moisture content as plant response indicator. *Atmospheric Environment*, 120, 182-190
- Döring, A.S., Pellegrini, E., Della Bartola, M., Nali, C., Lorenzini, G., Petersen, M., 2014a. How do background ozone concentrations affect the biosynthesis of rosmarinic acid in *Melissa officinalis*? *J. Pl. Physiol.*, 171: 35-41.
- Döring, A.S., Pellegrini, E., Campanella, A., Trivellini, A., Gennai, C., Petersen, M., Nali, C., Lorenzini, G., 2014b. How sensitive is *Melissa officinalis* to realistic ozone concentrations? *Pl. Phys. Biochem.*, 74: 156-164
- EC, 2008. Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. *Official Journal of the European Union*, L 152 of 11/6/2008.
- Fares, S., Mereu, S., Scarascia Mugnozza, G., Vitale, M., Manes, F., Frattoni, M., Ciccioli, P., Gerosa, G., Loreto, F., 2009. The ACCENT-VOCBAS field campaign on biosphere-atmosphere interactions in a Mediterranean ecosystem of Castelporziano (Rome): site characteristics, climatic and meteorological conditions, and eco-physiology of vegetation. *Biogeosciences* 6, 1043–1058.
- Fernández-Fernández, M.I., Gallego, M.C., García, J.A., Acero, F.J., 2011. A study of surface ozone variability over the Iberian Peninsula during the last fifty years. *Atmospheric Environment* 45, 1946-1959.
- Ferretti, M., Fagnano, M., Amoriello, T., Badiani, M., Ballarin-Denti, A., Buffoni, A., Bussotti, F., Castagna, A., Cieslik, S., Costantini, A., De Marco, A., Gerosa, G., Lorenzini, G., Manes, F., Merola, G., Nali, C., Paoletti, E., Petriccione, B., Racalbutto, S., Rana, G., Ranieri, A., Tagliaferro, A., Vialletto, G., Vitale, M., 2007. Measuring, modelling and testing ozone exposure, flux and effects on vegetation in southern European conditions – what does not work? A review from Italy. *Environmental Pollution* 146, 648–658.

- Flowers, M.D., Fiscus, E.L., Burkey, K.O., Booker, F.L., Dubois, J-J B., 2007. Photosynthesis, chlorophyll fluorescence, and yield of snap bean (*Phaseolus vulgaris* L.) genotypes differing in sensitivity to ozone. *Environmental and Experimental Botany* 61, 190–198.
- Fusaro, L., Gerosa, G., Salvatori, E., Marzuoli, R., Monga, R., Kuzminsky, E., Angelaccio, C., Quarato, D., Fares, S., 2016a. Early and late adjustments of the photosynthetic traits and stomatal density in *Quercus ilex* L. grown in an ozone-enriched environment. *Plant Biology* 18 (S1), 13-21.
- Fusaro, L., Salvatori, E., Manes, F., 2016b. Effects of nitrogen deposition, drought and their interaction, on functional and structural traits of *Fraxinus ornus* L. and *Quercus ilex* L. *Plant Biosystems*, in press, DOI: 10.1080/11263504.2016.1193070.
- Fusaro, L., Salvatori, E., Mereu, S., Silli, V., Bernardini, A., Tinelli, A., Manes, F., 2015. Researches in Castelporziano test site: ecophysiological studies on Mediterranean vegetation in a changing environment. *Rendiconti Lincei* 26(3), 473-481.
- Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R., et al., 2008. Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science* 320, 889–892
- Gerosa, G., 2002. Flussi di ozono in agroecosistemi: approcci ecofisiologici e modellistici. PhD thesis, University of Milan, Italy
- Gerosa, G., Cieslik, S., Ballarin-Denti, A., 2003. Micrometeorological determination of time-integrated stomatal ozone fluxes over wheat: a case study in Northern Italy. *Atmospheric Environment* 37 (6), 777-788.
- Gerosa, G., Derghi, F., Cieslik, S., 2007. Comparison of Different Algorithms for Stomatal Ozone Flux Determination from Micrometeorological Measurements. *Water Air & Soil Pollution* 179, 309-321
- Gerosa, G., Marzuoli, R., Finco, A., Ebone, A., Tagliaferro, F., 2008. Ozone effects on fruit productivity and photosynthetic response of two tomato cultivars in relation to stomatal fluxes. *Italian Journal of Agronomy* 3, 61-70
- Gerosa, G., Marzuoli, R., Finco, A., Monga, R., Fusaro, I., Faoro, F., 2014. Contrasting effects of water salinity and ozone concentration on two cultivars of durum wheat (*Triticum durum* Desf.) in Mediterranean conditions. *Environmental Pollution* 193, 13-21.
- Gerosa, G., Marzuoli, R., Rossini, M., Panigada, C., Meroni, M., Colombo, R., Faoro, F., Iriti M., 2009. A flux-based assessment of the effects of ozone on foliar injury, photosynthesis, and yield of bean (*Phaseolus vulgaris* L. cv. Borlotto Nano Lingua di Fuoco) in open-top chambers. *Environmental Pollution* 157, 1727-1736.
- González-Fernández, I., Calvo, E., Gerosa, G., Bermejo, V., Marzuoli, R., Calatayud, V., Alonso, R., 2014. Setting ozone critical levels for protecting horticultural Mediterranean crops: Case study of tomato. *Environmental Pollution* 185, 178-187.
- Goumenaki, E., Taybi, T., Borland, A., Barnes, J., 2010. Mechanisms underlying the impacts of ozone on photosynthetic performance. *Environmental and Experimental Botany* 69(3), 259-266.
- Guidi, L., Degl'Innocenti, E., Soldatini, G.F., 2002. Assimilation of CO₂, enzyme activation and photosynthetic electron transport in bean leaves, as affected by high light and ozone. *New Phytologist* 156, 377-388.
- Harmens, H., Mills, G., Hayes, F., Norris, D.A., Sharps, K., 2015a. Twenty eight years of ICP Vegetation: an overview of its activities. *Annali di Botanica* 5, 31-43.
- Heagle, A.S., Miller, J.E., Sherril, D., 1994. A white clover system to estimate effects of tropospheric ozone on plants. *Journal of Environmental Quality* 23, 613–621.
- Iriti, M., Belli, L., Nali, C., Lorenzini, G., Gerosa, G., Faoro F., 2006. Ozone sensitivity of currant tomato (*Lycopersicon pimpinellifolium*), a potential bioindicator specie. *Environmental Pollution* 141, 275-282.
- Karlsson, G.P., Karlsson, P.E., Danielsson, H., Pleijel, H., 2003. Clover as a tool for bioindication of phytotoxic ozone—5 years of experience from southern Sweden—consequences for the short-term critical levels. *Sci. Total Environ.* 301, 205-213.

- Lorenzini, G., Nali, C., Biagioni, M., 1995. An analysis of the distribution of surface ozone in Tuscany (central Italy) with the use of a new miniaturized bioassay with ozone-sensitive tobacco seedlings. *Environmental Monitoring and Assessment* 34, 59-72.
- Lorenzini, G., Nali, C., Dota, M.R., Martorana, F., 2000. Visual assessment of foliar injury induced by ozone on indicator tobacco plants: a data quality evaluation. *Environmental Monitoring and Assessment* 62, 175–191.
- Lorenzini, G., Pellegrini, E., Campanella, A., Nali, C., 2014. It's not just the heat and the drought: the role of ozone air pollution in the 2012 heat wave. *Agrochimica*, 58, Special issue: 40-52.
- Lorenzini, G., 1994. A miniaturized kit for ozone biomonitoring. *Appl. Biochem. Biotechnol.* 48, 1-4.
- Manes, F., Altieri, A., Tripodo, P., Booth, C.E., Unsworth, M.H., 1990b. Bioindication study of effects of ambient ozone on tobacco and radish plants using a protectant chemical (EDU). *Annali di Botanica* 48, 133-149.
- Manes, F., Federico, R., Cortiello, M., Angelini, R., 1990a. Ozone induced increase of peroxidase activity in Tobacco (*Nicotiana tabacum* L. cv. Burley 21) leaves. *Phytopathologia Mediterranea* 29, 101-106.
- Manes, F., De Santis, F., Giannini, M.A., Vazzana, C., Capogna, F., Allegrini, I., 2003. Integrated ambient ozone evaluation by passive samplers and clover biomonitoring mini-stations. *Sci. Total Environ.* 308, 133–141.
- Manes, F., Vitale, M., Di Traglia, M., 2005. Tropospheric ozone impact on plants and monitoring in natural and urban areas characterized by Mediterranean climate. *Plant Biosystems* 139 (3), 265-278.
- Marzuoli, R., Finco, A., Chiesa, M., Gerosa, G., 2016a. A dose-response relationship for marketable yield reduction of two lettuce (*Lactuca sativa* L.) cultivars exposed to tropospheric ozone in Southern Europe. *Environmental Science and Pollution Research*, doi:10.1007/s11356-016-8224-6.
- Marzuoli, R., Monga, R., Finco, A., Gerosa, G., 2016b. Biomass and physiological responses of *Quercus robur* (L.) young trees during 2 years of treatments with different levels of ozone and nitrogen wet deposition. *Trees* 30, 1995-2010.
- Mereu, S., Gerosa, G., Marzuoli, R., Fusaro, L., Salvatori, E., Finco, A., Spano, D., Manes, F., 2011. Gas exchange and JIP-test parameters of two Mediterranean maquis species are affected by sea spray and ozone interaction. *Environmental and Experimental Botany* 73, 80-88.
- Mereu, S., Gerosa, G., Finco, A., Fusaro, L., Muys, B., Manes, F., 2009. Improved sapflow methodology reveals considerable night-time ozone uptake by Mediterranean species. *Biogeosciences* 6, 3151–3162.
- Mills, G., Pleijel, H., Braun, S., Büker, P., Bermejo, V., Calvo, E., Danielsson, H., Emberson, L., González Fernández, I., Grünhage, L., Harmens, H., Hayes, F., Karlsson, P-E., Simpson, D., 2011. New stomatal flux-based critical levels for ozone effects on vegetation. *Atm. Envir.* 45, 5064-5068.
- Monga, R., 2015. Physiologic and ecological drivers and agronomical consequences of the ozone like syndrome in wheat. PhD thesis, University of Milan, Italy
- Monga, R., Marzuoli, R., Alonso, R., Bermejo, V., González-Fernández, I., Faoro, F., Gerosa, G., 2015. A varietal screening of different durum wheat cultivars (*Triticum durum*, Desf.) exposed to high levels of ozone in Open-Top Chambers. *Atmospheric Environment* 110,18-26
- Nali, C., Balducci, E., Frati, L., Paoli, L., Loppi, S., Lorenzini, G., 2007. Integrated biomonitoring of air quality with plants and lichens: A case study on ambient ozone from central Italy. *Chemosphere* 67 (2007) 2169–2176.
- Nali, C., Crocicchi, L., Lorenzini, G., 2004. Plants as indicators of urban air pollution (ozone and trace elements) in Pisa, Italy. *Journal of Environmental Monitoring*, 636-645.
- Nali, C., Francini, A., Lorenzini, G., 2006. Biological monitoring of ozone: the twenty-year Italian experience. *J. Environ. Monit* 8, 25–33.
- Nali, C., Lorenzini, G., 2007. Air Quality Survey Carried Out by Schoolchildren: An Innovative Tool for Urban Planning. *Environmental Monitoring and Assessment* 131(1), 201–210
- Nali, C., Pucciarello, C., Lorenzini, G., 2002. Ozone distribution in central Italy and its effect on crop productivity. *Agriculture, Ecosystems and Environment* 90, 277–289.

- Nali, C., Francini, A., Lorenzini, G., 2009. White clover clones as a cost-effective indicator of phytotoxic ozone: 10 years of experience from central Italy. *Environ. Pollut.* 157, 1421-1426.
- Ochoa-Hueso, R., Allen, E. B., Branquinho, C., Cruz, C., Dias, T., Fenn, M. E., Stock, W. D., 2011. Nitrogen deposition effects on Mediterranean-type ecosystems: an ecological assessment. *Environmental Pollution*, 159(10), 2265-2279.
- Oukarroum, A., Schansker, G., Strasser, R.J., 2009. Drought stress effects on Photosystem I content and Photosystem II thermotolerance analyzed using Chl a fluorescence kinetics in barley varieties differing in their drought tolerance. *Physiologia Plantarum* 137, 188–199.
- Pellegrini, E., 2014. PSII photochemistry is the primary target of oxidative stress imposed by ozone in *Tilia americana*. *Urban Forestry & Urban Greening* 13, 94–102.
- Pellegrini, E., Bertuzzi, S., Candotto Carniel, F., Lorenzini, G., Nali, C., Tretiach, M., 2014a. Ozone tolerance in lichens: a possible explanation from biochemical to physiological level using *Flavoparmelia caperata* as test organism. *J. Pl. Physiol.*, 171: 1514-1523.
- Pellegrini, E., Campanella, A., Lorenzini, G., Nali, C., 2014b. Biomonitoring of ozone: a tool to initiate the young people into the scientific method and environmental issues. A case study in Central Italy. *Urban Forestry & Urban Greening*, 13: 800-805.
- Pellegrini, E., Campanella, A., Paolucci, M., Trivellini, A., Gennai, C., Muganu, M., Nali, C., Lorenzini, G., 2015a. Functional leaf traits and diurnal dynamics of photosynthetic parameters predict the behavior of grapevine varieties towards ozone. *PLoS ONE*, 10 (8): e0135056. doi: 10.1371/journal.pone.0135056
- Pellegrini, E., Carucci, M.G., Campanella, A., Lorenzini, G., Nali, C., 2011. Ozone stress in *Melissa officinalis* plants assessed by photosynthetic function. *Env. Exp. Bot.*, 73: 94-101
- Pellegrini, E., Francini, A., Lorenzini, G., Nali, C., 2015b. Ecophysiological and antioxidant traits of *Salvia officinalis* under ozone stress. *Environ. Sci. Pollut. Res.* 22: 13083-13093.
- Pellegrini, E., Nali, C., Lorenzini, G., 2013a. Ecophysiology of *Tilia americana* under ozone fumigation. *Atmospheric Pollution Research*, 4: 142-146
- Pellegrini, E., Trivellini, A., Campanella, A., Francini, A., Lorenzini, G., Nali, C., Vernieri, P., 2013b. Signaling molecules and cell death in *Melissa officinalis* plants exposed to ozone. *Plant Cell Reports*, 32: 1965-1980.
- Phoenix, G.K., Hicks, W.K., Cinderby, S., Kuylenstierna, J.C., Stock, W.D., Dentener, F.J., et al., 2006. Atmospheric nitrogen deposition in world biodiversity hotspots: The need for a greater global perspective in assessing N deposition impacts. *Glob Change Biol* 12: 470–476.
- Pollastrini, M., Desotgiu, R., Camin, F., Ziller, L., Gerosa, G., Marzuoli, R., Bussotti, F., 2014. Severe drought events increase the sensitivity to ozone on poplar clones. *Environmental and Experimental Botany* 100, 94-104.
- Salvatori E., Fusaro L. Strasser R.J., Bussotti F., Manes F., 2015. Effects of acute O₃ stress on PSII and PSI photochemistry of sensitive and resistant snap bean genotypes (*Phaseolus vulgaris* L.), probed by Prompt Chlorophyll “a” fluorescence and 820 nm Modulated Reflectance. *Plant Physiology and Biochemistry* 97, 368-377.
- Salvatori, E., Fusaro, L., Mereu, S., Bernardini, A., Puppi, G., Manes, F., 2013. Different O₃ response of sensitive and resistant snap bean genotypes (*Phaseolus vulgaris* L.): The key role of growth stage, stomatal conductance, and PSI activity. *Environmental and Experimental Botany* 87, 79-91.
- Scalet, M., Federico, R., Guido, M.C., Manes F., 1995. Peroxidase activity and polyamine changes in response to ozone and simulated acid rain in Aleppo pine needles. *Environmental and Experimental Botany* 35, 417-425
- Schansker, G., Tóth, S.Z., Strasser, R.J., 2005. Methylviologen and dibromothymoquinone treatments of pea leaves reveal the role of photosystem I in the Chl a fluorescence rise OJIP. *Biochimica et Biophysica Acta* 1706, 250–261.
- Sicard, P., De Marco, A., Troussier, F., Renou, C., Vas, N., Paoletti, E., 2013. Decrease in surface ozone concentrations at Mediterranean remote sites and increase in the cities. *Atmospheric Environment* 79, 705-715.

- Strasser, R.J., Tsimilli-Michael, M., Qiang, S., Goltsev, V., 2010. Simultaneous in vivo recording of prompt and delayed fluorescence and 820-nm reflection changes during drying and after rehydration of the resurrection plant *Haberlea rhodopensis*. *Biochimica et Biophysica Acta* 179, 1313–1326.
- Strasser, R.J., Tsimilli-Michael, M., Srivastava, A., 2004. Analysis of the fluorescence transient. In: Papageorgiou, G., Govindjee, H. (Eds.), *Chlorophyll Fluorescence: A Signature of Photosynthesis*. Kluwer Academic Publishers, The Netherlands, p. 321–362.
- Tattini, M., Loreto, F., 2014. Plants in Mediterranean areas: “Living in the sun”. *Environ Exp Bot* 103: 1–2.
- The European Nitrogen Assessment: Sources, Effects and Policy Perspectives, 2011. Edited by Sutton M.A., Howard C.E., Erisman J.W., Billen G., Bleeker A., Grennfelt P., van Grinsven H., Grizzetti B. Cambridge University Press, Cambridge, UK.
- Tonelli, M., Pellegrini, E., D’Angiolillo, F., Petersen, M., Nali, C., Pistelli, L., Lorenzini G., 2015. Ozone-elicited secondary metabolites in shoot cultures of *Melissa officinalis* L. *Plant Cell, Tissue and Organ Culture*, 120: 617-629.
- Tuovinen, J.-P., Ashmore, M., Emberson, L., Simpson, D., 2004. Testing and improving the EMEP ozone deposition module. *Atmospheric Environment* 38, 2373-2385
- Tuovinen, J.-P., Simpson, D., Emberson, L., Ashmore, M., Gerosa, G., 2007. Robustness of modelled ozone exposures and doses. *Environmental Pollution* 146 (3), 578-586
- UN/ECE, 2010. Manual on the methodologies and criteria for modelling and mapping critical loads and levels and air pollution effects, risks and trends. Revision 2015. Available at: http://icpmapping.org/Mapping_Manual
- UN/ECE, 2015a. Chapter 3 of the Mapping Manual, “Mapping Critical Levels for Vegetation”, Revision June 2015. Available at: http://www.rivm.nl/media/documenten/cce/manual/binnenop17Juni/Ch3-MapMan-2016-05-03_vf.pdf
- UN/ECE, 2015b. Chapter 5 of the Mapping Manual, “MAPPING CRITICAL LOADS FOR ECOSYSTEMS”, Revision June 2015. Available at: <http://www.rivm.nl/media/documenten/cce/manual/Ch5-MapMan-2015-08-24.pdf>
- Vahisalu, T., Puzõrjova, I., Brosché, M., Valk, E., Lepiku, M., Moldau, H., Pechter, P., Wang, Y-S., Lindgren, O., Salojärvi, J., Loog, M., Kangasjärvi, J., Kollist, H., 2010. Ozone-triggered rapid stomatal response involves the production of reactive oxygen species, and is controlled by SLAC1 and OST1. *The Plant Journal* 62, 442-453.
- Vainonen, J., Kangasjärvi, J., 2015. Plant signalling in acute ozone exposure. *Plant, Cell and Environment* 38, 240–252.
- Valletta, A., Salvatori, E., Santamaria, A. R., Nicoletti, M., Toniolo, C., Caboni, E., Bernardini, A., Pasqua, G., Manes, F., 2016. Ecophysiological and phytochemical response to ozone of wine grape cultivars of *Vitis vinifera* L. *Natural Product Research* 30,22,
- Vitale, M., Salvatori, E., Loreto, F., Fares, S., Manes, F., 2008. Physiological responses of *Quercus ilex* leaves to water stress and acute ozone exposure under controlled conditions. *Water, Air and Soil Pollution*, 189, 113-125.

CHAPTER 6 - BIODIVERSITY AS AN IMPORTANT INDICATOR OF SOIL ACIDITY AND EUTROPHICATION: THE ROLE OF THE MODELLING IN PRESERVING IT.

- Attorre, F., Maggini, A., Di Traglia, M., De Sanctis, M., & Vitale, M., 2013. A methodological approach for assessing the effects of disturbance factors on the conservation status of Mediterranean coastal dune systems. *Applied Vegetation Science*, 16(2), 333-342.
- Belyazid, S. Westling, O., Sverdrup, H., 2006. Modelling and changes in forest soil chemistry at 16 Swedish coniferous forest sites following deposition reduction. *Environmental Pollution*, 144, 596-609.
- Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L., De Vries, W., 2010. Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Appl.* 20, 30-59.
- Bonten, T.C., Groenenberg, J.E., Meesenburg, H., De Vries, W., 2001/2011. Using advanced surface complexation models for modelling soil chemistry under forests: Solling forest, Germany. *Environmental Pollution*, 159, 2831-2839.

- Brumme, R. and Khanna, P. K. 2008. Ecological and site historical aspects of N dynamics and current N status in temperate forests. *Global Change Biology* 14, 125–141.
- Cosby, B.J., Ferrier, R.C., Jenkins, A., Wright, R.F., 2001. Modelling the effects of acid deposition: Refinements, adjustments and inclusion of nitrogen dynamics in the MAGIC model. *Hydrology and Earth System Sciences*, 5, 499-517.
- Cosby, B.J., Hornberger, G.M., Galloway, J.N., Wright, R.F., 1985. Modelling the effects of acid deposition: Assessment of a lumped parameter model of soil water and streamwater chemistry. *Water Resources Research*, 21, 51-63.
- De Luca, E., Novelli, C., Barbato, F., Menegoni, P., Iannetta, M., Nascetti, G., 2011. Coastal dune systems and disturbance factors: monitoring and analysis in central Italy. *Environmental Monitoring and Assessment* 183: 437–50.
- EEA, 2007. Air pollution in Europe 1990–2004. EEA Report, No. 2/2007.
- Ellenberg, H., 1979. Zeigerwerte von Gefäßpflanzen Mitteleuropas. *Scripta Geobotanica* 9, 1-122.
- Ellenberg, H., 1988. *Vegetation ecology of central Europe*. Cambridge University Press.
- Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W., Paulissen, D., 1991. Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica* 18, 1-248.
- Grunewald, R., Schubert, H., 2007. The definition of a new plant diversity index “HDune” for assessing human damage on coastal dunes – derived from the Shannon index of entropy H’. *Ecological Indicators* 7: 1–21.
- Lindner, M., Maroschek, M., Netherer, S., Kremer, A., Barbati, A., Garcia-Gonzalo, J., ..., Lexer, M. J., 2010. Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecology and Management*, 259(4), 698-709.
- Martinez, M.L., Psuty, N.P., Lubke, R.A., 2004. A perspective on coastal dunes. In: Martinez, M.L. & Psuty, N.P. (eds.) *Coastal dunes. Ecology and conservation*, pp. 3–10. Springer, Heidelberg, DE.
- Matyssek, R., Le Thiec, D., Löw, M., Dizengremel, P., Nunn, A.J., Häberle, K.H., 2006. Interactions between drought and O₃ stress in forest trees. *Plant Biology* 8, 11–17.
- Nillson, J. & Grennfelt, P., 1988. Critical loads for sulphur and nitrogen. *Miljörapport 1988: 15*; Nordic Council of Ministers, Copenhagen.
- Pielou, E. C., 1966. The measurement of diversity in different types of biological collections. *Journal of theoretical biology*, 13, 131-144.
- Posch, M., Reinds, G.J., 2009. A very simple dynamic soil acidification model for scenario analyses and target load calculations. *Environmental Modelling and Software*, 24, 329-340.
- Rehfuess, K.E., Ågren, G.I., Andersson, F., et al., 1999. Relationships between recent changes of growth and nutrition of Norway spruce, Scots pine and European beech forests in Europe-RECOGNITION. Working Paper 19, European Forest Institute, Joensuu, Finland, 94pp.
- Reinds, G.J., Posch, M., De Vries, W., 2001. A semi-empirical dynamic soil acidification model to use in spatially explicit integrated assessment models for Europe. (Alterra rapport84). Wageningen (Netherlands): Alterra.
- Shannon, C.E. and Weaver, W. (1949) *The Mathematical Theory of Communication*. University of Illinois Press, Urbana.
- Sutton, M.A., Simpson, D., Levy, P.E., Smith, R.I., Reis, S., Van Oijen, M. and de Vries, W. (2008), Uncertainties in the relationship between atmospheric nitrogen deposition and forest carbon sequestration. *Global Change Biology*, 14: 2057–2063.

CHAPTER 7 - The contribution of Italy to the ICP WATERS Programme

- Bobbink R., Hettelingh, J.-P. (eds.), 2011. Review and revision of empirical critical loads and dose-response relationships. Proceedings of an Expert Workshop, Noordwijkerhout, 23-25 June 2010. Bilthoven, National Institute for Public Health and the Environment (RIVM): 243 pp.
- Cosby, B.J., R.C. Ferrier, A. Jenkins & R.F. Wright., 2001. Modelling the effects of acid deposition – fifteen years of MAGIC: refinements, adjustments and inclusion of nitrogen dynamics. *Hydrol. Earth System Sci.*, 5: 499-517.
- de Wit, H., Lindholm, M., 2010. Nutrient enrichment effects of atmospheric N deposition on biology in oligotrophic surface waters - a review. ICP Waters report 101/2010: 39 pp.
- Escudero, C., 2015. Intercomparison 1529: pH, Conductivity, Alkalinity, NO₃-N, Cl, SO₄, Ca, Mg, Na, K, TOC, Al, Fe, Mn, Cd, Pb, Cu, Ni, and Zn. ICP Waters report 123/2015: 56 pp.
- Evans, C.D., Cullen, J.M., Alewell, C., Kopáček, J., Marchetto, A., Moldan, F., Prechtel, A., Rogora, M., Veselý, J. and Wright, R.F., 2001. Recovery from acidification in European surface waters. *Hydrol. Earth System Sci.*, 5, 283-297.
- Fjellheim, A., Johannessen, A., Landås, T., 2015. Biological intercalibration. Invertebrates 1915 - ICP Waters report 124/2015: 31 pp.
- Garmo, Ø.A., Skjelkvåle, B.L., de Wit, H.A., Colombo, L., Curtis, C., Fölster, J., Hoffmann, A., Hruška, J., Høgåsen, T., Jeffries, D.S., Keller, W., Krám, P., Majer, V., Monteith, D.T., Paterson, A., Rogora, M., Rzychon, D., Steingruber, S., Stoddard, J.L., Vuorenmaa, J., Worsztynowicz, A., 2014. Trends in Surface Water Chemistry in Acidified Areas in Europe and North America from 1990 to 2008. *Water Air Soil Pollut.* 225: 1880.
- Guilizzoni, P., Marchetto, A., Lami, A., Cameron, N.G., Appleby, P.G., Rose, N.L., Schnell, Ø. A., Belis, C.A., Giorgis, A., Guzzi, L., 1996. The environmental history of a mountain lake (Lago Paione Superiore, Central Alps, Italy) for the last c. 100 years: a multidisciplinary, paleolimnological study. *J. Paleolimnol.*, 15: 245-264.
- Helliwell, R., Wright, R., Jackson-Blake, L., Ferrier, R., Aherne, J., Cosby, B., Evans, C., Forsius, M., Hruska, J., Jenkins, A., Krám, P., Kopacek, J., Majer, V., Moldan, F., Posch, M., Potts, J., Rogora, M., Schoepp, W., 2014. Assessing recovery from acidification of European surface waters in the year 2010: An evaluation of projections made with the MAGIC model in 1995. *Envir. Sci. Technol.* 48: 13280–13288.
- Marchetto, A., Mosello, R., Psenner, R., Barbieri, A., Bendetta, G., Tait, D., Tartari, G. A., 1994. Evaluation of the level of acidification and the critical loads for Alpine lakes *Ambio.* 23: 150-154.
- Marchetto, A., Mosello, R., Rogora, M., Manca, M., Boggero, A., Morabito, G., Musazzi, S., Tartari, G.A., Nocentini, A.M., Pugnetti, A., Bettinetti, R., Panzani, P., Armiraglio, M., Cammarano, P., Lami, A., 2004. The chemical and biological response of two remote mountain lakes in the Southern Central Alps (Italy) to twenty years of changing physical and chemical climate. *J. Limnol.* 63: 77-89.
- Meteo Svizzera. (2012). Rapporto sul clima – Cantone Ticino 2012. Rapporto di lavoro Meteo Svizzera: 63 pp.
- Mosello, R., Marchetto, A., Brizzio, M.C., Rogora, M., Tartari, G.A., 2000. Results from the Italian Participation in the International Co-operative programme on Assessment and Monitoring of Acidification of Rivers and Lakes (ICP Waters). *J. Limnol.*: 59: 47-54. Climate control on sulphate and nitrate concentrations in alpine streams of Northern Italy along a nitrogen saturation gradient.
- Raddum, G. G., Skjelkvåle, B. L., 2001. Critical Limit of Acidifying Compounds to Invertebrates in Different Regions of Europe. *Water Air Soil Pollut.*, 130, 825-830.
- Rogora, M., Marchetto, A., Mosello, R., 2001. Trends in the chemistry of atmospheric deposition and surface waters in the Lago Maggiore watershed. *Hydrol. Earth System Sci.*, 5: 379-390.
- Rogora, M., Mosello, R., Arisci, S., 2003a. The effect of climate warming on the hydrochemistry of alpine lakes. *Water Air Soil Poll.*, 148: 347-361.
- Rogora, M., Marchetto, A., Mosello, R., 2003b. Modelling the effects of the deposition of acidity and nitrogen on selected lakes and streams in Central Alps (Italy). *Hydrol. Earth System Sci.* 7: 540-551.

- Rogora, M., 2004. Acidification and recovery at mountain lakes in Central Alps assessed by the MAGIC model. *J. Limnol.* 63: 133-142.
- Rogora, M., Mosello, R., Arisci, S., Brizzio, M.C., Barbieri, A., Balestrini, R., Waldner, P., Schmitt, M., Stähli, M., Thimonier, A., Kalina, M., Puxbaum, H., Nickus, U., Ulrich, E., Probst, A., 2006a. An overview of atmospheric deposition chemistry over the Alps: present status and long-term trends. *Hydrobiologia* 562: 17-40.
- Rogora, M., Mosello, R., Calderoni, A., Barbieri, A., 2006b. Nitrogen budget of a subalpine lake in North-Western Italy: the role of atmospheric input in the upward trend of nitrogen concentrations. *Verh. Internat. Verein. Limnol.*, 29 (4): 2027-2030.
- Rogora, M., 2007. Synchronous trends in N-NO₃ export from N-saturated river catchments in relation to climate. *Biogeochemistry* 86: 251–268.
- Rogora, M., Mosello, R., 2007. Climate as a confounding factor in the response of surface water to nitrogen deposition in an area South of the Alps. *Applied Geochem.* 22: 1122-1128.
- Rogora, M., Massafiero, J., Marchetto, A., Tartari, G.A., Mosello, R., 2008a. The water chemistry of Northern Patagonian lakes and their nitrogen status in comparison with remote lakes in different regions of the globe. *J. Limnol.* 67 (2): 75-86.
- Rogora, M., Arese, C., Balestrini, R., Marchetto, A., 2008b. Climate control on short- and long-term variations of sulphate and nitrate concentrations in alpine streams of Northern Italy along a N saturation gradient. *Hydrol. Earth Syst. Sci.*, 12: 371–381.
- Rogora, M., Arisci, S., Marchetto, A., 2012. The role of nitrogen deposition in the recent nitrate decline in lakes and rivers in Northern Italy. *Science of the Total Environment* 417-418C: 219-228.
- Rogora, M., Colombo, L., Lepori, F., Marchetto, A., Steingruber, S., Tornimbeni, O., 2013. Thirty years of chemical changes in alpine acid-sensitive lakes in the Alps. *Water Air Soil Pollut.* 224:1746.
- Rogora, M., Colombo, L., Marchetto, A., Mosello, R., Steingruber, S., 2016. Temporal and spatial patterns in the chemistry of wet deposition in Southern Alps. *Atm. Envir.* 146: 44-54.
- Romano D., Bernetti, A., Córdor, R.D., De Lauretis, R., Di Cristofaro, E., Lena, F., Gagna, A., Gonella, B., Pantaleoni, M., Peschi, E., Taurino, E., Vitullo, M., 2014. Italian Emission Inventory 1990-2012. Informative Inventory Report. Rome, Institute for Environmental Protection and Research (2014): 157 pp.
- Skjelkvåle, B.L., Stoddard, J., Jeffries, D., Tørseth, K., Høgåsen, T., Bowman, J., Mannio, J., Monteith, D., Mosello, R., Rogora, M., Rzychon, D., Vesely, J., Wieting, J., Wilander A., Worsztynowicz, A., 2005. Regional scale evidence for improvements in surface water chemistry 1990-2001. *Environ. Pollut.* 137: 165-76.
- Steingruber, S., 2015. Deposition of Acidifying and Eutrophying Pollutants in Southern Switzerland from 1988 to 2013. *Bollettino della Società ticinese di scienze naturali*, 103: 37-45.
- Stoddard J. L., Jeffries, D. S., Lükewille, A., Clair, T. A., Dillon, P. J., Driscoll, C. T., Forsius, M., Johannessen, M. J., Kahl, S., Kellogg, J. H., Kemp, A., Mannio, J., Monteith, D. T., Murdoch, P. S., Patrick, S., Rebsdorf, A., Skjelkvåle, B. L., Stainton, M. P., Traaen, T., van Dam, H., Webster, K. E., Wieting, J., Wilander, A., 1999. Regional trends in aquatic recovery from acidification in North America and Europe. *Nature* 401, 575-578.
- Thies, H., Nickus, U., Mair, V., Tessadri, R., Tait, D., Thaler, B., Psenner, R., 2007. Unexpected Response of High Alpine Lake Waters to Climate Warming. *Environ. Sci. Technol.*, 41 (21), 7424–7429.
- Tornimbeni, O., Rogora, M., 2012. An evaluation of trace metals in high altitude lakes of the Central Alps: present levels, origins and possible speciation in relation to pH values. *Wat. Air Soil Pollut.* 223: 1895-1909.
- Wright, R. F., Camarero, L., Cosby, B.J., Ferrier, R. C., Forsius, M., Helliwell, R., Jenkins, A., Kopacek, J., Majer, V., Moldan, F., Posch, M., Rogora, M., Schöpp, W., 2005. Recovery of acidified European surface waters. *Environ. Sci. Technol.*, 39: 64A-74A.
- Wright, R.F., Aherne, J., Bishop, K., Camarero, L., Cosby, B.J., Erlandsson, M., Evans, C.D., Hardekopf, D., Helliwell, R., Hruska, J., Jenkins, A., Moldan, F., Posch, M., Rogora, M., 2006. Modelling the effect of climate change on

recovery of acidified freshwaters: sensitivity of individual processes in the MAGIC and SMART models. *Sci. Tot. Environ.* 365: 154-166.

Wright R.F., Jenkins, A., 2001. Climate change as a confounding factor in reversibility of acidification: RAIN and CLIMEX projects. *Hydrol Earth Syst Sc* 5: 477–486.

Wright, R.F., Helliwell, R., Hruska, J., Larssen, T., Rogora, M., Rzychoń, D., Skjelkvåle, B.L., Worsztynowicz, A., 2011. Impacts of Air Pollution on Freshwater Acidification under Future Emission Reduction Scenarios; ICP Waters contribution to WGE report. ICP Waters report 108/2011: 27 pp.

CHAPTER 8 - EFFECTS OF AIR POLLUTION ON TECHNICAL MATERIALS AND CULTURAL HERITAGE. THE CONTRIBUTION OF ITALY TO ICP MATERIALS

Doytchinov, S., Screpanti, A., Leggeri, G., 2009. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 60: Combined stock at risk and mapping for selected urban areas of Italy. Italian National Agency for New Technologies, Energy and the Environment (ENEA), Rome, Italy.

Doytchinov, S., Screpanti, A., Leggeri, G., 2010. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 63: Combined stock at risk and mapping for Italy at the national level. Italian National Agency for New Technologies, Energy and the Environment (ENEA), Rome, Italy.

Doytchinov, S., Screpanti, A., Leggeri, G., Varotsos, C., 2011. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 68: Pilot study on inventory and condition of stock of materials at risk at United Nations Educational, Scientific and Cultural Organization (UNESCO) cultural heritage sites. Part I Methodology. Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy.

Doytchinov, S., Screpanti, A., Leggeri, G., 2012. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 70: Pilot study on inventory and condition of stock of materials at risk at United Nations Educational, Scientific and Cultural Organization (UNESCO) cultural heritage sites. Part II Determination of stock of materials at risk for individual monuments. Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy.

Doytchinov, S., Spezzano, P., Screpanti, A., Leggeri, G., 2014. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 73: Pilot study on inventory and condition of stock of materials at risk at United Nations Educational, Scientific and Cultural Organization (UNESCO) cultural heritage sites. Part III Economic evaluation. Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy.

Spezzano, P., Screpanti, A., Di Benedetto, E., Leggeri, G., 2015. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 77: Pilot study on inventory and condition of stock of materials at risk at United Nations Educational, Scientific and Cultural Organization (UNESCO) cultural heritage sites. Part IV The relationship between the environment and the artefact. Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Rome, Italy.

Tidblad, J., Kucera, V., Mikhailov, A.A., 1998. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 30: Statistical analysis of 8 year materials exposure and acceptable deterioration and pollution levels, Swedish Corrosion Institute, Stockholm, Sweden.

Tidblad, J., Yates, T., Doytchinov, S., Faller, M., Kreislova, K., 2010a. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 61: Assessment of stock of materials at risk including cultural heritage, Swerea KIMAB AB, Stockholm, Sweden.

Tidblad, J., Faller, M., Grøntoft, T., Kreislova, K., Varotsos, C., de la Fuente, D., Lombardo, T., Doytchinov, S., Brüggerhoff, S., Yates, T., 2010b. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 65: Economic assessment of corrosion and soiling of materials including cultural heritage, Swerea KIMAB AB, Stockholm, Sweden.

Tidblad, J., Grøntoft, T., Kreislova, K., Faller, M., de la Fuente, D., Yates, T., Verney-Carron, A., 2014. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 76: Trends in pollution, corrosion and soiling 1987-2012. Swerea KIMAB AB, Stockholm, Sweden.

Tidblad, J., Kreislova, K., Faller, M., de la Fuente, D., Yates, T., Verney-Carron, A., 2016. “UNECE international co-operative programme on effects on materials, including historic and cultural monuments”. Report no. 78: Results of corrosion and soiling from the 2011-2015 exposure programme for trend analysis. Swerea KIMAB AB, Stockholm, Sweden.

UNECE, 2015. Mapping of Effects on Materials. Updated 25 August 2015. Chapter IV of Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution effects, risks and trends. Convention on Long-range Transboundary Air Pollution. Accessed on 28 June 2016 on Web at www.icpmapping.org.

Watt, J., Doytchinov, S., Lefevre, R., Ionescu, A., de la Fuente, D., Kreislova, K., Screpanti, A., 2009. Stock at Risk. In: J. Watt, J. Tidblad, V. Kucera, and R. Hamilton (Eds), *The Effects of Air Pollution on Cultural Heritage* (pp. 147-187), Springer.

CHAPTER 9 - VEGETATION AND URBAN AIR QUALITY: RECENT FINDINGS

Amorim, J.H., Rodrigues, V., Tavares, R., et al., 2013. CFD modelling of the aerodynamic effect of trees on urban air pollution dispersion. *Sci Total Environ* 461–462(0): 541-551.

Bartra, J., Mullol, J., del Cuvillo, A., Dávila, I., Ferrer, M., Jáuregui, I., Montoro, J., Sastre, J., Valero, A., 2007. Air Pollution and Allergens. *J Investig Allergol Clin Immunol* 17(2): 3-8.

Beck, I., Jochner, S., Gilles, S., et al., 2013. High environmental ozone levels lead to enhanced allergenicity of birch pollen. *PLoS ONE* 8(11): e80147.

Benjamin MT and Winer AM. 1998. Estimating the ozone-forming potential of urban trees and shrubs. *Atmos Environ* 32(1): 53-68.

Bottalico, F., Chirici, G., Giannetti, F., De Marco, A., Nocentini, S., Paoletti, E., Salbitano, F., Sanesi, G., Serenelli, C., Travaglini, D., 2016. Air pollution removal by green infrastructures and urban forests in the city of Florence. *Agriculture and Agricultural Science Procedia*, 8: 243–251.

Calfapietra, C., Fares, S., Manes, F., et al., 2013. Role of biogenic volatile organic compounds (BVOC) emitted by urban trees on ozone concentration in cities: A review. *Environ Pollut* 183: 71-80.

Calfapietra, C., Morani, A., Sgrigna, G., Di Giovanni, S., Muzzini, V., Pallozzi, E., Guidolotti, G., Nowak, D., Fares, S., 2016. Removal of ozone by urban and peri-urban forests: Evidence from laboratory, field, and modeling approaches. *Journal of Environmental Quality* 45, 224-233.

Cariñanos, P., Adinolfi, C., Díaz de la Guardia, C., et al., 2016. Characterization of allergen-emission sources in urban areas. *J Environ Qual* 45: 244-252.

Churkina, G., Grote, R., Butler, T.M., et al., 2015. Natural selection? Picking the right trees for urban greening. *Environ Sci Policy* 47: 12-17.

Dzierzanowski, K., Popek, R., Gawronska, H., et al., 2011. Deposition of particulate matter of different size fractions on leaf surfaces and in waxes of urban forest species. *Int J Phytoremediation* 13(10): 1037-1046.

EFA, 2011. *EFA Book on Respiratory Allergies - Raise Awareness, Relieve the Burden*. European Federation of Allergy and Airway Diseases Patients Association (EFA). Valovirta E. (Ed.). Italy. 59 p.

Fares, S., McKay, M., Holzinger, R., Goldstein, A.H., 2010. Ozone fluxes in a *Pinus ponderosa* ecosystem are dominated by non-stomatal processes: Evidence from long-term continuous measurements. *Agric. For. Meteorol.* 150, 420–431.

Fares, S., Savi, F., Fusaro, L., Conte, A., Salvatori, E., Aromolo, R., Manes, F., 2016. Particle deposition in a peri-urban Mediterranean forest. *Environmental Pollution*, 2018, 1278-1286.

- Forsberg, B., Bråbäck, L., Keune, H., Kobernus, M., von Krauss, M.K., Yang, A., Bartonova, A., 2012. An expert assessment on climate change and health - with a European focus on lungs and allergies. *Environ Health* 11 Suppl 1:S4.
- Fusaro, L., Salvatori, E., Mereu, S., Marando, F., Scassellati, E., Abbate, G., Manes, F., 2015. Urban and peri-urban forests in the metropolitan area of Rome: ecophysiological response of *Quercus ilex* L. in two Green Infrastructures in an Ecosystem Services perspective. *Urban For. Urban Gree.* 14, 1147–1156.
- Gromke, C., Blocken, B., 2015. Influence of avenue-trees on air quality at the urban neighborhood scale. Part II: Traffic pollutant concentrations at pedestrian level. *Environmental Pollution* 196, 176-184.
- Grote, R., Samson, R., Alonso, R., Amorim, J.U., Calfapietra, C., Cariñanos, P., Churkina, G., Fares, S., Le Thiec, D., Niinemets, U., Mikkelsen, T.N., Paoletti, E., Tiwary, A., 2016. Functional traits of urban trees in relation to their air pollution mitigation potential: A holistic discussion. *Frontiers in Ecology and the Environment*, in press.
- Guidolotti, G., Salviato, M., Calfapietra, C., 2016 Comparing estimates of EMEP MSC-W and UFORE models in air pollutant reduction by urban trees *Environmental Science and Pollution Research* July, in press.
- Holzinger, R., Lee, A., McKay, M., Goldstein, A.H., 2006. Seasonal variability of monoterpene emission factors for a *Ponderosa* pine plantation in California, *Atmospheric Chemistry and Physics*, 6, 1267-1274.
- Irga, P.J., Burchett, M.D., Torpy, F.R., 2015. Does urban forestry have a quantitative effect on ambient air quality in an urban environment? *Atmospheric Environment* 120, 173-181.
- Kardel, F., Wuyts, K., Babanezhad, M., et al., 2012. Tree leaf wettability as passive bio-indicator of urban habitat quality. *Environ Exp Bot* 75: 277-285.
- Karl, M., Guenther, A., Köble, R., et al., 2009. A new European plant-specific emission inventory of biogenic volatile organic compounds for use in atmospheric transport models. *Biogeosciences* 6(6): 1059-1087.
- Manes, F., Chirici, G., Munafò, M., Marando, F., Capotorti, G., Blasi, C., Salvatori, E., Fusaro, L., Ciancarella, L., Mircea, M., Marchetti, M., 2016. Regulating Ecosystem Services of forests in ten Italian Metropolitan Cities: Air quality improvement by PM₁₀ and O₃ removal. *Ecological Indicators* 67, 425-440.
- Manes, F., Incerti, G., Salvatori, E., Vitale, M., Ricotta, C., Costanza, R., 2012. Urban ecosystem services: Tree diversity and stability of tropospheric ozone removal *Ecological Applications* 22, 349-360.
- Manes, F., Silli, V., Salvatori, E., Incerti, G., Galante, G., Fusaro, L., Perrino, C., 2014 Urban ecosystem services: Tree diversity and stability of PM₁₀ removal in the metropolitan area of Rome. *Annali di Botanica* 4, 19-26.
- Marando, F., Salvatori, E., Manes, F., 2016. Removal of PM₁₀ by Forests as a Nature-Based Solution for Air Quality Improvement in the Metropolitan City of Rome. *Forests* 7(7), 150.
- Nowak, D.J., Crane, D.E., 1998. The Urban Forest Effects (UFORE) model: quantifying urban forest structure and functions. In Hansen M, and Burk T. *Integrated tools for natural resources inventories in the 21st century*. St. Paul. 714-720.
- Nowak, D., Jovan, S., Branquinho, C., Augusto, S., Ribeiro, M.C., Kretsch, C.E. 2015. Chapter 4: Biodiversity, air quality and human health. In: Romanelli, C., Cooper, D., Campbell-Lendrum, D., Maiero, M., Karesh, W.B., Hunter, D., Golden, C.D. *Connecting Global Priorities - Biodiversity and Human Health: A State of Knowledge Review*. World Health Organization and Secretariat of the Convention on Biological Diversity. ISBN: ISBN 978 92 4 150853 7, p 63-74. <https://www.cbd.int/health/SOK-biodiversity-en.pdf>
- Nowak, D.J., Hirabayashi, S., Bodine, A., Hoehn, R., 2013. Modeled PM_{2.5} removal by trees in ten U.S. cities and associated health effects. *Environmental Pollution* 178, Pages 395-402.
- Nowak, D.J., 2006 Institutionalizing urban forestry as a “biotechnology” to improve environmental quality *Urban Forestry and Urban Greening* 5, 93-100.
- Paoletti, E., 2006. Impact of ozone on Mediterranean forests: A review. *Environmental Pollution*, 144, 463-474.
- Paoletti, E., 2009. Ozone and urban forests in Italy. *Environmental Pollution*, 157: 1506-1512.
- Paoletti, E., Bardelli, T., Giovannini, G., Pecchioli, L., 2011. Air quality impact of an urban park over time. *Procedia Environmental Sciences*, 4: 10-16.

- Paoletti, E., Karnosky, D.F., Percy, K.E., 2004. Urban trees and air pollution. In: *Forestry Serving Urbanised Societies* (Konijnendijk C.C., Schipperijn J., Hoyer K.K., Eds) IUFRO World Series Vol. 14, pp. 129-154.
- Pascal, M., Corso, M., Chanel, O., Declercq, C., Badaloni, C., Cesaroni, G., Henschel, S., Meister, K., Haluza, D., Martin-Olmedo, P., Medina, S., 2013. Assessing the public health impacts of urban air pollution in 25 European cities: results of the Aphekomp project. *Sci. Total Environ.* 449, 390-400.
- Ren, Y., Ge, Y., Gu, B., Min, Y., Tani, A., Chang, J., 2014. Role of Management Strategies and Environmental Factors in Determining the Emissions of Biogenic Volatile Organic Compounds from Urban Greenspaces. *Environ. Sci. Technol.*, 48 (11), 6237–6246.
- Roy, S., Byrne, J., Pickering, C., 2012. A systematic quantitative review of urban tree benefits, costs, and assessment methods across cities in different climatic zones. *Urban Forestry & Urban Greening* 11(4), 351–363.
- Salmond, J.A., Tadaki, M., Vardoulakis, S., et al., 2013. Health and climate related ecosystem services provided by street trees in the urban environment. *Environmental Health* 15(1): 95-111.
- Sieghardt, M., Mursch-Radlgruber, E., Paoletti, E., Couenberg, E., Dimitrakopoulos, A., Rego, F., Hatzistathis, A., Randrup, T.B., 2005. The abiotic urban environment: Impact of urban growing conditions on urban vegetation. In: *Urban Forests and Trees* (Konijnendijk C.C., Nilsson K., Randrup T.B., Schipperijn J. Eds) Springer-Verlag, Berlin Heidelberg, pp. 281-323.
- Silli, V., Salvatori, E., Manes, F., 2015. Removal of airborne particulate matter by vegetation in an urban park in the city of Rome (Italy): an ecosystem services perspective. *Ann. Bot.* 5, 53–62.
- Terzaghi, E., Wild, E., Zachello, G., Cerabolini, B.E.L., Jones, K.C., Di Guarda, A., 2013. Forest Filters Effects: Role of leaves in capturing/releasing air particulate matter and its associated PAHs. *Atmospheric Environment* 74, 378-384.
- Tiwary, A., Williams, I.D., Heidrich, O., et al., 2016. Development of multi-functional streetscape green infrastructure using a performance index approach. *Environ Pollut* 208A: 209-220.
- Traidl-Hoffman, C., Kashe, A., Menzel, A., Jakob, T., Thiel, M., Ring, J., Behrendt, H., 2003. Impact of pollen on human health: more than allergen carriers? *International Archives of Allergy and Immunology* 13(1): 1-13.
- Vos, P.E.J., Maiheu, B., Vankerkom, J., Janssen, S., 2013. Improving local air quality in cities: To tree or not to tree? *Environmental Pollution* 183, 113-122.
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kaźmierczak, A., Niemela, J., James, P., 2007. Promoting ecosystem and human health in urban areas using green infrastructure: a literature review. *Landscape Urban Plann.* 81, 167–178.
- Wang, H., Shi, H., Li, Y., et al., 2013. Seasonal variations in leaf capturing of particulate matter, surface wettability and micromorphology in urban tree species. *Front Environ Sci Eng* 7(4): 579-588.
- Weber, C., 2013. Ecosystem services provided by urban vegetation: A literature review. In: Rauch S, Morrison G, Norra S et al. (eds.) *Urban Environment*: Springer Netherlands, 119-131.
- Yang, J., Chang, Y., Yan, P., 2015. Ranking the suitability of common urban tree species for controlling PM2.5 pollution. *Atmos Pollut Res* 6(2): 267-277.

CHAPTER 10 - EFFECTS OF AIR POLLUTION ON HEALTH

- Ancona, C., Golini, M. N., Ciancarella, L., Demaria, M., Badaloni, C., Cesaroni, G., Cadum, E., Forastiere, F., 2015. Health impact assessment of PM2.5 and NO₂ in Italy. The VIIAS National Study. In: *Environmental Health Perspectives - Abstracts of the 27th Conference of the International Society of Environmental Epidemiology (ISEE)*, n. 1912.
- Ciancarella, L., Adani, M., Briganti, G., Cappelletti, A., Ciucci, A., Cremona, G., D’Elia, I., D’Isidoro, M., Mircea, M., Piersanti, A., Righini, G., Russo, F., Vitali, L., Zanini, G., 2016. La simulazione nazionale di AMS-MINNI relativa all’anno 2010. Simulazione annuale del Sistema Modellistico Atmosferico di MINNI e validazione dei risultati tramite confronto con i dati osservati. Technical Report RT/2016/12/ENEA, ENEA, ISSN 0393-3016 (in Italian).

- Cox, D., 1972. Regression models and life-tables. *Journal of the Royal Statistical Society, Series B* 34 (2), 187–220.
- D’Elia, I., Bencardino, M., Ciancarella, L., Contaldi, M., Vialetto, G., 2009. Technical and Non-Technical Measures for air pollution emission reduction: The integrated assessment of the regional Air Quality Management Plans through the Italian national model. *Atmospheric Environment* 43, 6182-6189.
- EC, 2008. Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. *Official Journal of the European Union*, L. 152 of 11/6/2008.
- Hupkens, CLH., van den Berg, J., van der Zee, J., 1999. National health interview surveys in Europe: an overview. *Health Policy* 47 (2), 145-168.
- Ignaccolo, R., Ghigo, S., Bande, S., 2012. Functional zoning for air quality. *Environmental and Ecological Statistics*, 20 (1), 109-127.
- IARC, 2015. *Outdoor Air Pollution*. IARC monographs on the evaluation of carcinogenic risk to humans. Volume 109. Lyon, France: IARC.
- Mircea, M., Ciancarella L., Briganti G., Calori G., Cappelletti A., Cionni I., Costa M., Cremona G., D’Isidoro M., Finardi S., Pace G., Piersanti A., Righini G., Silibello C., Vitali L., Zanini G., 2014. Assessment of the AMS-MINNI system capabilities to simulate air quality over Italy for the calendar year 2005. *Atmospheric Environment* 84, 178-188.
- Uccelli, R., Mastrantonio, M., Altavista, P., Caiaffa, E., Cattani, G., Belli, S., Comba, P., 2016. Female lung cancer mortality and long term exposure to particulate matter in Italy. *European Journal of Public Health*, in press.
- WHO, 2006. Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide. Global update 2005. Copenhagen: WHO.
- WHO Regional Office for Europe, 2013. Review of evidence on health aspects of air pollution – REVIHAAP Project. Technical Report. Copenhagen: WHO Regional Office for Europe.
- WHO, 2015. Landmark Resolution on “Health and the Environment: Addressing the health impact of air pollution”. 68th World Health Assembly, 26 May 2015, Geneva, Switzerland.
- WHO Regional Office for Europe, 2015. Economic cost of the health impact of air pollution in Europe: Clean air, health and wealth. Copenhagen: WHO Regional Office for Europe.

CHAPTER 11 - WHAT REMAINS TO BE DONE TO REDUCE AIR POLLUTION?

- Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N., Winiwarter, W., 2015. Adjusted historic emission data, projections, and optimized emission reduction targets for 2030 – A comparison with COM data 2013. Part A: Results for EU-28. TSAP Report #16A, Version 1.1, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria, January 2015.
- Amann, M., Borken-Kleefeld, J., Cofala, J., Hettelingh, J-P., Heyes, C., Hoglund, L., Holland, M., Kiesewetter, G., Klimont, Z., Rafaj, P., Posch, M., Sander, R., Schöpp, W., Wagner, F., Winiwarter, W., 2014. The Final Policy Scenarios of the EU Clean Air Policy Package. TSAP Report #11, Version 1.0, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria, February 2014.
- Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sander, R., Schöpp, W., Wagner, F., Winiwarter, W., 2011. Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications. *Environmental Modelling & Software* 26, 1489-1501.
- Carlsson-Kanyama A., 1998. Climate change and dietary choices: how can emissions of greenhouse gases from food consumption be reduced? *Food Policy* 23:277–293.
- Carlsson-Kanyama A., Ekstrom M.P., Shanahan H., 2003. Food and life cycle energy inputs: consequences of diet and ways to increase efficiency. *Ecological Economics* 44:293–307.

- Ciucci, A., D'Elia, I., Wagner, F., Sander, R., Ciancarella, L., Zanini, G., Schöpp, W., 2016. Cost-effective reductions of PM_{2.5} concentrations and exposure in Italy. *Atmospheric Environment*, 140, 84-93.
- COM, 2013. COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT, THE COUNCIL, THE EUROPEAN ECONOMIC AND SOCIAL COMMITTEE AND THE COMMITTEE OF THE REGIONS. A Clean Air Programme for Europe. Brussels, 18.12.2013, COM(2013) 918 final.
http://ec.europa.eu/environment/air/clean_air_policy.htm
- D'Elia, I., Bencardino, M., Ciancarella, L., Contaldi, M., Vialetto, G., 2009. Technical and Non-technical measures for air pollution emission reduction: The integrated assessment of the Regional Air Quality Management Plans through the Italian national model. *Atmospheric Environment*, 43, 6182-6189.
- D'Elia, I., Peschi, E., 2016. How national integrated air quality models can be used in defining environmental policies: the revision of the NEC Directive. ENEA Technical Report, RT/2016/30/ENEA.
- EC, 2001. Directive 2001/81/EC of the European Parliament and of the Council of 23 October 2001 on national emission ceilings for certain atmospheric pollutants. *Official Journal of the European Union*, L. 309 of 27/11/2001
- EC, 2016. Directive 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC. *Official Journal of the European Union*, L. 344 of 17/12/2016.
- ECE, 2012a. Decision 2012/1. Amendment of annex I to the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone. ECE/EB.AIR/111/Add.1.
http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE_EB.AIR_111_Add.1_ENG_DECISION_1.pdf
- ECE, 2012b. Decision 2012/2. Amendment of the text of and annexes II to IX to the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone and the addition of new annexes X and XI.
 ECE/EB.AIR/111/Add.1
http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE_EB.AIR_111_Add.1_ENG_DECISION_2.pdf
- Engstrom R., Wadeskog A., Finnveden G., 2007. Environmental assessment of Swedish agriculture. *Ecological Economics* 60:550–563
- Mircea M., Ciancarella L., Briganti G., Calori G., Cappelletti A., Cionni I., Costa M., Cremona G., D'Isidoro M., Finardi S., Pace G., Piersanti A., Righini G., Silibello C., Vitali L., Zanini G., 2014. Assessment of the AMS-MINNI system capabilities to simulate air quality over Italy for the calendar year 2005. *Atmospheric Environment* 84, 178-188.
- Westhoek H., Lesschen J.P., Rood T., Wagner S., De Marco A., Murphy-Bokern D., Leip A., van Grinsven H., Sutton M., Oenema O., 2014. Food choices, health and environment: Effects of cutting Europe's meat and dairy intake. *Global Environmental Change*, 26: 196-205.

ENEA
Promotion and Communication Service

www.enea.it

March 2017



Corpo Forestale dello Stato



ISPRA

Istituto Superiore per la Protezione e la Ricerca Ambientale



DIEP/Lazio

Dipartimento di Epidemiologia del Servizio Sanitario Regionale Regione Lazio



UNIVERSITÀ
DEGLI STUDI
FIRENZE



UNIVERSITÀ
CATTOLICA
del Sacro Cuore
sede di Brescia



SAPIENZA
UNIVERSITÀ DI ROMA



Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria



PIEMONTE
Arpa
Agenzia Regionale
per la Protezione Ambientale



UNIVERSITÀ DI PISA



TerraData
environmetrics



WSL



UNICAM
Università di Camerino

1336



Consiglio
Nazionale delle
Ricerche

ENEA

www.enea.it