

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/321529949>

Outdoor cultivation of a bioactive *Tolypothrix* sp. in Green Wall Panel (GWP®) photobioreactors

Conference Paper · December 2017

CITATIONS

0

READS

17

10 authors, including:



Nataszia Biondi

University of Florence

42 PUBLICATIONS 2,348 CITATIONS

[SEE PROFILE](#)



Niccolo Bassi

University of Florence

16 PUBLICATIONS 2,014 CITATIONS

[SEE PROFILE](#)



Alberto Niccolai

University of Florence

15 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)



Mario Tredici

University of Florence

121 PUBLICATIONS 4,845 CITATIONS

[SEE PROFILE](#)



ALGAE EUROPE

05-07 DEC 2017 | BERLIN | GERMANY

BOOK OF ABSTRACTS



Liliana Rodolfi^{1,2}, Natascia Biondi¹, Giacomo Sampietro², Marco De Vito¹, Niccolò Bassi², Alberto Niccolai¹, Gimena Arganaraz¹, João Morais³, Vitor Vasconcelos^{3,4}, Mario R. Tredici¹

¹Department of Agrifood Production and Environmental Sciences (DISPAA), University of Florence, Italy

²Fotosintetica & Microbiologica S.r.l., Florence, Italy

³Interdisciplinary Center of Marine and Environmental Research- CIIMAR, Matosinhos, Portugal

⁴Faculty of Sciences, University of Porto, Portugal

Abstract:

The work here described was performed within the EU funded project NOMORFILM, which aims to discover novel bioactive compounds from microalgae for medical applications in both treatment and prophylaxis against biofilm infections on implanted prosthetic devices.

Within this study, a *Tolypothrix* sp. strain from the culture collection of CIIMAR (Portugal) was selected for its interesting antimicrobial activity and was cultivated outdoors in reactors of increasing volume: 8-L bubble tubes, 40-L GWP®-III (50° tilted, south facing panels) and 290-L GWP®-II (vertical east-west facing panels). The highest daily productivity (44.2 g m⁻² of directly illuminated surface area, equivalent to 34.1 g m⁻² ground surface area) was obtained in the 40-L GWP®-III photobioreactor in late June (27.5 MJ m⁻² d⁻¹ of average global solar radiation on the horizontal). The daily productivity decreased to 34.6 g m⁻² of directly illuminated surface area, equivalent to 26.7 g m⁻² ground surface area in late August (18.7 MJ m⁻² d⁻¹ of average global solar radiation on the horizontal). In early September, with similar average global solar radiation on the horizontal (16.3 MJ m⁻² d⁻¹), the strain was cultivated in the 290-L GWP®-II reactor attaining much lower productivities (17.4 g m⁻² of directly illuminated surface area d⁻¹, equivalent to 16.5 g m⁻² ground surface area d⁻¹), mainly because of the lower irradiance at the panel's surfaces (11.8 MJ m⁻² d⁻¹ on the vertical reactor against 24.7 MJ m⁻² d⁻¹ on the 50° tilted one).

Acknowledgment: This research was funded by the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 634588 (Nomofilm).