



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Groups in which each subgroup is commensurable with a normal subgroup.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Groups in which each subgroup is commensurable with a normal subgroup / Carlo, Casolo; Ulderico, Dardano; Silvana, Rinauro. - In: JOURNAL OF ALGEBRA. - ISSN 0021-8693. - STAMPA. - 496:(2018), pp. 48-60. [10.1016/j.jalgebra.2017.11.016]

Availability:

This version is available at: 2158/1109988 since: 2018-02-06T12:19:06Z

Published version:

DOI: 10.1016/j.jalgebra.2017.11.016

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Groups in which each subgroup is commensurable with a normal subgroup

to the memory of Jim Wiegold

Carlo Casolo, Ulderico Dardano, Silvana Rinauro

Abstract

A group G is a CN-group if for each subgroup H of G there exists a normal subgroup N of G such that the index $|HN : (H \cap N)|$ is finite. The class of CN-groups contains properly both the well-known classes of core-finite groups and of finite-by-abelian groups. In the present paper it is shown that a CN-group whose periodic images are locally finite is finite-by-abelian-by-finite. Then such groups are described into some details by considering automorphisms of abelian groups. Finally, it is shown that if G is a locally graded group with the property that the above index is bounded independently of H , then G is finite-by-abelian-by-finite. ¹

1 Introduction

In a celebrated paper, B.H. Neumann [9] showed that for a group G the property that each subgroup H has finite index in a normal subgroup of G (i.e. $|H^G : H|$ is finite) is equivalent to the fact that G has finite derived subgroup (G is *finite-by-abelian*).

The class of groups with a dual property was considered in [1]. A group G is said a CF-group (*core-finite*) if each subgroup H contains a normal subgroup of G with finite index in H (i.e. $|H : H_G|$ is finite). As Tarski groups are CF, a complete classification of CF-groups seems to be much difficult. However, in [1] and [11] it has been proved that *a CF-group G whose periodic quotients are locally finite is abelian-by-finite and there exists an integer n such that $|H : H_G| \leq n$ for all $H \leq G$ (say that G is BCF, boundedly CF).*

¹Key words and phrases: locally finite, core-finite, subnormal, inert, CF-group.

2010 Mathematics Subject Classification: Primary 20F24, Secondary 20F18, 20F50, 20E15

Moreover a *locally graded BCF-group is abelian-by-finite*. Furthermore, an easy example of a metabelian (and even hypercentral) group which is CF but not BCF is given. It seems to be a still open question whether every locally graded CF-group is abelian-by-finite. Recall that a group is said abelian-by-finite if has an abelian subgroup with finite index and that a group is *locally finite* (*locally graded*, resp.) if each finitely generated subgroup is finite (has a proper subgroup with finite index, resp.).

With the aim of considering the above two classes in a common framework, recall that two subgroups H and K of a group G are said *commensurable* if and only if $H \cap K$ has finite index in both H and K . This is an equivalence relation and will be denoted by \sim . Clearly, if $H \sim K$, then $(H \cap L) \sim (K \cap L)$ and $HM \sim KM$ for each $L \leq G$ and $M \triangleleft G$.

In the present paper we consider the class of *CN-groups*, that is *groups in which each subgroup is commensurable to a normal subgroup*. Into details, for a subgroup H of a group G define $\delta_G(H)$ to be the minimum index $|HN : (H \cap N)|$ with $N \triangleleft G$. Then G is a CN-group if and only if $\delta_G(H)$ is finite for all $H \leq G$. Clearly both finite-by-abelian and CF groups are CN. Moreover, the class of CN-groups is both subgroup and quotient closed.

Note that if a subgroup H of a group G is commensurable with a normal subgroup N , then $S := (H \cap N)_N$ has finite index in H . Thus the class of CN-groups is contained in the class of *sbyf-groups*, that is, groups in which each subgroup H is *subnormal-by-finite*; that is to say that H contains a subnormal subgroup S of G such that the index $|H : S|$ is finite. It is known that *locally finite sbyf-groups are (locally nilpotent)-by-finite* (see [3]) and *nilpotent-by-Chernikov* (see [6]).

Recall also that from results in [4] it follows that *for an abelian-by-finite group properties CN and CF are equivalent*. However, for each prime p there is a nilpotent p -group with property CN which is neither finite-by-abelian nor abelian-by-finite, see Proposition 2.2 below.

Our main result is the following.

Theorem A *Let G be a CN-group such that every periodic image of G is locally finite. Then G is finite-by-abelian-by-finite.*

Here by finite-by-abelian-by-finite group we mean a group which has a subgroup which has finite index and is finite-by-abelian. The proof of Theorem A will be completed at the end of Sect. 5. Before, in Sect. 3, we study the action of a CN-group on its abelian sections, see Theorem 3.2 and Corollary 3.3. Then in Sect. 4 we consider also *BCN-groups*, that is, groups

G for which there is $n \in \mathbb{N}$ such that $\delta_G(H) \leq n$ for all $H \leq G$. We will show the following theorem.

Theorem B *Let G be a finite-by-abelian-by-finite group.*

- i) if G is CN, then the FC-center of G has finite index and is finite-by-abelian;*
- ii) G is CN if and only if it is finite-by-CF.*
- iii) G is BCN if and only if it is finite-by-BCF.*

It follows that that if the group G is periodic and finite-by-abelian-by-finite, then G is BCN if and only if it is CN. Then we consider non-periodic finite-by-abelian-by-finite BCF- and BCN-groups by Proposition 4.4.

The more restrictive property BCN reveals fruitful when we consider the wider class of locally graded groups.

Corollary *A locally graded BCN-group is finite-by-abelian-by-finite.*

Our notation is mostly standard and we refer to [10].

2 Preliminaries

We point out a sufficient condition for a group to be CN (or even BCN).

Proposition 2.1 *Let G be a group with a normal series $G_0 \leq G_1 \leq G$, where G_0 and G/G_1 have finite order m and n resp. If $H \leq G$, then H is commensurable with $H_1 := (H \cap G_1)G_0 \leq G_1$ and $\delta_G(H) \leq mn \cdot \delta_{G/G_0}(H_1/G_0)$.*

In particular, if each subgroup of G_1/G_0 is commensurable with a normal subgroup of G/G_0 , then G is a CN-group. \square

Now we give examples of non trivial CN-groups.

Proposition 2.2 *For each prime p there is a nilpotent p -group with property BCN, which is not abelian-by-finite nor finite-by-abelian.*

Proof. Consider a sequence P_n of isomorphic groups with order p^4 defined by $P_n := \langle x_n, y_n \mid x_n^{p^3} = y_n^p = 1, x_n^{y_n} = x_n^{1+p^2} \rangle = \langle x_n \rangle \rtimes \langle y_n \rangle$ where clearly $P'_n = \langle x_n^{p^2} \rangle$ has order p . Let $P := \text{Dr}_{n \in \mathbb{N}} P_n$ and consider the automorphism γ of P such that $x_n^\gamma = x_n^{1+p}$ and $y_n^\gamma = y_n$, for each $n \in \mathbb{N}$. Clearly, γ has order p^2 , acts as the automorphism $x \mapsto x^{1+p}$ on P/P' (which has exponent p^2) and acts trivially on P' (which is elementary abelian). Finally let $N := \langle x_0^{p^2} x_n^{p^2} \mid n \in \mathbb{N} \rangle$. Then N is a subgroup of P' with index p . Thus the p -group

$G := (P \rtimes \langle \gamma \rangle)/N$ is a BCN-group by Proposition 2.1 applied to the series $P'/N \leq P/N \leq G$.

We have that G' is infinite, since for each n we have $x_n^p = [x_n, \gamma] \in [P_n, \gamma] > P'_n$. Moreover, we have that $gN \in Z(P/N)$ if and only if $\forall i [g, P_i] \leq N$, and $N \cap P_i = 1$. Thus $Z(P/N) = Z(P)/N$ where $Z(P) = \text{Dr}_n \langle x_n^p \rangle$ has infinite index in P .

If, by contradiction, G is abelian-by-finite, then there is an abelian normal subgroup A/N of P/N with finite index. Then for some $m \in \mathbb{N}$ we have $P = AF$, where $F = \text{Dr}_{n < m} P_n$ is a finite normal subgroup of P . Therefore P/N is center-by-finite, a contradiction. \square

3 Automorphisms of abelian groups

As in [4], for the action of a group Γ on a group A , we consider the following properties:

- P) $\forall H \leq A \quad H = H^\Gamma$;
- AP) $\forall H \leq A \quad |H/H_\Gamma| < \infty$;
- BP) $\forall H \leq A \quad |H^\Gamma/H| < \infty$;
- CP) $\forall H \leq A \quad \exists K = K^\Gamma \leq A$ such that $H \sim K$, (H, K are commensurable).

When P holds, one says that Γ acts on A by means of *power automorphisms* or that A is Γ -*hamiltonian* ([10],[1]). Recall that if γ is a power automorphism of an abelian p -group A , then there exists a p -adic integer α such that $a^\gamma = a^\alpha$ for all $a \in A$ (see [10] for details). Here a^α stands for a^n , where $n \in \mathbb{N}$ is congruent to α modulo the order of a . On the other hand, a power automorphism of a non-periodic abelian group is either the identity or the inversion map.

Obviously both AP and BP imply CP. Moreover, *these three properties are equivalent, provided A is abelian and Γ is finitely generated, while they are in fact different in the general case even when A and Γ are elementary abelian p -groups* (see [4]). On the other hand, the properties AP and BP have previously characterized in [5] and [2] resp., as we are going to recall. To shorten statements we define a further property:

- \tilde{P}) Γ has P on the factors of a Γ -series $1 \leq V \leq D \leq A$ where
 - i) V is free abelian with finite rank,
 - ii) D/V is divisible periodic with finite total rank,
 - iii) A/D is periodic and has finite p -exponent for each prime $p \in \pi(D/V)$.

Theorem 3.1 [5],[2] *Let Γ be group acting on an abelian group A . Then:*
a) Γ has AP on A if and only if there is a Γ -subgroup A_1 such that A/A_1 is finite and Γ has either P or \tilde{P} on A_1 .
b) Γ has BP on A if and only if there is a Γ -subgroup A_0 such that A_0 is finite and Γ has either P or \tilde{P} on A/A_0 .

By next statement we give a characteration of the property CP along the same lines.

Theorem 3.2 *Let Γ be group acting on an abelian group A . Then:*
c) Γ has CP on A if and only if there are Γ -subgroups $A_0 \leq A_1 \leq A$ such that A_0 and A/A_1 are finite and Γ has either P or \tilde{P} on A_1/A_0 .

The proof of Theorem 3.2 is at the end of this section. Here we deduce a corollary.

Corollary 3.3 *For a group Γ acting on an abelian group A , the following are equivalent:*

- a) Γ has AP on A/A_0 for a finite Γ -subgroup A_0 of A ,
- b) Γ has BP on a finite index Γ -subgroup A_1 of A ,
- c) Γ has CP on A . □

Let us recall some basic facts from [4] where *inertial automorphisms* of abelian groups have been introduced. These are automorphisms γ of a group G such that $H^\gamma \sim H$ for all $H \leq G$. Clearly, if Γ has CP on G and $\gamma \in \Gamma$, then γ is inertial.

Proposition 3.4 *Let Γ be group acting on a locally nilpotent periodic group A . Then Γ has AP, BP, CP resp. on A if and only if Γ has AP, BP, CP resp. on finitely many primary components of A and P on all the other ones. □*

Lemma 3.5 *Let Γ be a group acting on an abelian group A . If Γ has CP, then:*

- i) Γ has P on the maximum periodic divisible subgroup of A .
- ii) if A is torsion-free, then each $\gamma \in \Gamma$ acts by conjugation on A by either the identity or the inversion map. □

Now we prove some lemmas. In the first one we do not require that the group A is abelian.

Lemma 3.6 *Let Γ be a group acting on a group A . If Γ has CP, then Γ has BP on the subgroup $X := \{a \in A \mid \langle a \rangle^\Gamma \text{ is finite}\}$ of A .*

Proof. For any $H \leq X$ there is K such that $H \sim K = K^\Gamma \leq A$. Then there is finite subgroup $F \leq X$ such that $H \leq KF$. Thus $H^\Gamma \leq KF^\Gamma$ and $|H^\Gamma : H| \leq |F^\Gamma| \cdot |HK : H|$ is finite. \square

Lemma 3.7 *Let Γ be a group acting on a p -group A which is the direct product of cyclic groups. If Γ has CP, then the following subgroup has finite index in A :*

$$X := \{a \in A \mid \langle a \rangle^\Gamma \text{ is finite}\}$$

Proof. Assume by contradiction that A/X is infinite.

Let us see that, by elementary facts, there is a sequence (a_n) of elements of A such that

- 1) $\langle a_n \mid n \in \mathbb{N} \rangle = \text{Dr}_{n \in \mathbb{N}} \langle a_n \rangle$,
- 2) $A_I / A_I \cap X$ is infinite, for each infinite subset I of \mathbb{N} , where $A_I := \langle a_i \mid i \in I \rangle$.

In fact, if A/X has finite rank, it has a Prüfer subgroup Q/X . Let Y be a countable subgroup such that $Q = YX$. By Kulikov Theorem (see [10]) Y is the direct product of cyclic groups, so that we may choose elements $a_n \in Y$ such that $\langle a_n \mid n \in \mathbb{N} \rangle = \text{Dr}_{n \in \mathbb{N}} \langle a_n \rangle \leq Y$ and $|a_n X| < |a_{n+1} X|$. The claim holds. Similarly, if A/X has infinite rank, we may consider its socle S/X and consider a countable subgroup Y such that $S = YX$. Then we may choose elements $a_n \in Y$ which are independent mod X and generate their direct product as in (1).

We claim now that there are sequences of infinite subsets I_n, J_n of \mathbb{N} and Γ -subgroups $K_n \leq A$ such that for each $n \in \mathbb{N}$:

- 3) $I_n \cap J_n = \emptyset$ and $I_{n+1} \subseteq J_n$
- 4) $K_n \sim A_{I_n}$
- 5) $(K_1 \dots K_i) \cap (A_{I_1} \dots A_{I_n}) \leq (A_{I_1} \dots A_{I_i}), \forall i \leq n$.

Proceed by induction on n . Choose an infinite subset I_1 of \mathbb{N} such that $J_1 := \mathbb{N} \setminus I_1$ is infinite. By CP-property there exists $K_1 = K_1^\Gamma$ commensurable with A_{I_1} .

Suppose we have defined I_j, J_j, K_j for $1 \leq j \leq n$ such that 3-5 holds. Since $(K_1 \dots K_n) \sim (A_{I_1} \dots A_{I_n})$, there is $m \in \mathbb{N}$ such that

- 6) $(K_1 K_2 \dots K_n) \cap A_\mathbb{N} \leq (A_{I_1} A_{I_2} \dots A_{I_n}) \langle a_1, \dots, a_m \rangle$.

Let I_{n+1} and J_{n+1} be disjoint infinite subsets of $J_n \setminus \{1, \dots, m\}$. By CP-property there exists $K_{n+1} = K_{n+1}^\Gamma$ commensurable with $A_{I_{n+1}}$. By the choice

of I_{n+1} it follows that

$$7) \quad (K_1 \dots K_i) \cap (A_{I_1} \dots A_{I_{n+1}}) \leq (K_1 \dots K_i) \cap (A_{I_1} \dots A_{I_n}) \quad \forall i \leq n$$

and so (5) holds for $n+1$, as wished. The claim is proved.

Note that by (2) and (5) it follows that $A_{I_n}/A_{I_n} \cap X$ is infinite for each $n \in \mathbb{N}$ and that also the following property holds

$$8) \quad (K_1 K_2 \dots K_n) \cap \bar{A} \leq (A_{I_1} A_{I_2} \dots A_{I_n}) \quad \forall n, \text{ where } \bar{A} := \text{Dr}_{n \in \mathbb{N}} A_{I_n}.$$

Now for each $n \in \mathbb{N}$, choose an element $b_n \in (A_{I_n} \cap K_n) \setminus X$. Then we have $B := \langle b_n \mid n \in \mathbb{N} \rangle = \text{Dr}_n \langle b_n \rangle$, where $\langle b_n \rangle^\Gamma$ is infinite and $\langle b_n \rangle^\Gamma \leq K_n \sim A_{I_n}$, so that

$$9) \quad \langle b_n \rangle^\Gamma \cap A_{I_n} \text{ is infinite for each } n.$$

Since there exists $B_0 = B_0^\Gamma \sim B$, we may take

$$- B_* := (B_0 \cap B)^\Gamma = (B_* \cap B)^\Gamma \leq B^\Gamma \text{ where } B_* \sim B.$$

Now $B_*/(B_* \cap B)$ and $B/(B_* \cap B)$ are both finite and there is $n \in \mathbb{N}$ such that if $B_n := \langle b_1, \dots, b_n \rangle$ we have

$$- (B_* \cap B)^\Gamma = B_* \leq (B_* \cap B) B_n^\Gamma \text{ and}$$

$$- B = (B_* \cap B) B_n.$$

Since $b_n \in K_n$ for each n , we have $B_n \leq \bar{K}_n := K_1 K_2 \dots K_n$ and

$$- B^\Gamma = (B_* \cap B)^\Gamma B_n^\Gamma \leq (B_* \cap B) B_n^\Gamma \leq (B_* \cap B) \bar{K}_n \leq B \bar{K}_n, \text{ so that}$$

$$- B^\Gamma \cap \bar{A} \leq B \bar{K}_n \cap \bar{A} = B(\bar{K}_n \cap \bar{A}) \leq B A_{I_1} A_{I_2} \dots A_{I_n} \text{ by (8) above.}$$

Thus

$$- \langle b_{n+1} \rangle^\Gamma \cap A_{I_{n+1}} \leq B^\Gamma \cap A_{I_{n+1}} \leq (B A_{I_1} A_{I_2} \dots A_{I_n}) \cap A_{I_{n+1}} = \langle b_{n+1} \rangle \text{ is finite,}$$

a contradiction with (9). \square

Lemma 3.8 *Let Γ be a group acting on an abelian periodic reduced group A . If Γ has CP, then there are Γ -subgroups $A_0 \leq A_1 \leq A$ such that A_0 and A/A_1 are finite and Γ has P on A_1/A_0 .*

Proof. By Proposition 3.4 it is enough to consider the case when A is a p -group. If A is the direct product of cyclic groups, by Lemma 3.7 we have that $A_1 := \{a \in A \mid \langle a \rangle^\Gamma \text{ is finite}\}$ has finite index in A . Further, by Lemma 3.6, Γ has BP on A_1 . Then the statement follows from Theorem 3.1.

Let A be any reduced p -group and B_* be a basic subgroup of A . Then there is $B = B^\Gamma \sim B_*$. Since A/B_* is divisible, then the divisible radical of A/B has finite index. Thus we may assume that A/B is divisible. By Kulikov Theorem (see [10]), also B is a direct product of cyclic groups, therefore by the above there are Γ -subgroups $B_0 \leq B_1 \leq B$ such that B_0 and B/B_1 are

finite and Γ has P on B_1/B_0 . We may assume $B_0 = 1$. Also, since A/B_1 is finite-by-divisible, it is divisible-by-finite and we may assume it is divisible.

Let $\gamma \in \Gamma$ and α be a p -adic integer such that $x^\gamma = x^\alpha$ for all $x \in B_1$. Consider the endomorphism $\gamma - \alpha$ of A and note that $B_1 \leq \ker(\gamma - \alpha)$. Thus $A/\ker(\gamma - \alpha) \simeq \text{im}(\gamma - \alpha)$ is both divisible and reduced, hence trivial. It follows $\gamma = \alpha$ on the whole A . \square

Proof of Theorem 3.2 For the sufficiency of the condition note that for any subgroup $H \leq A$ we have $H \sim H \cap A_1$ and the latter is in turn commensurable with a Γ -subgroup since Γ has BP on A_1 by Theorem 3.1.

Concerning necessity, we first prove the statement when A is periodic. Let $A = D \times R_1$, where D is divisible and R_1 is reduced. Then there is a $R = R^\Gamma \sim R_1$. Thus DR and $D \cap R$ are Γ -subgroups of A with finite index and order resp. Then we can assume $A = D \times R$. Let $X := \{a \in A \mid \langle a \rangle^\Gamma \text{ is finite}\}$. Clearly $D \leq X$, as Γ has P on D by Lemma 3.5. On the other hand, $X \cap R$ has finite index in R by Lemma 3.8. It follows A/X is finite and by Lemma 3.6 and Theorem 3.1 the statement holds.

In the non-periodic case, note that if V_0 is a maximal free subgroup of A (hence A/V_0 is periodic), then there is $V_1 = V_1^\Gamma \sim V_0$. Let $n := |V_1/(V_0 \cap V_1)|$. Thus by applying Lemma 3.5 we have

- *there is a free abelian Γ -subgroup $V := V_1^n$ such that A/V is periodic and each $\gamma \in \Gamma$ acts on V by either the identity or the inversion map.*

Suppose that V has finite rank. Consider now the action of Γ on the periodic group A/V and apply the above. Then there is a series $V \leq A_0 \leq A_1 \leq A$ such that A_0/V and A/A_1 are finite and Γ has either P or \tilde{P} on A_1/A_0 . Since A_0 has finite torsion subgroup T we can factor out T and assume $A_0 = V$. Then Γ has either P or \tilde{P} on A_1 as straightforward verification shows.

Suppose finally that V has infinite rank. Let $V_2 \leq V$ be such that V/V_2 is divisible periodic and its p -component has infinite rank for each prime p . We may assume $V := V_2$. By the above case when A is periodic, there is a Γ -series $V \leq A_0 \leq A_1 \leq A$ such that A_0/V and A/A_1 are finite and Γ has P on A_1/A_0 . We may factor out the torsion subgroup of A_0 , as it is finite, and assume $A_0 = V$.

Again let $V_2 \leq V$ be such that V/V_2 is divisible periodic and its p -component has infinite rank for each prime p . Let $\gamma \in \Gamma$ and α_p be a p -adic integer such that $x^\gamma = x^{\alpha_p}$ for all x in the p -component of A_1/V . Let $\epsilon = \pm 1$ be such that $x^\gamma = x^\epsilon$ for all $x \in V$. By Lemma 3.5, γ has P on

the maximum divisible subgroup D_p/V_2 of the p -component of A_1/V_2 . Thus $\alpha_p = \epsilon$ on D_p/V_2 . Therefore $x^\gamma = x^\epsilon$ for all $x \in V$ and for all $x \in A_1/V$. We claim that $a^\gamma = a^\epsilon$ for each $a \in A_1$. To see this, for any $a \in A_1$ consider $n \in \mathbb{N}$ such that $a^n \in V$. Then there is $v \in V$ such that $a^\gamma = a^\epsilon v$. Hence $a^{n\epsilon} = (a^n)^\gamma = (a^\gamma)^n = (a^\epsilon v)^n = a^{n\epsilon} v^n$. Thus $v^n = 1$. Therefore, as V is torsion-free, we have $v = 1$, as wished. \square

4 Abelian-by-finite CN-groups and Theorem B

Locally finite CF-groups are known to be abelian-by-finite and BCF (see [1]).

Proposition 4.1 *Let G be an abelian-by-finite group.*

- i) if G is CN, then G is CF;*
- ii) if G is BCN, then G is BCF.*

Proof. Let A be a normal abelian subgroup with finite index r . Then each $H \leq A$ has at most r conjugates in G . If $\delta_G(H) \leq n < \infty$ then for each $g \in G$ we have $|H : (H \cap H^g)| \leq 2\delta_G(H) \leq 2n$ hence $|H/H_G| \leq (2n)^r$. More generally, if H is any subgroup of G , then $|H/H_G| \leq r(2n)^r$. \square

We state now a key fact about non-periodic CN-groups.

Lemma 4.2 *Let G be a CN-group and $A = A(G)$ its subgroup generated by all infinite cyclic normal subgroups. Then G/A is periodic, A is abelian and each $g \in G$ acts on A by either the identity or the inversion map, hence $|G/C_G(A)| \leq 2$.*

Proof. For any $x \in G$ there is $N \triangleleft G$ which is commensurable with $\langle x \rangle$. Then $n := |N : (N \cap \langle x \rangle)|$ is finite. Thus $N^{n!} \leq \langle x \rangle$ where $N^{n!} \triangleleft G$. Hence G/A is periodic.

It is clear that A is abelian. Let $g \in G$. If $\langle a \rangle \triangleleft G$ and a has infinite order, then there is $\epsilon_a = \pm 1$ such that $a^g = a^{\epsilon_a}$. On the other hand, by Lemma 3.5, there is $\epsilon = \pm 1$ such that for each $a \in A$ there is a periodic element $t_a \in A$ such that $a^g = a^\epsilon t_a$. It follows $a^{\epsilon_a - \epsilon} = t_a$. Therefore $\epsilon_a = \epsilon$ is independent of a , as wished. \square

Lemma 4.3 *Let G be an FC-group. If G is a CN-group then G is finite-by-abelian.*

Proof. Let H be any subgroup of G . We shall prove that $|H^G : H|$ is finite. Consider $A = A(G)$ as in Lemma 4.2. Then $H \cap A \triangleleft G$ and $H/A \cap H$ is periodic. Hence we may assume H is periodic, that is, H contained in the torsion subgroup of the FC-group G . Our claim follows then from Lemma 3.6. \square

Proof of Theorem B Let G be a CN-group and $G_0 \leq G_1 \leq G$ be a normal series such that G_1/G_0 is abelian and both G_0 and G/G_1 are finite. Then G has CP on G_1/G_0 . By Corollary 3.3, the group G has BP on a subgroup $A_1/G_0 \leq G_1/G_0$ with finite index in G_1/G_0 . Thus A_1/G_0 is contained in the FC-centre of G/G_0 . Hence A_1 is contained in the FC-centre of F of G . So that G/F is finite. On the other hand, from Lemma 4.3 it follows that F' is finite.

Finally, (ii) and (iii) follow from Proposition 2.1 and Proposition 4.1. \square

Let us characterize BCF-groups among abelian-by-finite CF-groups.

Proposition 4.4 *Let G be a non-periodic group with an abelian normal subgroup A with finite index. Then the following are equivalent:*

- i) G is a BCF-group;*
- ii) G is a CF-group and there is $B \leq A$ such that B has finite exponent, $B \triangleleft G$ and each $g \in G$ acts by conjugation on A/B by either the identity or the inversion map.*

Proof. Let T be the torsion subgroup of A . By Lemma 3.5, each $g \in G$ acts on A/T as the automorphism $x \mapsto x^{\epsilon_g}$ where $\epsilon_g = \pm 1$. Then the equivalence of (i) and (ii) holds with $B := \langle A^{g^{-\epsilon_g}} \mid g \in G \rangle$, by Theorem 3 of [4]. \square

5 Proof of Theorem A

Our first statement in this section is a reduction to nilpotent groups.

Lemma 5.1 *A soluble p -group G with the property CN is nilpotent-by-finite.*

Proof. By Theorem 3.2, one may refine the the derived series of G to a finite G -series \mathcal{S} such that G has P on each infinite factor of \mathcal{S} . Recall that a p -group of power automorphisms of an abelian p -group is finite (see [10]). Then the stability group $S \leq G$ of the series \mathcal{S} , that is, the intersection of

the centralizers in G of the factors of the series, has finite index in G . On the other hand, by a theorem of Ph.Hall, S is nilpotent. \square

We recall now an elementary property of nilpotent groups.

Lemma 5.2 *Let G be a nilpotent group with class c . If G' has finite exponent e , then $G/Z(G)$ has finite exponent dividing e^c .*

Proof. Argue by induction on c , the statement being clear for $c = 1$. Assume $c > 1$ and that G/Z has exponent dividing e^{c-1} , where $Z/\gamma_c(G) := Z(G/\gamma_c(G))$. Then for all $g, x \in G$ we have $[g^{e^{c-1}}, x] \in \gamma_c(G) \leq G' \cap Z(G)$. Therefore $1 = [g^{e^{c-1}}, x]^e = [g^{e^c}, x]$, and $g^{e^c} \in Z(G)$, as claimed. \square

Next lemma follows easily from Lemma 6 in [8].

Lemma 5.3 *Let G be a nilpotent p -group and N a normal subgroup such that G/N is an infinite elementary abelian group. If H and U are finite subgroup of G such that $H \cap U = 1$, there exists a subgroup V of G such that $U \leq V$, $H \cap V = 1$ and $V/N \cap V$ is infinite. \square*

We deduce a technical lemma which is a tool for our purpose.

Lemma 5.4 *Let G be a nilpotent p -group and N be a normal subgroup such that G/N is an infinite elementary abelian group. If N contains the FC-center of G and G' is abelian with finite exponent, then there are subgroups H, U of G such that $H \cap U = 1$, with injective maps $n \mapsto h_n \in H$ and $(i, n) \mapsto u_{i,n} \in [G, h_i^{-1}h_n] \cap U$, where $i, n \in \mathbb{N}$, $i < n$.*

Proof. Let us show that for each $n \in \mathbb{N}$ there is an $(n + 1)$ -uple $v_n := (h_n, u_{0,n}, u_{1,n}, \dots, u_{n-1,n})$ of elements of G such that:

1. $\{h_1, \dots, h_n\}$ is linearly independent modulo N ;
2. $u_{i,n} \in [G, h_i^{-1}h_n] \quad \forall i \in \{0, \dots, n-1\}$;
3. $\{u_{j,h} \mid 0 \leq j < k \leq n\}$ is \mathbb{Z} -independent in G' ;
4. $H_n \cap U_n = 1$, where $H_n := \langle h_1, \dots, h_n \rangle$ and $U_n := \langle u_{j,h} \mid 0 \leq j < k \leq n \rangle$.

Then the statement is true for $H := \bigcup_{n \in \mathbb{N}} H_n$ and $U := \bigcup_{n \in \mathbb{N}} U_n$.

Let $h_0 := 1$ and choose $h_1 \in G \setminus N$. Since $N \geq F$, the FC-center of G , we have that y_1 has an infinite numbers of coniugates in G , hence $[G, h_1]$ is infinite and residually finite. Thus we may choose $u_{0,1} \in [G, h_1]$ such that $\langle u_{0,1} \rangle \cap \langle h_1 \rangle = 1$.

Assume then that we have defined v_i for $i \leq n$, that is, we have elements $h_0, \dots, h_n, u_{j,k}$, with $0 \leq j < k \leq n$ such that conditions 1-4 hold. To define an adequate v_{n+1} , note that by Lemma 5.3 we have that there exists $V_n \leq G$ such that $H_n \leq V_n$, $U_n \cap V_n = 1$ and $V_n N / N$ is infinite. Then choose

$$i) \quad h_{n+1} \in V_n \setminus N U_n H_n.$$

Note that $h_{n+1} \notin F H_n \leq N H_n$, so that $\{h_1, \dots, h_{n+1}\}$ is independent mod F . In particular $\forall i \in \{0, \dots, n\}$, $h_i^{-1} h_{n+1} \notin F$, hence also $[G, h_i^{-1} h_{n+1}]$ is infinite. Since G' is residually finite, we may recursively choose $u_{0,n+1}, \dots, u_{n,n+1}$ such that $\forall i \in \{0, \dots, n\}$

$$ii) \quad u_{i,n} \in [G, h_i^{-1} h_n]$$

$$iii) \quad \langle u_{i,n+1} \rangle \cap U_n \langle u_{h,n+1} \mid 0 \leq h < i \rangle H_{n+1} = 1$$

Then properties 1-3 hold for v_{n+1} . Finally suppose there are $h \in H_n, u \in U_n, s, t_0, \dots, t_n \in \mathbb{Z}$ such that

$$iii) \quad a = h h_{n+1}^s = u u_{0,n+1}^{t_1} \cdots u_{n,n+1}^{t_n} \in H_{n+1} \cap U_{n+1}.$$

Then from (iii) it follows $u_{n,n+1}^{t_n} = \dots = u_{0,n+1}^{t_1} = 1$. Hence $a = h h_{n+1}^s = u \in V_n \cap U_n = 1$ and 4 holds. \square

Lemma 5.5 *Let G be a nilpotent p -group. If G is CN, then G' has finite exponent.*

Proof. If, by contradiction, G' has infinite exponent, then the same happens to the abelian group $G'/\gamma_3(G)$ and there is N such that $G' \geq N \geq \gamma_3(G)$ and G'/N is a Prüfer group. We may assume $N = 1$, that is, G' itself is a Prüfer group and $G' \leq Z(G)$. Let us show that for any $H \leq G$ we have $|H^G : H| < \infty$, hence G' is finite, a contradiction. In fact we have that, by CN-property there is $K \triangleleft G$ such that $K \sim H$. Thus H has finite index in HK and we can also assume $H = HK$, that is, H/H_G is finite. Thus, we can assume $H_G = 1$ and $H \cap G' = 1$, that is, H is finite with order p^n and HG' is an abelian Chernikov group. It follows that H is contained in the n -th socle S of $HG' \triangleleft G$, where S is finite and normal in G , as wished. \square

Lemma 5.6 *Let G be a nilpotent p -group. If G is CN, then G is finite-by-abelian-by-finite.*

Proof. Assume, by contradiction, G is a counterexample. Then both G' and $G/Z(G)$ are infinite. However, they have finite exponent by Lemmas 5.5 and 5.2. Moreover, even the FC-center F of G has infinite index by Lemma 4.3. On the other hand, G/F has finite exponent, since $F \geq Z(G)$.

Then $N := FG^pG'$ has infinite index in G , otherwise the abelian group G/FG' has finite rank and finite exponent, hence it is finite. This implies that the nilpotent group G/F is finite, a contradiction.

If G' is abelian we are in a condition to apply Lemma 5.4 and get infinite elements and subgroups $h_n \in H$, $u_{i,n} \in U$ as in that statement. By CN-property there is K such that $H \sim K \triangleleft G$. So that the set $\{h_n(H \cap K) \mid n \in \mathbb{N}\}$ is finite. Hence there is $i \in \mathbb{N}$ and an infinite set $I \subseteq \mathbb{N}$ such that for each $n \in I$ we have $h_i^{-1}h_n \in H \cap K$ and $u_{i,n} \in U \cap [G, H \cap K] \leq U \cap K$. Therefore $U \cap K$ is infinite, in contradiction with $U \cap K \sim U \cap H = 1$.

For the general case, proceed by induction on the nilpotency class $c > 1$ of G and assume that the statement is true for $G/Z(G)$ and even that this is finite-by-abelian. Then there is a subgroup $L \leq G$ such that G/L is abelian and $L/Z(G)$ is finite. Thus L' is finite and, by the above, G/L' is finite-by-abelian-by-finite, a contradiction. \square

Proof of Theorem A. Recall from the Introduction that all subgroups of G are subnormal-by-finite. Thus, by above quoted results in [6] and [3] resp., we may assume that G is locally nilpotent and soluble

Assume first G is periodic. Then, by Lemma 3.4, only finitely many primary components are non-abelian. Thus we may assume G is a p -group and apply Lemma 5.1 and Lemma 5.6. It follows that G is finite-by-abelian-by-finite.

To treat the general case, consider $A = A(G)$ as in Lemma 4.2. We may assume A is central in G . Let V be a torsion-free subgroup of A such that A/V is periodic. Then G/V is locally finite and we may apply the above. Thus there is a series $V \leq F \leq G_1 \leq G$ such that G acts trivially on V , G_1/G_0 is abelian, while G_0/V and G/G_1 are finite. Then we can assume $G = G_1$ and note that the stabilizer S of the series has now finite index. Since S is nilpotent (by Ph.Hall Theorem) we can assume that $G = S$ is nilpotent. If T is the torsion subgroup of G , then VT/T is contained in the center of G/T . Since all factor of the upper central series of G/T are torsion-free we have G/T is abelian. Thus $G' \leq T \cap G_0$ is finite. \square

Proof of Corollary. If the statement is false, by Theorem A we may assume there is a counterexample G periodic and not locally finite. Also we may

assume G is finitely generated and infinite. Let R be the locally finite radical of G . By Theorem A again, R is finite-by-abelian-by-finite. By Theorem B(i), there is a finite subgroup $G_0 \triangleleft G$ such that R/G_0 is abelian-by-finite. We may assume $G_0 = 1$, so that R is abelian-by-finite.

We claim that $\bar{G} := G/R$ has finite exponent at most $(n+1)!$ where n is such that $n \geq \delta_G(H)$ for each $H \leq G$. In fact, for each $x \in \bar{G}$, there is $\bar{N} \triangleleft \bar{G}$ such that $|\bar{N} : (\bar{N} \cap \langle x \rangle)| \leq n$. Thus $\bar{N}^{n!} \leq \langle x \rangle$ and $\bar{N}^{n!} \triangleleft G$. Hence $\bar{N}^{n!} = 1$ and $x^{n!} = 1$.

By the positive answer (for all exponents) to the Restricted Burnside Problem, there is a positive integer k such that every finite image of \bar{G} has order at most k . Since \bar{G} is finitely generated, this means that the finite residual \bar{K} of \bar{G} has finite index and is finitely generated as well. Since also \bar{G} is locally graded (see [7]), we have $\bar{K} = 1$ and \bar{G} is finite. Therefore G is abelian-by-finite, a contradiction. \square

References

- [1] J. T. Buckley, J.C. Lennox, B. H. Neumann, H. Smith, J. Wiegold, Groups with all subgroups normal-by-finite. *J. Austral. Math. Soc. Ser. A* **59** (1995), no. 3, 384-398.
- [2] C. Casolo, Groups with finite conjugacy classes of subnormal subgroups, *Rend. Sem. Mat. Univ. Padova* **81** (1989), 107-149.
- [3] C. Casolo, Groups in which all subgroups are subnormal-by-finite, *Advances in Group Theory and Applications* **1** (2016), 3345 DOI: 10.4399/97888548908173
- [4] U. Dardano, S. Rinauro, Inertial automorphisms of an abelian group, *Rend. Sem. Mat. Univ. Padova* **127** (2012), 213-233. doi:10.4171/RSMUP/127-11
- [5] S. Franciosi, F. de Giovanni, M.L. Newell, Groups whose subnormal subgroups are normal-by-finite, *Comm. Alg.* **23(14)** (1995), 5483-5497.
- [6] H. Heineken, Groups with neighbourhood conditions for certain lattices. *Note di Matematica*, **1** (1996), 131143.
- [7] P. Longobardi, M. Maj, H. Smith. A note on locally graded groups. *Rend. Sem. Mat. Univ. Padova*, **94** (1995), 275-277.
- [8] W. Möhres, Torsionsgruppen, deren Untergruppen alle subnormal sind. *Geom. Dedicata*, **31** (1989), 237244.

- [9] B. H. Neumann, Groups with finite classes of conjugate subgroups, *Math. Z.* **63** (1955), 76-96.
- [10] D.J.S. Robinson, *A course in the theory of groups*, Graduate Texts in Mathematics, 80, Springer-Verlag, New York, 1996.
- [11] H. Smith, J. Wiegold, Locally graded groups with all subgroups normal-by-finite, *J. Austral. Math. Soc. Ser. A* **60** (1996), no. 2, 222-227.

Carlo Casolo, Dipartimento di Matematica U. Dini, Università di Firenze, Viale Morgagni 67A, I-50134 Firenze, Italy. email: casolo@math.unifi.it

Ulderico Dardano, Dipartimento di Matematica e Applicazioni "R. Caccioppoli", Università di Napoli "Federico II", Via Cintia - Monte S. Angelo, I-80126 Napoli, Italy. email: dardano@unina.it

Silvana Rinauro, Dipartimento di Matematica, Informatica ed Economia, Università della Basilicata, Via dell'Ateneo Lucano 10 - Contrada Macchia Romana, I-85100 Potenza, Italy. email: silvana.rinauro@unibas.it