

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/321530308>

# Microalgae biomass incorporation in salted cookies (“crackers”) increases their functional properties

Conference Paper · December 2017

CITATIONS

0

READS

16

8 authors, including:



Natascia Biondi

University of Florence

42 PUBLICATIONS 2,348 CITATIONS

[SEE PROFILE](#)



Mario Tredici

University of Florence

121 PUBLICATIONS 4,845 CITATIONS

[SEE PROFILE](#)



Isabel Sousa

University of Lisbon / Instituto Superior de Ag...

90 PUBLICATIONS 1,322 CITATIONS

[SEE PROFILE](#)



Anabela Raymundo

Instituto Piaget

61 PUBLICATIONS 1,266 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Using Freezing Technology in Sheep's Cheese Production [View project](#)



# ALGAE EUROPE

05-07 DEC 2017 | BERLIN | GERMANY

## BOOK OF ABSTRACTS



Ana Paula Batista<sup>1</sup>, Alberto Niccolai<sup>2</sup>, Ivana Bursic<sup>1</sup>, Liliana Rodolfi<sup>2,3</sup>, Natascia Biondi<sup>2</sup>, Mario R. Tredici<sup>2</sup>, Isabel Sousa<sup>1</sup>, Anabela Raymundo<sup>1</sup>

<sup>1</sup>LEAF – Linking Landscape, Environment, Agriculture and Food, Instituto Superior de Agronomia, Universidade de Lisboa, Portugal; <sup>2</sup>Department of Agrifood Production and Environmental Sciences (DISPAA), University of Florence, Italy; <sup>3</sup>Fotosintetica & Microbiologica S.r.l., Florence, Italy

#### Abstract:

In a previous work [1], microalgae sweet cookies were developed, but microalgae savoury foods might be more easily accepted due to their characteristic marine flavour. This study evaluated salted cookies (“crackers”) with 2% and 6% (w/w) of *Arthrospira platensis* F&M-C256, *Chlorella vulgaris* Allma, *Tetraselmis suecica* F&M-M33 and *Phaeodactylum tricornutum* F&M-M40 biomass, in terms of physicochemical properties, *in vitro* digestibility (IVD) and radical scavenging activity (RSA).

Microalgae crackers showed significantly higher ( $p<0.05$ ) protein content than the control (9.8% dw), with the highest values (>14% dw) for 6% *A. platensis* and *C. vulgaris*. The highest total dietary fiber (6.7% dw) was found for 6% *A. platensis* cracker while the control had 5.0% (dw). The addition of algae did not significantly increase the salt content in relation to the control (1.9% dw), except for 6% *T. suecica* (2.6% dw).

Microalgae crackers showed high IVD, with values ranging from 81.9% to 85.8%, with the highest value for *A. platensis*. These values are within the IVD range of commercial salted crackers.

The highest RSAs (24.4-24.6%, which corresponds to 0.25 mg/g<sub>alga</sub> of vitamin C equivalent antioxidant capacity) were found for *A. platensis* and *P. tricornutum* both at 6% incorporation. Total phenolics in microalgae crackers ranged from 1.1 to 2.4 mg of gallic acid equivalent per g of algal biomass, with the highest value for *T. suecica* 6%.

[1] Batista A.P. et al (2017) Algal Research, 26:161-171

*Acknowledgments:* Fundação para a Ciência e a Tecnologia (UID/AGR/04129/2013), COST Action ES1408.