


Special Publication 18

Hypogene Cave Morphologies



Edited by
Alexander Klimchouk
Ira D. Sasowsky
John Mylroie
Scott A. Engel
Annette Summers Engel

Special Publication 18



Hypogene Cave Morphologies

2014

Selected papers and abstracts of the
symposium held February 2 through 7, 2014,
San Salvador Island, Bahamas

Edited by
Alexander Klimchouk
Ira D. Sasowsky
John Mylroie
Scott A. Engel
Annette Summers Engel

Copyright © 2014 by the Karst Waters Institute, Inc. except where individual contributors to this volume retain copyright. All rights reserved with the exception of non-commercial photocopying for the purposes of scientific or educational advancement.

Published by: Karst Waters Institute, Inc.
P.O. Box 4142
Leesburg, Virginia 20177
<http://www.karstwaters.org>

Please visit our web page for ordering information.

The Karst Waters Institute is a non-profit 501 (c) (3) research and education organization incorporated in West Virginia. The mission of the Institute is to improve the fundamental understanding of karst water systems through sound scientific research and the education of professionals and the public.

ISBN Number: 978-0-9789976-7-0

Recommended citation for this volume:

A. Klimchouk, I. Sasowsky, J. Mylroie, S.A. Engel, and A.S. Engel, Eds., 2014, Hypogene Cave Morphologies. Selected papers and abstracts of the symposium held February 2 through 7, 2014, San Salvador Island, Bahamas. Karst Waters Institute Special Publication 18, Karst Waters Institute, Leesburg, Virginia. 111 p.

Published electronically.

Cover caption: Precision radial cave passage survey in progress at Altar Cave, San Salvador, Bahamas.
Photo credit: Ira D. Sasowsky

HYPOGENE CAVE MORPHOLOGIES

San Salvador, Bahamas

February 2-7, 2014

CONFERENCE ORGANIZERS

John Mylroie
Alexander Klimchouk
Ira D. Sasowsky

KARST WATERS INSTITUTE OFFICERS

President	Janet S. Herman, Ph.D.
Executive Vice-President	William B. White, Ph.D.
Comptroller	David C. Culver, Ph.D.
Secretary	Ira D. Sasowsky, Ph.D.
Treasurer of the Board	Harvey DuChene
Vice-President for Communications	Annette Summers Engel, Ph.D.
Vice-President for Education	Horton H. Hobbs, III, Ph.D.
Vice-President for Research	Carol Wicks, Ph.D.

KARST WATERS INSTITUTE BOARD OF DIRECTORS

Emily Davis	Horton H. Hobbs, III, Ph.D.	Ira D. Sasowsky, Ph.D.
Daniel Doctor, Ph.D.	William K. Jones (Chair)	Benjamin Schwartz, Ph.D.
Harvey R. DuChene	Jon B. Martin, Ph.D.	Steven Taylor, Ph.D.
John W. Hess, Ph.D.	Paul J. "PJ" Moore, Ph.D.	Dorothy Vesper, Ph.D. (Vice-Chair)
Matthew Covington, Ph.D.	Megan L. Porter, Ph.D.	William B. White, Ph.D.

TABLE OF CONTENTS

Preface.....	v
Keynote Papers	
Hypogene Speleogenesis – Discussion of Definitions <i>Dublyansky</i>	1
The Methodological Strength of the Hydrogeological Approach to Distinguishing Hypogene Speleogenesis <i>Klimchouk</i>	4
Contributed Papers	
Bahamian Caves and Blue Holes: Exquisitely Preserved Fossil Assemblages and Taphonomic Influences <i>Albury and Mylroie</i>	14
Hypogene Cave Patterns in Iron Ore Caves: Convergence of Forms or Processes? <i>Auler, Piló, Parker, Senko, Sasowsky, and Barton</i>	15
Earth Tide, A Potential Driver for Hypogenic Fluid Flow: Observations from a Submarine Cave in SW Turkey <i>Bayari and Ozyurt</i>	20
Preliminary Considerations on Hypogene Morphology in Toca da Boa Vista and Toca da Barriguda Caves, Northeastern Brazil <i>Borges, Casarin, Menezes, Srivastava, Silva, Bezerra, and Auler</i>	25
Facies Analyses and Diagenetic Evolution of the Hypogene Toca da Boa Vista/Toca da Barriguda Cave Complex, Salitre Formation (Irecê Basin, Brazil) – Preliminary Results <i>Cazarin, Menezes Santos, Srivastava, Vieira, Bezerra, and Borghi</i>	26
Modeling Speleogenesis using Computational Fluid Dynamics: Potential Applications to Hypogene Caves <i>Covington and Myre</i>	27
A Review on Hypogene Caves in Italy <i>De Waele, Galdenzi, Madonia, Menichetti, Parise, Piccini, Sanna, Sauro, Tognini, Vattano, and Vigna</i>	28
Sulfuric Acid Water Table Caves (Grotte du Chat / Acqua Fitusa / Bad Deutsch Altenburg + Kraushöhle) <i>De Waele, Plan, Audra, Vattano, and Madonia</i>	31
Morphological Effects of Condensation-Corrosion Speleogenesis at Devils Hole Ridge, Nevada <i>Dublyansky and Spötl</i>	36
Condensation Corrosion: Measurements and Geomorphic Evidence in the Frasassi Caves <i>Galdenzi, Jones, and Macalady</i>	44
Permian Hydrothermal Karst in Kraków Region (Southern Poland) and Its Peculiar Internal Sediments <i>Gradziński, Lewandowska, Paszkowski, Duliński, Nawrocki, and Żywiecki</i>	47
Hypogene Limestone Caves in Germany, Geochemical Background and Regionality <i>Kempe</i>	48
How Deep is Hypogene? Gypsum Caves in the South Harz <i>Kempe</i>	57
Reservoir Characteristics of the Complex Karst of the Lluçmajor Platform, Mallorca Island (Spain), Tool for Hydrocarbon Reservoir Appraisal <i>Lapointe</i>	65

TABLE OF CONTENTS

Hypogean Speleogenesis and CO ₂ : Suggestions from Karst of Italy <i>Menichetti</i>	72
Active Hypogene Speleogenesis in a Regional Karst Aquifer: Ayyalon Cave, Israel <i>Naaman, Dimentman, and Frumkin</i>	73
Hypogene vs Epigene Caves: The Sulfur and Oxygen Isotope Fingerprint <i>Onac</i>	75
Per Ascensum Cave Morphologies in Three Continents and One Island, Including Places Where They Shouldn't Occur <i>Osborne</i>	76
Helium Isotopes as Indicator of Current Hypogenic Karst Development in Taurids Karst Region, Turkey <i>Ozyurt and Bayari</i>	77
Evolving Interpretations of Hypogene Speleogenesis in the Black Hills, South Dakota <i>Palmer and Palmer</i>	82
Isotopic Studies of Byproducts of Hypogene Speleogenesis and Their Contribution to the Geologic Evolution of the Western United States <i>Polyak, Asmerom, Hill, Palmer, Provencio, Palmer, McIntosh, Decker, and Onac</i>	88
Devils Hole: Antikarst in the Mojave <i>Riggs</i>	95
Hypogene Paleokarst in the Triassic of the Dolomites (Northern Italy) <i>Riva</i>	97
Speleogenesis by the Sulfidic Springs at Northern Sierra de Chiapas, Mexico, Based on their Water Chemistry <i>Rosales-Lagarde and Boston</i>	101
Pondering the Importance of Subaerial Corrosion as a Speleogenetic Agent <i>Sasowsky</i>	102
Fingerprinting Water-Rock Interaction in Hypogene Speleogenesis: Potential and Limitations of Isotopic Depth-Profiling <i>Spötl and Dublyansky</i>	103

A REVIEW ON HYPOGENE CAVES IN ITALY

Jo De Waele¹, Sandro Galdenzi², Giuliana Madonia³, Marco Menichetti⁴, Mario Parise⁵, Leonardo Piccini⁶, Laura Sanna⁷, Francesco Sauro¹, Paola Tognini⁸, Marco Vattano³, and Bartolomeo Vigna⁹

Although hypogene cave systems have been described since the beginning of the 20th century, the importance in speleogenesis of ascending fluids that acquired their aggressiveness from in-depth sources has been fully realized only in the last decades. Aggressiveness of waters can be related to carbonic and sulfuric acids and the related corrosion-dissolution processes give rise to different types of caves and underground morphologies.

The abundance of hydrothermal springs and associated travertine deposits, and the widespread interaction between volcanic or sub-volcanic phenomena and karst in many sectors of the Italian peninsula are a strong evidence of hypogene speleogenesis. Furthermore, researches on secondary minerals have allowed to discover hypogene caves formed by highly acidic vapors in sub-aerial environments, also showing that most of these caves have extremely rich mineral associations.

Despite this, until the late 1980s the only known important cave systems of clear hypogene origin in Italy were considered to be the ones hosted in the Frasassi Canyon and Monte Cucco, in which important gypsum deposits undoubtedly showed that sulfuric acid played an important role in the creation of voids (Galdenzi, 1990, 2001; Galdenzi & Maruoka, 2003; Menichetti et al., 2007). Afterwards many other caves were categorized as formed by the sulfuric acid speleogenesis throughout the entire Apennines. Following the broad definition of hypogene caves by Palmer in 1991, and the even more general one of Klimchouk in the last decade (Klimchouk, 2007, 2009), the number of caves considered of hypogene origin in Italy has grown rapidly. Figure 1 shows the hypogene karst systems of Italy, including, besides the well-known and published ones, also the known and less studied, and presumed hypogene cave systems (see also Table 1).

More recently, in some of these caves detailed studies have been carried out including geomorphology, mineralogy, and geochemistry. Sulfuric acid caves are known from many regions along the Apennine chain (Tuscany, Umbria, Marche, Latium, Campania, Calabria) (Forti, 1985; Forti et al., 1989; Galdenzi and Menichetti, 1989, 1995; Galdenzi, 1997, 2001, 2009; Galdenzi et al., 2010; Piccini, 2000; Menichetti, 2009, 2011; Mecchia, 2012; De Waele et al., 2013b), but also from Piedmont, Apulia, Sicily (Vattano et al., 2013) and Sardinia (De Waele et al., 2013a). In this last region ascending fluids have also formed a hypogene cave in quartzite rock. Oxidation of sulfides can locally create hypogene cave morphologies in dominantly epigenic caves, such as in the Venetian forealps (this cave is not shown in Figure 1, being largely epigenic in origin) (Tisato et al., 2012). Ascending fluids have also created large solution voids in Messinian gypsum beds in Piedmont, and these can be defined hypogene caves according to the definition by Klimchouk (Vigna et al., 2010). Some examples of hypogene cave systems due to the rise of CO₂-rich fluids are also known in Liguria and Tuscany (Piccini, 2000). In the Alps and Prealps (Lombardy), some ancient high mountain karst areas exhibit evidences of an early hypogene origin, deeply modified and re-modeled by later epigenic processes. Hypogene morphologies are thus preserved as inactive features, and it is often difficult to distinguish them from epigenic ones.

At almost twenty years distance from the first review paper on hypogene cave systems in Central Italy by S. Galdenzi and M. Menichetti (1995), we give a review of the state-of-the-art knowledge on hypogene caves actually known from the whole of Italy.

¹Department of Biological, Geological and Environmental Sciences, Via Zamboni 67, 40126 Bologna, Italy, jo.dewaele@unibo.it, cescosauro@gmail.com

²Viale Verdi, 10 Jesi, Italy, galdenzi.sandro@tiscali.it

³Department of Earth and Marine Sciences, Via Archirafi 22, 90123 Palermo, Italy, marco.vattano@unipa.it, giuliana.madonia@unipa.it

⁴Department of Earth, Life and Environmental Sciences, Campus Scientifico Enrico Mattei, Via Cà Le Suore, 2/4 - 61029 Urbino, Italy, marco.menichetti@uniurb.it

⁵National Research Council, IRPI, Via Amendola 122-I, 70126 Bari, Italy, m.parise@ba.irpi.cnr.it

⁶Department of Earth Sciences, Via La Pira 4, 50121, Firenze, Italy, leonardo.piccini@unifi.it

⁷Institute for Biometeorology, National Research Council of Italy, Traversa La Crucca 3, 07100 Sassari, Italy, l.sanna@ibimet.cnr.it

⁸Gruppo Grotte Milano CAI-SEM; Progetto Speleologia Glaciale, paolatognini@iol.it

⁹Department of Territorial Engineering, of the Environment and of Geotechnologies, Politechnical University of Turin, Corso Duca degli Abruzzi 24, 10129 Torino, Italy, Bartolomeo.vigna@polito.it


Table 1. Hypogene cave systems of Italy ordered in well-known (studied), known and presumed ones. For location see Figure 1.

Cave system	Studied	Known	Presumed
1 Valle di Scerscen		X	
2 Valmalenco		X	
3 N Grigna		X	
4 Tremezzina		X	
5 Lezzeno			X
6 Mt Fenera			X
7 Moncalvo	X		
8 Grotta della Guerra			X
9 Scogli Neri		X	
10 Monsummano	X		
11 Mt Pisani		X	
12 Tassare-Mt Nerone	X		
13 Mt Rotondo	X		
14 Mt Cucco	X		
15 Frasassi	X		
16 Faggeto Tondo	X		
17 Mt di Campiglia			X
18 Mt. Civitello	X		
19 Mt Cetona		X	
20 Roselle		X	
21 Montecchio	X		
22 Parrano	X		
23 Pozzi della Piana	X		
24 Triponzo	X		
25 Acquasanta Terme	X		
26 Cittareale		X	
27 Buco del Pretaro	X		
28 Mt Soratte		X	
29 Pozzo del Merro	X		
30 Subiaco		X	
31 Fiume Coperto	X		
32 Mt Lattari		X	
33 S. Cesarea Terme	X		
34 Capo Palinuro	X		
35 Mt Sellaro	X		
36 Cassano allo Ionio	X		
37 Terme Luigiane		X	
38 Phaff	X		
39 S. Barbara	X		
40 Corona 'e Sa Craba	X		
41 Monte Inici	X		
42 Acqua Fitusa	X		
43 Mt Kronio	X		

REFERENCES

- De Waele J., Forti P., Naseddu A. 2013a. Speleogenesis of an ex-humed hydrothermal sulphuric acid karst in Cambrian carbonates (Mount San Giovanni, Sardinia). *Earth Surface Processes and Landforms*, 38: 1369-1379.
- De Waele J., Galli E., Piccini L., Rossi A. 2013b. Descrizione morfologica e mineralogica della grotta ipogenica sulfurea di Montecchio (Grosseto, Toscana). In: Cucchi F., Guidi P. (Eds.), *Atti del XXI Congresso Nazionale di Speleologia "Diffusione delle conoscenze"*, Trieste 2-5 giugno 2011: 380-386.
- Forti P. 1985. Le mineralizzazioni della grotta di Cala Fetente (Salerno, Campania). *Mondo Sotterraneo*, 1985(1-2): 41-50.
- Forti P., Menichetti M., Rossi A. 1989. Speleothems and speleogenesis of the Faggeto Tondo Cave (Umbria, Italy). In: Hazslinszky T., Takacsne B.K. (Eds.), *Proceedings of the 10th International Congress of Speleology*, Budapest, 1: 74-76.
- Galdenzi S. 1990. Un modello genetico per la Grotta Grande del Vento. *Memorie Istituto Italiano di Speleologia*, II(4): 123-142.
- Galdenzi S. 1997. Initial geological observations in caves bordering the Sibari plain (southern Italy). *Journal of Cave and Karst Studies*, 59: 81-86.
- Galdenzi S. 2001. L'azione morfogenetica delle acque sulfuree nelle Grotte di Frasassi, Acquasanta Terme (Appennino marchigiano-Italia) e di Movile (Dobrogea-Romania). *Le Grotte d'Italia*, V(2): 49-61.
- Galdenzi S., 2009. Hypogene caves in the Apennines (Italy). In: Klimchouk, A.B., Ford, D.C. (Eds.), *Hypogene Speleogenesis and Karst Hydrogeology of Artesian Basins*. Special Paper 1. Ukrainian Institute of Speleology and Karstology, Simferopol, pp. 101-116.
- Galdenzi S., Maruoka T. 2003. Gypsum deposits in the Frasassi caves, Central Italy. *Journal of Cave and Karst Studies*, 65: 111-125.
- Galdenzi S., Menichetti M. 1989. Evolution of underground karst systems in the Umbria-Marche Apennines in central Italy. In: Hazslinszky T., Takacsne K. (Eds.), *Proceedings of the 10th International Congress of Speleology*, Budapest, 3: 745-747.
- Galdenzi S., Menichetti M. 1995. Occurrence of hypogenic caves in a karst region: examples from central Italy. *Environmental Geology*, 26: 39-47.
- Galdenzi S., Cocchioni F., Filipponi G., Selvaggio R., Scuri S., Morichetti L., Cocchioni M. 2010. The sulfidic thermal caves of Acquasanta Terme (central Italy). *Journal of Cave and Karst Studies*, 72(1): 43-58.
- Klimchouk A.B. 2007. Hypogene speleogenesis. Hydrogeological and morphogenetic perspective. *National Cave and Karst Research Institute, Carlsbad, Special Paper Series 1*, 77 p.
- Klimchouk A.B. 2009. Morphogenesis of hypogenic caves. *Geomorphology*, 106: 100-117.

Figure 1. Italian karst areas and the hypogene karst systems (modified from Sivelli and De Waele, 2013, *Speleologia* 68, special issue printed for the 16th ICS Brno). wk = well known, k = known, p = presumed (GIS elaboration by M.L. Garberi).



Mecchia M. 2012. Indizi di speleogenesi ipogenica nelle grotte del Monte Soratte. *Notiziario dello Speleo Club Roma*, 16: 58-69.
 Menichetti M. 2009. Speleogenesis of the hypogenic caves in Central Italy. In: White W.B. (Ed.), *Proceedings of the 15th International Congress on Speleology*, Kerrville, Texas, pp. 909-915.

Menichetti M. 2011. Hypogenic caves in western Umbria (Central Italy). *Acta Carsologica*, 40(1): 129-145.

Menichetti M., Chirencio M.I., Onac B., Bottrell S. 2007. Depositi di gesso nelle grotte del Monte Cucco e della Gola di Frasassi. Considerazioni sulla speleogenesi. In: *Atti Congresso Nazionale di Speleologia*, Iglesias, pp. 308-325.

Palmer A.N. 1991. Origin and morphology of limestone caves. *Geological Society of America Bulletin*, 103: 1-21.

Piccini L. 2000. Il carsismo di origine idrotermale del Colle di Monsummano (Pistoia-Toscana). *Le Grotte d'Italia*, 1: 33-43

Sivelli M., De Waele J. (Eds.). 2013. A journey across speleological Italy. Map in Scale 1:1,500,000, Società Speleologica Italiana, attached to *Speleologia* 68.

Tisato N., Sauro F., Bernasconi S.M., Bruijn R.H.C., De Waele J. 2012. Hypogenic contribution to speleogenesis in a predominant epigenic karst system: A case study from the Venetian Alps, Italy. *Geomorphology*, 151/152: 156-163.

Vattano M., Audra P., Benvenuto F., Bigot J.-Y., De Waele J., Galli E., Madonia G., Nobécourt J.-C. 2013. Hypogenic caves of Sicily (southern Italy). In: Filippi M., Bosak P. (Eds.), *Proceedings of the 16th International Congress of Speleology*, Brno, 19-27 July 2013, 3: 144-149.

Vigna B., Fiorucci A., Banzato C., Forti P., De Waele J. 2010. Hypogenic gypsum karst and sinkhole formation at Moncalvo (Asti, Italy). *Zeitschrift für Geomorphologie N.F.*, 54, suppl. 2: 285-306.