



Dithiocarbamates with potent inhibitory activity against the *Saccharomyces cerevisiae* β -carbonic anhydrase

Murat Bozdag, Fabrizio Carta, Daniela Vullo, Semra Isik, Zeid AlOthman, Sameh M. Osman, Andrea Scozzafava & Claudiu T. Supuran

To cite this article: Murat Bozdag, Fabrizio Carta, Daniela Vullo, Semra Isik, Zeid AlOthman, Sameh M. Osman, Andrea Scozzafava & Claudiu T. Supuran (2016) Dithiocarbamates with potent inhibitory activity against the *Saccharomyces cerevisiae* β -carbonic anhydrase, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31:1, 132-136, DOI: [10.3109/14756366.2015.1010529](https://doi.org/10.3109/14756366.2015.1010529)

To link to this article: <https://doi.org/10.3109/14756366.2015.1010529>



Published online: 11 Feb 2015.



Submit your article to this journal [↗](#)



Article views: 201



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 8 View citing articles [↗](#)

RESEARCH ARTICLE

Dithiocarbamates with potent inhibitory activity against the *Saccharomyces cerevisiae* β -carbonic anhydrase

Murat Bozdogan¹, Fabrizio Carta¹, Daniela Vullo¹, Semra Isik¹, Zeid AlOthman², Sameh M. Osman², Andrea Scozzafava¹, and Claudiu T. Supuran^{1,3}

¹Laboratorio di Chimica Bioinorganica, Dipartimento di Chimica, Università degli Studi di Firenze, Sesto Fiorentino, Florence, Italy, ²Department of Chemistry, College of Science, King Saud University, Riyadh, Saudi Arabia, and ³Sezione di Scienze Farmaceutiche, Dipartimento NEUROFARBA, Università degli Studi di Firenze, Sesto Fiorentino, Florence, Italy

Abstract

Dithiocarbamates (DTCs) prepared from primary or secondary amines, which incorporated amino/hydroxyl-alkyl, mono-/bicyclic aliphatic/heterocyclic rings based on the quinuclidine, piperidine, hydroxy-/carboxy-/amino-substituted piperidine, morpholine and piperazine scaffolds, were investigated for the inhibition of α - and β -carbonic anhydrases (CAs, EC 4.2.1.1) of pharmacologic relevance, such as the human (h) isoform hCA I and II, as well as the *Saccharomyces cerevisiae* β -CA, scCA. The yeast and its β -CA were shown earlier to be useful models of pathogenic fungal infections. The DTCs investigated here were medium potency hCA I inhibitors (K_i s of 66.5–910 nM), were more effective as hCA II inhibitors (K_i s of 8.9–107 nM) and some of them showed excellent, low nanomolar activity against the yeast enzyme, with inhibition constants ranging between 6.4 and 259 nM. The detailed structure activity relationship for inhibition of the yeast and human enzymes is discussed. Several of the investigated DTCs showed excellent selectivity ratios for inhibiting the yeast over the human cytosolic CA isoforms.

Keywords

β -Class enzyme, carbonic anhydrase, dithiocarbamate, inhibitor, *Saccharomyces cerevisiae*

History

Received 16 December 2014
Revised 15 January 2015
Accepted 15 January 2015
Published online 6 February 2015

Introduction

The primary sulfonamides possessing the general formula RSO_2NH_2 (where R may be an aromatic, heterocyclic or aliphatic moiety, but also an even smaller groups such as OH and NH_2)^{1–7} dominated the landscape of carbonic anhydrase (CA, EC 4.2.1.1) inhibitors (CAIs) drug design for 70 years, since the discovery that such compounds inhibit the enzyme, in the 1940s^{8–15}. However, recently, new important chemotypes with such properties emerged, by combining structure-based drug discovery strategies (X-ray crystallography and fragment-based drug design) with massive campaigns of screening vast library of compounds for detecting such an activity^{16–38}. The interest in CAIs is mainly motivated by their pharmacological properties and clinical use as diuretics, antiglaucoma, antiobesity or antiepileptic agents, as well as anticancer agents/diagnostic tools^{1–3,29–39}. Among the new chemotypes with CA inhibitory properties ultimately reported, the dithiocarbamates (DTCs) are undoubtedly among the most interesting ones^{40–44}.

These compounds have been rationally discovered as CAIs after our report of trithiocarbonate (CS_3^{2-} , TTC), an inorganic anion similar to carbonate, as an interesting (milli–micromolar)

CAI⁴⁵. In the X-ray crystal structure of this anion bound to hCA II, a monodentate coordination of the inhibitor through one sulfur atom to the zinc ion from the enzyme active site has been observed. Furthermore, a hydrogen bond in which another sulfur of TTC and the OH of Thr199 (a crucial residue for the catalytic cycle of α -CAs)^{1–3} were involved, also stabilized the anion when bound within the enzyme active site. Thus, the CS_2^- moiety present in TTC was detected as a new zinc-binding group (ZBG) for CA inhibition. As DTCs incorporate this ZBG, a first series of such compounds was prepared and evaluated for their inhibitory activity against several mammalian, fungal and bacterial CAs by this group^{40–44}. Several low nanomolar and subnanomolar CAIs were thus detected against all these enzymes. X-ray crystal structures were also reported for three DTCs complexed to hCA II, i.e. compounds A–C (Figure 1)^{40,41}. DTCs A–C inhibited hCA II with K_i s of 25, 41 and 0.95 nM, respectively, and hCA IX (an isoform involved in cancerogenesis)^{1,3,5} with K_i s of 53, 757 and 6.2 nM, respectively^{40,41}. As seen from Figure 1, the binding mode of the new ZBG present in these compounds was identical to that of TTC, with one sulfur coordinated to the metal ion, but the organic scaffold present in the DTC was observed to make extensive contacts with many amino acid residues from the active site. This explains the wide range of inhibitory power of these derivatives, from the subnanomolar to the micromolar range, for the entire series of around 30 DTC reported so far^{40–44}. Interestingly, these compounds are highly water soluble (being salts), and the DTC C was also effective *in vivo* as an

Address for correspondence: Daniela Vullo, Dipartimento di Chimica, Laboratorio di Chimica Bioinorganica, Università degli Studi di Firenze, Rm. 188, Via della Lastruccia 3, Sesto Fiorentino, Florence 50019, Italy. Tel/Fax: +39 055 4573143. E-mail: daniela.vullo@unifi.it




Figure 1. (A) Structure of DTCs A–C used as lead molecules in the present study. (B) Electronic density for the adduct of dithiocarbamate B bound within the active site of hCA II^{40,41}. The zinc ion is shown as the central sphere and the amino acid residues involved in the binding are evidenced and numbered (hCA I numbering system)⁴⁰. The zinc ligands (except the sulfur from the DTC compound), His94, 96 and 119, are not shown for the sake of simplicity.

antiglaucoma agent when administered topically directly to the eye of hypertensive rabbits, a widely used animal model of this disease⁴¹. However, for the moment only one series of DTCs was reported and investigated as CAIs, and this is the reason why we extend here the earlier studies^{40,41}, reporting the CA inhibitory properties of a new series of such compounds against another CA of interest, the β -class enzyme from the yeast, *Saccharomyces cerevisiae* scCA^{46–48}, a model organism of great relevance when investigating fungal CAs as drug targets^{49–51}.


Materials and methods

Chemistry

The DTCs **1b–21b** investigated as scCA inhibitors were prepared by reaction of the amines **1a–21a** with CS₂ in the presence of bases, such as NaOH or KOH⁵².

CA inhibition

A stopped-flow instrument (SX.18MV-R Applied Photophysics model) was used for assaying the CA-catalyzed CO₂ hydration activity⁵³. Inhibitor and enzyme were preincubated for 15 min for allowing the complete formation of the enzyme-inhibitor adduct. IC₅₀ values were obtained from dose response curves working at seven different concentrations of test compound (from 0.1 nM to 50 μ M), by fitting the curves using PRISM (www.graphpad.com) and non-linear least squares methods, the obtained values representing the mean of at least three different determinations⁵⁴. The inhibition constants (K_i) were derived from the IC₅₀ values by using the Cheng–Prusoff equation, as follows: $K_i = IC_{50}/(1 + [S]/K_m)$, where $[S]$ represents the CO₂ concentration at which the measurement was carried out and K_m the concentration of substrate at which



Scheme 1. Preparation of DTCs **1b–21b**, by reaction of amines **1a–21a** with carbon disulfide in the presence of sodium/potassium hydroxide.

the enzyme activity is at half maximal. All enzymes used were recombinant, produced in *Escherichia coli* as reported earlier^{13,47}. The concentrations of enzymes used in the assay were: hCA I, 12.0 nM; hCA II, 9.2 nM and scCA, 15.3 nM.

Results and discussion




In the present article, we extended the series of DTCs investigated earlier^{40–44}, including both primary as well as secondary derivatives, which were obtained in such a way as to explore novel chemical space. The starting amines (**1a–21a**) used to synthesize DTCs **1b–21b** reported here (Scheme 1) included *N,N*-dimethylaminoethylenediamine **1a**, aminoalcohols with three to five carbon atoms in their molecule **2a–4a**, the bicyclic quinuclidine-3-amine (both the racemate as well as the R- and S-enantiomeric DTCs **5b–7b** incorporating this scaffold were obtained), piperidine **8a** and several of its derivatives with hydroxyl-, carboxy-, acetamido- and boc-amido functionalities in various positions of the heterocyclic ring, of types **9a–16a**; morpholine and piperazine derivatives **17a–19a**, as well as phenethylamine **20a** and its sulfamoylated derivative, 4-aminoethylbenzenesulfonamide **21a**, a well know CAI which binds to the enzyme through the sulfamoyl moiety^{1,3} (Scheme 1). The choice of these scaffolds was motivated by the fact that the structure–activity relationship (SAR) for the inhibition of CAs with the DTCs reported earlier^{7,8} was primarily influenced by the organic scaffold of the inhibitor. In fact, important differences of activity were observed between primary and secondary DTCs and between aliphatic or aromatic/heterocyclic derivatives, respectively^{40–44}. Here we investigated compounds belonging to these DTCs types, with a variety of substitution patterns, based on several scaffolds mentioned earlier (Table 1) among which primary, aliphatic aminoalkyl- or hydroxyalkyl derivatives (**1b–4b**); primary, bicyclic, bulky DTCs (**5b–7b**), with various stereochemistries; secondary, piperidine, morpholine and piperazine-based DTCs, of types **8b–19b**, for which the lead compound was the morpholine-DTC (compound **C**) reported earlier⁴⁰. Indeed, the largest number of the new DTCs investigated here belongs to this subgroup.

We investigated the CA inhibitory properties of DTCs **1b–21b**, as well as the sulfonamide CAI acetazolamide (**AAZ**) as standard, against three relevant CAs, the human (h) cytosolic CA isoforms, hCA I and II (widespread, off-target enzymes when non-mammalian CAs should be inhibited) as well as the yeast scCA, an enzyme belonging to the β -class, which we showed earlier to be a good model for studying fungal enzymes belonging to this class¹³. Indeed, several pathogenic fungi, such as *Cryptococcus neoformans*, *Candida albicans* and *C. glabrata*, encode β -CAs homologous to scCA^{49–51}.

The inhibition data of compounds **1b–21b** are shown in Table 1, and the following SAR can be delineated:


- The slow isoform hCA I was poorly inhibited by most of the DTCs investigated here. Few compounds, i.e. **1b**, **4b**, **18b** and **21b** showed inhibition constants ranging between 66.5 and 97.5 nM, being rather efficient hCA I inhibitors. The remaining derivatives were weaker inhibitors, with inhibition constants ranging between 109 and 910 nM, and were much less efficient compared to the clinically used

Table 1. hCA I, II and scCA inhibition data with DTCs **1b–21b** and acetazolamide (AAZ, 5-acetamido-1,3,4-thiadiazole-2-sulfonamide) as standard drug, by a stopped-flow CO₂ hydase assay.

No	R	R ¹	K _I (nM)*		
			hCA I	hCA II	scCA
1b	Me ₂ N(CH ₂) ₂	H	85.9	35.8	82.4
2b	HO(CH ₂) ₃	H	706	41.7	72.5
3b	HO(CH ₂) ₄	H	295	24.3	41.4
4b	HO(CH ₂) ₅	H	66.5	17.3	20.5
5b		H	494	48.7	216
6b (R)		H	240	18.9	278
7b (S)		H	615	65.9	181
8b	–(CH ₂) ₅ –		252	30.1	13.8
9b	–(CH ₂) ₃ –CH(OH)CH ₂ –		428	60.7	19.1
10b	–(CH ₂) ₄ –CH(COONa)–		485	80.1	10.5
11b	–(CH ₂) ₃ –CH(COONa)CH ₂ –		290	45.4	18.7
12b (R)	–(CH ₂) ₃ –CH(COONa)CH ₂ –		496	80.5	9.1
13b (S)	–(CH ₂) ₃ –CH(COONa)CH ₂ –		109	8.9	23.8
14b	–(CH ₂) ₂ –CH(COONa)(CH ₂) ₂ –		337	78.7	6.4
15b	–(CH ₂) ₃ –CH(NHAc)CH ₂ –		910	47.9	45.9
16b	–(CH ₂) ₃ –CH(NHBoc)CH ₂ –		683	13.2	24.1
17b	–CH(Me)CH ₂ –O–(CH ₂) ₂ –		434	60.2	16.5
18b	–CH(COONa)CH ₂ –O–(CH ₂) ₂ –		84.7	78.5	18.4
19b	–(CH ₂) ₂ N(CH ₂ CONHC ₆ H ₁₁)(CH ₂) ₂ –		415	67.2	259
20b	Ph(CH ₂) ₂	H	425	107	51.5
21b	H ₂ NO ₂ SC ₆ H ₄ (CH ₂) ₂	H	97.5	48.1	78.9
AAZ	–	–	250	12	82.6

*Mean from three different assays. Errors were in the range of ± 5 –10% of the reported values (data not shown).

sulfonamide acetazolamide (AAZ, K_I of 250 nM), Table 1. This is a positive feature for a CAI, since hCA I inhibition is associated with side effects of the systemically or topically acting sulfonamides, such as AAZ or dorzolamide DZA^{1,3,5}.



The SAR for the inhibition of this isoform is straightforward: primary DTCs were the most effective CAIs detected here (e.g. **1b**, **4b** and **21b**), whereas among the secondary DTCs only the 3-morpholine-carboxylic acid derivative **18b** was a good inhibitor. Among the hydroxyalkyl-substituted DTCs **2b–4b**, it may be observed that hCA I inhibitory power was very weak for the compound with three carbon atoms (**2b**), increased considerably for its homolog with four carbon atoms (**3b**) whereas for the 5-aminopentanol derivative **4b** it reached a good level, with a K_I of 66.5 nM

(a > 10 times increase of the inhibitory power with an increase of the aliphatic chain from 3 to 5 CH₂ groups).

- (ii) The physiologically dominant cytosolic isoform hCA II was more sensitive to inhibition by the compounds investigated here compared to hCA I. DTCs **1b–21b** showed inhibition constants ranging between 8.9 and 107 nM (AAZ has a K_I of 12 nM against this isoform, Table 1). Most of the new DTCs were medium potency hCA II inhibitors, with K_Is ranging between 30.1 and 80.5 nM. They include the following DTCs: **1b**, **2b**, **5b**, **7b–12b**, **14b**, **15b**, **17b–19b** and **21b**. The most ineffective hCA II inhibitor was the phenethyl derivative **20b** (K_I of 107 nM), whereas the sulfamoylated analog **21b** showed an improved efficacy, with a K_I of 48.1 nM. The best hCA II inhibitors (K_Is in the range of 8.9–24.3 nM), were **3b**, **4b**, **6b**, **13b** and **16b** (Table 1). These effective inhibitors belong to both classes of DTCs, primary (**3b**, **4b** and **6b**) and secondary ones (**13b** and **16b**), respectively. They incorporate the hydroxyalkyl moieties, and as for hCA I, the inhibition efficiency increases with the length of the aliphatic chain from 3 to 5. Another interesting case was observed for the quinuclidine derivatives. The racemate **5b** was a medium potency inhibitor (K_I of 48.7 nM) whereas the R-stereoisomer was 2.57 times a more effective inhibitor than the racemic compound, with a K_I of 18.9 nM. On the contrary, the S-stereoisomer was the least effective hCA II inhibitor among these derivatives, with a K_I of 65.9 nM (3.5 times less effective compared to its antipode **6b**). The same behavior was observed for the other racemate and the two enantiomers investigated here, the derivatives of nipecotic acid **11b–13b**. In this case the S-enantiomer **13b** was 9 times a better hCA II inhibitor compared to its antipode **12b**, whereas the racemate **11b** had an intermediate behavior between the two diastereoisomers, with a K_I of 45.4 nM (Table 1). The position of the other functional group on the heterocyclic ring on which the CS₂[–] moiety was attached was also an important factor influencing the activity of the heterocyclic DTCs investigated here. The regiomeric DTCs derived from pipecolic, nipecotic and isonipecotic acids (**10b**, **11b** and **14b**, respectively) had quite different activities: the nipecotic acid derivative **14b** was roughly two times more effective as a hCA II inhibitor compared to its isomers **10b** and **11b**. The two amine derivatives **15b** and **16b** were also different in their affinity for hCA II, with the bulkier, Boc derivative **16b** being a potent inhibitor (K_I of 13.2 nM), whereas the acetamido one **15b** having a weaker activity (K_I of 47.9 nM). All these data show that rather small differences in the scaffold of these DTCs lead to quite different inhibitory activities, probably due to the fact that as explained earlier, the scaffold of the inhibitor participates in many interactions with amino acid residues from the CA active site, which may stabilize or in case of clashes, weaken the inhibitory power of the corresponding compound.
- (iii) scCA was also effectively inhibited by DTCs **1b–21b**, which showed inhibition constants in the range of 6.4–278 nM. The SAR for the inhibition of the yeast enzyme is more complicated compared to the ones outlined above for the inhibition of the two human cytosolic isoforms. Thus, two DTCs, the quinuclidine derivatives **5b–7b** and the bulkier derivative **19b**, behaved as weak scCA inhibitors, with K_Is of 181–278 nM. It may be observed that all of them are rather bulky, sterically hindered compounds which probably cannot easily accommodate within the scCA active site which is a long and not very wide channel^{48–50}. Another group of DTCs, among which **1b–3b**, **15b**, and **20b**, **21b**, similarly to AAZ (K_I of 82.6 nM) showed medium potency

activity against scCA, with K_{1s} ranging between 41.1 and 82.4 nM (Table 1). These compounds incorporate the amino/hydroxyalkyl chains with two to four carbon atoms (**1b–3b**), arylalkyl chains (as in **20b** and **21b**) or the acetamido-piperidine scaffold present in **15b** (which is in fact an outlier, as all other piperidine, and morpholine DTCs investigated here showed a better inhibition profile against scCA compared to **15b**). Indeed, the most effective scCA inhibitors were just these derivatives, i.e. **8b–14b**, **16b–18b** (K_{1s} of 6.4–24.1 nM) together with the hydroxyl-pentyl DTC **4b** (K_{1s} of 20.5 nM). Thus, for the primary aliphatic derivatives **2b–4b**, scCA inhibitory activity increased with the length of the carbon atoms chain. For the piperidine derivatives activity was better when another functional group (such as OH, COOH, BocNH) was attached on the ring (compared to the parent derivative of the series, **8b**), but the substitution pattern and the stereochemistry were factors which influenced the activity considerably. For example the nipecotic acid **11b–13b** are again of interest to analyze: the R-enantiomer **12b** was highly effective (K_{1s} of 9.1 nM, the same potency as the hydroxyl-substituted piperidine **10b**), whereas the S-enantiomer **13b** had a 2.5 times lower activity (K_{1s} of 23.8 nM). The best scCA inhibitor was however the isonipecotic acid derivative **14b**, which had a K_{1s} as low as 6.4 nM, being the best scCA inhibitor reported to date. The morpholine DTCs **17b** and **18b** also had a good scCA inhibition profile, with K_{1s} of 16.5–18.4 nM. It should be noted that many DTCs investigated here were several times better scCA inhibitors compared to **AAZ**, a clinically used sulfonamide inhibitor, but a moderate scCA inhibitor. Another important feature, some of the DTCs investigated here showed selectivity for the inhibition of the yeast over the human enzymes. Examples of such derivatives are **8b–12b**, **14b**, **17b** and **20b**. In the case of **14b**, the selectivity ratio for inhibiting scCA over hCA I was of 52.6 and of inhibiting scCA over hCA II of 12.3, making this derivative one of the most selective inhibitor for the yeast over the human CA isoforms.

Conclusion

We investigated the inhibition of two human, off-target CA isoforms and the yeast β -class enzyme scCA with a series of DTCs prepared both from primary and secondary amines and carbon disulfide, in the presence of strong bases. These DTCs incorporated amino/hydroxyl-alkyl, mono-/bicyclic aliphatic/heterocyclic rings based on the quinuclidine, piperidine, hydroxy-/carboxy-/amino-substituted piperidine, morpholine and piperazine scaffolds. The investigated compounds were medium potency hCA I inhibitors (K_{1s} of 66.5–910 nM), they were more effective as hCA II inhibitors (K_{1s} of 8.9–107 nM), and some of them showed excellent, low nanomolar activity against the yeast enzyme, with inhibition constants ranging between 6.4 and 259 nM. The detailed structure activity relationship for inhibition of the yeast and human enzymes is discussed. Several of the investigated DTCs showed excellent selectivity ratios for inhibiting the yeast over the human cytosolic CA isoforms.

Declaration of interest

This research was financed in part by two EU projects (Dynano and Metoxia). The authors declare no conflict of interest.

References

- Supuran CT. Structure-based drug discovery of carbonic anhydrase inhibitors. *J Enzyme Inhib Med Chem* 2012;27:759–72.
- Alterio V, Di Fiore A, D'Ambrosio K, et al. Multiple binding modes of inhibitors to carbonic anhydrases: how to design specific drugs targeting 15 different isoforms? *Chem Rev* 2012;112:4421–68.
- Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nat Rev Drug Discov* 2008;7:168–81.
- Capasso C, Supuran CT. Sulfa and trimethoprim-like drugs-antimetabolites acting as carbonic anhydrase, dihydropteroate synthase and dihydrofolate reductase inhibitors. *J Enzyme Inhib Med Chem* 2014;29:379–87.
- Monti SM, Supuran CT, De Simone G. Anticancer carbonic anhydrase inhibitors: a patent review (2008–2013). *Expert Opin Ther Pat* 2013;23:737–49.
- Supuran CT. Carbonic anhydrase inhibitors. *Bioorg Med Chem Lett* 2010;20:3467–3474.
- Rummer JL, McKenzie DJ, Innocenti A, et al. Root effect hemoglobin may have evolved to enhance general tissue oxygen delivery. *Science* 2013;340:1327–9.
- Harju AK, Bootorabi F, Kuuslahti M, et al. Carbonic anhydrase III: a neglected isozyme is stepping into the limelight. *J Enzyme Inhib Med Chem* 2013;28:231–9.
- Supuran CT. Carbonic anhydrases: from biomedical applications of the inhibitors and activators to biotechnologic use for CO₂ capture. *J Enzyme Inhib Med Chem* 2013;28:229–30.
- Çavdar H, Ekinci D, Talaz O, et al. α -Carbonic anhydrases are sulfatases with cyclic diol monosulfate esters. *J Enzyme Inhib Med Chem* 2012;27:148–54.
- De Simone G, Alterio V, Supuran CT. Exploiting the hydrophobic and hydrophilic binding sites for designing carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2013;8:793–810.
- Durdagi S, Scozzafava G, Vullo D, et al. Inhibition of mammalian carbonic anhydrases I–XIV with grayanotoxin III: solution and in silico studies. *J Enzyme Inhib Med Chem* 2014;29:469–75.
- Bilginer S, Unluer E, Gul HI, et al. Carbonic anhydrase inhibitors. Phenols incorporating 2- or 3-pyridyl-ethenylcarbonyl and tertiary amine moieties strongly inhibit *Saccharomyces cerevisiae* β -carbonic anhydrase. *J Enzyme Inhib Med Chem* 2014;29:495–9.
- Sharma A, Tiwari M, Supuran CT. Novel coumarins and benzocoumarins acting as isoform-selective inhibitors against the tumor-associated carbonic anhydrase IX. *J Enzyme Inhib Med Chem* 2014;29:292–6.
- Pan J, Lau J, Mesak F, et al. Synthesis and evaluation of ¹⁸F-labeled carbonic anhydrase IX inhibitors for imaging with positron emission tomography. *J Enzyme Inhib Med Chem* 2014;29:249–55.
- Fabrizi F, Mincione F, Somma T, et al. A new approach to antiglaucoma drugs: carbonic anhydrase inhibitors with or without NO donating moieties. Mechanism of action and preliminary pharmacology. *J Enzyme Inhib Med Chem* 2012;27:138–47.
- Ekinci D, Karagoz L, Ekinci D, et al. Carbonic anhydrase inhibitors: *in vitro* inhibition of α isoforms (hCA I, hCA II, hCA III, hCA IV) by flavonoids. *J Enzyme Inhib Med Chem* 2013;28:283–8.
- Carta F, Supuran CT, Scozzafava A. Novel therapies for glaucoma: a patent review 2007–2011. *Expert Opin Ther Pat* 2012;22:79–88.
- Masini E, Carta F, Scozzafava A, Supuran CT. Antiglaucoma carbonic anhydrase inhibitors: a patent review. *Expert Opin Ther Pat* 2013;23:705–16.
- Supuran CT, Maresca A, Gregañ F, Remko M. Three new aromatic sulfonamide inhibitors of carbonic anhydrases I, II, IV and XII. *J Enzyme Inhib Med Chem* 2013;28:289–93.
- Alp C, Maresca A, Alp NA, et al. Secondary/tertiary benzenesulfonamides with inhibitory action against the cytosolic human carbonic anhydrase isoforms I and II. *J Enzyme Inhib Med Chem* 2013;28:294–8.
- Carta F, Scozzafava A, Supuran CT. Sulfonamides (RSO₂NH₂): a patent review 2008–2012. *Expert Opin Ther Pat* 2012;22:747–58.
- Supuran CT. Carbonic anhydrase inhibitors: an editorial. *Expert Opin Ther Pat* 2013;23:677–9.
- Wilkinson BL, Bornaghi LF, Houston TA, et al. Carbonic anhydrase inhibitors: inhibition of isozymes I, II and IX with triazole-linked O-glycosides of benzene sulfonamides. *J Med Chem* 2007;50:1651–7.
- Capasso C, Supuran CT. Antiinfective carbonic anhydrase inhibitors: a patent and literature review. *Expert Opin Ther Pat* 2013;23:693–704.
- Supuran CT. Carbonic anhydrase inhibitors as emerging drugs for the treatment of obesity. *Expert Opin Emerg Drugs* 2012;17:11–5.

27. Carta F, Supuran CT. Diuretics with carbonic anhydrase inhibitory action: a patent and literature review (2005–2013). *Expert Opin Ther Pat* 2013;23:681–91.
28. Scozzafava A, Supuran CT, Carta F. Antiobesity carbonic anhydrase inhibitors: a literature and patent review. *Expert Opin Ther Pat* 2013; 23:725–35.
29. D'Ascenzio M, Carradori S, Secci D, et al. Selective inhibition of human carbonic anhydrases by novel amide derivatives of probenecid: synthesis, biological evaluation and molecular modelling studies. *Bioorg Med Chem* 2014;22:3982–8.
30. Bonneau A, Maresca A, Winum JY, Supuran CT. Metronidazole-coumarin conjugates and 3-cyano-7-hydroxy-coumarin act as isoform-selective carbonic anhydrase inhibitors. *J Enzyme Inhib Med Chem* 2013;28:397–401.
31. Allouche F, Chabchoub F, Carta F, Supuran CT. Synthesis of aminocyanopyrazoles via a multi-component reaction and anti-carbonic anhydrase inhibitory activity of their sulfamide derivatives against cytosolic and transmembrane isoforms. *J Enzyme Inhib Med Chem* 2013;28:343–9.
32. Del Prete S, De Luca V, Scozzafava A, et al. Biochemical properties of a new α -carbonic anhydrase from the human pathogenic bacterium *Vibrio cholerae*. *J Enzyme Inhib Med Chem* 2014;29: 23–7.
33. Singh S, Supuran CT. 3D-QSAR CoMFA studies on sulfonamide inhibitors of the Rv3588c β -carbonic anhydrase from *Mycobacterium tuberculosis* and design of not yet synthesized new molecules. *J Enzyme Inhib Med Chem* 2014;29:449–55.
34. Gieling RG, Parker CA, De Costa LA, et al. Inhibition of carbonic anhydrase activity modifies the toxicity of doxorubicin and melphalan in tumour cells in vitro. *J Enzyme Inhib Med Chem* 2013;28:360–9.
35. Maresca A, Vullo D, Scozzafava A, Supuran CT. Inhibition of the alpha- and beta-carbonic anhydrases from the gastric pathogen *Helicobacter pylori* with anions. *J Enzyme Inhib Med Chem* 2013; 28:388–91.
36. Alp C, Özsoy S, Alp NA, et al. Sulfapyridine-like benzenesulfonamide derivatives as inhibitors of carbonic anhydrase isoenzymes I, II and VI. *J Enzyme Inhib Med Chem* 2012;27:818–24.
37. Maresca A, Scozzafava A, Vullo D, Supuran CT. Dihalogenated sulfanilamides and benzolamides are effective inhibitors of the three β -class carbonic anhydrases from *Mycobacterium tuberculosis*. *J Enzyme Inhib Med Chem* 2013;28:384–7.
38. Koz O, Ekinci D, Perrone A, et al. Analysis of saponins and phenolic compounds as inhibitors of α -carbonic anhydrase isoenzymes. *J Enzyme Inhib Med Chem* 2013;28:412–7.
39. Ebbesen P, Pettersen EO, Gorr TA, et al. Taking advantage of tumor cell adaptations to hypoxia for developing new tumor markers and treatment strategies. *J Enzyme Inhib Med Chem* 2009;24(S1):1–39.
40. Carta F, Aggarwal M, Maresca A, et al. Dithiocarbamates: a new class of carbonic anhydrase inhibitors. Crystallographic and kinetic investigations. *Chem Commun (Cambridge)* 2012;48:1868–70.
41. Carta F, Aggarwal M, Maresca A, et al. Dithiocarbamates strongly inhibit carbonic anhydrases and show antiglaucoma action in vivo. *J Med Chem* 2012;55:1721–30.
42. Monti SM, Maresca A, Viparelli F, et al. Dithiocarbamates strongly inhibit the beta-class fungal carbonic anhydrases from *Cryptococcus neoformans*, *Candida albicans* and *Candida glabrata*. *Bioorg Med Chem Lett* 2012;22:859–62.
43. Maresca A, Carta F, Vullo D, Supuran CT. Dithiocarbamates strongly inhibit the beta-class carbonic anhydrases from *Mycobacterium tuberculosis*. *J Enzyme Inhib Med Chem* 2013;28: 407–11.
44. Avram S, Milac AL, Carta F, Supuran CT. More effective dithiocarbamate derivatives inhibiting carbonic anhydrases, generated by QSAR and computational design. *J Enzyme Inhib Med Chem* 2013;28:350–9.
45. Temperini C, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors. X-ray crystal studies of the carbonic anhydrase II – trithiocarbonate adduct – an inhibitor mimicking the sulfonamide and urea binding to the enzyme. *Bioorg Med Chem Lett* 2010;20: 474–8.
46. Isik S, Kockar F, Arslan O, et al. Carbonic anhydrase inhibitors. Inhibition of the β -class enzyme from the yeast *Saccharomyces cerevisiae* with anions. *Bioorg Med Chem Lett* 2008;18:6327–31.
47. Isik S, Kockar F, Aydin M, et al. Carbonic anhydrase inhibitors. Inhibition of the β -class enzyme from the yeast *Saccharomyces cerevisiae* with sulfonamides and sulfamates. *Bioorg Med Chem* 2009;17:1158–63.
48. Isik S, Guler OO, Kockar F, et al. *Saccharomyces cerevisiae* β -carbonic anhydrase inhibition and activation studies. *Curr Pharm Des* 2010;16:3327–36.
49. Hall RA, Mühlischlegel FA. Fungal and nematode carbonic anhydrases: their inhibition in drug design. In: Supuran CT, Winum JY, eds. *Drug design of zinc-enzyme inhibitors: functional, structural, and disease applications*. Hoboken: John Wiley & Sons; 2009:301–22.
50. Ohndorf UM, Schlicker C, Steegborn C. Crystallographic studies on carbonic anhydrases from fungal pathogens for structure-assisted drug development. In: Supuran CT, Winum JY, eds. *Drug design of zinc-enzyme inhibitors: functional, structural, and disease applications*. Hoboken: John Wiley & Sons; 2009:323–34.
51. Carta F, Innocenti A, Hall RA, et al. Carbonic anhydrase inhibitors. Inhibition of the β -class enzymes from the fungal pathogens *Candida albicans* and *Cryptococcus neoformans* with branched aliphatic/aromatic carboxylates and their derivatives. *Bioorg Med Chem Lett* 2011;21:2521–6.
52. Bozdag M, Carta F, Vullo D, et al. New dithiocarbamates with effective carbonic anhydrase inhibitory activity. *J Med Chem*, submitted.
53. Khalifah RG. The carbon dioxide hydration activity of carbonic anhydrase. I. Stop-flow kinetic studies on the native human isoenzymes B and C. *J Biol Chem* 1971;246:2561–73.
54. Pacchiano F, Carta F, McDonald PC, et al. Ureido-substituted benzenesulfonamides potently inhibit carbonic anhydrase IX and show antimetastatic activity in a model of breast cancer metastasis. *J Med Chem* 2011;54:1896–902.