



Synthesis and carbonic anhydrase activating properties of a series of 2-amino-imidazolines structurally related to clonidine¹

Niccolò Chiaramonte, Soumia Maach, Caterina Biliotti, Andrea Angeli, Gianluca Bartolucci, Laura Braconi, Silvia Dei, Elisabetta Teodori, Claudiu T. Supuran & Maria Novella Romanelli

To cite this article: Niccolò Chiaramonte, Soumia Maach, Caterina Biliotti, Andrea Angeli, Gianluca Bartolucci, Laura Braconi, Silvia Dei, Elisabetta Teodori, Claudiu T. Supuran & Maria Novella Romanelli (2020) Synthesis and carbonic anhydrase activating properties of a series of 2-amino-imidazolines structurally related to clonidine¹, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 35:1, 1003-1010, DOI: [10.1080/14756366.2020.1749602](https://doi.org/10.1080/14756366.2020.1749602)

To link to this article: <https://doi.org/10.1080/14756366.2020.1749602>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 27 Apr 2020.



[Submit your article to this journal](#)



Article views: 192



[View related articles](#)



[View Crossmark data](#)

RESEARCH PAPER



Synthesis and carbonic anhydrase activating properties of a series of 2-amino-imidazolines structurally related to clonidine¹

Niccolò Chiaramonte, Soumia Maach, Caterina Biliotti, Andrea Angeli , Gianluca Bartolucci , Laura Braconi, Silvia Dei , Elisabetta Teodori , Claudiu T. Supuran and Maria Novella Romanelli

Section of Pharmaceutical and Nutraceutical Sciences, Department of Neuroscience, Psychology, Drug Research and Child's Health, University of Florence, Sesto Fiorentino, Italy

ABSTRACT

The Carbonic Anhydrase (CA, EC 4.2.1.1) activating properties of histamine have been known for a long time. This compound has been extensively modified but only in few instances the imidazole ring has been replaced with other heterocycles. It was envisaged that the imidazoline ring could be a bioisoster of the imidazole moiety. Indeed, we report that clonidine, a 2-aminoimidazoline derivative, was found able to activate several human CA isoforms (hCA I, IV, VA, VII, IX, XII and XIII), with potency in the micromolar range, while it was inactive on hCA II. A series of 2-aminoimidazoline, structurally related to clonidine, was then synthesised and tested on selected hCA isoforms. The compounds were inactive on hCA II while displayed activating properties on hCA I, VA, VII and XIII, with K_A values in the micromolar range. Two compounds (**10** and **11**) showed some preference for the hCA VA or VII isoforms.

ARTICLE HISTORY

Received 14 March 2020
Revised 26 March 2020
Accepted 26 March 2020

KEYWORDS

Carbonic anhydrase; activator; clonidine; histamine; imidazoline

1. Introduction

Carbonic anhydrases (CAs, EC 4.2.1.1) are metallo-enzymes widespread in all life kingdoms. These enzymes catalyse a plethora of reactions, among which the reversible hydration of CO_2 is the most important one¹. The active site contains the cofactor, a metal ion (usually Zn^{2+}) which coordinates a water molecule responsible, once activated as hydroxide ion, of the nucleophilic attack onto carbon dioxide. Eight genetically different families have been found (α -); 15 isoforms belonging to the α class have been characterised in humans^{1,2}.

CAs have been drug targets since more than 70 years; inhibitors of these enzyme are used for the treatment of oedema, glaucoma and epilepsy but several new therapeutic applications are under study³. In recent years, the attention has been focussed also on activators of these enzymes, despite the fact that CA are among the most efficient enzymes known. In fact, genetic deficiencies of several CA isoforms were reported in the last decades (reviewed in Refs. [3,4]), and in principle a loss of function of these enzymes could be treated with CA selective activators (CAAs). In addition, there is evidence that CA activation improves cognitive performance⁵⁻¹⁰. However, the influence of CA on these processes is complex since also inhibitors have been found to improve memory deficits in animal models (reviewed in Ref. [11]); these findings point out the need for isoform selective inhibitors or activators to elucidate the role of CA isoforms in cognitive processes. Other possible applications of CAAs could be in the formation of artificial tissues¹² and in CO_2 capture and sequestration processes¹³.

Histamine (HST, Chart 1) was among the first reported activators, whose interaction mode was elucidated by means of X-ray crystallography¹⁴. The adduct with hCA II revealed a complex network of H-bonds involving the Zn-bound water molecule, His64 and the imidazole ring of the activator, which is located far away from the metal ion, in a region approaching the edge of the active site cavity. X-ray crystallographic studies have later shown that also other activators bind in this area⁴.

As common structural feature, CAAs possess flexible tails decorated with protonable moieties, with $\text{p}K_a$ values spanning between 6 and 8. The molecule of histamine has been extensively modified, placing substituents on the imidazole C atoms and on the NH_2 group, showing that the latter is not essential, since it can be largely modified to keep or improve potency (reviewed in Ref. [4]). Only in few instances the imidazole ring has been replaced by another heterocycle, such as a thiaziazole ring¹⁵.




Chart 1. Chemical structure of CA activators.

In search for bioisosters of the imidazole moiety, our attention was attracted by the imidazoline ring. This feature is present in a well-known drug, Clonidine (CLO, Chart 1), which is clinically used

CONTACT Claudiu T. Supuran claudiu.supuran@unifi.it; Maria Novella Romanelli novella.romanelli@unifi.it Section of Pharmaceutical and Nutraceutical Sciences, Department of Neuroscience, Psychology, Drug Research and Child's Health, University of Florence, Via Ugo Schiff 6, Sesto Fiorentino, 50019, Italy

¹Dedicated to the memory of Professor Fulvio Gualtieri, our incomparable master who recently passed away.

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

as an antihypertensive agent being an agonist at the central α_2 -adrenergic receptor, but it is able to interact with other targets, such as the imidazoline binding sites and the hyperpolarization-activated cyclic nucleotide gated channels^{16,17}. Therefore, we decided to measure the potential CA activating properties of this compound, finding that CLO behaves as CAA on several CA isoforms (Table 1). Encouraged by this positive outcome, we synthesised a series of 2-substituted imidazolines (compounds **1–24**, Chart 1) and tested their activity on five different hCA isoforms. The ubiquitous cytosolic enzymes CA I and II, the mitochondrial CA VA, which is associated with the glucose homeostasis,¹⁸ the cytosolic CA VII which is particularly abundant in the CNS and has been recently demonstrated to have a protective role against oxidative damage,¹⁹ and the cytosolic CA XIII, which is particularly expressed in the reproductive organs^{20,21} were selected.

2. Materials and methods

2.1. Chemistry

All melting points were taken on a Büchi apparatus and are uncorrected. NMR spectra were recorded on a Bruker Avance 400 spectrometer (400 MHz for ¹H NMR, 100 MHz for ¹³C). Chromatographic separations were performed on a silica gel column by gravity chromatography (Kieselgel 40, 0.063–0.200 mm; Merck) or flash chromatography (Kieselgel 40, 0.040–0.063 mm; Merck). Yields are given after purification, unless differently stated. When reactions were performed under anhydrous conditions, the mixtures were maintained under nitrogen. High-resolution mass spectrometry (HR-MS) analyses were performed with a Thermo Finnigan LTQ Orbitrap mass spectrometer equipped with an electrospray ionisation source (ESI). Analyses were carried out in positive ion mode monitoring protonated molecules, [M + H]⁺ species, and a proper dwell time acquisition was used to achieve 60,000 units of resolution at Full Width at Half Maximum (FWHM). Elemental composition of compounds were calculated on the basis of their measured accurate masses, accepting only results with an attribution error less than 5 ppm and a not integer RDB (double bond/ring equivalents) value, in order to consider only the protonated species²². Compounds were named following IUPAC rules by means of ChemDraw 14.0.

2.1.1. General procedure for the preparation of 2-amino-imidazoline 2–20

A mixture of the appropriate intermediate (**1a**²³ or **1b**²⁴, 0.05 g) and the amine (1 eq) was suspended in THF (5 ml) and the mixture was heated at 70 °C until reaction completion (TLC monitoring, eluent CH₂Cl₂/CH₃OH/NH₄OH 90:10:1); alternatively, an excess of amine (5 eq) was used as solvent. Volatiles were evaporated under vacuum and the residue was triturated with Et₂O to remove the unreacted amine, affording the desired imidazoline derivative. In some instances, additional purification by means of flash chromatography was necessary. The conditions for each compound are reported in Table 2. With this method, the following compounds were prepared.

N-benzyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **2**²³. White solid; m.p. 143–146 °C ESI-LC-MS (*m/z*) 176.0 [M + H]⁺ [¹H]-NMR (D₂O) δ : 3.56 (s, 4H, CH₂CH₂); 4.31 (s, 2H, CH₂); 7.24–7.35 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ : 42.70 (CH₂Ph); 45.65 (2CH₂); 126.93 (CH_{Ar}); 128.00 (CH_{Ar}); 128.95 (CH_{Ar}); 136.25 (C_{Ar}); 159.91 (C=N) ppm.

N-Phenethyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **3**²⁵. White solid; m.p. 82–83 °C; ESI-LCMS (*m/z*) 189.7 [M + H]⁺.

Table 1. Activation constants of Clonidine (CLO) and histamine (HST) on selected human CA isoforms, measured by means of a stopped-flow, CO₂ hydratase assay.^a

Compound	hCA K _A (μM)							
	I	II	IV	VA	VII	IX	XII	XIII
CLO	73.6	>200	132	42.6	8.4	54.1	126	7.8
HST ^b	2.1	125	25.3	0.01	37.5	35.1	27.9	4.6

^aErrors (data not shown) are in the range of ± 5 –10% of the reported values from three different assays.

^bFrom Ref. [4].

Table 2. Synthetic details for the preparation of compounds 2–20 (see general formula in Scheme 1) starting from **1a** (R₁ = H) or **1b** (R₁ = Me) and amines R₂R₃NH.

N	Starting material	R ₂	R ₃	Solvent	Reaction time (h) ^a	Yields (%) ^b
2	1a	H	–CH ₂ Ph	THF	7	94
3	1a	H	–(CH ₂) ₂ Ph	THF	16	90
(<i>S</i>)- 4	1a	H	(<i>S</i>)–CH(Me)Ph	Amine ^c	2	36 ^d
(<i>R</i>)- 4	1a	H	(<i>R</i>)–CH(Me)Ph	Amine ^c	2	74 ^d
5	1a	H	–(CH ₂) ₃ Ph	THF	3	88
6	1a	Me	–CH ₂ Ph	THF	22	32 ^d
7	1a	Me	–(CH ₂) ₂ Ph	THF	2	88
8	1b	H	–(CH ₂) ₂ Ph	Amine ^c	2	86
9	1b	Me	–(CH ₂) ₂ Ph	Amine ^c	2	97
10	1b	Me	–CH ₂ Ph	Amine ^c	3	70 ^d
11	1b	H	–CH ₂ Ph	Amine ^c	2	81
12	1b	H	–(CH ₂) ₃ Ph	Amine ^c	2	45 ^d
13	1b	H	–CH ₂ C ₆ H ₄ Cl(4)	THF	16	97
14	1b	H	–CH ₂ C ₆ H ₄ OCH ₃ (4)	THF	4	89 ^d
15	1b	H	–CH ₂ C ₆ H ₄ F(4)	Amine ^c	6	89
16	1b	H	–CH ₂ C ₆ H ₄ Cl(3)	Amine ^c	6	88 ^d
17	1b	H	–CH ₂ C ₆ H ₄ OCH ₃ (3)	Amine ^c	24	92
18	1b	H	–CH ₂ C ₆ H ₄ F(3)	Amine ^c	16	87
19	1a	H	–(CH ₂) ₂ NHCH ₂ Ph	Amine ^c	3	56 ^d
20	1b	H	–(CH ₂) ₂ NHCH ₂ Ph	Amine ^c	4	57 ^d

^aHeating at 70 °C; all amines were commercially available.

^bUnless otherwise stated, yields are given after trituration with diethyl ether.

^cThe amine was used in excess (5 eq).

^dAfter chromatographic separation.

[¹H]-NMR (D₂O) δ : 2.75 (t, *J* = 6.6 Hz, 2H, CH₂); 3.33 (t, *J* = 6.6 Hz, 2H, CH₂); 3.43 (s, 4H, CH₂CH₂); 7.14–7.31 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ : 34.76 (CH₂); 42.63 (2CH₂); 43.78 (CH₂); 126.94 (CH_{Ar}); 128.84 (CH_{Ar}); 129.13 (CH_{Ar}); 138.50 (C_{Ar}); 159.83 (C=N) ppm.

(*S*) *N*-(1-Phenylethyl)-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide (*S*)-**4**²⁶. Purification by flash chromatography (CH₂Cl₂/CH₃OH/NH₃ 87:13:1.3 as eluent); m.p. 103–107 °C; ESI-LCMS (*m/z*) 190.2 [M + H]⁺. [¹H]-NMR (D₂O) δ : 1.41 (d, *J* = 6.8 Hz, 3H, CH₃); 3.49–3.53 (m, 4H, 2CH₂); 4.55 (q, *J* = 6.8 Hz, 1H, CH); 7.23–7.35 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ : 22.52 (CH₃); 42.62 (2CH₂); 52.79 (CH); 125.59 (CH_{Ar}); 128.03 (CH_{Ar}); 129.08 (CH_{Ar}); 142.03 (C_{Ar}); 158.98 (C=N) ppm.

(*R*) *N*-(1-Phenylethyl)-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide (*R*)-**4**²⁶. Purification by flash chromatography (CH₂Cl₂/CH₃OH/NH₃ 87:13:1.3 as eluent); m.p. 105–109 °C; ESI-LCMS (*m/z*) 190.2 [M + H]⁺. [¹H]-NMR (D₂O) δ : 1.40 (d, *J* = 6.8 Hz, 3H, CH₃); 3.47–3.52 (m, 4H, 2CH₂); 4.54 (q, *J* = 6.8 Hz, 1H, CH); 7.21–7.40 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ : 22.54 (CH₃); 42.67 (2CH₂); 52.84 (CH); 125.63 (CH_{Ar}); 128.07 (CH_{Ar}); 129.13 (CH_{Ar}); 142.06 (C_{Ar}); 159.03 (C=N) ppm.

N-(3-Phenylpropyl)-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **5**. White solid; m.p. 85–87 °C. [¹H]-NMR (CDCl₃) δ : 1.90 (p, *J* = 7.2 Hz, 2H, CH₂); 2.70 (t, 2H, *J* = 7.9 Hz, PhCH₂); 3.34 (apparent q, *J* = 6.4 Hz, 2H, NCH₂); 3.60 (s, 4H, CH₂CH₂); 7.07 (bs, 1H, NH), 7.12–7.26 (m, 5H, Ar); 7.63 (s, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ :

30.68 (CH₂); 32.63 (CH₂); 42.57 (CH₂); 43.39 (CH₂); 126.22 (CH_{Ar}); 128.54 (CH_{Ar}); 140.62 (C_{Ar}); 159.67 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃⁺ 204.1495; found 204.1498.

N-Benzyl-*N*-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **6**. Purification by flash chromatography (CH₂Cl₂/CH₃OH/NH₃ 80:20:1 as eluent). White solid; m.p. 150–153 °C. [¹H]-NMR (D₂O) δ: 2.89 (s, 3H, NCH₃); 3.61 (s, 4H, CH₂CH₂); 4.42 (s, 2H, CH₂Ph); 7.15–7.40 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ: 36.38 (CH₃); 43.14 (2CH₂); 54.29 (CH₂Ph); 127.05 (CH_{Ar}); 128.25 (CH_{Ar}); 129.11 (CH_{Ar}); 134.71 (C_{Ar}); 160.548 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₁H₁₆N₃⁺ 190.1339; found 190.1342.

N-Methyl-*N*-phenethyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **7**. White solid; m.p. 200–204 °C. [¹H]-NMR (D₂O) δ: 2.80–2.89 (m, 5H, NCH₃ + PhCH₂); 3.42 (s, 4H, 2CH₂); 3.46 (t, *J* = 6.6 Hz, 2H, NCH₂); 7.19–7.33 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ: 32.62 (CH₂Ph); 36.19 (NCH₃); 42.83 (N-CH₂); 52.53 (CH₂CH₂); 126.98 (CH_{Ar}); 128.82 (CH_{Ar}); 129.05 (CH_{Ar}); 138.13 (C_{Ar}); 159.89 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃⁺ 204.1495; found 204.1499.

1-Methyl-*N*-phenethyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **8**. White solid; m.p. 118–220 °C. [¹H]-NMR (CDCl₃) δ: 3.00–3.08 (m, 5H, PhCH₂ + NCH₃); 3.48 (s, 4H, 2CH₂); 3.65 (apparent q, *J* = 6.8 Hz, 2H, NCH₂); 6.96 (s, 1H, NH), 7.12–7.35 (m, 5H, Ar), 7.49 (t, *J* = 5.6, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 33.78 (CH₃); 35.49 (CH₂); 41.04 (CH₂); 45.22 (CH₂); 50.11 (CH₂); 126.79 (CH_{Ar}); 128.67 (CH_{Ar}); 129.32 (CH_{Ar}); 137.87 (C_{Ar}); 158.22 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃⁺ 204.1495; found 204.1494.

N,1-Dimethyl-*N*-phenethyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **9**. White solid; m.p. 109–110 °C. [¹H]-NMR (CDCl₃) δ: 2.82 (s, 3H, CH₃); 2.97 (t, *J* = 6.8 Hz, 2H, PhCH₂); 3.13 (s, 3H, CH₃); 3.55–3.62 (m, 4H, CH₂CH₂); 3.77 (t, *J* = 6.9 Hz, 2H, CH₂N); 7.32 (t, *J* = 7.5 Hz, 5H, Ar), 8.34 (bs, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 33.66 (CH₂); 36.97 (CH₃); 39.70 (CH₃); 40.33 (CH₂); 52.89 (CH₂); 54.71 (CH₂); 127.23 (CH-Ar); 128.92 (CH-Ar); 136.96 (C-Ar); 162.47 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₃H₂₀N₃⁺ 218.1652; found 218.1649.

N-Benzyl-*N*,1-dimethyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **10**. Purification by flash chromatography (CH₂Cl₂/CH₃OH/NH₃ 90:10:1 as eluent). Oil. [¹H]-NMR (CDCl₃) δ: 3.01 (s, 3H, CH₃); 3.07 (s, 3H, CH₃); 3.70–3.87 (m, 4H, CH₂CH₂); 4.63 (s, 2H, CH₂Ph); 7.22–7.35 (m, 5H, Ar), 8.10 (s, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 37.03 (CH₃); 39.72 (CH₃); 40.53 (CH₂); 53.14 (CH₂); 56.60 (CH₂); 127.20 (CH_{Ar}); 128.44 (CH_{Ar}); 129.25 (CH_{Ar}); 133.94 (C_{Ar}); 162.63 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃⁺ 204.1495; found 204.1497.

N-Benzyl-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **11**. White solid; m.p. 130–133 °C. [¹H]-NMR (CDCl₃) δ: 3.16 (s, 3H, CH₃); 3.58–3.66 (m, 4H, CH₂CH₂); 4.67 (d, *J* = 5.6 Hz, 2H, CH₂Ph); 6.87 (s, 1H, NH); 7.25–7.38 (m, 3H, Ar); 7.49 (d, *J* = 7.2 Hz, 2H, Ar); 8.21 (s, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 33.76 (CH₃); 41.31 (CH₂); 46.38 (CH₂); 50.22 (CH₂); 128.13 (CH_{Ar}); 128.34 (CH_{Ar}); 129.04 (CH_{Ar}); 135.63 (C_{Ar}); 158.35 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₁H₁₆N₃⁺ 190.1339; found 190.1338.

1-Methyl-*N*-(3-phenylpropyl)-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **12**. Purification by flash chromatography (CH₂Cl₂/CH₃OH/NH₃ 90:10:1 as eluent). Gum. [¹H]-NMR (CDCl₃) δ: 2.02 (p, *J* = 7.6 Hz, 2H); 2.69 (t, *J* = 8.0 Hz, 2H, CH₂); 2.97 (s, 3H, CH₃); 3.41–3.54 (m, 4H, CH₂CH₂); 3.60 (t, *J* = 8.0 Hz, 2H, CH₂); 7.10–7.25 (m, 5H, Ar) ppm. [¹³C]-NMR (CDCl₃) δ: 30.62 (CH₂), 32.93 (CH₂), 33.77 (NCH₃), 40.97 (CH₂), 43.83 (CH₂), 50.11 (CH₂), 126.01 (CH_{Ar}), 128.44 (CH_{Ar}), 128.58 (CH_{Ar}), 141.25 (C_{Ar}), 157.91 (C=N) ppm. ESI-

HRMS (*m/z*) [M + H]⁺: calculated for C₁₃H₂₀N₃⁺ 218.1652; found 218.1654.

N-(4-Chlorobenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **13**. White solid, m.p. 84 °C. [¹H]-NMR (MeOD) δ: 2.96 (s, 3H, CH₃), 3.61–3.76 (m, 4H, CH₂CH₂), 4.41 (s, 2H, CH₂), 7.31 (d, *J* = 8.4 Hz, 2H, Ar), 7.35 (d, *J* = 8.4 Hz, 2H, Ar), 7.42 (s, 2H, NH) ppm. [¹³C]-NMR (MeOD) δ: 30.68 (CH₃), 40.95 (CH₂), 45.41 (CH₂), 50.09 (CH₂), 128.92 (CH_{Ar}), 130.37 (CH_{Ar}), 133.46 (C-Cl), 135.00 (C_{Ar}), 158.68 (C=N), ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₁H₁₅ClN₃⁺ 224.0949; found 224.0946.

N-(4-Methoxybenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **14**. White solid, m.p. 167 °C. [¹H]-NMR (CDCl₃) δ: 3.18 (s, 3H, CH₃), 3.64 (s, 4H, CH₂CH₂), 3.76 (s, 3H, OCH₃), 4.58 (d, *J* = 5.7 Hz, 2H, CH₂), 6.38 (s, 1H, NH), 6.85 (d, *J* = 8.6 Hz, 2H, Ar), 7.39 (d, *J* = 8.6 Hz, 2H, Ar), 8.10 (s, 1H, NH), ppm. [¹³C]-NMR (CDCl₃) δ: 33.77 (NCH₃), 41.29 (CH₂), 45.86 (CH₂), 50.14 (CH₂), 55.36 (OCH₃), 114.38 (CH_{Ar}), 127.62 (C_{Ar}), 129.69 (CH_{Ar}), 158.19 (C_{Ar}), 159.59 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃O⁺ 220.1444; found 220.1443.

N-(4-Fluorobenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **15**. White solid, m.p. 157 °C. [¹H]-NMR (MeOD) δ: 2.97 (s, 3H, CH₃), 3.61–3.75 (m, 4H, CH₂CH₂), 4.42 (s, 2H, CH₂), 7.08 (t, *J* = 8.5 Hz, 2H, Ar), 7.42 (dd, *J* = 8.5, 5.4 Hz, 2H, Ar), ppm. [¹³C]-NMR (MeOD) δ: 30.81 (CH₃), 40.98 (CH₂), 45.44 (CH₂), 50.11 (CH₂), 115.26 (d, *J*_{C-F} = 22 Hz, CH_{Ar}), 129.12 (d, *J*_{C-F} = 8 Hz, CH_{Ar}), 132.25 (C_{Ar}), 158.62 (C=N), 162.48 (d, *J*_{C-F} = 244 Hz, CF), ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₁H₁₅FN₃⁺ 208.1245; found 208.1245.

N-(3-Chlorobenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **16**. White solid, m.p. 190 °C. [¹H]-NMR (MeOD) δ: 2.96 (s, 3H, CH₃), 3.58–3.71 (m, 4H), 4.41 (s, 2H, CH₂), 7.20–7.25 (m, 1H, Ar), 7.25–7.37 (m, 3H, Ar) ppm. [¹³C]-NMR (MeOD) δ: 30.56 (CH₃), 40.93 (CH₂), 45.45 (CH₂), 50.06 (CH₂), 125.25 (CH_{Ar}), 126.91 (CH_{Ar}), 127.74 (CH_{Ar}), 130.07 (CH_{Ar}), 134.35 (C-Cl), 138.59 (C_{Ar}), 154.10 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₁H₁₅ClN₃⁺ 224.0949; found 224.0949.

N-(3-Methoxybenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **17**. White solid, m.p. 147 °C. [¹H]-NMR (CDCl₃) δ: 2.95 (s, 3H, CH₃), 3.61 (s, 4H, CH₂CH₂), 3.76 (s, 3H, OCH₃), 4.59 (s, 2H, CH₂), 6.77 (d, *J* = 6.4 Hz, 1H, NH), 7.06–7.66 (m, 4H, Ar), 8.31 (s, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 33.70 (CH₃), 41.32 (CH₂), 46.20 (CH₂), 50.22 (CH₂), 55.68 (OCH₃), 113.67 (CH_{Ar}), 114.00 (CH_{Ar}), 120.27 (CH_{Ar}), 129.96 (CH_{Ar}), 137.43 (C_{Ar}), 158.22 (C_{Ar}), 159.94 (C=N), ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₈N₃O⁺ 220.1444; found 220.1447.

N-(3-Fluorobenzyl)-1-methyl-4,5-dihydro-1*H*-imidazol-2-amine hydroiodide **18**. White solid, m.p. 195 °C. [¹H]-NMR (CDCl₃) δ: 3.19 (s, 3H, CH₃), 3.61–3.71 (m, 4H, CH₂CH₂), 4.68 (s, 2H, CH₂), 6.95 (t, *J* = 7.7 Hz, 1H, NH), 7.09–7.41 (m, 4H, Ar), 8.27 (bs, 1H, NH) ppm. [¹³C]-NMR (CDCl₃) δ: 33.86 (CH₃), 41.19 (CH₂), 45.42 (CH₂), 50.20 (CH₂), 115.05 (d, *J*_{CF} = 25 Hz, CH_{Ar}), 123.94 (CH_{Ar}), 130.45 (d, *J* = 8 Hz, CH_{Ar}), 138.48 (C_{Ar}), 158.11 (C=N), 162.78 (d, *J* = 246 Hz, CF) ppm.

*N*¹-Benzyl-*N*²-(4,5-dihydro-1*H*-imidazol-2-yl)ethane-1,2-diamine hydroiodide **19**. White solid, m.p. 230 °C (dec). [¹H]-NMR (MeOD) δ: 3.37 (t, *J* = 6.2 Hz, 2H, CH₂), 3.64 (t, *J* = 6.2 Hz, 2H, CH₂), 3.70 (s, 4H, CH₂CH₂), 4.34 (s, 2H, PhCH₂), 7.40–7.47 (m, 3H, Ar), 7.59–7.64 (m, 2H, Ar) ppm. [¹³C]-NMR (MeOD) δ: 38.87 (CH₃), 42.82 (CH₂CH₂), 45.73 (CH₂), 51.20 (PhCH₂), 128.92 (CH_{Ar}), 129.49 (CH_{Ar}), 129.99 (CH_{Ar}), 130.69 (C_{Ar}), 159.88 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₂H₁₉N₄⁺ 219.1604; found 219.1606.

*N*¹-Benzyl-*N*²-(1-methyl-4,5-dihydro-1*H*-imidazol-2-yl)ethane-1,2-diamine hydroiodide **20**. White solid, m.p. 235 °C (dec). [¹H]-NMR (MeOD) δ: 2.93 (s, 3H, CH₃), 3.33 (t, *J* = 6.0 Hz, 2H, CH₂), 3.63–3.72

(m, 6H, 3CH₂), 4.28 (s, 2H, PhCH₂), 7.38–7.44 (m, 3H, Ar), 7.48 – 7.54 (m, 2H, Ar) ppm. [¹³C]-NMR (MeOD) δ: 30.59 (CH₃), 39.07 (CH₃), 40.94 (CH₂), 45.61 (CH₂), 50.04 (PhCH₂), 51.17 (CH₂), 128.56 (CH_{Ar}), 129.37 (CH_{Ar}), 129.80 (CH_{Ar}), 130.81 (C_{Ar}), 156.18 (C=N) ppm. ESI-HRMS (*m/z*) [M + H]⁺: calculated for C₁₃H₂₁N₄⁺ 233.1761; found 233.1760.

2.1.2. *N*-(4,5-Dihydro-1*H*-imidazol-2-yl)ethane-1,2-diamine **21**²⁷

The compound was prepared following the procedure reported by A. Bucio-Cano et al.²⁷ Yellow oil, isolated then as oxalate salt. Yield: 61%; m.p. (oxalate salt): 153–154 °C. [¹H] NMR (D₂O) δ: 3.10 (t, *J* = 6 Hz, 2H, CH₂); 3.44 (t, *J* = 6 Hz, 2H, CH₂); 3.59 (s, 4H, NHCH₂CH₂NH₂) ppm. [¹³C] NMR (D₂O) δ: 38.28 (CH₂); 39.70 (CH₂); 42.82 (2CH₂); 159.80 (C); 165.0 (CO Oxalate) ppm.

2.1.3. *N*-Phenyl-4,5-dihydro-1*H*-imidazol-2-amine **22**²⁸

1-(2-Aminoethyl)-3-phenylthiourea²⁹ (110 mg, 0.56 mmol) was dissolved in THF (5 ml) and a 1 M sodium hydroxide/water solution (1 eq) was added while stirring. After 0.5 h, a solution of TsCl (1–1.1 equiv) in THF (5 ml) was slowly added and the mixture was stirred for 5 h at room T. The solvent was evaporated and diethyl ether and brine were added to the residue. The layers were separated and the aqueous layer was extracted three times with diethyl ether. The combined organic layers were dried (Na₂SO₄), filtered and evaporated, leaving a residue which was purified by flash chromatography (CH₂Cl₂: MeOH: NH₃ 80:20:1 as eluent). Compound **22** was obtained as a white solid, m.p. 137–140 °C, in 50% yields. [¹H]-NMR (D₂O) δ: 3.40 (s, 4H, CH₂CH₂), 6.94 (d, *J* = 7.2 Hz, 2H), 7.03 (t, *J* = 7.2 Hz, 1H), 7.27 (t, *J* = 8.0 Hz, 2H) ppm. [¹³C]-NMR (D₂O) δ: 42.37 (CH₂), 123.34 (CH_{Ar}), 123.40 (CH_{Ar}), 129.52 (CH_{Ar}), 146.90 (C_{Ar}), 161.03 (C=N) ppm.

2.1.4. 1-(2-(Methylthio)-4,5-dihydro-1*H*-imidazol-1-yl) ethanone **23**³⁰

The imidazoline **1a** was acylated on one imidazoline-nitrogen atom to obtain **23** as reported by Gomez-San Juan et al.³⁰ Yellow solid, yield: 100%; m.p. = 107–110 °C. [¹H]-NMR (D₂O) δ: 2.07 (s, 3H, SCH₃); 2.23 (s, 3H, CH₃); 3.76 (t, *J* = 8.8 Hz, 2H, CH₂); 3.92 (t, *J* = 8.8 Hz, 2H, CH₂) ppm. [¹³C]-NMR (D₂O) δ: 14.5 (SCH₃); 23.5 (CH₃); 48.0 (2CH₂); 161.5 (C=N); 171.9 (CO) ppm.

2.1.5. 2-(Benzylthio)-4,5-dihydro-1*H*-imidazole **24** hydrobromide³¹

A mixture of commercially-available 2-imidazolidinethione (0.1 g, 0.98 mmol) and benzyl bromide (1.1 eq) in anhydrous methanol (5 ml) was heated at 65 °C for 2 h. After cooling, the solvent was removed under vacuum and the residue was treated with Et₂O until it became solid; filtration and drying under vacuum gave the desired compound as a white solid, m.p. 181–183 °C. ESI-LC-MS (*m/z*) 193.1 [M + H]⁺ [¹H]-NMR (D₂O) δ: 3.81 (s, 4H, CH₂CH₂); 4.35 (s, 2H, CH₂); 7.31–7.42 (m, 5H, Ar) ppm. [¹³C]-NMR (D₂O) δ: 35.15 (CH₂Ph); 45.24 (2CH₂); 128.31 (CH_{Ar}); 128.72 (CH_{Ar}); 129.10 (CH_{Ar}); 134.12 (C_{Ar}); 169.69 (C=N) ppm.

2.2. CA activation

An Sx.18Mv-R Applied Photophysics (Oxford, UK) stopped-flow instrument has been used to assay the catalytic activity of various CA isozymes for CO₂ hydration reaction³². Phenol red (at a concentration of 0.2 mM) was used as indicator, working at the

absorbance maximum of 557 nm, with 10 mM Hepes (pH 7.5, for α-CAs)^{33–37} as buffers, 0.1 M NaClO₄ (for maintaining constant ionic strength), following the CA-catalysed CO₂ hydration reaction for a period of 10 s at 25 °C. The CO₂ concentrations ranged from 1.7 to 17 mM for the determination of the kinetic parameters and inhibition constants. For each activator at least six traces of the initial 5–10% of the reaction have been used for determining the initial velocity. The uncatalysed rates were determined in the same manner and subtracted from the total observed rates. Stock solutions of activators (at 0.1 mM) were prepared in distilled-deionised water and dilutions up to 1 nM were made thereafter with the assay buffer. Enzyme and activator solutions were pre-incubated together for 15 min prior to assay, in order to allow for the formation of the enzyme-activator complexes. The activation constant (K_A), defined similarly with the inhibition constant K_i, can be obtained by considering the classical Michaelis–Menten equation (Equation (1)), which has been fitted by non-linear least squares by using PRISM 3:

$$v = v_{\max} / \{1 + (K_M/[S])(1 + [A]_f/K_A)\} \quad (1)$$

where [A]_f is the free concentration of activator.

Working at substrate concentrations considerably lower than K_M ([S] ≪ K_M), and considering that [A]_f can be represented in the form of the total concentration of the enzyme ([E]_t) and activator ([A]_t), the obtained competitive steady-state equation for determining the activation constant is given by Equation (2):

$$v = v_0 \cdot K_A / \{K_A + ([A]_t - 0.5 \{([A]_t + [E]_t + K_A) - ([A]_t + [E]_t + K_A)^2 - 4[A]_t \cdot [E]_t\}^{1/2})\} \quad (2)$$


where v₀ represents the initial velocity of the enzyme-catalysed reaction in the absence of activator^{33–39}. This type of approach to measure enzyme–ligand interactions is in excellent agreement with recent results from native mass spectrometry measurements⁴⁰.

3. Results and discussion


3.1. Chemistry

2-Amino imidazolines **2–20** were prepared by reacting primary and secondary amines with 2-(methylthio)-4,5-dihydro-1*H*-imidazole hydroiodide **1a**²³ or its *N*¹-methyl derivative **1b**²⁴ using tetrahydrofuran or an excess of amine as solvents (Scheme 1). The final compounds were obtained as hydroiodide salts in different yields. Synthetic details are reported in the experimental section. Reactions with 3- or 4-nitrobenzyl amine or with dibenzyl amine were unsuccessful.

This method failed also to give compound **22** (step b, Scheme 2), whose synthesis was therefore attempted according to the procedure of Gomez Saint-Juan (step c, Scheme 2);³⁰ however, also



Scheme 1. Synthesis of compounds 2-20.



Scheme 2. Synthesis of compounds 22 and 23.

this method was abandoned since the reaction of 1-acetyl-2-(methylthio)-4,5-dihydro-1*H*-imidazole **23** with aniline was not successful. Finally, **22** was prepared from 1-(2-aminoethyl)-3-phenylthiourea²⁹ according to Heinelt et al.²⁸ Compound **21**²⁷ was synthesised through condensation of guanidine and ethylenediamine while **24**³¹ was prepared by reaction of imidazoline-2-thione with benzyl bromide.

3.2. CA activating profile

The stopped-flow method³² has been used for assaying the CO_2 hydration activity catalysed by different CA isoforms; the results are expressed as K_A (activation constant, μM). The activating profile of CLO is reported in Table 1, in comparison with HST. CLO behaved as an activator on several CA isoforms (I, IV, VA, VII, IX, XII and XIII), with K_A values in the range 7.8–136 μM , while on CA II it was inactive up to a dose of 200 μM . In particular, the isoforms most sensitive to CLO were CA VII and CA XIII.

The K_A values of the synthesised compounds are reported in Table 3. All compounds have been tested as hydroiodide salts, with the exception of **21** (oxalate), **22** and **23** (free bases), and **24** (as HBr salt). As CLO, none of the compounds was active on hCA II at the highest tested concentration, while on the other isoforms the compounds showed K_A values mainly in the low-medium range, allowing to derive the following structure–activity relationships.

3.2.1. hCA I

The K_A value of CLO on this isoform was 76.3 μM . Removal of both chlorine atoms abolished activity, since **22** ($R_2 = \text{Ph}$) was inactive when tested up to a 150 μM concentration. On the contrary, inserting a CH_2 unit between the exocyclic N atom and the Ph ring of **22** improved the activity: in fact, **2** (K_A 4.18 μM), 18 times more potent than CLO, was one of the most potent compounds on this isoform among the newly synthesised analogues. The elongation of the methylene chain gave compounds less active than **2**; interestingly, a chain formed by 3 CH_2 units was tolerated (**5**, K_A 36.7 μM) while a CH_2CH_2 chain was not (**3**, $K_A > 150 \mu\text{M}$). Side-chain branching abolished activity (**S-4**, **R-4**: $K_A > 150 \mu\text{M}$). Methylation of the exocyclic N_x atom was tolerated when R_2 was a phenethyl group (**7**: K_A 68.6 μM) but not when R_2 was a benzyl moiety (**6**: $K_A > 150 \mu\text{M}$). Also the methylation of the endocyclic N_1 atom gave contrasting results, since compounds **8** and **12** were more active than their non-methylated analogues **3** and **5**, but **11** ($R_2 = \text{benzyl}$) showed a fourfold decrease of activity with respect to **2** (**11**: K_A 16.9; **2**: 4.18 μM). Double methylation was productive for the phenethyl analogues (compare **9** with **3**, **7** and **8**); for the benzyl analogue **10** this modification improved the

activity only with respect to **6**. The aminoethyl derivative **21** (K_A 3.87 μM) was the most potent compound on the hCA I isoform; the addition of a benzyl moiety on the primary amine group abolished activity (**19**, $K_A > 100 \mu\text{M}$), and also the N^1 -methyl analogue **20** was inactive.


Aromatic substitution on the benzyl moiety did not improve the potency: in fact, while the 4-Cl derivative **13** was equipotent with **11**, a 4-OMe (**14**) or 4-F substituent (**15**) increased from 2 to 3.5 times the K_A values. The same substituents in the meta position reduced to a higher extent or abolished the activity. As far as the sulphur analogues **1a**, **23** and **24** are concerned, the small methyl group seemed tolerated, not the bulkier benzyl moiety (**24**, $K_A > 150 \mu\text{M}$). The basicity of the amidine group appeared to be not crucial, since the NH and the *N*-acetyl derivatives (**1a** and **23**, respectively) were equipotent.

3.2.2. hCA VA

The K_A value of CLO on this isoform was 42.6 μM . The removal of both chlorine atoms did not affect activity, since **22** ($R_2 = \text{Ph}$, K_A 52.7 μM) was almost equipotent with CLO. Also the insertion of a CH_2 unit between the exocyclic N atom and the Ph ring of **22** did not substantially modify potency (**2**, K_A 45.7 μM). On this isoform, the majority of the compounds showed good activating properties, with potency higher than CLO: the K_A values of **24**, **23**, **3–5**, **7–9**, **11** and **12** were in the range 3.7–17.2 μM .

The most potent compound was **10** (K_A 0.9 μM), a benzyl derivative carrying a methyl group on both N_1 and N_x atoms; this compound was 47 times more active than CLO. The removal of the exocyclic $N_x\text{-Me}$ group decreased 4 times the activity (**11**, K_A 3.7 μM), while the removal of the N_1 -methyl group was more detrimental: as a matter of fact, compounds **6** (K_A 40.5 μM) and **2** (K_A 45.5 μM) were about 40 times less potent than **10** (K_A 0.9 μM). On the contrary, the degree of methylation did not substantially affect the potency of the phenylethyl and phenyl propyl derivatives, since compounds **3**, **5**, **7**, **8** and **12** had K_A values in the range 9.9–17.2 μM . Similarly, aromatic substitution on the benzyl moiety slightly decreased the potency without substantial modulation, the K_A values of compounds **13–18** being 2–4 times higher than **11**. Side-chain branching (compounds **S-4** and **R-4**) improved the activity on this isoform, and a small enantioselectivity was observed: the *R*-enantiomer was twice more potent as the *S*-isomer. A benzyl moiety on the terminal amino group of **21** (K_A 31.2 μM) increased the activity, as analogue **19** and its N^1 -methyl derivative **20** were about 3 times more potent than the parent compound.

As far as the sulphur derivatives are concerned, the replacement of the $N_x\text{H}$ moiety of **2** (K_A 45.5 μM) with S (**24**, K_A 11.1 μM) brought a fourfold improvement in activity. Acetylation of the N^1

Table 3. Activation constant (K_A) of the synthesised compounds and Clonidine (CLO) for human I, II, VA, VII and XIII Carbonic Anhydrase isoforms.

N	R ₁	X	R ₂	K_A (μM) ^b				
				hCA I	hCA II	hCA VA	hCA VII	hCA XIII
1a	H	S	-CH ₃	9.61	>150	38.3	41.9	>100
2	H	NH	-CH ₂ Ph	4.18	>150	45.7	35.2	>100
3	H	NH	-(CH ₂) ₂ Ph	>150	>150	16.7	18.9	>100
S-4	H	NH	(S)-CH(Me)Ph	>150	>150	12.4	31.5	>100
R-4	H	NH	(R)-CH(Me)Ph	>150	>150	4.92	24.2	>100
5	H	NH	-(CH ₂) ₃ Ph	36.7	>100	9.9	11.4	24.3
6	H	NMe	-CH ₂ Ph	>150	>150	40.5	11.0	>100
7	H	NMe	-(CH ₂) ₂ Ph	68.6	>100	14.9	2.4	6.5
8	CH ₃	NH	-(CH ₂) ₂ Ph	95.4	>100	14.6	16.2	31.0
9	CH ₃	NMe	-(CH ₂) ₂ Ph	20.2	>100	14.9	2.6	36.9
10	CH ₃	NMe	-CH ₂ Ph	30.2	>100	0.9	6.5	17.4
11	CH ₃	NH	-CH ₂ Ph	16.9	>100	3.7	0.9	19.1
12	CH ₃	NH	-(CH ₂) ₃ Ph	10.9	>100	17.2	3.1	10.9
13	CH ₃	NH	-CH ₂ C ₆ H ₄ Cl(4)	17.7	>100	10.5	22.7	22.9
14	CH ₃	NH	-CH ₂ C ₆ H ₄ OCH ₃ (4)	37.6	>100	15.3	30.2	19.3
15	CH ₃	NH	-CH ₂ C ₆ H ₄ F(4)	59.7	>100	9.3	17.4	24.4
16	CH ₃	NH	-CH ₂ C ₆ H ₄ Cl(3)	99.2	>100	10.4	24.0	37.0
17	CH ₃	NH	-CH ₂ C ₆ H ₄ OCH ₃ (3)	>100	>100	13.1	29.4	14.7
18	CH ₃	NH	-CH ₂ C ₆ H ₄ F(3)	>100	>100	14.9	19.4	23.3
19	H	NH	-(CH ₂) ₂ NHCH ₂ Ph	>100	>100	11.0	41.7	20.1
20	CH ₃	NH	-(CH ₂) ₂ NHCH ₂ Ph	>100	>100	11.9	29.0	16.3
21	H	NH	-CH ₂ CH ₂ NH ₂	3.87	>150	31.2	91.6	>100
22	H	NH	-Ph	>150	>150	52.7	32.6	>100
23	COCH ₃	S	-CH ₃	12.7	>150	15.0	30.9	>100
24	H	S	-CH ₂ Ph	>150	>150	11.1	46.7	>100
CLO	H	NH	(2,6-dichloro)Ph	76.3	>200	42.6	8.4	7.8

^aAll compounds have been tested as HI salts, with the exception of **21** (oxalate), **22** and **23** (free bases), **24** (HBr), and CLO (HCl).

^bMean from 3 different determinations (errors in the range of 5–10% of the reported values, data not shown).

nitrogen was also favourable, as **23** was twice more potent than **1a**.

3.2.3. hCA VII

The K_A value of CLO on this isoform was 8.4 μM . All the tested compounds showed activation properties on this isoform, with K_A values between 0.9 and 91.6 μM . The least potent compound was the primary amine **21** (K_A 91.6 μM), whose activity was however improved by adding a benzyl group on the terminal NH₂ moiety (**19**, K_A 41.7 μM) and a methyl group on the endocyclic N atom (**20**, K_A 29.0 μM). The removal of chlorine atoms of CLO reduced 4 times the activity (**22**, $R_2 = \text{Ph}$, K_A 32.6 μM) while the separation of the phenyl and N_xH moieties by means of a CH₂ unit did not substantially modified potency (**3**, K_A 35.2 μM). On the contrary, the potency increased by elongating the chain from 1 to 3 CH₂ units (**3**, K_A 35.2 μM ; **5**, K_A 11.4 μM) and by adding a methyl group on the N_xH moiety: with the latter modification the potency of **3** (K_A 18.9 μM) and of **2** (K_A 35.2 μM) were increased 3 (**6**, K_A 11.0 μM) and 8 times (**7**, K_A 2.4 μM), respectively. Side-chain branching did not substantially affect activity, since **S-4** and **R-4** were equipotent with **2**. Methylation on the endocyclic N atom was the most effective modification in this set of molecules: as a matter of fact, with this structural change the K_A value of **2** (K_A 35.2 μM) was reduced 39 times, and **11** (K_A 0.9 μM) resulted in the most potent compound on this isoform. The same modification was also effective on the phenylpropyl derivative **5** (K_A 11.4 μM), whose activity was increased 4 times (**12**, K_A 3.1 μM). The double methylation on the N₁ and N_x atoms gave potent compounds (**10**, K_A 6.5 μM and **9**, K_A 2.6 μM) even if the K_A values are, respectively,

7 and 3 times lower than that of **11**. Aromatic substitution on the benzyl moiety of **11** was detrimental for activity, as compounds **13–18** were 19–33 times less potent than **11**. As far as the sulphur analogues **1a**, **23** and **24** are concerned, their K_A values were in the range 30.9–46.7 μM , not better than the other tested 2-aminoimidazoline derivatives. Attempts to crystallise adducts of **7**, **11** and **12** with hCA VII are ongoing.

3.2.4. hCA XIII

This is the isoform most sensitive to CLO among those studied (K_A 7.8 μM). As it happened on the hCA I isoform, the removal of both chlorine atoms, to give **22**, abolished activity. Several other compounds resulted inactive when tested at concentrations up to 100 μM , i.e. the sulphur derivatives, the polar aminoethyl derivative **21**, and all the compounds having both the N₁ and N_x atoms as secondary amines, with the exception of the lipophilic phenylpropyl derivative **5** (K_A 24.3 μM). The activity of **21** could be restored by adding a benzyl moiety on the primary amino-group (**19**, K_A 20.1 μM) and a methyl group on the endocyclic N atom (**20**, K_A 16.3 μM). Also the methylation of the phenethyl analogue **3** on the N_x atom re-established activity, giving **7** (K_A 6.5 μM) which resulted the most potent compound of the series on this isoform. Methylation on the endocyclic N₁ atom gave compounds **8**, **11** and **12** whose potency ranged from 10.9 to 31.0 μM , the most potent being the derivative carrying a phenylpropylamino side chain (**12**). Methylation on both N₁ and N_x atoms or aromatic substitution on the benzyl moiety did not improve activity.

As far as selectivity is concerned, the two compounds showing submicromolar K_A values displayed also interesting selectivity

profiles: **10** was more active on hCA VA with respect to hCA I (33 times), II (>100 times), VII (7 times), and XIII (19 times), while **11** showed a preference for hCA VII over hCA I (19 times), II (>100 times), VA (4 times), and XII (21 times).

4. Conclusions

We have synthesised a series of 2-aminoimidazolines, structurally related to Clonidine, and tested them on five different hCA isoforms (I, II, VA, VII and XIII). As the lead compound, none of the newly synthesised molecules was active on the ubiquitously expressed CA II; on the contrary, the compounds showed activity in the micromolar range on the other tested CA isoforms. Structure–activity relationships were derived, which were different on the various isoforms, suggesting that it could be possible, in this class of compounds, to find molecules, selective for a particular CA isoform. Indeed, from these preliminary modifications it has been possible to find two compounds, **10** and **11**, with a promising preference towards, respectively, CA VA and VII. Work is underway to improve both potency and selectivity, in order to find new pharmacological tool to activate specific CA isoforms in pathologies characterised by their loss of functionality.

Disclosure statement

The authors report no conflict of interest.

Funding

This work was supported by grants from the University of Florence [Fondo Ricerca Ateneo RICATEN18].

ORCID

Andrea Angeli  <http://orcid.org/0000-0002-1470-7192>
 Gianluca Bartolucci  <http://orcid.org/0000-0002-5631-8769>
 Silvia Dei  <http://orcid.org/0000-0003-0898-7148>
 Elisabetta Teodori  <http://orcid.org/0000-0002-9705-3875>
 Claudiu T. Supuran  <http://orcid.org/0000-0003-4262-0323>
 Maria Novella Romanelli  <http://orcid.org/0000-0002-5685-3403>

References

- Supuran CT, De Simone G, Carbonic anhydrases: an overview. In: Claudiu T, Supuran GDS, eds. Carbonic anhydrases as biocatalysts – from theory to medical and industrial applications. Amsterdam: Elsevier; 2015.
- Jensen EL, Clement R, Kosta A, et al. A new widespread subclass of carbonic anhydrase in marine phytoplankton. *ISME J* 2019;13:2094–106.
- Bozdag M, Altamimi ASA, Vullo D, et al. State of the art on carbonic anhydrase modulators for biomedical purposes. *Curr Med Chem* 2019;26:2558–73.
- Supuran CT. Carbonic anhydrase activators. *Future Med Chem* 2018;10:561–73.
- Sun MK, Alkon DL. Carbonic anhydrase gating of attention: memory therapy and enhancement. *Trends Pharmacol Sci* 2002;23:83–9.
- Canto de Souza L, Provensi G, Vullo D, et al. Carbonic anhydrase activation enhances object recognition memory in mice through phosphorylation of the extracellular signal-regulated kinase in the cortex and the hippocampus. *Neuropharmacology* 2017;118:148–56.
- Temperini C, Scozzafava A, Supuran CT. Carbonic anhydrase activators and the drug design. *Curr Pharm. Des* 2008;14:708–15.
- Akokcak S, Supuran CT. Activation of α -, β -, γ -, δ -, ζ - and η -class of carbonic anhydrases with amines and amino acids: a review. *J Enz Inhib Med Chem* 2019;34:1652–9.
- Supuran CT, Vullo D, Manole G, et al. Designing of novel carbonic anhydrase inhibitors and activators. *Curr Med Chem Cardiovasc Hematol Agents* 2004;2:49–68.
- Akokcak S, Lolak N, Bua S, et al. α -Carbonic anhydrases are strongly activated by spinaceamine derivatives. *Bioorg Med Chem* 2019;27:800–4.
- Provensi G, Carta F, Nocentini A, et al. A new kid on the block? Carbonic anhydrases as possible new targets in Alzheimer's disease. *Int J Mol Sci* 2019;20:4724.
- Wang X, Schröder HC, Schlossmacher U, et al. Modulation of the initial mineralization process of SaOS-2 cells by carbonic anhydrase activators and polyphosphate. *Calcified Tissue Int* 2014;94:495–509.
- Capasso C, Supuran CT, Targeting carbonic anhydrases in biotechnology. In: Capasso C, Supuran CT eds. Targeting carbonic anhydrases. *Future Medicine*; 2014:158–69.
- Briganti F, Mangani S, Orioli P, et al. Carbonic anhydrase activators: X-ray crystallographic and spectroscopic investigations for the interaction of isozymes I and II with histamine. *Biochemistry* 1997;36:10384–92.
- Supuran CT, Barboiu M, Luca C, et al. Carbonic anhydrase activators. Part 14. Syntheses of mono and bis pyridinium salt derivatives of 2-amino-5-(2-aminoethyl)- and 2-amino-5-(3-aminopropyl)-1,3,4-thiadiazole and their interaction with isozyme II. *Eur J Med Chem* 1996;31:597–606.
- Bousquet P, Hudson A, García-Sevilla JA, Li J-X. Imidazoline receptor system: the past, the present, and the future. *Pharmacol Rev* 2020;72:50–79.
- Romanelli MN, Sartiani L, Masi A, et al. HCN channels modulators: the need for selectivity. *Curr Top Med Chem* 2016;16:1764–91.
- Price TO, Sheibani N, Shah GN. Regulation of high glucose-induced apoptosis of brain pericytes by mitochondrial CA VA: a specific target for prevention of diabetic cerebrovascular pathology. *Biochim Biophys Acta* 2017;1863:929–35.
- Di Fiore A, Monti DM, Scaloni A, et al. Protective role of carbonic anhydrases III and VII in cellular defense mechanisms upon redox unbalance. *Oxid Med Cell Longev* 2018;2018:2018306.
- José O, Torres-Rodríguez P, Forero-Quintero LS, et al. Carbonic anhydrases and their functional differences in human and mouse sperm physiology. *Biochem Biophys Res Commun* 2015;468:713–8.
- Karim K, Giribabu N, Muniandy S, Salleh N. Estrogen and progesterone differentially regulate carbonic anhydrase II, III, IX, XII, and XIII in ovariectomized rat uteri. *Syst Biol Reprod Med* 2016;62:57–68.
- Marshall AG, Hendrickson CL. High-resolution mass spectrometers. *Ann Rev Anal Chem* 2008;1:579–99.
- Aoyagi N, Endo T. Synthesis of five- and six-membered cyclic guanidines by guanylation with isothiuronium iodides and amines under mild conditions. *Synth Commun* 2017;47:442–8.
- McKay AF, Kreling ME. Nitration of 1-substituted-2-iminoimidazolines. *J Org Chem* 1957;22:1581–3.

25. Tronche P, Amelot A, Bayard J, Laroussinie C. Synthesis of some N-substituted 2-aminoimidazolines. *Ann Pharm Fr* 1960; 18:726–35.
26. Genc M, Servi S. Microwave-induced synthesis of 2-aminoimidazolines under neat conditions. *Synth Commun* 2009;39: 3263–77.
27. Bucio-Cano A, Reyes-Arellano A, Correa-Basurto J, et al. Targeting quorum sensing by designing azoline derivatives to inhibit the N-hexanoyl homoserine lactone-receptor CviR: synthesis as well as biological and theoretical evaluations. *Bioorg Med Chem* 2015;23:7565–77.
28. Heinelt U, Lang HJ. Process for synthesizing heterocyclic compounds, Patent US2004/0242560 A1.
29. Tilley JW, Levitan P, Kierstead RW, Cohen M. Antihypertensive (2-aminoethyl)thiourea derivatives. 1. *J Med Chem* 1980;23:1387–92.
30. Gómez-SanJuan A, Botija JM, Méndez A, et al. C-N bond forming reactions in the synthesis of substituted 2-aminoimidazole derivatives. *Arkivoc* 2014;44–56.
31. Aspinnall SR, Bianco EJ. A synthesis of 2-alkylamino-4,5-dihydroimidazoles. *J Am Chem Soc* 1951;73:602–3.
32. Khalifah RG. The carbon dioxide hydration activity of carbonic anhydrase. I. Stop-flow kinetic studies on the native human isoenzymes B and C. *J Biol Chem* 1971;246:2561–73.
33. Temperini C, Scozzafava A, Vullo D, Supuran CT. Carbonic anhydrase activators. Activation of isozymes I, II, IV, VA, VII, and XIV with L- and D-histidine and crystallographic analysis of their adducts with isoform II: engineering proton-transfer processes within the active site of an enzyme. *Chem Eur J* 2006;12:7057–66.
34. Akocak S, Lolak N, Vullo D, et al. Synthesis and biological evaluation of histamine Schiff bases as carbonic anhydrase I, II, IV, VII, and IX activators. *J Enz Inhib Med Chem* 2017;32: 1305–12.
35. Akocak S, Lolak N, Bua S, et al. Activation of human α -carbonic anhydrase isoforms I, II, IV and VII with bis-histamine schiff bases and bis-spinaceamine substituted derivatives. *J Enz Inhib Med Chem* 2019;34:1193–8.
36. Angeli A, Del Prete S, Osman SM, et al. Activation studies with amines and amino acids of the β -carbonic anhydrase encoded by the Rv3273 gene from the pathogenic bacterium *Mycobacterium tuberculosis*. *J Enz Inhib Med Chem* 2018;33:364–9.
37. Angeli A, Alasmay FAS, Del Prete S, et al. The first activation study of a δ -carbonic anhydrase: TweCA δ from the diatom *Thalassiosira weissflogii* is effectively activated by amines and amino acids. *J Enz Inhib Med Chem* 2018;33: 680–5.
38. Angeli A, Buonanno M, Donald WA, et al. The zinc – but not cadmium – containing ζ -carbonic from the diatom *Thalassiosira weissflogii* is potently activated by amines and amino acids. *Bioorg Chem* 2018;80:261–5.
39. Vistoli G, Aldini G, Fumagalli L, et al. Activation effects of carnosine- and histidine-containing dipeptides on human carbonic anhydrases: a comprehensive study. *Int J Mol Sci* 2020;21:1761.
40. Nguyen GTH, Tran TN, Podgorski MN, et al. Nanoscale ion emitters in native mass spectrometry for measuring ligand–protein binding affinities. *ACS Central Sci* 2019;5: 308–18.