



ELSEVIER

Contents lists available at ScienceDirect

Data in brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Myosin V fluorescence imaging dataset for single-molecule localization and tracking

Lucia Gardini ^{a, b}, Claudia Arbore ^a,
 Francesco Saverio Pavone ^{a, b, c}, Marco Capitanio ^{a, c, *}

^a LENS - European Laboratory for Non-linear Spectroscopy, Via Nello Carrara 1, 50019 Sesto Fiorentino, Italy

^b National Institute of Optics—National Research Council, Largo Fermi 6, 50125 Florence, Italy

^c Department of Physics and Astronomy, University of Florence, Via Sansone 1, 50019 Sesto Fiorentino, Italy



ARTICLE INFO

Article history:

Received 22 March 2019

Received in revised form 18 April 2019

Accepted 26 April 2019

Available online 23 May 2019

Keywords:

Total internal reflection fluorescence (TIRF) microscopy

Myosin

Single molecule biophysics

ABSTRACT

Myosin-5B is one of three members of the myosin-5 family of actin-based molecular motors fundamental in recycling endosome trafficking and collective actin network dynamics. Through single-molecule motility assays, we recently demonstrated that myosin-5B can proceed in 36-nm steps along actin filaments as single motor. By analyzing trajectories of single myosin-5B along actin filaments we showed that its velocity is dependent on ATP concentration, while its run length is independent on ATP concentration, as a landmark of processivity.

Here, we share image stacks acquired under total internal reflection fluorescence (TIRF) microscopy and representative trajectories of single myosin-5B molecules labelled with Quantum Dots (QD-myosin-5B) moving along actin filaments at different ATP concentrations (0.3–1000 μ M). Localization of QD-myosin-5B was performed with the PROOF software, which is freely available [1]. The data can be valuable for researchers interested in molecular motors motility, both from an experimental and modeling point of view, as well as to researchers developing single particle tracking algorithms. The data is related to the research article “Dissecting myosin-5B mechanosensitivity and calcium regulation at the single molecule level” Gardini et al., 2015.

© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. LENS - European Laboratory for Non-linear Spectroscopy, Via Nello Carrara 1, 50019 Sesto Fiorentino, Italy.

E-mail address: capitanio@lens.unifi.it (M. Capitanio).

Specifications table

Subject area	Biophysics
More specific subject area	Single molecule imaging
Type of data	Fluorescence images, Tiff and graph
How data was acquired	Total internal reflection fluorescence (TIRF) microscope
Data format	Raw and analysed
Experimental factors	Recombinant HMM myosin-5B was biotinylated at its C terminal. Streptavidin-QDs (emission ~655 nm) were selectively attached to biotinylated myosin-5B at a 1:5 M ratio to ensure that QD was conjugated to a single motor [3]. F-actin filaments were polymerized, labelled with rhodamine-phalloidin (emission ~580 nm) and immobilized on the coverslip surface. No pretreatment of data
Experimental features	Image stacks were acquired with an EMCCD camera at 50–150 ms integration times, EM gain = 300 under TIRF illumination. Pixel size was 91 nm. Moving QD-myosin-5B were isolated from immobile ones by visual inspection and regions of interest were cut from the original stacks to speed up the localization analysis
Data source location	Sesto Fiorentino, Italy, LENS - European Laboratory for Non-linear Spectroscopy
Data accessibility	All data is with this article
Related research article	Gardini et al. "Dissecting myosin-5B mechanosensitivity and calcium regulation at the single molecule level", Nature Communications 9, 2844 (2018)

Value of the data

- Single molecule techniques allow tracking the movements of molecular motors with nanometer accuracy to investigate their mechanochemical and biophysical properties [4–6]. Here we share raw image stacks of QD-labelled myosin-5B molecules while they move on actin filaments immobilized on a coverslip surface. The data can serve as a reference to develop novel single particle tracking algorithms and analysis methods to precisely derive the position and movement of single molecular motors.
- Generation of molecular motor trajectories using our freely available tracking software (PROOF, available as supplementary material in Gardini et al. [1]) or different tracking methods [7,8] can provide a benchmark for testing molecular motor models and, in particular, walking models for the myosin-5 family of motor proteins [9–11].
- The shared data can be useful to test automated methods for detection of moving vs immobile particles, step detection methods, and trajectory fitting [12].

1. Data

The shared data are image stacks acquired with a single-molecule motility assay in which actin filaments were immobilized on a coverslip surface and single myosin-5B proteins, bound to a Quantum Dot (QD-myosin-5B), were free to interact and possibly move along an actin filament in the presence of ATP (Fig. 1). Images were acquired under total internal reflection fluorescence (TIRF) microscopy, in which the sample is illuminated only in the vicinity of the coverslip surface. Table 1 lists the image stacks attached to this article in supplementary information, including ATP concentration and integration time. Measurements 0–12 are selected regions of interests, the FullField.tif stack contains 1000 frames of a full field (512×512 pixels) acquisition. We used the PROOF software described in Gardini et al. [1] to track the position of the QD-myosin-5B and reconstruct QD x-y trajectories as described in the next section (Fig. 2). The trajectories were then used to compute myosin run length (the distance traveled by the motor before detaching from the actin filament) and velocity, listed in the last two columns of Table 1, as described in the next section.

2. Experimental design, materials, and methods

Single molecule motility experiments were performed on an inverted fluorescence microscope (Nikon ECLIPSE TE300) equipped with a 532 nm laser (Coherent Sapphire) for rhodamine excitation (~3 mW on the sample) and a 488 nm Laser Physics argon laser for QDs-655 nm excitation (~3 mW on the sample) [13].

The sample was loaded into a flow chamber of about 20 μ L volume coated with 1% w/v nitrocellulose/amyl acetate on the coverslip surface. 2 mg/ml of inactivated NEM-myosin-2 was loaded into the




Fig. 1. **a)** Single molecule motility assay scheme. A QD-myo-5B is observed while proceeding along a single actin filament attached to a glass coverslip in the presence of ATP. **b)** Examples of QD-myo-5B motors moving along an actin filament (as indicated by the arrows). The images are superpositions of fluorescence images of actin filaments labelled with rhodamine phalloidin (green) and QD-myo-5B (red). Scale bar is 2 μm .

Table 1

Image stack number in supplementary materials is listed with the ATP concentration, the integration time of the measurement, the calculated run length and velocity.

measure	[ATP] (μM)	Integration time (ms)	Run length (nm)	Velocity (nm/s)
0	0.3	150	835	6
1	25	70	800	497
2	25	70	914	344
3	100	100	871	512
4	100	100	842	547
5	300	100	822	433
6	300	100	632	632
7	500	70	748	763
8	500	70	1077	699
9	700	70	854	718
10	700	70	995	618
11	1000	50	720	757
12	1000	50	625	781
FullField	100	100	–	–




Fig. 2. Example of a trajectory obtained by x, y localization of QD-myo-5B at 100 μM [ATP] through the PROOF program. The trajectory is calculated from attached stack 3 (Table 1) and the coordinates values of this trajectory are listed in the supplementary table.

flow chamber and incubated for 1 min, followed by rhodamine phalloidin F-actin. After careful washing, 0.1 nM QD-myosin-5B was introduced. Experiments were performed in the presence of 0.3–1000 μ M ATP.

Images were acquired in TIRF microscopy using a Nikon Plan Apo TIRF, 1.45 \times NA, 60X oil immersion objective. The image pixel size was 91 nm, obtained by projecting the fluorescence image onto an iXon 3 EMCCD camera, after an additional 3 \times magnification through an achromatic doublet telescope. Depending on the ATP concentration, different integration times (50–150 ms) were used at constant EM gain = 300 (Table 1) (see Gardini et al. [2] for detailed methods).

High precision localization of single QDs was performed by a custom-made Matlab algorithm, named “PROOF”, that automatically detects the fluorescent emitter within a region of interest (ROI) and fits a bidimensional Gaussian function to its intensity profile [1]. Myosin trajectories were derived from x,y coordinates of QDs moving along actin filaments for more than ten frames. The run length was calculated from x–y trajectories (Fig. 2) [2].

Acknowledgments

This work was supported by the European Union's Horizon 2020 research and innovation program under grant agreement no. 654148 Laserlab-Europe and by Ente Cassa di Risparmio di Firenze.

Transparency document

Transparency document associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2019.103973>.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.dib.2019.103973>.

References

- [1] L. Gardini, M. Capitanio, F.S. Pavone, 3D tracking of single nanoparticles and quantum dots in living cells by out-of-focus imaging with diffraction pattern recognition, *Sci. Rep.* 5 (2015) 16088, <https://doi.org/10.1038/srep16088>.
- [2] L. Gardini, S.M. Heissler, C. Arbore, Y. Yang, J.R. Sellers, F.S. Pavone, M. Capitanio, Dissecting myosin-5B mechanosensitivity and calcium regulation at the single molecule level, *Nat. Commun.* 9 (2018) 2844, <https://doi.org/10.1038/s41467-018-05251-z>.
- [3] S.R. Nelson, M.Y. Ali, K.M. Trybus, D.M. Warshaw, Random walk of processive, quantum dot-labeled myosin Va molecules within the actin cortex of COS-7 cells, *Biophys. J.* 97 (2009) 509–518. S0006-3495(09)00968-0 [pii]10.1016/j.bpj.2009.04.052.
- [4] P. Pierobon, S. Achouri, S. Courty, A.R. Dunn, J.A. Spudich, M. Dahan, G. Cappello, Velocity, processivity, and individual steps of single myosin V molecules in live cells, *Biophys. J.* 96 (2009) 4268–4275. S0006-3495(09)00603-1 [pii]10.1016/j.bpj.2009.02.045.
- [5] L. Gardini, A. Tempestini, F.S. Pavone, M. Capitanio, High-speed optical tweezers for the study of single molecular motors, in: C. Lavelle (Ed.), *Mol. Mot., Methods in Molecular Biology*, vol. 1805, Humana Press, New York, NY, 2018, pp. 151–184, https://doi.org/10.1007/978-1-4939-8556-2_9.
- [6] L. Melli, N. Billington, S.A. Sun, J.E. Bird, A. Nagy, T.B. Friedman, Y. Takagi, J.R. Sellers, Bipolar filaments of human nonmuscle myosin 2-A and 2-B have distinct motile and mechanical properties, *Elife* 7 (2018), <https://doi.org/10.7554/eLife.32871>.
- [7] M.P. Bruchez, Quantum dots find their stride in single molecule tracking, *Curr. Opin. Chem. Biol.* 15 (2011) 776–780, <https://doi.org/10.1016/j.cbpa.2011.10.011>.
- [8] A. von Diezmann, Y. Shechtman, W.E. Moerner, Three-Dimensional localization of single molecules for super-resolution imaging and single-particle tracking, *Chem. Rev.* 117 (2017) 7244–7275, <https://doi.org/10.1021/acs.chemrev.6b00629>.
- [9] A. Vilfan, Five models for myosin V, *Front. Biosci.* 14 (2009) 2269–2284, <https://doi.org/10.2741/3378>.
- [10] A. Vilfan, Elastic lever-arm model for myosin V, *Biophys. J.* 88 (2005) 3792–3805, <https://doi.org/10.1529/biophysj.104.046763>.
- [11] A.T. Lombardo, S.R. Nelson, M.Y. Ali, G.G. Kennedy, K.M. Trybus, S. Walcott, D.M. Warshaw, Myosin Va molecular motors manoeuvre liposome cargo through suspended actin filament intersections in vitro, *Nat. Commun.* 8 (2017) 15692, <https://doi.org/10.1038/ncomms15692>.
- [12] B.C. Carter, M. Vershinin, S.P. Gross, A comparison of step-detection methods: how well can you do? *Biophys. J.* 94 (2008) 306–319, <https://doi.org/10.1529/biophysj.107.110601>.
- [13] L. Gardini, M. Calamai, H. Hatakeyama, M. Kanzaki, M. Capitanio, F.S. Pavone, Three-dimensional tracking of quantum dot-conjugated molecules in living cells, in: Y.L. Lyubchenko (Ed.), *Nanoscale Imaging Methods Protoc., Methods in Molecular Biology*, vol. 1814, Humana Press, New York, NY, 2018, pp. 425–448, https://doi.org/10.1007/978-1-4939-8591-3_26.