

Some notions of subharmonicity over the quaternions

Caterina Stoppato

Dipartimento di Matematica e Informatica “U. Dini”, Università di Firenze
Viale Morgagni 67/A, I-50134 Firenze, Italy
stoppato@math.unifi.it

Abstract

This work introduces several notions of subharmonicity for real-valued functions of one quaternionic variable. These notions are related to the theory of slice regular quaternionic functions introduced by Gentili and Struppa in 2006. The interesting properties of these new classes of functions are studied and applied to construct the analogs of Green’s functions.

Acknowledgements. This work stems from a question posed by Filippo Bracci at Università di Roma “Tor Vergata”, where the author was the recipient of the “Michele Cuozzo” prize. The author warmly thanks the Cuozzo family, Università di Roma “Tor Vergata” and Filippo Bracci for the remarkable research opportunity. The author is partly supported by GNSAGA of INdAM and by Finanziamento Premiale FOE 2014 “Splines for accUrate NumeRics: adaptIve models for Simulation Environments” of MIUR. The authors wishes to thank the anonymous referee, whose precious suggestions significantly improved the presentation of this work.

1 Introduction

Let $\mathbb{H} = \mathbb{R} + i\mathbb{R} + j\mathbb{R} + k\mathbb{R}$ denote the real algebra of quaternions and let

$$\mathbb{S} := \{q \in \mathbb{H} : q^2 = -1\} = \{\alpha i + \beta j + \gamma k : \alpha^2 + \beta^2 + \gamma^2 = 1\}$$

denote the 2-sphere of quaternionic imaginary units. For each $I \in \mathbb{S}$, the subalgebra $L_I = \mathbb{R} + I\mathbb{R}$ generated by 1 and I is isomorphic to \mathbb{C} . In recent years, this elementary fact has been the basis for the introduction of a theory of quaternionic functions.

Definition 1.1 ([7]). *Let f be a quaternion-valued function defined on a domain Ω . For each $I \in \mathbb{S}$, let $\Omega_I = \Omega \cap L_I$ and let $f_I = f|_{\Omega_I}$ be the restriction of f to Ω_I . The restriction f_I is called holomorphic if it has continuous partial derivatives and*

$$\frac{1}{2} \left(\frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f_I(x + yI) \equiv 0. \quad (1)$$

The function f is called (slice) regular if, for all $I \in \mathbb{S}$, f_I is holomorphic.

The study of regular quaternionic functions has then grown into a full theory, described in the monograph [6]. It resembles the theory of holomorphic complex functions, but in a many-sided way that reflects the richness of the non-commutative setting.

In the present work, we consider several notions of subharmonicity related to the class of regular quaternionic functions. This study is distinct from the one performed in [10]. Indeed, that work studied the relation between regularity and real harmonicity; moreover, it introduced the notion of *slice harmonic function*: a quaternion-valued (or Clifford-valued) slice functions induced by a harmonic stem function. The present work searches instead for new notions of subharmonicity for real-valued functions of a quaternionic variable, compatible with composition with regular functions.

The first attempt is \mathbb{J} -*plurisubharmonicity*. However, this property is quite restrictive, besides being preserved by composition with a regular function f only if f is *slice preserving*, that is, if $f(\Omega_I) \subseteq L_I$ for all $I \in \mathbb{S}$.

For this reason, the alternative notions of *weakly subharmonic* and *strongly subharmonic* function are introduced. Composition with regular functions turns out to map strongly subharmonic functions into weakly subharmonic ones. Moreover, composition with slice preserving regular functions is proven to preserve weak subharmonicity.

These new notions of subharmonicity turn out to have many nice properties that recall the complex and pluricomplex cases, including mean-value properties and versions of the maximum modulus principle.

These results are finally applied to construct quaternionic analogs of Green's functions, which reveal many peculiarities due to the non-commutative setting.

An appendix comprises the classic properties of subharmonic and plurisubharmonic functions used for our new constructions.

2 Prerequisites

Let us recall a few properties of the algebra of quaternions \mathbb{H} , on which we consider the standard Euclidean metric and topology.

- For each $I \in \mathbb{S}$, the couple $1, I$ can be completed to a (positively oriented) orthonormal basis $1, I, J, K$ by choosing $J \in \mathbb{S}$ with $I \perp J$ and setting $K = IJ$.
- The coordinates of any $q \in \mathbb{H}$ with respect to such a basis can be recovered as

$$\begin{aligned} x_0(q) &= \frac{1}{4}(q - IqI - JqJ - KqK) \\ x_1(q) &= \frac{1}{4I}(q - IqI + JqJ + KqK) \\ x_2(q) &= \frac{1}{4J}(q + IqI - JqJ + KqK) \\ x_3(q) &= \frac{1}{4K}(q + IqI + JqJ - KqK). \end{aligned}$$

- Mapping each $v \in T_{q_0}\mathbb{H} \cong \mathbb{H}$ to Iv for all $q_0 \in \mathbb{H}$ defines an (orthogonal) complex structure on \mathbb{H} , called *constant*. A biholomorphism between (\mathbb{H}, I) and $(L_I^2, I) \cong (\mathbb{C}^2, i)$ can be constructed by mapping each q to $(z_1(q), z_2(q))$, where

$$\begin{aligned} z_1(q) &= x_0(q) + Ix_1(q) = (q - IqI)\frac{1}{2}, \\ z_2(q) &= x_2(q) + Ix_3(q) = (q + IqI)\frac{1}{2J}, \end{aligned}$$

are such that $z_1(q) + z_2(q)J = q$. Both z_1 and z_2 depend on the choice of I ; z_2 also depends on J , but only up to a multiplicative constant $c \in L_I$.

For every domain Ω and every function $f : \Omega \rightarrow \mathbb{H}$, let us denote by $f = f_1 + f_2J$ the corresponding decomposition with f_1, f_2 ranging in L_I . Furthermore $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2 : C^1(\Omega, L_I) \rightarrow C^0(\Omega, L_I)$ will denote the corresponding complex derivatives. In other words,

$$\begin{aligned} \partial_1 &= \frac{1}{2} \left(\frac{\partial}{\partial x_0} - I \frac{\partial}{\partial x_1} \right) \\ \bar{\partial}_1 &= \frac{1}{2} \left(\frac{\partial}{\partial x_0} + I \frac{\partial}{\partial x_1} \right) \\ \partial_2 &= \frac{1}{2} \left(\frac{\partial}{\partial x_2} - I \frac{\partial}{\partial x_3} \right) \\ \bar{\partial}_2 &= \frac{1}{2} \left(\frac{\partial}{\partial x_2} + I \frac{\partial}{\partial x_3} \right). \end{aligned}$$

We notice that these derivatives commute with each other, and that $\partial_1, \bar{\partial}_1$ depend only on I , while $\partial_2, \bar{\partial}_2$ depend on both I and J .

The definition of regular function (Definition 1.1) amounts to requiring that the restriction to Ω_I be holomorphic from (Ω_I, I) to (\mathbb{H}, I) for all $I \in \mathbb{S}$. Curiously, if the domain is carefully chosen then a stronger property holds.

Definition 2.1. *Let Ω be a domain in \mathbb{H} . Ω is a slice domain if it intersects the real axis \mathbb{R} and if, for all $I \in \mathbb{S}$, the intersection Ω_I with the complex plane L_I is connected. Moreover, Ω is termed symmetric if it is axially symmetric with respect to the real axis \mathbb{R} .*

If we denote by $\partial_c f$ the slice derivative

$$\partial_c f(x + Iy) = \frac{1}{2} \left(\frac{\partial}{\partial x} - I \frac{\partial}{\partial y} \right) f_I(x + yI)$$

introduced in [7] and by $\partial_s f$ the spherical derivative

$$\partial_s f(q) = (q - \bar{q})^{-1} (f(q) - f(\bar{q}))$$

introduced in [8], then the aforementioned property can be stated as follows.

Theorem 2.2 ([12]). *Let Ω be a symmetric slice domain, let $f : \Omega \rightarrow \mathbb{H}$ be a regular function and let $q_0 \in \Omega$. Chosen $I, J \in \mathbb{S}$ so that $q_0 \in L_I$ and $I \perp J$, let $z_1, z_2, \bar{z}_1, \bar{z}_2$ be the induced coordinates and let $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2$ be the corresponding derivations. Then*

$$\begin{pmatrix} \bar{\partial}_1 f_1 & \bar{\partial}_2 f_1 \\ \bar{\partial}_1 f_2 & \bar{\partial}_2 f_2 \end{pmatrix} \Big|_{q_0} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}. \quad (2)$$

Furthermore, if $q_0 \notin \mathbb{R}$ then

$$\begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 \\ \partial_1 f_2 & \partial_2 f_2 \end{pmatrix} \Big|_{q_0} = \begin{pmatrix} \partial_c f_1(q_0) & -\overline{\partial_s f_2(q_0)} \\ \partial_c f_2(q_0) & \overline{\partial_s f_1(q_0)} \end{pmatrix}. \quad (3)$$

If, on the contrary, $q_0 \in \mathbb{R}$ then

$$\begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 \\ \partial_1 f_2 & \partial_2 f_2 \end{pmatrix} \Big|_{q_0} = \begin{pmatrix} \partial_c f_1(q_0) & -\overline{\partial_c f_2(q_0)} \\ \partial_c f_2(q_0) & \overline{\partial_c f_1(q_0)} \end{pmatrix}. \quad (4)$$

We point out that we have not proven that f is holomorphic with respect to the constant structure I : with respect to the basis $1, I, J, IJ$, equality (2) only holds at those points q_0 that lie in L_I . In fact, regularity is related to a different notion of holomorphy, which involves non-constant orthogonal complex structures. Let us recall the notations $Re(q) = x_0(q)$, $Im(q) = q - Re(q)$ for $q \in \mathbb{H}$ and let us set

$$\mathbb{J}_{q_0} v := \frac{Im(q_0)}{|Im(q_0)|} v \quad \forall v \in T_{q_0}(\mathbb{H} \setminus \mathbb{R}) \cong \mathbb{H}.$$

Then $\pm \mathbb{J}$ are orthogonal complex structures on

$$\mathbb{H} \setminus \mathbb{R} = \bigcup_{I \in \mathbb{S}} (\mathbb{R} + I\mathbb{R}^+)$$

and they are induced by the natural identification with the complex manifold $\mathbb{C}\mathbb{P}^1 \times (\mathbb{R} + i\mathbb{R}^+)$.

Theorem 2.3 ([5]). *Let Ω be a symmetric slice domain, and let $f : \Omega \rightarrow \mathbb{H}$ be an injective regular function. Then the real differential of f is invertible at each $q \in \Omega$ and the push-forward of \mathbb{J} via f , that is,*

$$\mathbb{J}_{f(q)}^f v = \frac{Im(q)}{|Im(q)|} v \quad \forall v \in T_{f(q)} f(\Omega \setminus \mathbb{R}) \cong \mathbb{H}, \quad (5)$$

is an orthogonal complex structure on $f(\Omega \setminus \mathbb{R})$.

In the hypotheses of the previous theorem, f is (obviously) a holomorphic map from $(\Omega \setminus \mathbb{R}, \mathbb{J})$ to $(f(\Omega \setminus \mathbb{R}), \mathbb{J}^f)$. Furthermore, there is a special class of regular functions such that $\mathbb{J}^f = \mathbb{J}$.

Remark 2.4. *Let $f : \Omega \rightarrow \mathbb{H}$ be a slice preserving regular function, namely a regular function such that $f(\Omega_I) \subseteq L_I$ for all $I \in \mathbb{S}$. Then $f(\Omega \setminus \mathbb{R}) = f(\Omega) \setminus \mathbb{R}$ and f is a holomorphic map from $(\Omega \setminus \mathbb{R}, \mathbb{J})$ to $(f(\Omega) \setminus \mathbb{R}, \mathbb{J})$*

3 Quaternionic notions of subharmonicity

Let Ω be a domain in \mathbb{H} and let

$$\text{us}(\Omega) = \{u : \Omega \rightarrow [-\infty, +\infty), u \text{ upper semicontinuous}, u \not\equiv -\infty\}.$$

For $u \in \text{us}(\Omega)$, we aim at defining some notion of subharmonicity that behaves well when we compose u with a regular function. Remark 2.4 encourages us to consider \mathbb{J} -plurisubharmonic and \mathbb{J} -harmonic functions, i.e., functions that are pluri(sub)harmonic with respect to the complex structure \mathbb{J} . However, the notion of \mathbb{J} -plurisubharmonicity on a symmetric slice domain Ω is induced by plurisubharmonicity in $\mathbb{C}\mathbb{P}^1 \times D_\Omega$ with

$$D_\Omega = \{x + iy \in \mathbb{R} + i\mathbb{R}^+ : x + y\mathbb{S} \subset \Omega\},$$

which amounts to constance in the first variable and subharmonicity in the second variable. We conclude:

Proposition 3.1. *Let Ω be a symmetric domain in \mathbb{H} . A function $u \in \text{us}(\Omega)$ is \mathbb{J} -pluri(sub)harmonic in $\Omega \setminus \mathbb{R}$ if, and only if, there exists a (sub)harmonic function $v : D_\Omega \rightarrow \mathbb{R}$ such that $u(x + Iy) = v(x + iy)$ for all $I \in \mathbb{S}$ and for all $x + iy \in D_\Omega$. In particular, a function $u \in C^2(\Omega, \mathbb{R})$ is \mathbb{J} -plurisubharmonic in $\Omega \setminus \mathbb{R}$ if, and only if, $u(x + Iy)$ does not depend on I and*

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) u(x + Iy) \geq 0; \quad (6)$$

it is \mathbb{J} -harmonic if, and only if, equality holds at all points.

Clearly, if we complete $1, I$ to a basis $1, I, J, IJ$ with $I \perp J$, consider the induced coordinates $z_1, z_2, \bar{z}_1, \bar{z}_2$ and let $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2$ be the corresponding derivations, then inequality (6) is equivalent to $\bar{\partial}_1 \partial_1 u_I \geq 0$.

In order to get a richer class of functions, we need to suitably weaken the notion of subharmonicity considered. We are thus encouraged to give the following definition.

Definition 3.2. *Let Ω be a domain in \mathbb{H} and let $u \in \text{us}(\Omega)$. We call u weakly subharmonic if for all $I \in \mathbb{S}$ the restriction $u_I = u|_{\Omega_I}$ is subharmonic (after the natural identification between L_I and \mathbb{C}). We say that u is weakly harmonic if, for all $I \in \mathbb{S}$, u_I is harmonic.*

Remark 3.3. *A function $u \in C^2(\Omega, \mathbb{R})$ is weakly subharmonic if, and only if, inequality (6) holds for all $I \in \mathbb{S}$; u is weakly harmonic if, and only if, equality holds at all points.*

By construction:

Proposition 3.4. *Let Ω be a symmetric domain in \mathbb{H} and let $u \in \text{us}(\Omega)$. If u is \mathbb{J} -pluri(sub)harmonic in $\Omega \setminus \mathbb{R}$ then it is weakly (sub)harmonic in $\Omega \setminus \mathbb{R}$. If, moreover, u is continuous at all points of $\Omega \cap \mathbb{R}$ then u is weakly (sub)harmonic in Ω .*

The converse implication is not true, as shown by the next example. Here, $\langle \cdot, \cdot \rangle$ denotes the Euclidean scalar product on $\text{Im}(\mathbb{H}) \cong \mathbb{R}^3$.

Example 3.5. *All real affine functions $u : \mathbb{H} \rightarrow \mathbb{R}$ are weakly harmonic, including the coordinates x_0, x_1, x_2, x_3 with respect to any basis $1, I, J, IJ$ with $I, J \in \mathbb{S}, I \perp J$. On the other hand, x_1, x_2, x_3 are not \mathbb{J} -plurisubharmonic in $\mathbb{H} \setminus \mathbb{R}$, as $x_1(x + Iy) = y\langle I, i \rangle, x_2(x + Iy) = y\langle I, j \rangle, x_3(x + Iy) = y\langle I, k \rangle$ are not constant in I .*

Actually, a stronger property holds for real affine functions $u : \mathbb{H} \rightarrow \mathbb{R}$: they are pluriharmonic with respect to any constant orthogonal complex structure. This motivates the next definition.

Definition 3.6. *Let Ω be a domain in \mathbb{H} and let $u \in \text{us}(\Omega)$. We say that u is strongly (sub)harmonic if it is pluri(sub)harmonic with respect to every constant orthogonal complex structure on Ω .*

Remark 3.7. *A function $u \in C^2(\Omega, \mathbb{R})$ is strongly subharmonic if, for all $I, J \in \mathbb{S}$ with $I \perp J$,*

$$H_{I,J}(u) = \begin{pmatrix} \bar{\partial}_1 \partial_1 u & \bar{\partial}_1 \partial_2 u \\ \bar{\partial}_2 \partial_1 u & \bar{\partial}_2 \partial_2 u \end{pmatrix} \quad (7)$$

is a positive semidefinite matrix at each $q \in \Omega$. The function u is strongly harmonic if for all $I, J \in \mathbb{S}$ with $I \perp J$ the matrix $H_{I,J}(u)$ has constant rank 0.

Clearly, if the matrix $H_{I,J}(u)$ is positive semidefinite then its $(1,1)$ -entry $\bar{\partial}_1 \partial_1 u$ is non-negative. Similarly, if $H_{I,J}(u)$ has constant rank 0 then $\bar{\partial}_1 \partial_1 u \equiv 0$. This leads to the next result, which, however, is not only true for $u \in C^2(\Omega, \mathbb{R})$ but also for $u \in \text{us}(\Omega)$.

Proposition 3.8. *Let $u \in \text{us}(\Omega)$. If u is strongly (sub)harmonic then it is weakly (sub)harmonic.*

Proof. Let $u \in \text{us}(\Omega)$ be strongly (sub)harmonic, let $I \in \mathbb{S}$ and let us prove that u_I is (sub)harmonic. By construction, u is pluri(sub)harmonic with respect to the constant orthogonal complex structure I . Moreover, the inclusion map $\text{incl} : \Omega_I \rightarrow \Omega$ is a holomorphic map from (Ω_I, I) to (Ω, I) . As a consequence, $u_I = u \circ \text{incl}$ is (sub)harmonic, as desired. \square


Example 3.5 shows that strong (sub)harmonicity does not imply \mathbb{J} -pluri(sub)harmonicity. The converse implication does not hold, either:

Example 3.9. *The function $u(q) = \text{Re}(q^2)$ is \mathbb{J} -pluriharmonic in $\mathbb{H} \setminus \mathbb{R}$, as $u(x + Iy) = x^2 - y^2$ does not depend on I and $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)u(x + Iy) \equiv 0$. On the other hand, u is not strongly subharmonic. Actually, it is not plurisubharmonic with respect to any constant orthogonal complex structure I : after choosing $J \in \mathbb{S}$ with $J \perp I$, we compute*

$$u(z_1 + z_2 J) = \text{Re}(z_1^2 - z_2 \bar{z}_2 + (z_1 z_2 + z_2 \bar{z}_1)J) = \frac{z_1^2 + \bar{z}_1^2}{2} - z_2 \bar{z}_2$$

for all $z_1, z_2 \in L_I$, so that $H_{I,J}(u) \equiv \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$.

We have proven the following implications (none of which can be reversed):



A similar scheme can be drawn for the quaternionic notions of harmonicity. We show with a further example that a strongly subharmonic function is not necessarily strongly *harmonic* when it is weakly harmonic, or even \mathbb{J} -pluriharmonic.

Example 3.10. *Consider the function $u : \mathbb{H} \rightarrow \mathbb{R}$ with $u(q) := \log |q|$ for all $q \in \mathbb{H} \setminus \{0\}$ and $u(0) := -\infty$. u is \mathbb{J} -pluriharmonic in $\mathbb{H} \setminus \mathbb{R}$, as $u(x + Iy) = \frac{1}{2} \log(x^2 + y^2)$. As a consequence, u is also weakly harmonic. On the other hand, for any choice of $I, J \in \mathbb{S}$, the fact that $u(z_1 + z_2 J) = \frac{1}{2} \log(z_1 \bar{z}_1 + z_2 \bar{z}_2)$ implies that*

$$\begin{aligned}
 H_{I,J}(u)|_{z_1 + z_2 J} &= \frac{1}{2(z_1 \bar{z}_1 + z_2 \bar{z}_2)^2} \begin{pmatrix} z_2 \bar{z}_2 & -z_1 \bar{z}_2 \\ -z_2 \bar{z}_1 & z_1 \bar{z}_1 \end{pmatrix} \\
 &= \frac{1}{2(|z_1|^2 + |z_2|^2)^2} \begin{pmatrix} |z_2|^2 & -z_1 \bar{z}_2 \\ -z_2 \bar{z}_1 & |z_1|^2 \end{pmatrix}.
 \end{aligned}$$

Hence, u is strongly subharmonic but it is not strongly harmonic.

Let us review a few classical constructions in our new environment.

Remark 3.11. *On a given domain Ω , let us denote by $\text{wsh}(\Omega)$ the set of weakly subharmonic functions, by $\text{ssh}(\Omega)$ that of strongly subharmonic functions, and by $\text{psh}_{\mathbb{J}}(\Omega)$ that of \mathbb{J} -plurisubharmonic functions on Ω (if Ω equals a symmetric domain minus \mathbb{R}). If S is any of these sets then:*

1. S is a convex cone;
2. for all $u \in S$, if φ is a real-valued C^2 function on a neighborhood of $u(\mathbb{R})$ and if φ is increasing and convex then $\varphi \circ u : \Omega \rightarrow \mathbb{R}$ also belongs to S ;
3. for all $u_1, u_2 \in S$, the function $u(q) = \max\{u_1(q), u_2(q)\}$ belongs to S ;

4. if $\{u_\alpha\}_{\alpha \in A}$ (with $A \neq \emptyset$) is a family in S , locally bounded from above, and if $u(q) = \sup_{\alpha \in A} u_\alpha(q)$ for all $q \in \Omega$ then the upper semicontinuous regularization u^* belongs to S .

Example 3.12. For any $\alpha > 0$, the function $u : \mathbb{H} \rightarrow \mathbb{R} \ q \mapsto |q|^\alpha$ is strongly subharmonic in \mathbb{H} and it is \mathbb{J} -plurisubharmonic in $\mathbb{H} \setminus \mathbb{R}$.

Example 3.13. The functions $Re^2(q) = x_0^2(q)$ and $|Im(q)|^2 = x_1^2(q) + x_2^2(q) + x_3^2(q)$ are strongly subharmonic in \mathbb{H} . (They are also \mathbb{J} -plurisubharmonic in $\mathbb{H} \setminus \mathbb{R}$, as $Re^2(x+Iy) = x^2$ and $|Im(x+Iy)|^2 = y^2$).

We conclude this section showing that any given subharmonic function on an axially symmetric planar domain extends to a weakly subharmonic function on the corresponding symmetric domain of \mathbb{H} .

Remark 3.14. If we start with a domain $D \subseteq \mathbb{C}$ that is symmetric with respect to the real axis and a (sub)harmonic function v on D , we may define a weakly (sub)harmonic function u on the symmetric domain $\Omega = \bigcup_{x+iy \in D} x + y\mathbb{S}$ by setting

$$u(x + Iy) := \frac{1 + \langle I, i \rangle}{2} v(x + iy) + \frac{1 - \langle I, i \rangle}{2} v(x - iy)$$

for all $x \in \mathbb{R}, I \in \mathbb{S}, y > 0$ such that $x + Iy \in \Omega$ and $u(x) := v(x)$ for all $x \in \Omega \cap \mathbb{R}$.

4 Composition with regular functions

We now want to understand the behavior of the different notions of subharmonicity we introduced, under composition with regular functions. For \mathbb{J} -pluri(sub)harmonicity, Remark 2.4 immediately implies:

Proposition 4.1. Let Ω be a symmetric domain in \mathbb{H} and let $u \in \text{us}(\Omega)$. u is \mathbb{J} -pluri(sub)harmonic in $\Omega \setminus \mathbb{R}$ if, and only if, for every symmetric domain Ω' in \mathbb{H} and every slice preserving regular function $f : \Omega' \setminus \mathbb{R} \rightarrow \Omega \setminus \mathbb{R}$, the composition $u \circ f$ is \mathbb{J} -pluri(sub)harmonic in $\Omega' \setminus \mathbb{R}$.

It is essential to restrict to slice preserving regular functions. If we compose u with an injective regular function f then the only sufficient condition we know in order for $u \circ f$ to be \mathbb{J} -pluri(sub)harmonic is, that u be \mathbb{J}^f -pluri(sub)harmonic (see Theorem 2.3).

For weak (sub)harmonicity, we can prove the next result.

Theorem 4.2. Let $u \in \text{us}(\Omega)$. u is weakly (sub)harmonic in Ω if, and only if, for every symmetric domain Ω' in \mathbb{H} and every slice preserving regular function $f : \Omega' \rightarrow \Omega$, the composition $u \circ f$ is weakly (sub)harmonic in Ω' .

Proof. If $u \circ f$ is weakly (sub)harmonic for all slice preserving regular f then in particular $u = u \circ id$ is weakly (sub)harmonic.

Conversely, let $u \in \text{us}(\Omega)$ be weakly (sub)harmonic, let $f : \Omega' \rightarrow \Omega$ be a slice preserving regular function and let us prove that $u \circ f$ is weakly (sub)harmonic. For each $I \in \mathbb{S}$, u_I is (sub)harmonic in Ω_I and the restriction f_I is a holomorphic map from (Ω'_I, I) to (Ω_I, I) . As a consequence, $(u \circ f)_I = u_I \circ f_I$ is (sub)harmonic in Ω'_I . \square

As for strong harmonicity and subharmonicity, they are preserved under composition with quaternionic affine transformations, as the latter are holomorphic with respect to any constant structure $I \in \mathbb{S}$:

Remark 4.3. If u is a strongly (sub)harmonic function on a domain $\Omega \subseteq \mathbb{H}$ then, for any $a, b \in \mathbb{H}$ with $b \neq 0$, the function $v(q) = u(a + qb)$ is strongly (sub)harmonic in $\Omega b^{-1} - a$.

However, strong harmonicity and subharmonicity are not preserved by composition with other regular functions, not even slice preserving regular functions (see Example 3.9). For this reason, we address the study of their composition with regular functions by direct computation, starting with the C^2 case.

Lemma 4.4. *Let $I, J \in \mathbb{S}$ with $I \perp J$ and let us consider the associated $\partial_1, \partial_2, \bar{\partial}_1, \bar{\partial}_2$. Let Ω be a domain in \mathbb{H} and let $u \in C^2(\Omega, \mathbb{R})$. For every symmetric slice domain Ω' and for every regular function $f : \Omega' \rightarrow \Omega$, we have*

$$\bar{\partial}_1 \partial_1(u \circ f)|_{q_0} = (\overline{\partial_1 f_1}, \overline{\partial_1 f_2})|_{q_0} \cdot H_{I, J}(u)|_{f(q_0)} \cdot \begin{pmatrix} \partial_1 f_1 \\ \partial_1 f_2 \end{pmatrix} \Big|_{q_0} \quad (8)$$

at each $q_0 \in \Omega'_I$. If, moreover, f is slice preserving then

$$\bar{\partial}_1 \partial_1(u \circ f)|_{q_0} = |\partial_1 f_1|_{q_0}^2 \cdot \bar{\partial}_1 \partial_1 u|_{f(q_0)} \quad (9)$$

at each $q_0 \in \Omega'_I$. The same is true if there exists a constant $c \in \mathbb{H}$ such that $f_I + c$ maps Ω'_I to L_I .

Proof. We compute:

$$\partial_1(u \circ f) = (\partial_1 u) \circ f \cdot \partial_1 f_1 + (\partial_2 u) \circ f \cdot \partial_1 f_2 + (\bar{\partial}_1 u) \circ f \cdot \partial_1 \bar{f}_1 + (\bar{\partial}_2 u) \circ f \cdot \partial_1 \bar{f}_2$$

and

$$\begin{aligned} \bar{\partial}_1 \partial_1(u \circ f) &= \bar{\partial}_1((\partial_1 u) \circ f) \cdot \partial_1 f_1 + (\partial_1 u) \circ f \cdot \bar{\partial}_1 \partial_1 f_1 + \\ &\quad + \bar{\partial}_1((\partial_2 u) \circ f) \cdot \partial_1 f_2 + (\partial_2 u) \circ f \cdot \bar{\partial}_1 \partial_1 f_2 + \\ &\quad + \bar{\partial}_1((\bar{\partial}_1 u) \circ f) \cdot \partial_1 \bar{f}_1 + (\bar{\partial}_1 u) \circ f \cdot \bar{\partial}_1 \partial_1 \bar{f}_1 + \\ &\quad + \bar{\partial}_1((\bar{\partial}_2 u) \circ f) \cdot \partial_1 \bar{f}_2 + (\bar{\partial}_2 u) \circ f \cdot \bar{\partial}_1 \partial_1 \bar{f}_2. \end{aligned}$$

If we evaluate the previous expression at a point $q \in L_I$, equality (2) guarantees the vanishing of all terms but the first and the third. Hence,

$$\bar{\partial}_1 \partial_1(u \circ f)|_q = \bar{\partial}_1((\partial_1 u) \circ f)|_q \cdot \partial_1 f_1|_q + \bar{\partial}_1((\bar{\partial}_1 u) \circ f)|_q \cdot \partial_1 \bar{f}_1|_q$$

where

$$\begin{aligned} \bar{\partial}_1((\partial_1 u) \circ f)|_q &= \\ &= \partial_1 \partial_1 u|_{f(q)} \cdot \bar{\partial}_1 f_1|_q + \partial_2 \partial_1 u|_{f(q)} \cdot \bar{\partial}_1 f_2|_q + \bar{\partial}_1 \partial_1 u|_{f(q)} \cdot \bar{\partial}_1 \bar{f}_1|_q + \bar{\partial}_2 \partial_1 u|_{f(q)} \cdot \bar{\partial}_1 \bar{f}_2|_q = \\ &= \bar{\partial}_1 \partial_1 u|_{f(q)} \cdot \overline{\partial_1 f_1}|_q + \bar{\partial}_2 \partial_1 u|_{f(q)} \cdot \overline{\partial_1 f_2}|_q \end{aligned}$$

and

$$\begin{aligned} \bar{\partial}_1((\bar{\partial}_1 u) \circ f)|_q &= \\ &= \partial_1 \bar{\partial}_1 u|_{f(q)} \cdot \bar{\partial}_1 f_1|_q + \partial_2 \bar{\partial}_1 u|_{f(q)} \cdot \bar{\partial}_1 f_2|_q + \bar{\partial}_1 \bar{\partial}_1 u|_{f(q)} \cdot \bar{\partial}_1 \bar{f}_1|_q + \bar{\partial}_2 \bar{\partial}_1 u|_{f(q)} \cdot \bar{\partial}_1 \bar{f}_2|_q = \\ &= \bar{\partial}_1 \bar{\partial}_1 u|_{f(q)} \cdot \overline{\partial_1 f_1}|_q + \bar{\partial}_2 \bar{\partial}_1 u|_{f(q)} \cdot \overline{\partial_1 f_2}|_q. \end{aligned}$$

Thus,

$$\begin{aligned} \bar{\partial}_1 \partial_1(u \circ f)|_q &= \bar{\partial}_1 \partial_1 u|_{f(q)} \cdot |\partial_1 f_1|_q^2 + \bar{\partial}_2 \partial_1 u|_{f(q)} \cdot \overline{\partial_1 f_2}|_q \cdot \partial_1 f_1|_q + \\ &\quad + \bar{\partial}_1 \bar{\partial}_1 u|_{f(q)} \cdot \overline{\partial_1 f_1}|_q \partial_1 f_2|_q + \bar{\partial}_2 \bar{\partial}_1 u|_{f(q)} \cdot |\partial_1 f_2|_q^2; \end{aligned}$$

that is,

$$\bar{\partial}_1 \partial_1(u \circ f)|_q = (\overline{\partial_1 f_1}, \overline{\partial_1 f_2})|_q \cdot \begin{pmatrix} \bar{\partial}_1 \partial_1 u & \bar{\partial}_1 \bar{\partial}_1 u \\ \bar{\partial}_2 \partial_1 u & \bar{\partial}_2 \bar{\partial}_1 u \end{pmatrix} \Big|_{f(q)} \cdot \begin{pmatrix} \partial_1 f_1 \\ \partial_1 f_2 \end{pmatrix} \Big|_q.$$

Finally, if there exists $c = c_1 + c_2 J \in \mathbb{H}$ such that $f_I + c$ maps Ω'_I to L_I , then $f_2 \equiv -c_2$ in Ω'_I so that $\partial_1 f_2$ vanishes identically in Ω'_I and

$$\bar{\partial}_1 \partial_1(u \circ f)|_q = |\partial_1 f_1|_q^2 \cdot \bar{\partial}_1 \partial_1 u|_{f(q)},$$

as desired. \square

We are now ready to study the composition of strongly (sub)harmonic C^2 functions with regular functions.

Theorem 4.5. *Let $u \in C^2(\Omega, \mathbb{R})$. u is strongly (sub)harmonic if, and only if, for every symmetric slice domain Ω' and for every regular function $f : \Omega' \rightarrow \Omega$, the composition $u \circ f$ is weakly (sub)harmonic.*

Proof. If $u : \Omega \rightarrow \mathbb{R}$ is strongly subharmonic then, for all $I, J \in \mathbb{S}$ (with $I \perp J$), the matrix $H_{I,J}(u)$ is positive semidefinite. For every regular function $f : \Omega' \rightarrow \Omega$ and for all $I \in \mathbb{S}$, Lemma 4.4 implies $\bar{\partial}_1 \partial_1 (u \circ f)|_{z_1} \geq 0$ at each $z_1 \in \Omega_I$. Hence, $u \circ f$ is weakly subharmonic.

Conversely, if $u \circ f$ is weakly subharmonic for every regular function $f : \Omega' \rightarrow \Omega$ then we can prove that u is strongly subharmonic in the following way. Let us fix $I, J \in \mathbb{S}, p \in \Omega$ (with $I \perp J$) and prove that $H_{I,J}(u)$ is positive semidefinite at p , i.e., that

$$(\bar{v}_1, \bar{v}_2) \cdot H_{I,J}(u)|_p \cdot \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \geq 0$$

for arbitrary $v_1, v_2 \in L_I$. Let us set $v := v_1 + v_2 J$ and $f(q) := qv + p$ for $q \in B(0, R)$ (with $R > 0$ small enough to guarantee the inclusion of $f(B(0, R)) = B(p, |v|R)$ into Ω). By direct computation, $\partial_c f \equiv v$. Formula (3) yields the equalities $\partial_1 f|_0 = v_1, \partial_1 f|_0 = v_2$. Taking into account Lemma 4.4 and the fact that $f(0) = p$, we conclude that

$$(\bar{v}_1, \bar{v}_2) \cdot H_{I,J}(u)|_p \cdot \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \bar{\partial}_1 \partial_1 (u \circ f)|_0.$$

Since $u \circ f$ is weakly subharmonic, $\bar{\partial}_1 \partial_1 (u \circ f)|_0 \geq 0$ and we have proven the desired inequality.

Analogous reasonings characterize strong *harmonicity*. \square

The previous result allows us to construct a large class of examples of weakly subharmonic functions.

Example 4.6. *For any regular function $f : \Omega \rightarrow \mathbb{H}$ on a symmetric slice domain Ω , the components of f with respect to any basis $1, I, J, IJ$ with $I, J \in \mathbb{S}, I \perp J$ are weakly harmonic. Furthermore, for all $\alpha > 0$ the functions $\log |f|, |f|^\alpha, \operatorname{Re}^2 f, |\operatorname{Im} f|^2$ are weakly subharmonic.*

5 Mean-value property and consequences

We can characterize weak and strong (sub)harmonicity of $u \in \operatorname{us}(\Omega)$ in terms of mean-value properties. For each $I \in \mathbb{S}, a \in \Omega, b \in \mathbb{H} \setminus \{0\}$ such that Ω includes the disc $\Gamma_{I,a,b} := \{a + \lambda b : \lambda \in L_I, |\lambda| \leq 1\}$, we will use the notation

$$l_I(u; a, b) := \frac{1}{2\pi} \int_0^{2\pi} u(a + e^{I\vartheta} b) d\vartheta. \quad (10)$$

Proposition 5.1. *Let Ω be a domain in \mathbb{H} and let $u \in \operatorname{us}(\Omega)$. u is weakly subharmonic if, and only if, the inequality*

$$u(a) \leq l_I(u; a, b) \quad (11)$$

holds for all $I \in \mathbb{S}, a \in \Omega_I, b \in L_I \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega_I$. u is weakly harmonic if, and only if, (u does not take the value $-\infty$ and) equality always holds in formula (11).

Proof. Fix any $I \in \mathbb{S}$. By the mean-value characterization of subharmonic functions (see Theorem 8.1), u_I is subharmonic in Ω_I if, and only if, $u(a) \leq l_I(u; a, b)$ for all $a \in \Omega_I, b \in L_I \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega_I$. The corresponding equalities characterize harmonicity because u_I is harmonic if, and only if, u_I and $-u_I$ are both subharmonic (see Corollary 8.3). \square

Proposition 5.2. *Let Ω be a domain in \mathbb{H} and let $u \in \operatorname{us}(\Omega)$. u is strongly subharmonic if, and only if, the inequality*

$$u(a) \leq l_I(u; a, b) \quad (12)$$

holds for all $I \in \mathbb{S}, a \in \Omega, b \in \mathbb{H} \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega$. u is strongly harmonic if, and only if, (u does not take the value $-\infty$ and) equality always holds in formula (12).

Proof. For each $I \in \mathbb{S}$, let us apply the mean-value characterization of plurisubharmonic functions (see Theorem 8.6) to establish whether u is I -plurisubharmonic. This happens if, and only if, $u(a) \leq l_I(u; a, b)$ for all $a \in \Omega, b \in \mathbb{H} \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega$. The corresponding equalities characterize I -pluriharmonicity. \square

As an application of the previous results, we can extend Theorem 4.5 to all $u \in \text{us}(\Omega)$.

Theorem 5.3. *Let $u \in \text{us}(\Omega)$. u is strongly (sub)harmonic if, and only if, for every regular function $f : \Omega' \rightarrow \Omega$ the composition $u \circ f$ is weakly (sub)harmonic.*

Proof. Let us suppose the composition $u \circ f$ with any regular function $f : \Omega' \rightarrow \Omega$ to be weakly subharmonic and let us prove that u is strongly subharmonic. By Proposition 5.2, it suffices to prove that, for any $I \in \mathbb{S}$, $a \in \Omega$, $b \in \mathbb{H} \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega$, the inequality

$$u(a) \leq l_I(u; a, b)$$

holds. If we set $f(q) := a + qb$, then $f(0) = a$ and f maps the disc $\Gamma_{I,0,1}$ into the disc $\Gamma_{I,a,b}$. Thus, it suffices to prove that

$$u(f(0)) \leq l_I(u \circ f; 0, 1).$$

But this inequality is true by Proposition 5.1, since $u \circ f$ is weakly subharmonic in a domain Ω' such that $\Gamma_{I,0,1} \subset \Omega'_I$. Analogous considerations can be made for the harmonic case.

Conversely, let $u \in \text{us}(\Omega)$ be strongly (sub)harmonic, let $f : \Omega' \rightarrow \Omega$ be a regular function and let us prove that $u \circ f$ is weakly (sub)harmonic. For each $I \in \mathbb{S}$, u is I -pluri(sub)harmonic and the restriction f_I is a holomorphic map from (Ω'_I, I) to (Ω, I) . As a consequence, $(u \circ f)_I = u \circ f_I$ is (sub)harmonic in Ω'_I , as desired. \square

A form of maximum modulus principle holds for weakly or strongly plurisubharmonic functions.

Proposition 5.4. *Let Ω be a domain in \mathbb{H} and suppose $u \in \text{us}(\Omega)$ to be weakly subharmonic. If u has a local maximum point $p \in \Omega_I$ then u_I is constant in the connected component of Ω_I that includes p . If, moreover, u is strongly subharmonic, then u is constant in Ω .*

Proof. In our hypotheses, u_I is a subharmonic function with a local maximum point $p \in \Omega_I$. Thus, u_I is constant in the connected component of Ω_I that includes p by the maximum modulus principle for subharmonic functions (see Theorem 8.2).

If, moreover, u is strongly subharmonic then it is I -plurisubharmonic. Since we assumed Ω to be connected, u is constant in Ω by the maximum modulus principle for plurisubharmonic functions (see Theorem 8.8). \square

Let us now consider maximality.

Definition 5.5. *Let S be a class of real-valued functions on an open set D and let v be an element of S . Suppose that, for any relatively compact subset G of D and for all $\nu \in S$ with $\nu \leq v$ in ∂G , the inequality $\nu \leq v$ holds throughout G . In this situation, we say that v is maximal in S (or among the elements of S).*

Bearing in mind that harmonic functions of one complex variable are the maximal elements of the class of plurisubharmonic functions (see Remark 8.9), we can characterize weak harmonicity as follows.

Remark 5.6. *Let Ω be a domain in \mathbb{H} and let $u \in \text{wsh}(\Omega)$. u is weakly harmonic if, and only if, for all $I \in \mathbb{S}$, the restriction u_I is maximal among subharmonic functions on Ω_I . As a consequence, if u is weakly harmonic then u is maximal in $\text{wsh}(\Omega)$.*

Let us now consider strongly subharmonic functions. Since they are plurisubharmonic with respect to all constant structures, we can make the following observation.

Remark 5.7. *Let Ω be a domain in \mathbb{H} and let $u \in \text{ssh}(\Omega)$. If u is strongly harmonic then it is maximal in $\text{ssh}(\Omega)$. Furthermore, if $u \in C^2(\Omega)$ then u is maximal in $\text{ssh}(\Omega)$ if and only if $\det H_{I,J}(u) \equiv 0$ for all $I, J \in \mathbb{S}$ with $I \perp J$.*

It is easy to exhibit a maximal element of $\text{ssh}(\Omega)$ that is not strongly harmonic.

Example 5.8. *The function $u(q) = \log |q|$ is a strongly subharmonic function on \mathbb{H} . The explicit computations in Example 3.10 show that u is maximal but not strongly harmonic.*

6 Approximation

An approximation result holds for strongly subharmonic functions. For all $\varepsilon > 0$, let

$$\Omega_\varepsilon := \begin{cases} \{q \in \Omega : \text{dist}(q, \partial\Omega) > \varepsilon\} & \text{if } \Omega \neq \mathbb{H} \\ \mathbb{H} & \text{if } \Omega = \mathbb{H} \end{cases}$$

and let $u * \chi_\varepsilon$ denote the convolution of u with the standard smoothing kernels χ_ε of $\mathbb{H} \cong \mathbb{R}^4$ (see Definition 8.4).

Proposition 6.1. *Let $u \in \text{ssh}(\Omega)$. If $\varepsilon > 0$ is such that Ω_ε is not empty, then $u * \chi_\varepsilon \in C^\infty \cap \text{ssh}(\Omega_\varepsilon)$. Moreover, $u * \chi_\varepsilon$ monotonically decreases with decreasing ε and*

$$\lim_{\varepsilon \rightarrow 0^+} u * \chi_\varepsilon(q) = u(q) \quad (13)$$

for each $q \in \Omega$.

Proof. Fix any $I \in \mathbb{S}$, so that u is I -plurisubharmonic. The fact that $u * \chi_\varepsilon \in C^\infty \cap \text{ssh}(\Omega_\varepsilon)$, as well as our second statement, follow from the corresponding classic properties of plurisubharmonic functions (see Theorem 8.7). \square

On the other hand, convolution with the standard smoothing kernel χ_ε does not preserve weak subharmonicity. This can be shown with an example, which uses the notation $l_I(u; a, b)$ established in formula (10). As customary, $l_I(u; \cdot, b)$ denotes the function $a \mapsto l_I(u; a, b)$.

Example 6.2. *The function $u(q) = \text{Re}(q^2)$ is in $\text{wsh}(\mathbb{H})$, but $u * \chi_\varepsilon$ does not belong to $\text{wsh}(\mathbb{H})$. Indeed, we saw that for each orthonormal basis $1, I, J, IJ$ of \mathbb{H} we have $H_{I,J}(u) \equiv \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$. Hence, $-u$ is strongly subharmonic and the same is true for $-u * \chi_\varepsilon$ by the previous proposition. In particular, $-u * \chi_\varepsilon \in \text{wsh}(\mathbb{H})$ so that $u * \chi_\varepsilon$ can only be in $\text{wsh}(\mathbb{H})$ if it is weakly harmonic. This amounts to requiring that for each $I \in \mathbb{S}$, $a \in \Omega_I, b \in L_I \setminus \{0\}$ such that $\Gamma_{I,a,b} \subset \Omega$ the equality*

$$u * \chi_\varepsilon(a) = l_I(u * \chi_\varepsilon; a, b) = l_I(u; \cdot, b) * \chi_\varepsilon(a),$$

holds. But this happens if, and only if, $u(a - q) = l_I(u; a - q, b)$ for all q in the support $\overline{B(0, \varepsilon)}$ of χ_ε . This cannot be true, since (if z_1, z_2 denote the complex variables with respect to the orthonormal basis $1, I, J, IJ$) the function $-u$ is strictly subharmonic in z_2 .

7 Green's functions

We now consider the analogs of Green's functions in the context of weakly and strongly subharmonic functions.

Definition 7.1. *Let Ω be a domain in \mathbb{H} , let $q_0 \in \Omega$, and set*

$$\begin{aligned} \text{wsh}_{q_0}(\Omega) &:= \left\{ u \in \text{wsh}(\Omega) : u < 0, \limsup_{q \rightarrow q_0} |u(q) - \log |q - q_0|| < \infty \right\} \\ \text{ssh}_{q_0}(\Omega) &:= \text{wsh}_{q_0}(\Omega) \cap \text{ssh}(\Omega). \end{aligned}$$

For all $q \in \Omega$, let us define

$$\begin{aligned} w(q) &:= \begin{cases} -\infty & \text{if } \text{wsh}_{q_0}(\Omega) = \emptyset \\ \sup\{u(q) : u \in \text{wsh}_{q_0}(\Omega)\} & \text{otherwise} \end{cases} \\ s(q) &:= \begin{cases} -\infty & \text{if } \text{ssh}_{q_0}(\Omega) = \emptyset \\ \sup\{u(q) : u \in \text{ssh}_{q_0}(\Omega)\} & \text{otherwise} \end{cases} \end{aligned}$$

The Green function of Ω with logarithmic pole at q_0 , denoted $g_{q_0}^\Omega$, is the upper semicontinuous regularization w^* of w . The strongly subharmonic Green function of Ω with logarithmic pole at q_0 , denoted $G_{q_0}^\Omega$, is the upper semicontinuous regularization s^* of s .

Remark 7.2. By construction, $G_{q_0}^\Omega(q) \leq g_{q_0}^\Omega(q)$ for all $q \in \Omega$. Moreover, any inclusion $\Omega' \subseteq \Omega$ implies $g_{q_0}^\Omega(q) \leq g_{q_0}^{\Omega'}(q)$ and $G_{q_0}^\Omega(q) \leq G_{q_0}^{\Omega'}(q)$.

Let us construct a basic example. We will use the notations $\mathbb{B} := B(0, 1)$, where

$$B(q_0, R) := \{q \in \mathbb{H} : |q - q_0| < R\}$$

for all $q_0 \in \mathbb{H}$, $R > 0$, and $\mathbb{B}_I := \mathbb{B} \cap L_I$.

Example 7.3. We can easily prove that

$$G_0^{\mathbb{B}}(q) = g_0^{\mathbb{B}}(q) = \log |q|$$

for all $q \in \mathbb{B}$. Indeed, $q \mapsto \log |q|$ is clearly an element of $\text{ssh}_0(\mathbb{B}) \subseteq \text{wsh}_0(\mathbb{B})$. Furthermore, for each $u \in \text{wsh}_0(\mathbb{B})$, the inequality $u(q) \leq \log |q|$ holds throughout \mathbb{B} . Indeed, for all $I \in \mathbb{S}$ it holds $u_I(z) \leq \log |z|$ for all $z \in \mathbb{B}_I$ because $z \mapsto \log |z|$ is the (complex) Green function of the disc \mathbb{B}_I .

Further examples can be derived by means of the next results.

Lemma 7.4. Let f be any affine transformation of \mathbb{H} , let Ω be a domain in \mathbb{H} and fix $q_0 \in \Omega$. Then

$$G_{f(q_0)}^{f(\Omega)}(f(q)) = G_{q_0}^\Omega(q)$$

for all $q \in \Omega$.

Proof. By repeated applications of Remark 4.3, we conclude that

$$\text{ssh}_{q_0}(\Omega) = \{u \circ f : u \in \text{ssh}_{f(q_0)}(f(\Omega))\}.$$

Thanks to this equality, the statement immediately follows from Definition 7.1. \square

Lemma 7.5. Let Ω be a symmetric slice domain in \mathbb{H} , fix $q_0 \in \Omega$ and take a regular function $f : \Omega \rightarrow \mathbb{H}$. Then

$$G_{f(q_0)}^{f(\Omega)}(f(q)) \leq g_{q_0}^\Omega(q)$$

for all $q \in \Omega$. If, moreover, f is slice preserving then

$$g_{f(q_0)}^{f(\Omega)}(f(q)) \leq g_{q_0}^\Omega(q)$$

for all $q \in \Omega$. If, additionally, f admits a regular inverse $f^{-1} : f(\Omega) \rightarrow \Omega$, then the last inequality becomes an equality at all $q \in \Omega$.

Proof. By Theorem 5.3,

$$\text{wsh}_{q_0}(\Omega) \supseteq \{u \circ f : u \in \text{ssh}_{f(q_0)}(f(\Omega))\}.$$

If f is a slice preserving regular function then, by Theorem 4.2,

$$\text{wsh}_{q_0}(\Omega) \supseteq \{u \circ f : u \in \text{wsh}_{f(q_0)}(f(\Omega))\}.$$

The last inclusion is actually an equality if f admits a regular inverse $f^{-1} : f(\Omega) \rightarrow \Omega$, which is necessarily slice preserving. The three statements now follow from Definition 7.1. \square

In the last statement, we assumed Ω to be a symmetric slice domain for the sake of simplicity. The result could, however, be extended to all slice domains.

Lemmas 7.4 and 7.5, along with the preceding example, yield what follows.

Example 7.6. For each $x_0 \in \mathbb{R}$ and each $R > 0$, the equalities

$$\log \frac{|q - x_0|}{R} = G_{x_0}^{B(x_0, R)}(q) = g_{x_0}^{B(x_0, R)}(q)$$

hold for all $q \in B(x_0, R)$. We point out that this function is strongly subharmonic in $B(x_0, R)$ and weakly harmonic in $B(x_0, R) \setminus \{x_0\}$.

Example 7.7. For each $q_0 \in \mathbb{H}$ and each $R > 0$ it holds

$$\log \frac{|q - q_0|}{R} = G_{q_0}^{B(q_0, R)}(q) \leq g_{q_0}^{B(q_0, R)}(q)$$

for all $q \in B = B(q_0, R)$. We point out that $G_{q_0}^B$, though strongly and weakly subharmonic, is not weakly harmonic if $q_0 \notin \mathbb{R}$. Indeed, if we fix an orthonormal basis $1, I, J, IJ$ and write $q_0 = q_1 + q_2 J$ with $q_1, q_2 \in L_I$, then by Lemma 4.4

$$(\bar{\partial}_1 \partial_1 G_{q_0}^B)|_z = \frac{1}{R^2} (\bar{\partial}_1 \partial_1 G_0^{\mathbb{B}})|_{z - \frac{q_0}{R}} = \frac{|q_2|^2}{2(|z - q_1|^2 + |q_2|^2)^2}$$

for $z \in L_I$. This expression only vanishes when $q_2 = 0$, that is, when $q_0 \in L_I$.

We are now in a position to make the next remarks.

Remark 7.8. Let Ω be a bounded domain in \mathbb{H} and let $q_0 \in \Omega$. For all $r, R > 0$ such that $B(q_0, r) \subseteq \Omega \subseteq B(q_0, R)$, we have that

$$\log \frac{|q - q_0|}{R} \leq G_{q_0}^\Omega(q) \leq \log \frac{|q - q_0|}{r}.$$

As a consequence, $G_{q_0}^\Omega$ is not identically equal to $-\infty$, it belongs to $\text{ssh}_{q_0}(\Omega)$ and it coincides with the supremum s appearing in Definition 7.1.

Remark 7.9. Let Ω be a bounded domain in \mathbb{H} and let $q_0 \in \Omega$. For all $R > 0$ such that $\Omega \subseteq B(q_0, R)$, we have that

$$\log \frac{|q - q_0|}{R} \leq g_{q_0}^\Omega(q),$$

whence $g_{q_0}^\Omega$ is not identically equal to $-\infty$. If, moreover, $q_0 = x_0 \in \mathbb{R}$ then for all $r > 0$ such that $B(x_0, r) \subseteq \Omega$ we have that

$$g_{x_0}^\Omega(q) \leq \log \frac{|q - x_0|}{r}.$$

In this case, $g_{x_0}^\Omega$ belongs to $\text{wsh}_{x_0}(\Omega)$ and it coincides with the supremum w appearing in Definition 7.1.

When Ω admits a well-behaved exhaustion function, we can prove a few further properties.

Theorem 7.10. Let Ω be a bounded domain in \mathbb{H} . Suppose there exists $\rho \in C^0(\Omega, (-\infty, 0)) \cap \text{ssh}(\Omega)$ such that $\{q \in \Omega : \rho(q) < c\} \subset\subset \Omega$ for all $c < 0$. Then for all $q_0 \in \Omega$ and for all $p \in \partial\Omega$

$$\lim_{q \rightarrow p} G_{q_0}^\Omega(q) = 0.$$

Moreover, $G_{q_0}^\Omega$ is continuous in $\Omega \setminus \{q_0\}$.

Proof. Let $B(q_0, r) \subseteq \Omega \subseteq B(q_0, R)$ and let $C > 0$ be such that $C\rho < \log \frac{r}{R}$ in $\overline{B(q_0, r)}$. If we set

$$u(q) = \begin{cases} \log \frac{|q - q_0|}{R} & q \in \overline{B(q_0, r)} \\ \max \left\{ C\rho(q), \log \frac{|q - q_0|}{R} \right\} & q \in \Omega \setminus B(q_0, r) \end{cases}$$

then $u \in \text{ssh}_{q_0}(\Omega)$. Thus, $u \leq G_{q_0}^\Omega \leq 0$, where $\lim_{q \rightarrow p} u(q) = 0$ for all $p \in \partial\Omega$. This prove the first statement.

To prove the second statement, we only need to prove the lower semicontinuity of $G_{q_0}^\Omega$. Let us choose $\lambda \in (0, 1)$ such that $\rho < -\lambda$ in $\overline{B(q_0, \lambda)}$. For any $\varepsilon \in (0, \lambda)$ such that $\log \frac{\varepsilon}{R} > (1 - \varepsilon) \log \varepsilon^2$ (whence $\log \frac{\varepsilon}{R} > \varepsilon - \frac{1}{\varepsilon}$), let us set

$$\alpha(q) = (1 - \varepsilon) \log(\varepsilon|q - q_0|) - \varepsilon$$

on $\overline{B(q_0, \varepsilon)}$.

By Proposition 6.1, for each sufficiently small $\delta > 0$, the convolution $G_{q_0}^\Omega * \chi_\delta$ is an element of $C^\infty \cap \text{ssh}(\Omega_\delta)$. If we choose $\eta \in (0, \varepsilon)$ so that $(1 - \varepsilon) \log(\varepsilon\eta) > \log \frac{\eta}{R}$ then we may choose $\delta = \delta_\varepsilon > 0$ so that:

- $(1 - \varepsilon) \log(\varepsilon\eta) > G_{q_0}^\Omega * \chi_\delta$ in $\partial B(q_0, \eta)$,

- Ω_δ includes $\rho^{-1}([-\infty, -\varepsilon^3])$, and
- $G_{q_0}^\Omega * \chi_\delta < 0$ in $\rho^{-1}(-\varepsilon^3)$.

We may then set

$$\beta(q) := G_{q_0}^\Omega * \chi_{\delta_\varepsilon}(q) - \varepsilon$$

for all $q \in \Omega_{\delta_\varepsilon}$ and

$$\gamma(q) := \varepsilon^{-2} \rho(q)$$

for all $q \in \Omega$. By construction, α, β, γ can be patched together in a continuous strongly subharmonic function defined on Ω , namely

$$u_\varepsilon := \begin{cases} \alpha & \text{in } \overline{B(q_0, \eta)} \\ \max\{\alpha, \beta\} & \text{in } \overline{B(q_0, \varepsilon)} \setminus B(q_0, \eta) \\ \beta & \text{in } \rho^{-1}([-\infty, -\varepsilon]) \setminus B(q_0, \varepsilon) \\ \max\{\beta, \gamma\} & \text{in } \rho^{-1}([-\varepsilon, -\varepsilon^3]) \\ \gamma & \text{in } \Omega \setminus \rho^{-1}([-\infty, -\varepsilon^3]) \end{cases}$$

We remark that $\rho^{-1}([-\infty, -\varepsilon]) \setminus B(q_0, \varepsilon)$ increases as $\varepsilon \rightarrow 0$ and that

$$\bigcup_{\varepsilon \in (0, \lambda)} (\rho^{-1}([-\infty, -\varepsilon]) \setminus B(q_0, \varepsilon)) = \Omega \setminus \{q_0\}.$$

Thus, for all $q \in \Omega \setminus \{q_0\}$,

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(q) = \lim_{\varepsilon \rightarrow 0} G_{q_0}^\Omega * \chi_{\delta_\varepsilon}(q) - \varepsilon = G_{q_0}^\Omega(q).$$

Moreover, for each $\varepsilon \in (0, \lambda)$ it holds $\frac{u_\varepsilon}{1-\varepsilon} \in \text{ssh}_{q_0}(\Omega)$, whence $\frac{u_\varepsilon}{1-\varepsilon} \leq G_{q_0}^\Omega$ in Ω . Thus,

$$G_{q_0}^\Omega(q) = \sup_{\varepsilon \in (0, \lambda)} \frac{u_\varepsilon(q)}{1-\varepsilon},$$

whence the lower semicontinuity of $G_{q_0}^\Omega$ immediately follows. \square

Similarly:

Proposition 7.11. *Let Ω be a bounded domain in \mathbb{H} . Suppose there exists $\rho \in C^0(\Omega, (-\infty, 0)) \cap \text{wsh}(\Omega)$ such that $\{q \in \Omega : \rho(q) < c\} \subset\subset \Omega$ for all $c < 0$. Then for all $q_0 \in \Omega$ and for all $p \in \partial\Omega$*

$$\lim_{q \rightarrow p} g_{q_0}^\Omega(q) = 0.$$

Proof. Let $B(q_0, r) \subseteq \Omega \subseteq B(q_0, R)$ and let $C > 0$ be such that $C\rho < \log \frac{r}{R}$ in $\overline{B(q_0, r)}$. If we set

$$u(q) = \begin{cases} \log \frac{|q-q_0|}{R} & q \in \overline{B(q_0, r)} \\ \max\{C\rho(q), \log \frac{|q-q_0|}{R}\} & q \in \Omega \setminus B(q_0, r) \end{cases}$$

then $u \in \text{wsh}_{q_0}(\Omega)$. Thus, $u \leq g_{q_0}^\Omega \leq 0$, where $\lim_{q \rightarrow p} u(q) = 0$ for all $p \in \partial\Omega$. \square

For a special class of domains Ω and points q_0 , the Green function $g_{q_0}^\Omega$ can be easily determined, as follows.

Theorem 7.12. *Let Ω be a bounded symmetric slice domain and let $x_0 \in \Omega \cap \mathbb{R}$. Consider the slice $\Omega_i = \Omega \cap \mathbb{C}$ of the domain and the (complex) Green function of Ω_i with logarithmic pole at x_0 , which we will denote as $\gamma_{x_0}^{\Omega_i}$. If we set*

$$u(x + Iy) := \gamma_{x_0}^{\Omega_i}(x + iy)$$

for all $x, y \in \mathbb{R}$ and $I \in \mathbb{S}$ such that $x + Iy \in \Omega$, then:

- u is a well-defined function on Ω ;

- u is \mathbb{J} -plurisubharmonic in $\Omega \setminus \mathbb{R}$ and it belongs to $\text{wsh}_{x_0}(\Omega)$;
- u coincides with $g_{x_0}^\Omega$.

Proof. The slice Ω_i of the domain is a bounded domain in \mathbb{C} . Hence, the function $\gamma_{x_0}^{\Omega_i}$ is a negative plurisubharmonic function on Ω_i with a logarithmic pole at x_0 (see Proposition 8.10). Moreover, since Ω_i is symmetric with respect to the real axis, it holds $\gamma_{x_0}^{\Omega_i}(x + iy) = \gamma_{x_0}^{\Omega_i}(x - iy)$ for all $x + iy \in \Omega_i$. It follows at once that u is well-defined, that it belongs to $\text{wsh}_{x_0}(\Omega)$, and that it is \mathbb{J} -plurisubharmonic in $\Omega \setminus \mathbb{R}$.

Moreover, let us fix any other $v \in \text{wsh}_0(\mathbb{B})$: we can prove that $v(q) \leq u(q)$ for all $q \in \Omega$, as follows. For each $I \in \mathbb{S}$, the inequality $v_I \leq u_I$ holds in Ω_I because (up to identifying L_I with \mathbb{C}) the function u_I is the (complex) Green function of Ω_I with logarithmic pole at x_0 and v_I is a negative subharmonic function on Ω_I with a logarithmic pole at x_0 . As a consequence, u coincides with $g_{x_0}^\Omega$. \square

7.1 A significant example

An interesting example to consider is that of the unit ball \mathbb{B} with a pole other than 0. It is natural to address it by means of the *classical Möbius transformations* of \mathbb{B} , namely the conformal transformations $v^{-1}M_{q_0}u$, where u, v are constants in $\partial\mathbb{B}$, $q_0 \in \mathbb{B}$ and M_{q_0} is the transformation of \mathbb{B} defined as

$$M_{q_0}(q) := (1 - q\bar{q}_0)^{-1}(q - q_0).$$

The transformation M_{q_0} has inverse $M_{q_0}^{-1} = M_{-q_0}$. It is regular if, and only if, $q_0 = x_0 \in \mathbb{R}$; in this case, it is also slice preserving. For more details, see [2, 11].

Our first observation can be derived from either Lemma 7.5 or Theorem 7.12.

Example 7.13. *For each $x_0 \in \mathbb{B} \cap \mathbb{R}$, we have*

$$G_{x_0}^{\mathbb{B}}(q) \leq g_{x_0}^{\mathbb{B}}(q) = \log \frac{|q - x_0|}{|1 - qx_0|}$$

for all $q \in \mathbb{B}$.

The same techniques do not work when the logarithmic pole q_0 is not real, as a consequence of the fact that M_{q_0} is not a regular function. Nevertheless, we can make the following observation.

Example 7.14. *Let us fix $q_0 \in \mathbb{B} \setminus \mathbb{R}$. We will prove that*

$$\log \frac{|q - q_0|}{|1 - q\bar{q}_0|} \leq g_{q_0}^{\mathbb{B}}(q)$$

by showing that $u(q) = \log \frac{|q - q_0|}{|1 - q\bar{q}_0|}$ is weakly subharmonic. We will also prove that: (a) the restriction u_I to $\mathbb{B}_I \setminus \{q_0\}$ is harmonic if, and only if, $q_0 \in \mathbb{B}_I$; (b) the function u is not strongly subharmonic.

- We first observe that

$$u(q) = \log |q - q_0| - \log |q - \tilde{q}_0| - \log |\bar{q}_0|,$$

with $\tilde{q}_0 := \bar{q}_0^{-1} = q_0|q_0|^{-2}$. With respect to any orthonormal basis $1, I, J, IJ$ and to the associated coordinates $z_1, z_2, \bar{z}_1, \bar{z}_2$, if we split q_0, \tilde{q}_0 as $q_0 = q_1 + q_2J, \tilde{q}_0 = \tilde{q}_1 + \tilde{q}_2J$ with $q_1, q_2, \tilde{q}_1, \tilde{q}_2 \in \mathbb{B}_I$ then

$$H_{I,J}(u)|_q = \frac{1}{2|q - q_0|^4} \begin{pmatrix} |z_2 - q_2|^2 & -(z_1 - q_1)(\bar{z}_2 - \bar{q}_2) \\ -(z_2 - q_2)(\bar{z}_1 - \bar{q}_1) & |z_1 - q_1|^2 \end{pmatrix} + \\ - \frac{1}{2|q - \tilde{q}_0|^4} \begin{pmatrix} |z_2 - \tilde{q}_2|^2 & -(z_1 - q_1)(\bar{z}_2 - \bar{q}_2) \\ -(z_2 - \tilde{q}_2)(\bar{z}_1 - \bar{q}_1) & |z_1 - \tilde{q}_1|^2 \end{pmatrix}.$$

- For all $z \in \mathbb{B}_I$ (that is, $z_1 = z, z_2 = 0$) it holds

$$(\bar{\partial}_1 \partial_1 u)|_z = \frac{|q_2|^2}{2(|z - q_1|^2 + |q_2|^2)^2} - \frac{|q_2|^2|q_0|^4}{2(|z|q_0|^2 - |q_1|^2 + |q_2|^2)^2}.$$

If $q_0 \in \mathbb{B}_I$ then $q_2 = 0$ and $(\bar{\partial}_1 \partial_1 u)$ vanishes identically in \mathbb{B}_I . Otherwise, $(\bar{\partial}_1 \partial_1 u)|_z > 0$ for all $z \in \mathbb{B}_I$ because

$$\begin{aligned} & |z|q_0|^2 - q_1|^2 + |q_2|^2 - |q_0|^2(|z - q_1|^2 + |q_2|^2) \\ &= |z|^2|q_0|^4 + |q_0|^2 - |q_0|^2|z|^2 - |q_0|^4 \\ &= |q_0|^2(1 - |q_0|^2)(1 - |z|^2) > 0. \end{aligned}$$

- In general, $H_{I,J}(u)$ is not positive semidefinite. To see this, let us choose a basis $1, I, J, IJ$ so that $q_1 \neq 0 \neq q_2$ and let us choose $z_1 = 0, z_2 = q_2$. We get

$$\begin{aligned} H_{I,J}(u)|_{q_2 J} &= \frac{1}{2|q_1|^4} \begin{pmatrix} 0 & 0 \\ 0 & |q_1|^2 \end{pmatrix} + \\ &\quad - \frac{1}{2(|\tilde{q}_1|^2 + |q_2 - \tilde{q}_2|^2)^2} \begin{pmatrix} |q_2 - \tilde{q}_2|^2 & q_1(\bar{q}_2 - \tilde{q}_2) \\ (q_2 - \tilde{q}_2)\tilde{q}_1 & |\tilde{q}_1|^2 \end{pmatrix}, \end{aligned}$$

where $q_2 - \tilde{q}_2 \neq 0$ by construction.

We would now like to consider a different approach, through regular transformations of \mathbb{B} . Indeed, the work [11] proved the following facts.

- The only regular bijections $\mathbb{B} \rightarrow \mathbb{B}$ are the so-called *regular Möbius transformations* of \mathbb{B} , namely the transformations $\mathcal{M}_{q_0} * u = \mathcal{M}_{q_0} u$ with $u \in \partial\mathbb{B}, q_0 \in \mathbb{B}$ and

$$\mathcal{M}_{q_0}(q) := (1 - q\bar{q}_0)^{-*} * (q - q_0).$$

Here, the symbol $*$ denotes the multiplicative operation among regular functions and f^{-*} is the inverse of f with respect to this multiplicative operation.

- For all $q \in \mathbb{B}$, it holds

$$\mathcal{M}_{q_0}(q) = M_{q_0}(T_{q_0}(q)),$$

where $T_{q_0} : \mathbb{B} \rightarrow \mathbb{B}$ is defined as $T_{q_0}(q) = (1 - qq_0)^{-1}q(1 - qq_0)$ and has inverse $T_{q_0}^{-1}(q) = T_{\bar{q}_0}(q)$.

- \mathcal{M}_{q_0} is slice preserving if, and only if, $q_0 = x_0 \in \mathbb{R}$ (in which case, $T_{q_0} = id_{\mathbb{B}}$ and $\mathcal{M}_{x_0} = M_{x_0}$).

If we fix $q_0 \in \mathbb{B} \setminus \mathbb{R}$ then, by Lemma 7.5,

$$\log |\mathcal{M}_{q_0}(q)| \leq g_{q_0}^{\mathbb{B}}(q) \tag{14}$$

for all $q \in \mathbb{B}$. The work [3] proved the quaternionic Schwarz-Pick Lemma and, in particular, the inequality

$$\log |f(q)| \leq \log |\mathcal{M}_{q_0}(q)|$$

valid for all regular $f : \mathbb{B} \rightarrow \mathbb{B}$ with $f(q_0) = 0$. It is therefore natural to ask ourselves whether an equality may hold in (14). However, this is not the case: as a consequence of the next result, inequality (14) is strict at all q not belonging to the same slice \mathbb{B}_I as q_0 . As a byproduct, we conclude that the set

$$\{\log |f| : f : \mathbb{B} \rightarrow \mathbb{H} \text{ regular, } f(q_0) = 0\}$$

is not a dense subset of $\text{wsh}_{q_0}(\mathbb{B})$.

Theorem 7.15. *If $q_0 \in \mathbb{B}_I$, then*

$$|M_{q_0}(T_{q_0}(q))| = |\mathcal{M}_{q_0}(q)| \leq |M_{q_0}(q)| \tag{15}$$

for all $q \in \mathbb{B}$. Equality holds if, and only if, $q \in \mathbb{B}_I$.

Proof. Inequality (15) is equivalent to

$$|T_{q_0}(q) - q_0| |1 - q\bar{q}_0| \leq |q - q_0| |1 - T_{q_0}(q)\bar{q}_0|.$$

Since $|T_{q_0}(q)| = |q|$, the last inequality is equivalent to

$$\begin{aligned} 0 &\leq (|q|^2 - 2\langle q, q_0 \rangle + |q_0|^2) (1 - 2\langle T_{q_0}(q), q_0 \rangle + |q|^2|q_0|^2) + \\ &\quad - (|q|^2 - 2\langle T_{q_0}(q), q_0 \rangle + |q_0|^2) (1 - 2\langle q, q_0 \rangle + |q|^2|q_0|^2) \\ &= 2(-|q|^2 - |q_0|^2 + 1 + |q|^2|q_0|^2) \langle T_{q_0}(q), q_0 \rangle + \\ &\quad 2(-1 - |q|^2|q_0|^2 + |q|^2 + |q_0|^2) \langle q, q_0 \rangle \\ &= 2(1 - |q_0|^2)(1 - |q|^2) \langle T_{q_0}(q) - q, q_0 \rangle. \end{aligned}$$

Thus, inequality (15) holds for $q \in \mathbb{B}$ if, and only if, $0 \leq \langle T_{q_0}(q) - q, q_0 \rangle$. This is equivalent to the non-negativity of the real part of $(T_{q_0}(q) - q)\bar{q}_0 = ((1 - qq_0)^{-1}q(1 - qq_0) - q)\bar{q}_0$ or, equivalently, of the real part of

$$\begin{aligned} &\left(\overline{(1 - qq_0)}q(1 - qq_0) - |1 - qq_0|^2q \right) \bar{q}_0 \\ &= \overline{(1 - qq_0)}(q - q^2q_0 - q + qq_0q)\bar{q}_0 \\ &= (1 - \bar{q}_0\bar{q})(qq_0q\bar{q}_0 - q^2|q_0|^2) \\ &= qq_0q\bar{q}_0 - q^2|q_0|^2 + |q|^2|q_0|^2(\bar{q}_0q - q\bar{q}_0) \\ &= (|q_0|^2 - \bar{q}_0^2)(|z_2|^2 - z_1z_2J) + (\bar{q}_0 - q_0)z_2J, \end{aligned}$$

where the last equality can be obtained by direct computation after splitting q as $q = z_1 + z_2J$, with $z_1, z_2 \in \mathbb{B}_I$ and $J \perp I$. If $q_0 = x_0 + Iy_0$ then the real part of the last expression equals $(x_0^2 + y_0^2 - x_0^2 + y_0^2)|z_2|^2 = 2y_0^2|z_2|^2$, which is clearly non-negative. Moreover, it vanishes if, and only if, $z_2 = 0$, i.e., $q \in \mathbb{B}_I$. \square

An example wherein inequality (15) holds and is strict had been constructed in [4]. That construction was used to prove that regular Möbius transformations are not isometries for the Poincaré distance of \mathbb{B} , defined as

$$\delta_{\mathbb{B}}(q, q_0) := \frac{1}{2} \log \left(\frac{1 + |M_{q_0}(q)|}{1 - |M_{q_0}(q)|} \right)$$

for all $q, q_0 \in \mathbb{B}$. The subsequent work [1] proved that, for each $q_0 \in \mathbb{B} \setminus \mathbb{R}$, there exists no Riemannian metric on \mathbb{B} having \mathcal{M}_{q_0} as an isometry.

Our new inequality (15) is equivalent to

$$\delta_{\mathbb{B}}(T_{q_0}(q), q_0) \leq \delta_{\mathbb{B}}(q, q_0).$$

In other words, we have proven the following property of the transformation T_{q_0} of \mathbb{B} : while all points $q \in \mathbb{B}_I$ are fixed, all points $q \in \mathbb{B} \setminus \mathbb{B}_I$ are attracted to q_0 with respect to the Poincaré distance.

8 Appendix

For the reader's convenience, we include in this appendix some classical results and definitions, which are used in the present work. We begin with some theorems concerning subharmonic functions of m real variables, along with some instrumental definitions. Let λ denote the Lebesgue measure on \mathbb{R}^m and σ denote the surface area measure.

Theorem 8.1 ([9, Theorem 2.4.1 (iii)]). *Let D be an open subset of \mathbb{R}^m and let $v : D \rightarrow [-\infty, +\infty)$ be an upper semicontinuous function which is not identically $-\infty$ on any connected component of D . The function v is subharmonic in D if, and only if, for any Euclidean ball $B(a, R)$ such that $\overline{B(a, R)} \subset D$, it holds*

$$v(a) \leq \mathbf{L}(v; a, R),$$

where

$$\mathbf{L}(v; a, R) := \frac{1}{s_m R^{m-1}} \int_{\partial B(a, R)} u(x) d\sigma(x), \quad s_m := \sigma(\partial B(0, 1)).$$

Theorem 8.2 ([9, Theorem 2.4.2]). *Let D be a bounded connected open subset of \mathbb{R}^m and let $v : D \rightarrow [-\infty, +\infty)$ be subharmonic in D . Then either v is constant or, for each $x \in D$,*

$$v(x) < \sup_{z \in \partial D} \left\{ \limsup_{D \ni y \rightarrow z} v(y) \right\}.$$

Corollary 8.3 ([9, Corollary 2.4.3]). *u is harmonic if, and only if, u and $-u$ are both subharmonic.*

Definition 8.4 ([9, §2.5]). *Define $h : \mathbb{R} \rightarrow \mathbb{R}$ by the formula*

$$h(t) := \begin{cases} \exp(-1/t) & \text{if } t > 0 \\ 0 & \text{if } t \leq 0 \end{cases}$$

and define $\chi : \mathbb{R}^m \rightarrow \mathbb{R}$ by the formula

$$\chi(x) := \frac{1}{c} h(1 - \|x\|^2), \quad c := \int_{B(0,1)} h(1 - \|x\|^2) d\lambda(x).$$

The standard smoothing kernels $\chi_\varepsilon : \mathbb{R}^m \rightarrow \mathbb{R}$ are defined, for all $\varepsilon > 0$, by the formula

$$\chi_\varepsilon(x) := \frac{1}{\varepsilon^m} \chi\left(\frac{x}{\varepsilon}\right).$$

Given a function v on a open subset D of \mathbb{R}^m , the convolution

$$v * \chi_\varepsilon(x) = \chi_\varepsilon * v(x) := \int_{\mathbb{R}^m} \chi_\varepsilon(x - y) v(y) d\lambda(y)$$

is well-defined on

$$D_\varepsilon := \begin{cases} \{x \in D : \text{dist}(x, \partial D) > \varepsilon\} & \text{if } D \neq \mathbb{R}^m \\ \mathbb{C}^n & \text{if } D = \mathbb{R}^m \end{cases}$$

Theorem 8.5 ([9, Theorem 2.5.5]). *Let D be an open subset of \mathbb{R}^m and let $v : D \rightarrow [-\infty, +\infty)$ be subharmonic. For all $\varepsilon > 0$ such that D_ε is not empty, $v * \chi_\varepsilon$ is C^∞ and subharmonic in D_ε . Moreover, $v * \chi_\varepsilon$ monotonically decreases with decreasing ε and*

$$\lim_{\varepsilon \rightarrow 0^+} v * \chi_\varepsilon(x) = v(x) \tag{16}$$

for each $x \in D$.

We now recall some properties of plurisubharmonic functions of n complex variables.

Theorem 8.6 ([9, Theorem 2.9.1]). *Let D be an open subset of \mathbb{C}^n and let $v : D \rightarrow [-\infty, +\infty)$ be an upper semicontinuous function which is not identically $-\infty$ on any connected component of D . v is plurisubharmonic in D if, and only if, for any $a \in D, b \in \mathbb{C}^n$ such that $\{a + \lambda b : \lambda \in \mathbb{C}, |\lambda| \leq 1\} \subset D$, it holds*

$$v(a) \leq l(v; a, b),$$

where

$$l(v; a, b) := \frac{1}{2\pi} \int_0^{2\pi} v(a + e^{it}b) dt.$$

Moreover, plurisubharmonicity is a local property.

Theorem 8.7 ([9, Theorem 2.9.2]). *Let D be an open subset of \mathbb{C}^n and let $v : D \rightarrow [-\infty, +\infty)$ be plurisubharmonic. For all $\varepsilon > 0$ such that D_ε is not empty, $v * \chi_\varepsilon$ is C^∞ and plurisubharmonic in D_ε . Moreover, $v * \chi_\varepsilon$ monotonically decreases with decreasing ε and*

$$\lim_{\varepsilon \rightarrow 0^+} v * \chi_\varepsilon(z) = v(z) \tag{17}$$

for each $z \in D$.

Theorem 8.8 ([9, Corollary 2.9.9]). *Let D be a bounded connected open subset of \mathbb{C}^n and let v be a plurisubharmonic function on D . Then either v is constant or, for each $x \in D$,*

$$v(z) < \sup_{w \in \partial D} \left\{ \limsup_{D \ni y \rightarrow w} v(y) \right\}.$$

Remark 8.9 ([9, §3.1]). *Let D be an open subset of \mathbb{C} and let v be a plurisubharmonic function on D . The function v is harmonic if, and only if, it is maximal among plurisubharmonic functions on D .*

Proposition 8.10 ([9, Proposition 6.1.1 (iv)]). *Let D be a bounded domain in \mathbb{C}^n , let $w \in D$ and let γ_w^D denote the pluricomplex Green function of D with pole at w . Then γ_w^D is a negative plurisubharmonic function with a logarithmic pole at w .*

References

- [1] N. Arcozzi and G. Sarfatti. Invariant metrics for the quaternionic Hardy space. *J. Geom. Anal.*, 25(3):2028–2059, 2015.
- [2] C. Bisi and G. Gentili. Möbius transformations and the Poincaré distance in the quaternionic setting. *Indiana Univ. Math. J.*, 58(6):2729–2764, 2009.
- [3] C. Bisi and C. Stoppato. The Schwarz-Pick lemma for slice regular functions. *Indiana Univ. Math. J.*, 61(1):297–317, 2012.
- [4] C. Bisi and C. Stoppato. Regular vs. classical Möbius transformations of the quaternionic unit ball. In G. Gentili, I. Sabadini, M. V. Shapiro, F. Sommen, and D. C. Struppa, editors, *Advances in Hypercomplex Analysis*, Springer INdAM Series. Springer, Milan, 2013.
- [5] G. Gentili, S. Salamon, and C. Stoppato. Twistor transforms of quaternionic functions and orthogonal complex structures. *J. Eur. Math. Soc. (JEMS)*, 16(11):2323–2353, 2014.
- [6] G. Gentili, C. Stoppato, and D. C. Struppa. *Regular functions of a quaternionic variable*. Springer Monographs in Mathematics. Springer, Heidelberg, 2013.
- [7] G. Gentili and D. C. Struppa. A new theory of regular functions of a quaternionic variable. *Adv. Math.*, 216(1):279–301, 2007.
- [8] R. Ghiloni and A. Perotti. Slice regular functions on real alternative algebras. *Adv. Math.*, 226(2):1662–1691, 2011.
- [9] M. Klimek. *Pluripotential theory*, volume 6 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 1991. Oxford Science Publications.
- [10] A. Perotti. Slice regularity and harmonicity on Clifford algebras. arXiv:1801.03045 [math.CV], 2018.
- [11] C. Stoppato. Regular Moebius transformations of the space of quaternions. *Ann. Global Anal. Geom.*, 39(4):387–401, 2011.
- [12] C. Stoppato. A new series expansion for slice regular functions. *Adv. Math.*, 231(3-4):1401–1416, 2012.