



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

A degree associated to linear eigenvalue problems in Hilbert spaces and applications to nonlinear spectral theory

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

A degree associated to linear eigenvalue problems in Hilbert spaces and applications to nonlinear spectral theory / PIERLUIGI BENEVIERI, ALESSANDRO CALAMAI, MASSIMO FURI, MARIA PATRIZIA PERA. - In: JOURNAL OF DYNAMICS AND DIFFERENTIAL EQUATIONS. - ISSN 1040-7294. - STAMPA. - 34:(2022), pp. 555-581. [10.1007/s10884-020-09921-9]

Availability:

This version is available at: 2158/1204859 since: 2024-04-08T10:18:43Z

Published version:

DOI: 10.1007/s10884-020-09921-9

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

(Article begins on next page)

A DEGREE ASSOCIATED TO LINEAR EIGENVALUE PROBLEMS IN HILBERT SPACES AND APPLICATIONS TO NONLINEAR SPECTRAL THEORY

PIERLUIGI BENEVIERI, ALESSANDRO CALAMAI, MASSIMO FURI,
AND MARIA PATRIZIA PERA

*Dedicated to the memory of our friend and
outstanding mathematician Russell Johnson*

ABSTRACT. We extend to the infinite dimensional context the link between two completely different topics recently highlighted by the authors: the classical eigenvalue problem for real square matrices and the Brouwer degree for maps between oriented finite dimensional real manifolds. Thanks to this extension, we solve a conjecture regarding global continuation in nonlinear spectral theory that we have formulated in a recent article. Our result (the ex conjecture) is applied to prove a Rabinowitz type global continuation property of the solutions to a perturbed motion equation containing an air resistance frictional force.

1. INTRODUCTION

Given a linear operator $L: \mathbb{R}^k \rightarrow \mathbb{R}^k$, consider the Classical Eigenvalue Problem

$$(1.1) \quad \begin{cases} Lx = \lambda x, \\ x \in S, \end{cases}$$

where S is the unit sphere of \mathbb{R}^k and $\lambda \in \mathbb{R}$. The solutions of (1.1) are pairs $(\lambda, x) \in \mathbb{R} \times S$, hereafter called *eigenpoints*, where λ is a real eigenvalue of L and x is one of the corresponding unit eigenvectors. Since the eigenpoints of (1.1) are the zeros of the C^∞ -map

$$\Psi: \mathbb{R} \times S \rightarrow \mathbb{R}^k, \quad (\lambda, x) \mapsto Lx - \lambda x,$$

in [8] we have shown that there is a link between the above purely algebraic problem and the Brouwer degree, $\deg(\Psi, U, 0)$, of Ψ with target $0 \in \mathbb{R}^k$ on convenient open subsets U of the cylinder $\mathbb{R} \times S$, which is a smooth k -dimensional real manifold with a natural orientation.

Roughly speaking, in [8] we have shown that

Date: 18th January 2021.

1991 *Mathematics Subject Classification.* 47J10, 47A75, 47H11, 55M25, 34C23.

Key words and phrases. eigenvalues, eigenvectors, nonlinear spectral theory, topological degree, bifurcation, differential equations.

The first, second and fourth authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

A. Calamai is partially supported by GNAMPA - INdAM (Italy).

- if $\lambda_* \in \mathbb{R}$ is a simple eigenvalue of L , and x_* and $-x_*$ are the two corresponding unit eigenvectors, then the “twin” eigenpoints $p_* = (\lambda_*, x_*)$ and $\bar{p}_* = (\lambda_*, -x_*)$ are isolated zeros of Ψ and give the same contribution to the Brouwer degree, which is either 1 or -1 , depending on the sign jump at λ_* of the (real) characteristic polynomial of L .

Still roughly speaking, here we extend this fact to the infinite dimensional case by considering a problem of the type

$$(1.2) \quad \begin{cases} Lx = \lambda Cx, \\ x \in S, \end{cases}$$

in which $\lambda \in \mathbb{R}$, L and C are bounded linear operators acting between two real Hilbert spaces G and H , C is compact, $L - \lambda C$ is invertible for some $\lambda \in \mathbb{R}$, and S is the unit sphere of the source space G .

Even in the special case $G = H$ and L the identity, Problem (1.2) cannot be treated using the degree of Leray and Schauder which, unlike that of Brouwer, does not hold in the context of smooth manifolds. Therefore, we apply the degree introduced in [9] for oriented C^1 Fredholm maps of index zero between real differentiable Banach manifolds, which extends the Brouwer degree for maps between oriented finite dimensional smooth manifolds, as well as the Leray-Schauder degree for C^1 compact vector fields in Banach spaces. To apply this degree we need the unit sphere S to be a smooth manifold. This is the reason of our restriction to Hilbert spaces instead of the more general Banach environment.

Here the degree regards the smooth map

$$\Psi: \mathbb{R} \times S \rightarrow H, \quad (\lambda, x) \mapsto Lx - \lambda Cx,$$

acting between the 1-codimensional submanifold $\mathbb{R} \times S$ of the Hilbert space $\mathbb{R} \times G$ and the target space H , whose zeros are called the *eigenpoints* of (1.2). The result obtained here, Theorem 3.6 below, extends, to the infinite dimensional case, the one in [8] concerning the Classical Eigenvalue Problem (1.1), provided that one calls $\lambda_* \in \mathbb{R}$ a *simple eigenvalue* of (1.2) if there exists $x_* \in S$ such that $\text{Ker}(L - \lambda_* C) = \mathbb{R}x_*$ and $H = \text{Im}(L - \lambda_* C) \oplus \mathbb{R}Cx_*$. In fact, we obtain that

- if λ_* is a simple eigenvalue of (1.2) and x_* and $-x_*$ are the two corresponding unit eigenvectors, then the “twin” eigenpoints $p_* = (\lambda_*, x_*)$ and $\bar{p}_* = (\lambda_*, -x_*)$ are isolated zeros of Ψ and give the same contribution to the degree, which is either 1 or -1 , depending on the orientation of Ψ .

As in [8], this crucial result regarding the “fair contribution to the degree of the twin eigenpoints” is applied to the study of the behaviour of the solution triples (s, λ, x) of the following perturbation of (1.2):

$$(1.3) \quad \begin{cases} Lx + sN(x) = \lambda Cx, \\ x \in S, \end{cases}$$

in which $N: S \rightarrow H$ is a compact C^1 -map. Precisely, if we denote by Σ the subset of $\mathbb{R} \times \mathbb{R} \times S$ of the solutions (s, λ, x) of (1.3) and we call *trivial* those having $s = 0$, our main result, Theorem 4.5 below, yields the following Rabinowitz type global continuation result that was conjectured in [7].

- If $q_* = (0, \lambda_*, x_*)$ is a trivial solution of (1.3) corresponding to a simple eigenvalue λ_* of the unperturbed problem (1.2), then the connected component of Σ containing q_* is either unbounded or encounters a trivial solution $q^* = (0, \lambda^*, x^*)$ with $\lambda^* \neq \lambda_*$.

The result obtained in [7] differs from this one in the final assertion and the additional assumption that G and H are separable. In fact, in [7], without any degree theory and with arguments of differential topology, we proved that

- the connected component of Σ containing $q_* = (0, \lambda^*, x^*)$ is either unbounded or encounters a trivial solution $q^* = (0, \lambda_*, x_*)$ different from q_* .

Notice that this assertion does not exclude the case $\lambda^* = \lambda_*$ and, consequently, $x^* = -x_*$, since λ_* is simple.

We believe that our global continuation result, Theorem 4.5, could be extended to the more general situation in which $\text{Ker}(L - \lambda_* C)$ is odd dimensional and

$$H = \text{Img}(L - \lambda_* C) \oplus C(\text{Ker}(L - \lambda_* C)).$$

Our belief is supported by the fact that in [5], under these assumptions, we have shown that

- the projection of Σ into the $s\lambda$ -plane has a connected set which contains $(0, \lambda_*)$ and is either unbounded or includes a “trivial eigenpair” $(0, \lambda^*)$ different from $(0, \lambda_*)$.

Moreover, our belief is also based on the fact that in [8], by means of the Brouwer degree for maps between oriented smooth manifolds, we obtained that

- if $G = H = \mathbb{R}^k$, C is the identity, and x_* is a unit eigenvector of L corresponding to an eigenvalue λ_* with odd algebraic multiplicity, then the connected component of Σ containing $(0, \lambda_*, x_*)$ is either unbounded or includes a trivial solution $(0, \lambda^*, x^*)$ with $\lambda^* \neq \lambda_*$.

So far, our effort to obtain the supposed extension of Theorem 4.5 has proved unsuccessful.

Theorem 4.5 falls into the subject of *nonlinear spectral theory*, which finds applications to differential equations (see e.g. [3, 16] and references therein).

Here we mention the work of R. Chiappinelli [14], which inspired some of our recent articles. In [14] a sort of “local persistence property” for a perturbed eigenvalue problem similar to (1.3) was proved. Precisely, under the assumptions

- $L: G \rightarrow G$ is a *self-adjoint* operator,
- $C = I$ is the identity of G ,
- $N: S \rightarrow G$ is Lipschitz continuous,
- $\lambda_* \in \mathbb{R}$ is an isolated simple eigenvalue of L ,
- $x_* \in S$ is an eigenvector corresponding to λ_* ,

it was shown that

- in a neighborhood V of $0 \in \mathbb{R}$ a Lipschitz continuous function $\varepsilon \mapsto (\lambda_\varepsilon, x_\varepsilon) \in \mathbb{R} \times S$ is defined with the properties that $(\lambda_0, x_0) = (\lambda_*, x_*)$ and that $Lx_\varepsilon + \varepsilon N(x_\varepsilon) = \lambda_\varepsilon x_\varepsilon$ for any $\varepsilon \in V$.

When G is infinite dimensional, the hypotheses of our global continuation result seem incompatible with the assumptions of Chiappinelli, due to the fact that

the identity is not a compact operator. However, Theorem 4.5 does apply to the equation

$$Lx + \varepsilon N(x) = \lambda x,$$

provided that N is C^1 and compact, and L is of the type $\lambda_* I + C$, with $\lambda_* \in \mathbb{R}$ and C compact. In fact, putting $\varepsilon = -\sigma/\mu$ and $\lambda = \lambda_* + 1/\mu$, the above equation becomes $x + \sigma N(x) = \mu Cx$, which is as in our problem (1.3) with $L = I$.

For results regarding the local persistence property when the eigenvalue λ_* is not necessarily simple we mention [4–8, 15, 17–21].

The last section of the paper contains three examples illustrating our main result, as well as an application to the study of the solutions (s, λ, x) of the boundary value problem

$$(1.4) \quad \begin{cases} x''(t) + sg(x'(t)) + \lambda x(t) = 0, \\ x(0) = 0 = x(\pi), \quad x \in S, \end{cases}$$

in which S is the unit sphere of the Hilbert space $H^2(0, \pi)$, and $g: \mathbb{R} \rightarrow \mathbb{R}$ is an odd increasing C^1 -function (such as the air resistance force $g(v) = v|v|$). From our result, with the help of the well-known notion of winding number of a self-map of the circle S^1 , we deduce that, given any trivial solution q_* of (1.4), the connected component of Σ containing q_* is unbounded and does not encounter other trivial solutions.

For pioneering articles regarding the use of the winding number in order to study the behavior of solutions to ordinary differential equations we mention [13, 22, 23].

2. NOTATION AND PRELIMINARIES

We introduce some notation, preliminaries, and known or unknown concepts that we will need in subsequent sections. In particular, we will outline the main notions related to the topological degree for oriented C^1 Fredholm maps of index zero between real differentiable Banach manifolds introduced in [9] (see also [10, 11] for additional details). Actually, some notions and results are new: we consider them necessary for a better understanding of the topics in Sections 3 and 4, the proof of Theorem 3.6 in particular.

2.1. Algebraic preliminaries. Let, hereafter, E and F be two real vector spaces. By $\mathcal{L}(E, F)$ we shall denote the vector space of the linear operators from E into F . The same notation will be used if, in addition, E and F are normed. In this case, however, we will tacitly assume that all the operators of $\mathcal{L}(E, F)$ are bounded, and that this space is endowed with the usual operator norm. If $E = F$, we will write $\mathcal{L}(E)$ instead of $\mathcal{L}(E, E)$. By $\text{Iso}(E, F)$ we shall mean the subset of $\mathcal{L}(E, F)$ of the invertible operators, and we will write $\text{GL}(E)$ instead of $\text{Iso}(E, E)$. The subspace of $\mathcal{L}(E, F)$ of the operators having finite dimensional image will be denoted by $\mathcal{F}(E, F)$, or simply by $\mathcal{F}(E)$ when $E = F$.

Let $I \in \mathcal{L}(E)$ indicate the identity of E . If $T \in \mathcal{L}(E)$ has the property that $I - T \in \mathcal{F}(E)$, we shall say that T is an *admissible operator (for the determinant)*. The symbol $\mathcal{A}(E)$ will stand for the affine subspace of $\mathcal{L}(E)$ of the admissible operators.

It is known (see [25]) that the determinant of an operator $T \in \mathcal{A}(E)$ is well defined as follows: $\det T := \det T|_{\hat{E}}$, where $T|_{\hat{E}}$ is the restriction (as domain and as

codomain) to any finite dimensional subspace \hat{E} of E containing $\text{Img}(I - T)$, with the understanding that $\det T|_{\hat{E}} = 1$ if $\hat{E} = \{0\}$.

As one can easily check, the function $\det: \mathcal{A}(E) \rightarrow \mathbb{R}$ inherits most of the properties of the classical determinant. Some of them are stated in the following

Remark 2.1. *Let $T, T_1, T_2 \in \mathcal{A}(E)$. Then*

- $\det T \neq 0$ if and only if T is invertible;
- $R \in \text{Iso}(E, F)$ implies $RT R^{-1} \in \mathcal{A}(F)$ and $\det(RT R^{-1}) = \det T$;
- $T_2 T_1 \in \mathcal{A}(E)$ and $\det(T_2 T_1) = \det(T_2) \det(T_1)$.

See, for example, [12] for a discussion about other properties of the determinant.

We will need the following remark, whose easy proof is left to the reader:

Remark 2.2. *Let $T \in \mathcal{L}(E)$ and let $E = E_1 \oplus E_2$ with $\dim E_2 < +\infty$. Assume that, with respect to the above decomposition, T can be represented in a block matrix form*

$$T = \begin{pmatrix} I_{11} & T_{12} \\ 0 & T_{22} \end{pmatrix},$$

where I_{11} is the identity of E_1 . Then $T \in \mathcal{A}(E)$ and $\det T = \det T_{22}$.

Recall that an operator $T \in \mathcal{L}(E, F)$ is said to be (*algebraic*) *Fredholm* if its kernel, $\text{Ker } T$, and its cokernel, $\text{coKer } T = F/T(E)$, are both finite dimensional.

The *index* of a Fredholm operator T is the integer

$$\text{ind } T = \dim(\text{Ker } T) - \dim(\text{coKer } T).$$

In particular, any invertible linear operator is Fredholm of index zero. Observe also that, if $T \in \mathcal{L}(\mathbb{R}^k, \mathbb{R}^s)$, then $\text{ind } T = k - s$.

The subset of $\mathcal{L}(E, F)$ of the Fredholm operators will be denoted by $\Phi(E, F)$; while $\Phi_n(E, F)$ will stand for the set $\{T \in \Phi(E, F) : \text{ind } T = n\}$. Obviously, $\Phi(E)$ and $\Phi_n(E)$ designate, respectively, $\Phi(E, E)$ and $\Phi_n(E, E)$.

One can easily check that $\mathcal{A}(E)$ is a subset of $\Phi_0(E)$. This is also a consequence of a well known property regarding Fredholm operators. Namely,

- (1) *if $T \in \Phi_n(E, F)$ and $K \in \mathcal{F}(E, F)$, then $T + K \in \Phi_n(E, F)$.*

Another fundamental property states that

- (2) *the composition of Fredholm operators is Fredholm and its index is the sum of the indices of all the composite operators.*

An useful consequence of property (2) is the following:

- If $T \in \Phi_n(E, F)$ and $k \in \mathbb{N}$, then the restriction of T to a k -codimensional subspace of E is Fredholm of index $n - k$.

Let $T \in \Phi_0(E, F)$. In [9], an operator $K \in \mathcal{F}(E, F)$ was called a *corrector* of T if $T + K$ is invertible. Since, during a conference, someone has critically observed that it is not necessary to correct an invertible operator, hereafter we will use the more appropriate word *companion* instead of *corrector*.

Notice that any $T \in \text{Iso}(E, F)$ has a *natural companion*: the trivial element of $\mathcal{L}(E, F)$. This fact was crucial in [9] for the construction of the degree theory that we will apply here.

Given $T \in \Phi_0(E, F)$, let us denote by $\mathcal{C}(T)$ the subset of $\mathcal{F}(E, F)$ of all the companions of T . As one can easily check, this set is nonempty. Moreover, $\mathcal{C}(T)$ admits a partition in two equivalence classes according to the following

Definition 2.3 (Equivalence relation). Two companions K_1 and K_2 of an operator $T \in \Phi_0(E, F)$ are *equivalent* (more precisely, *T-equivalent*) if the admissible operator $(T + K_2)^{-1}(T + K_1)$ has positive determinant.

Given two companions K_1 and K_2 of $T \in \Phi_0(E, F)$, the admissible automorphism $(T + K_2)^{-1}(T + K_1)$ is not the unique one that can be used to check whether or not K_1 and K_2 are equivalent. In fact, one has the following

Remark 2.4. Let $T \in \Phi_0(E, F)$ and $K_1, K_2 \in \mathcal{C}(T)$. Then, the determinants of the invertible operators

$$(T + K_2)^{-1}(T + K_1), (T + K_1)(T + K_2)^{-1}, (T + K_1)^{-1}(T + K_2), (T + K_2)(T + K_1)^{-1}$$

have the same sign. In fact, from the second property of Remark 2.1 one gets that the first two operators have the same determinant, while the third property implies the statement regarding the last two operators, being the inverses of the first two.

Thanks to the above equivalence relation, the following definition was introduced in [9].

Definition 2.5 (Orientation). An *orientation* of $T \in \Phi_0(E, F)$ is one of the two equivalence classes of $\mathcal{C}(T)$, denoted by $\mathcal{C}_+(T)$ and called the class of *positive companions* of the *oriented operator* T . The set $\mathcal{C}_-(T) = \mathcal{C}(T) \setminus \mathcal{C}_+(T)$ of the *negative companions* is the *opposite orientation* of T .

Some further definitions are in order.

Definition 2.6 (Natural orientation). Any $T \in \text{Iso}(E, F)$ admits the *natural orientation*: the one given by considering the trivial operator of $\mathcal{L}(E, F)$ as a positive companion.

Definition 2.7 (Canonical orientation). Any admissible operator $T \in \mathcal{A}(E)$ admits the *canonical orientation*: the one given by choosing as a positive companion any $K \in \mathcal{F}(E)$ such that $\det(T + K) > 0$. In particular, this applies for any $T \in \mathcal{L}(E)$, with $\dim E < \infty$.

Definition 2.8 (Associated orientation). Let E and F have the same finite dimension. Assume that they are oriented up to an inversion of both the orientations or, equivalently, assume that $E \times F$ has an orientation, say \mathcal{O} . Then any $T \in \mathcal{L}(E, F)$ admits the orientation *associated with* \mathcal{O} : the one given by choosing as a positive companion any $K \in \mathcal{F}(E, F)$ such that $T + K$ is orientation preserving.

Definition 2.9 (Oriented composition). The *oriented composition* of two oriented operators, $T_1 \in \Phi_0(E_1, E_2)$ and $T_2 \in \Phi_0(E_2, E_3)$, is the operator $T_2 T_1$ with the orientation given by considering $K = (T_2 + K_2)(T_1 + K_1) - T_2 T_1$ as a positive companion, where K_1 and K_2 are positive companions of T_1 and T_2 , respectively.

Observe that the oriented composition is associative. Indeed, if $T_1 \in \Phi_0(E_1, E_2)$, $T_2 \in \Phi_0(E_2, E_3)$ and $T_3 \in \Phi_0(E_3, E_4)$, and K_1, K_2 and K_3 are, respectively, companions of T_1, T_2 and T_3 , one has

$$\begin{aligned} & ((T_3 + K_3)(T_2 + K_2))(T_1 + K_1) - (T_3 T_2) T_1 \\ &= (T_3 + K_3)((T_2 + K_2)(T_1 + K_1)) - T_3(T_2 T_1). \end{aligned}$$

The following result implies an important property of the oriented composition (see Corollary 2.13 below). Moreover, it shows that Definition 2.9 is well-posed.

Lemma 2.10. *Given $T_1 \in \Phi_0(E_1, E_2)$, $T_2 \in \Phi_0(E_2, E_3)$, $K_1, K'_1 \in \mathcal{C}(T_1)$ and $K_2, K'_2 \in \mathcal{C}(T_2)$, consider the following companions of T_2T_1 :*

$$K = (T_2 + K_2)(T_1 + K_1) - T_2T_1 \quad \text{and} \quad K' = (T_2 + K'_2)(T_1 + K'_1) - T_2T_1.$$

Then K is equivalent to K' if and only if K_1 and K_2 are both equivalent or both not equivalent to K'_1 and K'_2 , respectively.

Proof. According to Definition 2.5, we need to compute the sign of

$$\det((T_2T_1 + K)^{-1}(T_2T_1 + K')).$$

We have

$$(T_2T_1 + K)^{-1}(T_2T_1 + K') = ((T_2 + K_2)(T_1 + K_1))^{-1}((T_2 + K'_2)(T_1 + K'_1)).$$

Thus, because of the second property in Remark 2.1, we obtain

$$\begin{aligned} & \det((T_2T_1 + K)^{-1}(T_2T_1 + K')) \\ &= \det\left((T_1 + K'_1)((T_2 + K_2)(T_1 + K_1))^{-1}((T_2 + K'_2)(T_1 + K'_1))(T_1 + K'_1)^{-1}\right) \\ &= \det((T_1 + K'_1)(T_1 + K_1)^{-1}(T_2 + K_2)^{-1}(T_2 + K'_2)). \end{aligned}$$

Therefore, applying the third property of the same remark, we get

$$\begin{aligned} & \det((T_2T_1 + K)^{-1}(T_2T_1 + K')) \\ &= \det((T_1 + K'_1)(T_1 + K_1)^{-1}) \det((T_2 + K_2)^{-1}(T_2 + K'_2)), \end{aligned}$$

and the assertion follows. \square

Definition 2.11 (Sign of an oriented operator). Let $T \in \Phi_0(E, F)$ be an oriented operator. Its *sign* is the integer

$$\text{sign } T = \begin{cases} +1 & \text{if } T \text{ is invertible and naturally oriented,} \\ -1 & \text{if } T \text{ is invertible and not naturally oriented,} \\ 0 & \text{if } T \text{ is not invertible.} \end{cases}$$

As a straightforward consequence of Remark 2.4, and taking into account Definitions 2.3, 2.5, 2.6, 2.11, we get the following

Remark 2.12. *Let $T \in \text{Iso}(E, F)$ be oriented. Then,*

$$\begin{aligned} \text{sign } T &= \text{sign } \det((T + K)^{-1}T) = \text{sign } \det(T(T + K)^{-1}) \\ &= \text{sign } \det(T^{-1}(T + K)) = \text{sign } \det((T + K)T^{-1}), \end{aligned}$$

where K is a positive companion of T .

Lemma 2.10 shows that, in the oriented composition, the inversion of the orientation of one (and only one) of the operators yields the inversion of the orientation of the composition. Hence, one gets the following

Corollary 2.13. *Let $T_1 \in \Phi_0(E_1, E_2)$ and $T_2 \in \Phi_0(E_2, E_3)$ be oriented. Then, $\text{sign}(T_2T_1) = \text{sign } T_2 \text{sign } T_1$, where T_2T_1 is the oriented composition of T_1 and T_2 .*

Proof. If one of the two operators is not invertible, then the assertion is obvious. Assume, therefore, that T_1 and T_2 are isomorphisms. If both the operators are naturally oriented, then the assertion follows from the definition of oriented composition. The other cases are a consequence of Lemma 2.10. \square

Given $R_1 \in \text{Iso}(E_1, F_1)$ and $R_2 \in \text{Iso}(E_2, F_2)$, observe that the function

$$\Lambda: \mathcal{L}(E_1, E_2) \rightarrow \mathcal{L}(F_1, F_2), \quad T \mapsto R_2 T R_1^{-1}$$

is a linear isomorphism (whose inverse is given by $\tilde{T} \mapsto R_2^{-1} \tilde{T} R_1$). One can see that Λ establishes a correspondence between certain pairs of subsets of $\mathcal{L}(E_1, E_2)$ and $\mathcal{L}(F_1, F_2)$. For example, $\text{Iso}(E_1, E_2)$ and $\text{Iso}(F_1, F_2)$, $\mathcal{F}(E_1, E_2)$ and $\mathcal{F}(F_1, F_2)$, $\Phi_0(E_1, E_2)$ and $\Phi_0(F_1, F_2)$. Moreover if, in particular, $T \in \Phi_0(E_1, E_2)$, then Λ sends the set $\mathcal{C}(T)$ onto the set $\mathcal{C}(\Lambda(T))$, and if $K_1, K_2 \in \mathcal{C}(T)$ are equivalent (according to Definition 2.3), so are $\Lambda(K_1), \Lambda(K_2) \in \mathcal{C}(\Lambda(T))$.

Since the oriented composition is associative, this notion can be extended to the composition of three (or more) oriented operators.

2.2. Topological preliminaries. Let, hereafter, X denote a metric space. We recall that a subset A of X is *locally compact* if any point of A admits a neighborhood, in A , which is compact. Therefore, any compact subset of X is locally compact, as is any relatively open subset of a locally compact set. However, the union of two locally compact subsets of X may not be locally compact.

We recall also that a continuous map between metric spaces is said to be *proper* if the inverse image of any compact set is a compact set, while it is called *locally proper* if its restriction to a convenient closed neighborhood of any point of its domain is proper. Thus, level sets of locally proper maps are locally compact.

One can check that proper maps are *closed*, in the sense that the image of any closed set is a closed set.

Notation 2.14. Let D be a subset of the product $X \times Y$ of two metric spaces. Given $x \in X$, we call *x-slice of D* the set $D_x = \{y \in Y : (x, y) \in D\}$.

Assume, from now on, that the vector spaces E and F are actually Banach. In this framework, any Fredholm operator is assumed to be bounded. Therefore, in addition to the algebraic properties (1) and (2) stated in Subsection 2.1, one has the following topological ones (see e.g. [29]):

- (3) if $T \in \Phi(E, F)$, then $\text{Img } E$ is closed in F ;
- (4) if $T \in \Phi(E, F)$, then T is proper on any bounded closed subsets of E ;
- (5) for any $n \in \mathbb{Z}$, the set $\Phi_n(E, F)$ is open in $\mathcal{L}(E, F)$;
- (6) if $T \in \Phi_n(E, F)$ and $K \in \mathcal{L}(E, F)$ is compact, then $T + K \in \Phi_n(E, F)$.

Let us now sketch the construction and summarize the main properties of the degree introduced in [9].

The basic fact is that, in the context of Banach spaces, the orientation of an operator $T_* \in \Phi_0(E, F)$ induces an orientation to the operators in a neighborhood of T_* . Indeed, due to the fact that $\text{Iso}(E, F)$ and $\Phi_0(E, F)$ are open in $\mathcal{L}(E, F)$, any companion of T_* remains a companion of all T sufficiently close to T_* .

Therefore, the following definition makes sense.

Definition 2.15. Let $\Gamma: X \rightarrow \Phi_0(E, F)$ be a continuous map defined on a metric space X . A *pre-orientation* of Γ is a function that to any $x \in X$ assigns an orientation $\omega(x)$ of $\Gamma(x)$. A pre-orientation (of Γ) is an *orientation* if it is *continuous*, in the sense that, given any $x_* \in X$, there exist $K \in \omega(x_*)$ and a neighborhood W of x_* such that $K \in \omega(x)$ for all $x \in W$. The map Γ is said to be *orientable* if it admits an orientation, and *oriented* if an orientation has been chosen. In particular, a subset Y of $\Phi_0(E, F)$ is orientable or oriented if so is the inclusion map $Y \hookrightarrow \Phi_0(E, F)$.

Observe that the set $\hat{\Phi}_0(E, F)$ of the oriented operators of $\Phi_0(E, F)$ has a natural topology, and the natural projection $\pi: \hat{\Phi}_0(E, F) \rightarrow \Phi_0(E, F)$ is a 2-fold covering space (see [10] for details). Therefore, an orientation of a map Γ as in Definition 2.15 could be regarded as a lifting $\hat{\Gamma}$ of Γ . This implies that, if the domain X of Γ is simply connected and locally path connected, then Γ is orientable.

Let $f: U \rightarrow F$ be a C^1 -map defined on an open subset of E , and denote by $df_x \in \mathcal{L}(E, F)$ the Fréchet differential of f at a point $x \in U$.

We recall that f is said to be *Fredholm of index n* , from now on written $f \in \Phi_n$, if $df_x \in \Phi_n(E, F)$ for all $x \in U$. Therefore, if $f \in \Phi_0$, Definition 2.15 and the continuity of the differential map $df: U \rightarrow \Phi_0(E, F)$ suggest the following

Definition 2.16 (Orientation of a Φ_0 -map in Banach spaces). Let U be an open subset of E and $f: U \rightarrow F$ a Fredholm map of index zero. A *pre-orientation* or an *orientation* of f are, respectively, a pre-orientation or an orientation of df , according to Definition 2.15. The map f is said to be *orientable* if it admits an orientation, and *oriented* if an orientation has been chosen.

Remark 2.17. A very special Φ_0 -map is given by an operator $T \in \Phi_0(E, F)$. Thus, for T there are two different notions of orientations: the algebraic one, according to Definition 2.5; and the one regarding T as a C^1 -map (according to Definition 2.16). In each case T admits exactly two orientations (in the second one this is due to the connectedness of the domain E). Hereafter, we shall tacitly assume that the two notions agree. Namely, T has an algebraic orientation ω if and only if its differential $dT_x: \dot{x} \mapsto T\dot{x}$ has the ω orientation for all $x \in E$.

We will show how the notion of orientation in Definition 2.16 can be extended to the case of maps acting between real Banach manifolds. To this purpose, we need some further notation.

For short, by a *manifold* we shall mean a smooth Banach manifold embedded in a real Banach space.

Given a manifold \mathcal{M} and a point $x \in \mathcal{M}$, the tangent space of \mathcal{M} at x will be denoted by $T_x\mathcal{M}$. If \mathcal{M} is embedded in a Banach space \tilde{E} , $T_x\mathcal{M}$ will be identified with a closed subspace of \tilde{E} , for example by regarding any tangent vector of $T_x\mathcal{M}$ as the derivative $\gamma'(0)$ of a smooth curve $\gamma: (-1, 1) \rightarrow \mathcal{M}$ such that $\gamma(0) = x$.

Assume that $f: \mathcal{M} \rightarrow \mathcal{N}$ is a C^1 -map between two manifolds, respectively embedded in \tilde{E} and \tilde{F} and modelled on E and F . As in the flat case, f is said to be *Fredholm of index n* (written $f \in \Phi_n$) if so is the differential $df_x: T_x\mathcal{M} \rightarrow T_{f(x)}\mathcal{N}$, for any $x \in \mathcal{M}$.

Given $f \in \Phi_0$, suppose that to any $x \in \mathcal{M}$ it is assigned an orientation $\omega(x)$ of df_x (also called *orientation of f at x*). As above, the function ω is called a *pre-orientation* of f , and an *orientation* if it is continuous, in a sense to be specified (see Definition 2.19).

Definition 2.18. The pre-oriented composition of two (or more) pre-oriented maps between manifolds is given by assigning, at any point x of the domain of the composite map, the composition of the orientations (according to Definition 2.9) of the differentials in the chain representing the differential at x of the composite map.

Assume that $f: \mathcal{M} \rightarrow \mathcal{N}$ is a C^1 -diffeomorphism. Then, in particular, given any $x \in \mathcal{M}$, the differential df_x is an isomorphism. Thus, for any $x \in \mathcal{M}$, we may take as $\omega(x)$ the natural orientation of df_x (recall Definition 2.6). This pre-orientation of f turns out to be continuous according to Definition 2.19 below (it is, in some sense, constant). From now on, unless otherwise stated, **any diffeomorphism will be considered oriented with the natural orientation**. In particular, in a composition of pre-oriented maps, all charts and parametrizations of a manifold will be tacitly assumed to be naturally oriented.

Definition 2.19 (Orientation of a Φ_0 -map between manifolds). Let $f: \mathcal{M} \rightarrow \mathcal{N}$ be a Φ_0 -map between two manifolds modelled on E and F , respectively. A pre-orientation of f is an *orientation* if it is *continuous* in the sense that, given any two charts, $\varphi: U \rightarrow E$ of \mathcal{M} and $\psi: V \rightarrow F$ of \mathcal{N} , such that $f(U) \subseteq V$, the pre-oriented composition

$$\psi \circ f \circ \varphi^{-1}: U \rightarrow V$$

is an oriented map according to Definition 2.16.

The map f is said to be *orientable* if it admits an orientation, and *oriented* if an orientation has been chosen.

Perhaps, the simplest example of non-orientable Φ_0 -map is given by a constant map from the 2-dimensional projective space into \mathbb{R}^2 (see [10]).

Remark 2.20. *One can check that the pre-oriented composition of orientations is an orientation.*

Remark 2.21. *Regarding the attribution that we will assign to some particular orientations of Φ_0 -maps between manifolds, whenever it makes sense, we will adapt the terminology for Φ_0 -operators, such as “natural orientation”, “associated orientation”, “canonical orientation”.*

For example any local diffeomorphism $f: \mathcal{M} \rightarrow \mathcal{N}$ admits the *natural orientation*, given by assigning the natural orientation to the operator df_x , for any $x \in \mathcal{M}$ (see Definition 2.6). As another example, assume the manifolds \mathcal{M} and \mathcal{N} have the same finite dimension and are oriented, then any C^1 -map between them admits the *associated orientation* (see Definition 2.8). A third example is given by a C^1 -map $f: \mathbb{R}^k \rightarrow \mathbb{R}^k$: it can be given the *canonical orientation* (see Definition 2.7).

The concept of canonical orientation of a C^1 -map $f: \mathbb{R}^k \rightarrow \mathbb{R}^k$ can be extended to a more general situation that we shall need in the next section. In fact, if E is a real Banach space, in spite of the fact that the function $\det: \mathcal{A}(E) \rightarrow \mathbb{R}$ can be discontinuous (see e.g. [12]), one has the following

Remark 2.22. Let X be a metric space and $E = E_1 \times E_2$ a real Banach space, with $\dim E_2 < +\infty$. Assume that $\Gamma: X \rightarrow \mathcal{A}(E)$ is a continuous map that can be represented in a block matrix form as follows:

$$\Gamma = \begin{pmatrix} I_{11} & \Gamma_{12} \\ 0 & \Gamma_{22} \end{pmatrix},$$

where I_{11} is the identity of E_1 , $\Gamma_{12}: X \rightarrow \mathcal{L}(E_2, E_1)$, and $\Gamma_{22}: X \rightarrow \mathcal{L}(E_2)$. Then, according to Remark 2.2, one has $\det \Gamma(x) = \det \Gamma_{22}(x)$, for all $x \in X$. Moreover, the pre-orientation of Γ given by assigning, to any $x \in X$, the canonical orientation of the operator $\Gamma(x)$ is actually an orientation, and has the property that $\text{sign } \Gamma(x) = \text{sign } \det \Gamma(x)$ for all $x \in X$.

Similarly to the case of a single map, one can define a notion of orientation of a continuous family of Φ_0 -maps depending on a parameter $s \in [0, 1]$. To be precise, one has the following

Definition 2.23 (Oriented Φ_0 -homotopy). A Φ_0 -homotopy between two Banach manifolds \mathcal{M} and \mathcal{N} is a C^1 -map $h: [0, 1] \times \mathcal{M} \rightarrow \mathcal{N}$ such that, for any $s \in [0, 1]$, the partial map $h_s = h(s, \cdot)$ is Fredholm of index zero. An orientation of h is a continuous function ω that to any $(s, x) \in [0, 1] \times \mathcal{M}$ assigns an orientation $\omega(s, x)$ to the differential $d(h_s)_x \in \Phi_0(T_x \mathcal{M}, T_{h(s,x)} \mathcal{N})$. Where “continuous” means that, given any chart $\varphi: U \rightarrow E$ of \mathcal{M} , a subinterval J of $[0, 1]$, and a chart $\psi: V \rightarrow F$ of \mathcal{N} such that $h(J \times U) \subseteq V$, the pre-orientation of the map $\Gamma: J \times U \rightarrow \Phi_0(E, F)$ that to any $(s, x) \in J \times U$ assigns the pre-oriented composition

$$d(\psi \circ h_s \circ \varphi^{-1})_x = d\psi_{h(s,x)} d(h_s)_x (d\varphi_x)^{-1}$$

is an orientation, according to Definition 2.15.

The homotopy h is said to be *orientable* if it admits an orientation, and *oriented* if an orientation has been chosen.

If a Φ_0 -homotopy h has an orientation ω , then any partial map $h_s = h(s, \cdot)$ has a *compatible* orientation $\omega(s, \cdot)$. Conversely, one has the following

Proposition 2.24 ([9, 10]). Let $h: [0, 1] \times \mathcal{M} \rightarrow \mathcal{N}$ be a Φ_0 -homotopy, and assume that one of its partial maps, say h_s , has an orientation. Then, there exists and is unique an orientation of h which is compatible with that of h_s . In particular, if two maps from \mathcal{M} to \mathcal{N} are Φ_0 -homotopic, then they are both orientable or both non-orientable.

As a consequence of Proposition 2.24, one gets that any C^1 -map $f: \mathcal{M} \rightarrow \mathcal{M}$ which is Φ_0 -homotopic to the identity is orientable, since so is the identity (even when \mathcal{M} is finite dimensional and not orientable).

The degree for oriented Φ_0 -maps defined in [9] satisfies the three fundamental properties stated below and called *Normalization*, *Additivity* and *Homotopy Invariance*. By an axiomatic approach similar to the one due to Amann-Weiss in [2] for the Leray-Schauder degree, in [11] it is shown that the degree constructed in [9] is the only possible integer valued function that satisfies these three properties.

To be more explicit, let us define, first, the domain of this degree function. Given an oriented Φ_0 -map $f: \mathcal{M} \rightarrow \mathcal{N}$, an open (possibly empty) subset U of \mathcal{M} , and a target value $y \in \mathcal{N}$, the triple (f, U, y) is said to be *admissible* for the degree

provided that $U \cap f^{-1}(y)$ is compact. From the axiomatic point of view, the degree is an integer valued function, \deg , defined on the class of all the admissible triples, that satisfies the following three *fundamental properties*:

- (Normalization) *If $f: \mathcal{M} \rightarrow \mathcal{N}$ is a naturally oriented diffeomorphism onto an open subset of \mathcal{N} , then*

$$\deg(f, \mathcal{M}, y) = 1, \quad \forall y \in f(\mathcal{M}).$$

- (Additivity) *Let (f, U, y) be an admissible triple. If U_1 and U_2 are two disjoint open subsets of U such that $U \cap f^{-1}(y) \subseteq U_1 \cup U_2$, then*

$$\deg(f, U, y) = \deg(f|_{U_1}, U_1, y) + \deg(f|_{U_2}, U_2, y).$$

- (Homotopy Invariance) *Let $h: [0, 1] \times \mathcal{M} \rightarrow \mathcal{N}$ be an oriented Φ_0 -homotopy, and $\gamma: [0, 1] \rightarrow \mathcal{N}$ a continuous path. If the set*

$$\{(s, x) \in [0, 1] \times \mathcal{M} : h(s, x) = \gamma(s)\}$$

is compact, then $\deg(h(s, \cdot), \mathcal{M}, \gamma(s))$ does not depend on $s \in [0, 1]$.

Other properties can be deduced from the fundamental ones (see [11] for details). We mention only some of them. One of these is the

- (Localization) *If (f, U, y) is an admissible triple, then*

$$\deg(f, U, y) = \deg(f|_U, U, y).$$

Another one is the

- (Excision) *If (f, U, y) is admissible and V is an open subset of U such that $f^{-1}(y) \cap U \subseteq V$, then*

$$\deg(f, U, y) = \deg(f, V, y).$$

A significant one is the

- (Existence) *If (f, U, y) is admissible and $\deg(f, U, y) \neq 0$, then the equation $f(x) = y$ admits at least one solution in U .*

Roughly speaking, given an admissible triple (f, U, y) , the integer $\deg(f, U, y)$ is an algebraic count of the solutions in U of the equation $f(x) = y$. More precisely, as a consequence of the fundamental properties, one gets the following

- (Computation Formula) *If (f, U, y) is admissible and y is a regular value for f in U , then the set $U \cap f^{-1}(y)$ is finite and*

$$\deg(f, U, y) = \sum_{x \in U \cap f^{-1}(y)} \text{sign}(df_x).$$

Another property that can be deduced from the fundamental ones is a generalization of the Homotopy Invariance Property, that we will need in Section 4. This requires the following extension of the concept of Φ_0 -homotopy:

Definition 2.25 (Extended Φ_0 -homotopy). An *extended Φ_0 -homotopy* from \mathcal{M} to \mathcal{N} is a C^1 -map $h: \mathcal{I} \times \mathcal{M} \rightarrow \mathcal{N}$, where \mathcal{I} is an arbitrary (nontrivial) real interval, such that any partial map $h_s = h(s, \cdot)$ of h is a Φ_0 -map.

The notion of orientation for an extended Φ_0 -homotopy is practically identical to the one in Definition 2.23 and its formulation is left to the reader.

As a consequence of the Excision and the Homotopy Invariance properties of the degree we get the following

- (Generalized Homotopy Invariance) *Let $h: \mathcal{I} \times \mathcal{M} \rightarrow \mathcal{N}$ be an oriented extended Φ_0 -homotopy, $\gamma: \mathcal{I} \rightarrow \mathcal{N}$ a continuous path, and W an open subset of $\mathcal{I} \times \mathcal{M}$. Given any $s \in \mathcal{I}$, denote by $W_s = \{x \in \mathcal{M} : (s, x) \in W\}$ the s -slice of W . If the set $\{(s, x) \in W : h(s, x) = \gamma(s)\}$ is compact, then $\deg(h_s, W_s, \gamma(s))$ does not depend on $s \in \mathcal{I}$.*

The easy proof of this property can be performed by showing that the integer valued function $s \in \mathcal{I} \mapsto \nu(s) := \deg(h_s, W_s, \gamma(s))$ is locally constant. In fact, given any $s_* \in \mathcal{I}$, because of the compactness of the set

$$\{(s, x) \in W : h(s, x) = \gamma(s)\},$$

one can find a box $J \times V \subseteq W$, with V open in \mathcal{M} and J an open interval containing s_* , such that $W_s \cap h_s^{-1}(\gamma(s)) \subseteq V$ for all $s \in J \cap \mathcal{I}$. Thus, from the Excision Property, one gets $\nu(s) = \deg(h_s, V, \gamma(s))$ for all $s \in J \cap \mathcal{I}$. Moreover, because of the Homotopy Invariance Property, $\nu(s)$ does not depend on $s \in J \cap \mathcal{I}$. Hence, since \mathcal{I} is connected and $s_* \in \mathcal{I}$ is arbitrary, one gets the assertion.

3. THE EIGENVALUE PROBLEM AND THE ASSOCIATED TOPOLOGICAL DEGREE

Hereafter, G and H will be two real Hilbert spaces, with inner product and norm denoted by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively.

Consider the eigenvalue problem

$$(3.1) \quad \begin{cases} Lx = \lambda Cx \\ x \in S, \end{cases}$$

where $\lambda \in \mathbb{R}$, $L, C \in \mathcal{L}(G, H)$, C is compact, and S is the unit sphere of G . We assume that the operator $L - \lambda C \in \mathcal{L}(G, H)$ is invertible for some $\lambda \in \mathbb{R}$. Therefore, because of the compactness of C , $L - \lambda C$ is Fredholm of index zero for any $\lambda \in \mathbb{R}$ (recall property (6) of Fredholm operators in Subsection 2.2). In particular, $\text{Ker}(L - \lambda C)$ is always finite dimensional, and nontrivial if and only if $\lambda \in \mathbb{R}$ is an eigenvalue of (3.1). Moreover, the set of all the real eigenvalues of (3.1) is discrete.

A pair (λ, x) belonging to the cylinder $\mathbb{R} \times S$ will be called an *eigenpoint* of (3.1) if it satisfies the equation $Lx = \lambda Cx$. In this case, x is a *unit eigenvector* of (3.1) corresponding to the eigenvalue λ .

The set of the eigenpoints of (3.1) will be denoted by \mathcal{S} . Hence, given any $\lambda \in \mathbb{R}$, the λ -slice $\mathcal{S}_\lambda = \{x \in S : (\lambda, x) \in \mathcal{S}\}$ of \mathcal{S} coincides with $S \cap \text{Ker}(L - \lambda C)$.

Observe that \mathcal{S}_λ is nonempty if and only if λ is an eigenvalue of (3.1). In this case \mathcal{S}_λ will be called the *eigensphere* of (3.1) corresponding to λ . In fact, it is a sphere whose (finite) dimension equals that of $\text{Ker}(L - \lambda C)$ minus one.

If $\lambda \in \mathbb{R}$ is an eigenvalue of (3.1), the nonempty subset $\{\lambda\} \times \mathcal{S}_\lambda$ of \mathcal{S} will be called the λ -*eigenet* of (3.1).

It is convenient to regard $\mathbb{R} \times S$ as the subset of the Hilbert space $\mathbb{R} \times G$ satisfying the equation $g(\lambda, x) = 1$, where $g: \mathbb{R} \times G \rightarrow \mathbb{R}$ is defined by $g(\lambda, x) = \langle x, x \rangle$. The

differential $dg_p \in \mathcal{L}(\mathbb{R} \times G, \mathbb{R})$ of g at a point $p = (\lambda, x)$ is given by $(\dot{\lambda}, \dot{x}) \mapsto 2\langle x, \dot{x} \rangle$. Therefore, the set of the critical points of g is the λ -axis $x = 0$ and, consequently, the number 1 is a regular value for g . This shows that $\mathbb{R} \times S$ is a smooth manifold of codimension one in $\mathbb{R} \times G$ and, given any $p = (\lambda, x) \in g^{-1}(1)$, the tangent space of $\mathbb{R} \times S$ at p is the kernel of dg_p , namely

$$T_{(\lambda, x)}(\mathbb{R} \times S) = \{(\dot{\lambda}, \dot{x}) \in \mathbb{R} \times G : \langle x, \dot{x} \rangle = 0\} = \mathbb{R} \times x^\perp.$$

Observe that, if $\dim G = 1$ and $(\lambda, x) \in \mathbb{R} \times S$, then $x^\perp = \{0\}$ and the tangent space $T_{(\lambda, x)}(\mathbb{R} \times S)$ is the subspace $\mathbb{R} \times \{0\}$ of $\mathbb{R} \times \mathbb{R}$. Moreover, the cylinder $\mathbb{R} \times S$ is disconnected: it is the union of two horizontal lines, $\mathbb{R} \times \{-1\}$ and $\mathbb{R} \times \{1\}$. Due to this fact, in order to write some statements in a simpler form, **hereafter, unless otherwise stated, we will assume that the dimensions of the Hilbert spaces G and H are bigger than 1**, so that the cylinder $\mathbb{R} \times S$ is connected.

Define the smooth map

$$\bar{\Psi}: \mathbb{R} \times G \rightarrow H \quad \text{by} \quad (\lambda, x) \mapsto Lx - \lambda Cx$$

and observe that it is Fredholm of index one. Therefore, its restriction

$$\Psi: \mathbb{R} \times S \rightarrow H$$

to the 1-codimensional submanifold $\mathbb{R} \times S$ of $\mathbb{R} \times G$ is Fredholm of index zero. To see this, recall that the differential of Ψ at $p \in \mathbb{R} \times S$ is the restriction of $d\bar{\Psi}_p$ to the 1-codimensional subspace $T_p(\mathbb{R} \times S)$ of $\mathbb{R} \times G$.

The map Ψ will play a fundamental role in this paper. Notice that its zeros are the eigenpoints of (3.1). That is, $S = \Psi^{-1}(0)$.

We point out that Ψ is orientable and, because of the connectedness of the manifold $\mathbb{R} \times S$, admits exactly two orientations. In fact, in the finite dimensional case, an orientation of Ψ is equivalent to a pair of orientations, one of the domain and one of the codomain, up to an inversion of both of them (see Definition 2.8); while, if $\dim G = +\infty$, the orientability of Ψ is a consequence of the fact that the cylinder $\mathbb{R} \times S$ is simply connected (it is actually contractible). Therefore, **from now on, we shall assume that Ψ is oriented**. No matter which one of the two orientations one selects, all the statements in this paper hold true.

Definition 3.1. Let X be a metric space and $\mathcal{K} \subseteq \mathcal{A} \subseteq X$. We shall say that \mathcal{K} is an *isolated subset* of \mathcal{A} if it is compact and relatively open in \mathcal{A} . Thus, there exists an open subset U of X such that $U \cap \mathcal{A} = \mathcal{K}$, which will be called an *isolating neighborhood* of \mathcal{K} among (the elements of) \mathcal{A} .

Definition 3.2. Let $\mathcal{K} \subset \mathbb{R} \times S$ be an isolated subset of $\Psi^{-1}(0)$. By the Ψ -degree of \mathcal{K} we mean the integer $\Psi\text{-deg}(\mathcal{K}) := \deg(\Psi, U, 0)$, where $U \subseteq \mathbb{R} \times S$ is any isolating neighborhood of \mathcal{K} among $\Psi^{-1}(0)$. If p is an isolated zero of Ψ , we shall simply write $\Psi\text{-deg}(p)$ instead of $\Psi\text{-deg}(\{p\})$.

Notice that this definition is well-posed, thanks to the Excision Property of the degree.

Remark 3.3. If $p \in \Psi^{-1}(0)$ is such that the differential $d\Psi_p$ is an isomorphism, then, as a consequence of the Local Inverse Function Theorem, Ψ maps diffeomorphically a neighborhood of p in $\mathbb{R} \times S$ onto a neighborhood of 0 in H . Thus, $\{p\}$ is isolated among $\Psi^{-1}(0)$ and, because of the Computation Formula of the degree (see

Section 2), $\Psi\text{-deg}(p) = \text{sign}(d\Psi_p)$, which is either 1 or -1 , depending on whether or not the orientation of $d\Psi_p$ is the natural one.

Definition 3.4. An eigenpoint (λ_*, x_*) of (3.1) will be called *simple* if the associated Φ_0 -operator $T = L - \lambda_* C$ satisfies the following conditions:

- (1) $\text{Ker } T = \mathbb{R}x_*$,
- (2) $Cx_* \notin \text{Img } T$.

Notice that, if $p_* = (\lambda_*, x_*)$ is a simple eigenpoint, then the eigenset $\{\lambda_*\} \times \mathcal{S}_{\lambda_*}$ has only two elements: p_* and its *twin eigenpoint* $\bar{p}_* = (\lambda_*, -x_*)$, which is as well simple. Moreover, since $T = L - \lambda_* C$ is Fredholm of index zero, its image has codimension one in H . Therefore one has the following

Remark 3.5. If (λ_*, x_*) is a simple eigenpoint of (3.1), then $H = \text{Img } T \oplus \mathbb{R}Cx_*$. Thus, λ_* is a simple eigenvalue of the equation $Lx = \lambda Cx$.

The following result is the key to the proof of Theorem 4.5. Despite the fact that we have assumed $\dim G > 1$, Theorem 4.5 holds true in any dimension: its assertion in the 1-dimensional case will be verified in Example 5.1.

Theorem 3.6. Let $p_* = (\lambda_*, x_*)$ and $\bar{p}_* = (\lambda_*, -x_*)$ be two simple twin eigenpoints of (3.1). Then, $\Psi\text{-deg}(p) = \Psi\text{-deg}(\bar{p}) = \pm 1$. Consequently, the Ψ -degree of the λ_* -eigenset $\{\lambda_*\} \times \mathcal{S}_{\lambda_*}$ is non-zero.

Proof. It is enough to prove that $\Psi\text{-deg}(p) = \Psi\text{-deg}(\bar{p}) = \pm 1$: the last assertion follows from the Additivity Property of the degree.

Since p_* is simple, the λ_* -eigensphere \mathcal{S}_{λ_*} of (3.1) consists of two antipodal points: x_* and $-x_*$. Both the tangent spaces of S at these points coincide with the 1-codimensional subspace x_*^\perp of G . Thus, the tangent spaces of the cylinder $\mathbb{R} \times S$ at the twin eigenpoints p_* and \bar{p}_* are equal to the 1-codimensional subspace $\mathbb{R} \times x_*^\perp$ of the Hilbert space $\mathbb{R} \times G$. The differentials $d\Psi_{p_*}$ and $d\Psi_{\bar{p}_*}$ (acting from $\mathbb{R} \times x_*^\perp$ to H) are given, respectively, by

$$(\dot{\lambda}, \dot{x}) \mapsto T\dot{x} - \dot{\lambda}Cx_* \quad \text{and} \quad (\dot{\lambda}, \dot{x}) \mapsto T\dot{x} + \dot{\lambda}Cx_*,$$

where T , as in Definition 3.4, denotes the operator $L - \lambda_* C$.

As one can check (see, for example, [7, Lemma 3.2]), the fact that the eigenpoints p_* and \bar{p}_* are simple implies that the differentials $d\Psi_{p_*}$ and $d\Psi_{\bar{p}_*}$ are invertible. Consequently, according to Remark 3.3, the Ψ -degrees of p_* and \bar{p}_* coincide, respectively, with $\text{sign}(d\Psi_{p_*})$ and $\text{sign}(d\Psi_{\bar{p}_*})$, which are both non-zero.

Thus, it remains to prove that these two signs are equal, which means that the orientations of Ψ at these points are both natural or both not natural.

For this purpose, it is convenient to fix an orientation of Ψ at one of the two eigenpoints p_* and \bar{p}_* (for example by choosing the natural orientation of $d\Psi_{\bar{p}_*}$) and to transport it, continuously, along a curve, up to the other eigenpoint.

A suitable curve is a λ_* -meridian. That is, a geodesic in $\mathbb{R} \times S$ of the type

$$\mathcal{G} = \{(\lambda_*, x) \in \mathbb{R} \times S : x = \sin \theta x_* + \cos \theta x_e, \theta \in [-\pi/2, \pi/2]\},$$

where x_e is an element of the *equator* $S \cap x_*^\perp$ of S (recall that $\dim G > 1$) and θ may be regarded as a *latitude*.

Having chosen x_e , and the consequent λ_* -meridian, we will “observe” the differential of Ψ , along \mathcal{G} , from the point of view of a self-map defined on a convenient “flat space”; namely, the Hilbert space $G_* \times \mathbb{R} \times \mathbb{R}$, where G_* is the 2-codimensional

subspace $x_*^\perp \cap x_e^\perp$ of G , which is nonempty because of the assumption $\dim G > 1$, even if trivial when $\dim G = 2$.

We will “observe” the map Ψ by means of a convenient composition $\tilde{\Psi} = \sigma^{-1} \circ \Psi \circ \eta$, where $\eta: W \rightarrow \mathbb{R} \times S$ is a parametrization (i.e. the inverse of a chart) defined on an open subset W of $G_* \times \mathbb{R} \times \mathbb{R}$ and $\sigma: G_* \times \mathbb{R} \times \mathbb{R} \rightarrow H$ is an invertible bounded linear operator (a global parametrization of H).

Let us define η , first. Consider the open subset

$$W = \{(y, \theta, \lambda) \in B \times (-\pi, \pi) \times \mathbb{R}\}$$

of $G_* \times \mathbb{R} \times \mathbb{R}$, where B stands for the open unit ball of G_* , and let $\eta: W \rightarrow \mathbb{R} \times S$ be the map given by

$$\eta(y, \theta, \lambda) = (\lambda, y + \sqrt{1 - \|y\|^2}(\sin \theta x_* + \cos \theta x_e)).$$

Notice that, under η , the eigenpoints \bar{p}_* and p_* correspond to $\bar{u}_* = (0, -\pi/2, \lambda_*)$ and $u_* = (0, \pi/2, \lambda_*)$, respectively. One can check that η is a diffeomorphism onto an open subset of $\mathbb{R} \times S$ containing the meridian \mathcal{G} .

We now define σ . From Remark 3.5 we get the splitting

$$(3.2) \quad H = T(x_*^\perp) \oplus \mathbb{R}Cx_* = (T(G_*) \oplus \mathbb{R}Tx_e) \oplus \mathbb{R}Cx_*.$$

Thus, H can be identified with $G_* \times \mathbb{R} \times \mathbb{R}$ by means of the isomorphism

$$\sigma: G_* \times \mathbb{R} \times \mathbb{R} \rightarrow H, \quad (y, a, b) \mapsto Ty + aTx_e + bCx_*.$$

We assume that η and σ^{-1} are naturally oriented and that $\tilde{\Psi}$ has the composite orientation. Therefore, recalling Corollary 2.13,

$$(3.3) \quad \text{sign}(d\tilde{\Psi}_{\bar{u}_*}) = \text{sign}(d\Psi_{\bar{p}_*}) \quad \text{and} \quad \text{sign}(d\tilde{\Psi}_{u_*}) = \text{sign}(d\Psi_{p_*}),$$

whatever the orientation of Ψ .

Hence, it remains to prove that $\text{sign}(d\tilde{\Psi}_{\bar{u}_*}) = \text{sign}(d\tilde{\Psi}_{u_*})$, no matter what is the orientation of $\tilde{\Psi}$.

To this purpose, consider the straight path $\gamma: [-\pi/2, \pi/2] \rightarrow W$ defined by $\gamma(\theta) = (0, \theta, \lambda_*)$. This path joins $\gamma(-\pi/2) = \bar{u}_*$ with $\gamma(\pi/2) = u_*$, therefore it is suitable for the continuous transport of the orientation of $\tilde{\Psi}$ from \bar{u}_* to u_* . Notice that the image of the simple arc $\theta \mapsto \eta(\gamma(\theta))$ is the λ_* -meridian \mathcal{G} .

Taking into account that $\Psi(\lambda, x) = Tx - (\lambda - \lambda_*)Cx$ and that $Tx_* = 0$, given any $\theta \in [-\pi/2, \pi/2]$, we get

$$d(\Psi \circ \eta)_{\gamma(\theta)}(\dot{y}, \dot{\theta}, \dot{\lambda}) = T\dot{y} - \dot{\theta} \sin \theta Tx_e - \dot{\lambda} \sin \theta Cx_* - \dot{\lambda} \cos \theta Cx_e.$$

Therefore, recalling that σ^{-1} , being linear, coincides with its differential, we obtain

$$d(\tilde{\Psi})_{\gamma(\theta)}(\dot{y}, \dot{\theta}, \dot{\lambda}) = \sigma^{-1}(T\dot{y} - \dot{\theta} \sin \theta Tx_e - \dot{\lambda} \sin \theta Cx_* - \dot{\lambda} \cos \theta Cx_e).$$

Since, according to the splitting (3.2), Cx_e can be written as $Ty_* + \alpha Tx_e + \beta Cx_*$ for some $y_* \in G_*$ and $\alpha, \beta \in \mathbb{R}$, we have

$$d(\tilde{\Psi})_{\gamma(\theta)}(\dot{y}, \dot{\theta}, \dot{\lambda}) = (\dot{y}, -\dot{\theta} \sin \theta, -\dot{\lambda} \sin \theta) - \dot{\lambda} \cos \theta (y_*, \alpha, \beta),$$

which can be represented as

$$d\tilde{\Psi}_{\gamma(\theta)}(\dot{y}, \dot{\theta}, \dot{\lambda}) = \begin{pmatrix} I_{11} & 0 & -\cos \theta y_* \\ 0 & -\sin \theta & -\alpha \cos \theta \\ 0 & 0 & -(\sin \theta + \beta \cos \theta) \end{pmatrix} \begin{pmatrix} \dot{y} \\ \dot{\theta} \\ \dot{\lambda} \end{pmatrix},$$

where I_{11} is the identity on G_* .

Thus, the continuous map $\Gamma: [-\pi/2, \pi/2] \rightarrow \mathcal{A}(G_* \times \mathbb{R} \times \mathbb{R})$, given by $\theta \mapsto d\tilde{\Psi}_{\gamma(\theta)}$, is in block matrix form as in Remark 2.22. Consequently, up to an inversion of the orientation of Ψ , we may assume that Γ has the canonical orientation, which is such that

$$\text{sign}(d\tilde{\Psi}_{\gamma(\theta)}) = \text{sign} \det(d\tilde{\Psi}_{\gamma(\theta)}) = \text{sign}(\sin \theta(\sin \theta + \beta \cos \theta)).$$

Recalling that $\bar{u}_* = \gamma(-\pi/2)$ and $u_* = \gamma(\pi/2)$, we finally obtain $\text{sign}(d\tilde{\Psi}_{\bar{u}_*}) = \text{sign}(d\tilde{\Psi}_{u_*})$, and the assertion “ $\Psi\text{-deg}(p) = \Psi\text{-deg}(\bar{p}) = \pm 1$ ” follows from (3.3). \square

4. THE PERTURBED EIGENVALUE PROBLEM AND GLOBAL CONTINUATION

Given, as before, two real Hilbert spaces G and H , consider the problem

$$(4.1) \quad \begin{cases} Lx + sN(x) = \lambda Cx \\ x \in S, \end{cases}$$

where $s, \lambda \in \mathbb{R}$, $L, C \in \mathcal{L}(G, H)$, C is compact, and $N: S \rightarrow H$ is a C^1 compact map defined on the unit sphere of G . As in the unperturbed problem (3.1), we assume that $L - \lambda C$ is invertible for some $\lambda \in \mathbb{R}$.

A triple $(s, \lambda, x) \in \mathbb{R} \times \mathbb{R} \times S$ is a *solution* of (4.1) if it satisfies the equation $Lx + sN(x) = \lambda Cx$. The third element $x \in S$ is said to be a *unit eigenvector* corresponding to the *eigenpair* (s, λ) . The set of all the solutions of (4.1) is denoted by Σ , while \mathcal{E} stands for the subset of \mathbb{R}^2 of the eigenpairs of (4.1).

Observe that \mathcal{E} is the projection of Σ into the $s\lambda$ -plane, and that the $s = 0$ slice Σ_0 of Σ coincides with the set $\mathcal{S} = \Psi^{-1}(0)$ of the eigenpoints of (3.1).

A solution of (4.1) is said to be *trivial* if it is of the type $(0, \lambda, x)$. In this case $p = (\lambda, x)$ is the *corresponding eigenpoint* (of the unperturbed problem (3.1)). When p is simple, the solution $(0, \lambda, x)$ will be as well said to be *simple*. Therefore, any simple solution is trivial, but not viceversa.

Consider the C^1 -map

$$\Psi^+: \mathbb{R} \times \mathbb{R} \times S \rightarrow H, \quad (s, \lambda, x) \mapsto \Psi(\lambda, x) + sN(x),$$

where, we recall, $\Psi: \mathbb{R} \times S \rightarrow H$ is defined by $\Psi(\lambda, x) = Lx - \lambda Cx$. Notice that the zeros of Ψ^+ are the solutions of (4.1); that is, $\Sigma = (\Psi^+)^{-1}(0)$.

Since Ψ is Fredholm of index zero, because of the compactness of N , any partial map $\Psi_s^+ = \Psi^+(s, \cdot, \cdot)$ of Ψ^+ is a Φ_0 -map from $\mathbb{R} \times S$ to H . In fact, in Banach spaces, the differential at any point of the domain of a compact C^1 -map is a compact linear operator, and this holds true also in Banach manifolds, due to the fact that they are locally diffeomorphic to open sets of Banach spaces.

Therefore, according to Definition 2.25, Ψ^+ is an extended Φ_0 -homotopy from $\mathbb{R} \times S$ into H .

Due to the fact that the partial map Ψ_0^+ of Ψ^+ coincides with the oriented map Ψ , thanks to Proposition 2.24, the extended Φ_0 -homotopy Ψ^+ admits an orientation (a unique one) which is compatible with that of Ψ . Therefore, from now on, Ψ^+ will be considered an oriented extended Φ_0 -homotopy.

Since the set $\Sigma = (\Psi^+)^{-1}(0)$ has a distinguished (trivial) subset, namely $\{0\} \times \Sigma_0$, it makes sense to consider the notion of bifurcation point. A trivial solution $q_* = (0, \lambda_*, x_*)$ of (4.1) is a *bifurcation point* provided that any neighborhood of q_* contains nontrivial solutions.

A bifurcation point q_* is said to be *global* (in the sense of Rabinowitz [27]) if there exists a connected set of nontrivial solutions whose closure contains q_* and it is either unbounded or meets a bifurcation point q^* different from q_* .

Particularly meaningful is the study of bifurcation points belonging to a set of trivial solutions of the type $\{0\} \times \{\lambda_*\} \times \mathcal{S}_{\lambda_*}$, whose eigensphere \mathcal{S}_{λ_*} is *nontrivial* (that is, with positive dimension). Since, in this case, 0 and λ_* are given, one can simply say that a point $x_* \in \mathcal{S}_{\lambda_*}$, regarded as an alias of $q_* = (0, \lambda_*, x_*)$, is a *bifurcation point* if so is q_* .

For a necessary condition and some sufficient conditions for a point x_* of a nontrivial eigensphere to be a bifurcation point see [17]. Results regarding the existence of (global or non-global) bifurcation points belonging to even-dimensional eigenspheres can be found in [4–8, 18, 19, 21].

Theorem 4.2 below, which is crucial for our main result (Theorem 4.5), provides, in particular, a sufficient condition for an isolated subset of trivial solutions of (4.1) to contain at least one bifurcation point. To prove it, we need the following lemma of point-set topology, which is particularly suitable to our purposes and is obtained from general results by C. Kuratowski (see [26], Chapter 5, Vol. 2). For an interesting paper on connectivity theory we also recommend [1].

Lemma 4.1 ([24]). *Let \mathcal{K} be a compact subset of a locally compact metric space X . If every compact subset of X containing \mathcal{K} has nonempty boundary, then $X \setminus \mathcal{K}$ contains a connected set whose closure in X is non-compact and intersects \mathcal{K} .*

Recall that, according to Notation 2.14, given a subset D of $\mathbb{R} \times \mathbb{R} \times S$ and $s \in \mathbb{R}$, the symbol D_s stands for the s -slice of D . Namely,

$$D_s = \{(\lambda, x) \in \mathbb{R} \times S : (s, \lambda, x) \in D\}.$$

Theorem 4.2. *Let Ω be an open subset of $\mathbb{R} \times \mathbb{R} \times S$. If $\deg(\Psi, \Omega_0, 0) \neq 0$, then Ω has a connected set of nontrivial solutions whose closure in Ω is non-compact and contains at least one bifurcation point.*

Proof. Since, by definition, a trivial solution of (4.1) is a bifurcation point if it is in the closure of the set of nontrivial solutions, the assertion is the same as that of Lemma 4.1 provided that X is the set of the solutions in Ω and \mathcal{K} is its subset of the trivial ones. Namely, $X = (\Psi^+)^{-1}(0) \cap \Omega$ and $\mathcal{K} = \{0\} \times X_0$, where X_0 is the $s = 0$ slice of X , which coincides with $\Psi^{-1}(0) \cap \Omega_0 = \Sigma_0 \cap \Omega_0$.

Thus, it is enough to prove that the metric pair (X, \mathcal{K}) satisfies the assumptions of Lemma 4.1.

Let us show first that X is locally compact. Recall that $\Psi: \mathbb{R} \times S \rightarrow H$ is a Fredholm map of index zero. Therefore, its extension $(s, \lambda, x) \mapsto \Psi(\lambda, x)$ is Fredholm of index one, being obtained by composing Ψ with the projection $(s, \lambda, x) \mapsto (\lambda, x)$, which is a Φ_1 -map (recall the property about the index of the composition of Fredholm operators in Subsection 2.1). Since Ψ^+ is obtained by adding to this extension of Ψ the compact C^1 -map $(s, \lambda, x) \mapsto sN(x)$, we get that Ψ^+ is as well Fredholm of index one. Therefore, Ψ^+ , being Fredholm, is a locally proper map (see [28]). This implies that the set $\Sigma = (\Psi^+)^{-1}(0)$ is locally compact, and so is its relatively open subset X . Moreover, $\mathcal{K} = \{0\} \times X_0$ is compact, since X_0 coincides with the set $\Psi^{-1}(0) \cap \Omega_0$, whose compactness is implicit in the assumption that $\deg(\Psi, \Omega_0, 0)$ is defined.

It remains to prove that any compact subset of X containing \mathcal{K} has nonempty boundary in X . Assume, by contradiction, that this is not the case. Hence there exists a compact subset D of X containing \mathcal{K} whose boundary, in X , is empty. Therefore, D is relatively open in X and, consequently, there exists an open subset W of Ω such that $X \cap W = D$. Incidentally we observe that, according to Definition 3.1, the compact set D is isolated among the elements of X .

Notice that, since $W \subseteq \Omega$, one has $D = \{(\lambda, s, x) \in W : \Psi^+(\lambda, s, x) = 0\}$. Thus, according to the Generalized Homotopy Invariance Property, $\deg(\Psi_s^+, W_s, 0)$ does not depend on $s \in \mathbb{R}$.

Because of the Excision Property, for $s = 0$ one has

$$\deg(\Psi_0^+, W_0, 0) = \deg(\Psi_0^+, \Omega_0, 0).$$

Therefore, since the partial map Ψ_0^+ of Ψ^+ coincides with Ψ , one gets

$$\deg(\Psi_s^+, W_s, 0) = \deg(\Psi, \Omega_0, 0) \neq 0, \quad \forall s \in \mathbb{R}.$$

Now, the compactness of D implies that there exists $s_* \in \mathbb{R}$ such that the set $D_{s_*} = (\Psi_{s_*}^+)^{-1}(0) \cap W_{s_*}$ is empty. Consequently, because of the Existence Property one gets $\deg(\Psi_{s_*}^+, W_{s_*}, 0) = 0$, and the assertion follows from the contradiction. \square

Corollary 4.3 below provides a sufficient condition for the existence of bifurcation of a global branch emanating from a point in an isolated subset of trivial solutions.

In order to deduce it from Theorem 4.2, we need to show that the map Ψ^+ is more than locally proper. Actually,

- Ψ^+ is proper on any bounded and closed subset of its domain.

To check this, observe that Ψ^+ is the sum of two maps: one is the restriction \hat{L} to the manifold $\mathbb{R} \times \mathbb{R} \times S$ of the linear operator

$$\bar{L}: \mathbb{R} \times \mathbb{R} \times G \rightarrow F, \quad (s, \lambda, x) \mapsto Lx,$$

which is Fredholm of index two; the other one is the map $(s, \lambda, x) \mapsto sN(x) - \lambda Cx$, which sends bounded sets into relatively compact sets. The linear operator \bar{L} , being Fredholm, is proper on bounded and closed subsets of its domain. Therefore, the same property is inherited by \hat{L} to the closed subset $\mathbb{R} \times \mathbb{R} \times S$ of $\mathbb{R} \times \mathbb{R} \times G$. One can check that this property is preserved by adding to \hat{L} a compact map.

From Theorem 4.2 we get a sufficient condition for the existence of a global bifurcation point. Recall that the slice Σ_0 of the set $\Sigma = (\Psi^+)^{-1}(0)$ of the solutions of (4.1) coincides with the set $\mathcal{S} = \Psi^{-1}(0)$ of the eigenpoints of (3.1).

Corollary 4.3. *Let \mathcal{K} be an isolated subset of Σ_0 such that $\Psi\text{-deg}(\mathcal{K}) \neq 0$. Then, there exists a connected set of nontrivial solutions of (4.1) whose closure contains a bifurcation point $q_* \in \{0\} \times \mathcal{K}$ and it is either unbounded or encounters a bifurcation point $q^* \notin \{0\} \times \mathcal{K}$. Consequently, q_* is a global bifurcation point.*

Proof. Let Ω be the open subset of $\mathbb{R} \times \mathbb{R} \times S$ obtained by removing the closed set of the elements of $\{0\} \times \Sigma_0$ which are not in $\{0\} \times \mathcal{K}$ (recall that \mathcal{K} , according to Definition 3.1, is relatively open in Σ_0). Thus, Ω_0 is an isolating neighborhood of \mathcal{K} among Σ_0 and, consequently, $\deg(\Psi, \Omega_0, 0) = \Psi\text{-deg}(\mathcal{K}) \neq 0$. Because of Theorem 4.2, there exists a connected set \mathcal{C} of nontrivial solutions whose closure in Ω , call it \mathcal{C}^+ , is non-compact and contains at least one bifurcation point q_* , which, necessarily, belongs to Ω .

It is enough to prove that \mathcal{C} satisfies the first assertion: the second one is a consequence of the fact that the closure of a connected set is as well connected. To this purpose, we need to show, first, that $q_* \in \{0\} \times \mathcal{K}$ and, after this, we may assume that \mathcal{C}^+ is bounded.

The point q_* belongs to $\{0\} \times \mathcal{K}$, since, because of the definition of Ω , one has $\Omega \cap (\{0\} \times \Sigma_0) = \{0\} \times \mathcal{K}$.

Assume now that \mathcal{C}^+ is bounded. Then, so is the closure $\bar{\mathcal{C}}$ of \mathcal{C} (in $\mathbb{R} \times \mathbb{R} \times \mathbb{S}$). It remains to show that $\bar{\mathcal{C}}$ contains a trivial solution q^* which does not belong to $\{0\} \times \mathcal{K}$.

Recall that Ψ^+ is proper on bounded closed subsets of its domain. Therefore, the subset $\bar{\mathcal{C}}$ of $(\Psi^+)^{-1}(0)$ is compact. Moreover, $\bar{\mathcal{C}}$ contains \mathcal{C}^+ , which is not compact. This implies that $\bar{\mathcal{C}}$ has at least one point q^* which is not in \mathcal{C}^+ . The fact that q^* is a bifurcation point not in $\{0\} \times \mathcal{K}$ follows from the definition of Ω . \square

Corollary 4.4. *If D is a compact component of Σ , then $\Psi\text{-deg}(D_0) = 0$.*

Proof. Observe that D_0 is an isolated subset of Σ_0 , due to the fact that the set of all the eigenvalues of (3.1) is discrete.

Suppose, by contradiction, that $\Psi\text{-deg}(D_0) \neq 0$. Then, Corollary 4.3 applies ensuring the existence of a connected set \mathcal{C} of nontrivial solutions of (4.1) whose closure $\bar{\mathcal{C}}$, which is as well connected, contains at least two trivial solutions: one, say q_* , belonging to $\{0\} \times D_0$, and one, call it q^* , outside $\{0\} \times D_0$.

Since q_* belongs to both the connected set $\bar{\mathcal{C}}$ and the component D , one gets $\bar{\mathcal{C}} \subseteq D$. Therefore, q^* belongs to D and, consequently, being trivial, belongs as well to $\{0\} \times D_0$, which is a contradiction yielding the assertion. \square

We are ready to prove our main achievement (Theorem 4.5). Its proof is based on previous results requiring the notion of degree for the oriented map Ψ and the convenient hypothesis $\dim G > 1$. In spite of this, its assertion is still valid when the space G has dimension one, as Example 5.1 shows.

Theorem 4.5. *Let (λ_*, x_*) be a simple eigenpoint of problem (3.1). Then, in the set Σ of the solutions of (4.1), the connected component containing $(0, \lambda_*, x_*)$ is either unbounded or includes a trivial solution $(0, \lambda^*, x^*)$ with $\lambda^* \neq \lambda_*$.*

Proof. We may assume that the connected component D of Σ containing $(0, \lambda_*, x_*)$ is bounded, and we need to show that its slice D_0 is not contained in $\{\lambda_*\} \times \mathcal{S}_{\lambda_*}$.

Recalling that $\Sigma = (\Psi^+)^{-1}(0)$ and that Ψ^+ is proper on bounded closed subsets of its domain, we get that D is compact. Then, Corollary 4.4 implies $\Psi\text{-deg}(D_0) = 0$ and, consequently, because of Theorem 3.6, $D_0 \not\subseteq \{\lambda_*\} \times \mathcal{S}_{\lambda_*}$. \square

5. SOME ILLUSTRATING EXAMPLES AND AN APPLICATION

We give now three examples illustrating the assertion of Theorem 4.5. The last one shows also that, in this theorem, the assumption that the solution $(0, \lambda_*, x_*)$ is simple cannot be removed.

After the examples, we will give an application of Theorem 4.5 to a motion equation containing a nonlinearity like an air resistance force.

5.1. Examples. The first example regards an exhaustive discussion of the solutions of problem (4.1) in the case when $\dim G = 1$. As we shall see, the assertion of Theorem 4.5 holds true also in this minimal dimension.

Example 5.1. Let $G = H = \mathbb{R}$ and consider the problem

$$(5.1) \quad \begin{cases} lx + sN(x) = \lambda cx, \\ |x| = 1, \end{cases}$$

in which l and c are two given real numbers, and N is an arbitrary real function.

We assume $c \neq 0$, so that the unperturbed problem (obtained by putting $s = 0$) has a unique eigenvalue, $\lambda_* = l/c$, and two corresponding twin eigenpoints:

$$p = (\lambda_*, x_*) = (l/c, 1), \quad \bar{p} = (\lambda_*, -x_*) = (l/c, -1),$$

both simple. We will interpret the assertion of Theorem 4.5 in this extreme situation.

For a solution (s, λ, x) of problem (5.1) we have two possibilities: $x = 1$ or $x = -1$.

For $x = 1$ one has $\lambda = (l + sN(1))/c$. Thus, the set of solutions of this type is given by the straight line

$$\Sigma_+ = \{(s, (l + sN(1))/c, 1) \in \mathbb{R}^3 : s \in \mathbb{R}\}.$$

Analogously, for $x = -1$ one gets

$$\Sigma_- = \{(s, (l - sN(-1))/c, -1) \in \mathbb{R}^3 : s \in \mathbb{R}\}.$$

Therefore, the set Σ of all the solutions of (5.1) is $\Sigma_+ \cup \Sigma_-$, and the assertion of Theorem 4.5 is satisfied for both the simple eigenpoints (λ_*, x_*) and $(\lambda_*, -x_*)$.

The following example concerns a differential equation with an evident physical meaning, and the parameter $2s$, when positive, can be regarded as a frictional coefficient. Its abstract formulation has infinitely many eigenpoints, all of them simple. The set Σ of the solution triples (s, λ, x) is the union of infinitely many unbounded components, each of them corresponding to one and only one eigenpoint.

Example 5.2. Let us show how Theorem 4.5 agrees with the structure of the non-zero solutions of the following boundary value problem:

$$(5.2) \quad \begin{cases} x''(t) + 2sx'(t) + \lambda x(t) = 0, \\ x(0) = 0 = x(\pi). \end{cases}$$

For this purpose, we will interpret it as an abstract problem of the type (4.5) by specifying what are here the spaces G and H , the linear operators L and C , and the map N .

Let $H^1(0, \pi)$ be the Hilbert space of the absolutely continuous real functions defined in $[0, \pi]$ with derivative in $L^2(0, \pi)$, and denote by $H^2(0, \pi)$ the Hilbert space of the C^1 real functions in $[0, \pi]$ with derivative in $H^1(0, \pi)$.

Clearly $H^1(0, \pi)$ is a subset of the Banach space $C[0, \pi]$, and the inclusion, we recall, is a compact operator. Therefore the injection of $H^1(0, \pi)$ into $L^2(0, \pi)$ is as well compact, due to the bounded injection of $C[0, \pi]$ into $L^2(0, \pi)$. Analogously, the inclusion of $H^2(0, \pi)$ into $C^1[0, \pi]$ is compact and the inclusion of $C^1[0, \pi]$ into $H^1[0, \pi]$ is continuous.

As a source space G we take the 2-codimensional closed subspace of $H^2(0, \pi)$ consisting of the functions x satisfying the boundary condition $x(0) = 0 = x(\pi)$. The target space H is $L^2(0, \pi)$.

Observe that the second derivative $x \mapsto x''$, as a linear operator from $H^2(0, \pi)$ to $L^2(0, \pi)$, is bounded and Fredholm of index 2, being surjective with 2-dimensional

kernel. Therefore, its restriction $L \in \mathcal{L}(G, H)$ is a Φ_0 -operator (recall the property about the composition of Fredholm operators).

Here C associates to any $x \in G$ the element $-x \in H$. Thus, C is a compact linear operator, since so is the injection of $H^2(0, \pi)$ into $L^2(0, \pi)$. The map N transforms $x \in G$ in $2x' \in H$ and, therefore, it is as well compact, as composition of a bounded linear operator into $H^1(0, \pi)$ with the compact injection into $L^2(0, \pi)$.

Among the infinitely many equivalent norms in $H^2(0, \pi)$ and, consequently, in G we choose the one associated with the inner product

$$\langle x, y \rangle = \frac{1}{\pi} \int_0^\pi (x(t)y(t) + x''(t)y''(t)) dt.$$

As in the previous sections, S denotes the unit sphere of G . Since we are interested in the non-zero solutions of (5.2), the linearity of N justifies the condition $x \in S$ in the following abstract formulation of our problem:

$$(5.3) \quad \begin{cases} Lx + sN(x) = \lambda Cx \\ x \in S. \end{cases}$$

Elementary computations show that the eigenvalues of the unperturbed problem (obtained with $s = 0$) are $\lambda_1 = 1, \lambda_2 = 4, \dots, \lambda_n = n^2, \dots$ and to any λ_n corresponds the 1-dimensional eigenspace $\mathbb{R}x_n = \text{Ker}(L - \lambda_n C)$, with $x_n \in S$ given by

$$x_n(t) = \sqrt{\frac{2}{1+n^4}} \sin(nt).$$

Let us show that $p_n = (\lambda_n, x_n)$ and $\bar{p}_n = (\lambda_n, -x_n)$ are simple eigenpoints, according to Definition 3.4. Since C is compact, the operator $T_n = L - \lambda_n C$ is Fredholm of index zero. Therefore, we need only to prove that Cx_n does not belong to $T_n(G)$, which means that the equation $T_n(x) = Cx_n$ has no solutions in G . In fact, there are no solutions of the resonant problem

$$\begin{cases} x''(t) + n^2 x(t) = \sin(nt), \\ x(0) = 0 = x(\pi). \end{cases}$$

With standard computations one can prove that, given $s \in \mathbb{R}$, the differential equation $x''(t) + 2sx'(t) + \lambda x(t) = 0$ has a non-zero solution verifying the boundary condition $x(0) = 0 = x(\pi)$ if and only if $\lambda = n^2 + s^2$, with $n \in \mathbb{N}$. Therefore, the subset \mathcal{E} of the $s\lambda$ -plane of the eigenpairs of (5.3) is composed of the disjoint union of infinitely many parabolas of equation $\lambda = n^2 + s^2$, $n \in \mathbb{N}$. Moreover, given $(s, n^2 + s^2) \in \mathcal{E}$, any solution in G of the differential equation

$$x''(t) + 2sx'(t) + (s^2 + n^2)x(t) = 0$$

belongs to the straight line $\mathbb{R}x_{s,n}$, where $x_{s,n}(t) = \exp(-st) \sin(nt)$.

As a consequence of this, given any eigenpoint $p_n = (\lambda_n, x_n)$, the connected component, in Σ , containing the corresponding trivial solution $(0, \lambda_n, x_n)$ is the unbounded curve

$$\{(s, s^2 + n^2, x_{s,n}/\|x_{s,n}\|) : s \in \mathbb{R}\}.$$

Obviously, for the twin eigenpoint $\bar{p}_n = (\lambda_n, -x_n)$, one gets

$$\{(s, s^2 + n^2, -x_{s,n}/\|x_{s,n}\|) : s \in \mathbb{R}\}.$$

In conclusion, for any eigenpoint the assertion of Theorem 4.5 is verified.

Example 5.3 below has already been considered in [7] in relation to the conjecture formulated there and solved by Theorem 4.5 above. It concerns a system of two ordinary differential equations with periodic boundary conditions, and the set Σ of its solutions (s, λ, x) has a component which is diffeomorphic to a circle and contains exactly four trivial solutions, all of them simple. These four solutions are associated with two eigenvalues of the unperturbed problem: a pair of twins for each eigenvalue. The other components of Σ are infinitely many 1-dimensional spheres (geometric circles). The projection of each of them into the $s\lambda$ -plane is a singleton $\{(0, \lambda)\}$, with λ an eigenvalue of the unperturbed problem.

Example 5.3. We are interested in the non-zero solutions of the following system of coupled differential equations with 2π -periodic boundary conditions:

$$(5.4) \quad \begin{cases} x_1'(t) + x_1(t) - sx_1(t) = \lambda x_2(t), \\ x_2'(t) - x_2(t) - sx_2(t) = -\lambda x_1(t), \\ x_1(0) = x_1(2\pi), \quad x_2(0) = x_2(2\pi). \end{cases}$$

As in Example 5.2, we interpret our problem in the abstract form

$$(5.5) \quad \begin{cases} Lx + sN(x) = \lambda Cx \\ x \in S, \end{cases}$$

where L , C and N are operators to be defined below, together with the source and the target spaces G and H .

Let $H^1((0, 2\pi), \mathbb{R}^2)$ be the Hilbert space of the absolutely continuous functions $x = (x_1, x_2): [0, 2\pi] \rightarrow \mathbb{R}^2$ with derivative in $L^2((0, 2\pi), \mathbb{R}^2)$.

We take as G the closed subspace of $H^1((0, 2\pi), \mathbb{R}^2)$ of the functions satisfying the periodic condition $x(0) = x(2\pi)$, and as H the space $L^2((0, 2\pi), \mathbb{R}^2)$. Observe that G has codimension 2 in $H^1((0, 2\pi), \mathbb{R}^2)$. Therefore, the operator $L: G \rightarrow H$, given by $(x_1, x_2) \mapsto (x_1' + x_1, x_2' - x_2)$, is Fredholm of index zero.

The operators N and C , given by $(x_1, x_2) \mapsto (-x_1, -x_2)$ and $(x_1, x_2) \mapsto (x_2, -x_1)$ respectively, are compact, since so is the injection

$$H^1((0, 2\pi), \mathbb{R}^2) \hookrightarrow L^2((0, 2\pi), \mathbb{R}^2).$$

The norm in G is the one associated with the inner product

$$\langle x, y \rangle_1 = \frac{1}{2\pi} \int_0^{2\pi} (x(t) \cdot y(t) + x'(t) \cdot y'(t)) dt,$$

where, given two vectors $a = (a_1, a_2)$ and $b = (b_1, b_2)$ in \mathbb{R}^2 , $a \cdot b$ denotes the standard dot product. As in the previous sections, S is the unit sphere in G .

The eigenvalues of the unperturbed problem (obtained by putting $s = 0$ in (5.5)) are $\lambda = \pm\sqrt{1+n^2}$, $n = 0, 1, 2, \dots$ and the set \mathcal{E} of the eigenpairs of (5.5) is the disjoint union of two sets: the circle

$$\mathcal{C} = \{(s, \lambda) \in \mathbb{R}^2 : s^2 + \lambda^2 = 1\}$$

and the isolated eigenpairs

$$\{(0, \lambda) \in \mathbb{R}^2 : \lambda = \pm\sqrt{1+n^2}, n = 1, 2, 3, \dots\}.$$

Notice in fact that, if (s, λ, x) satisfies (5.4) with $s \neq 0$, then x must be constant. Thus, the non-zero solutions of the periodic boundary value problem (5.4) are of two types: the constant ones, corresponding to the eigenpairs of the circle \mathcal{C} , and the oscillating ones associated with the isolated eigenpairs.

Let us examine first the case $(s, \lambda) \in \mathcal{C}$. One has $(s, \lambda) = (\cos \theta, \sin \theta)$, with $\theta \in [0, 2\pi]$, and the kernel of the linear operator

$$L + (\cos \theta)N - (\sin \theta)C \in \mathcal{L}(G, H)$$

is the straight line $\mathbb{R}x^\theta$, where $x^\theta \in S$ is the constant function

$$x^\theta: [0, 2\pi] \rightarrow \mathbb{R}^2, \quad t \mapsto (\cos(\theta/2), \sin(\theta/2)).$$

Therefore, the connected component D of Σ containing the trivial solution

$$q_* = (0, \lambda_*, x_*) = (0, 1, x^{\pi/2})$$

of (5.5) is diffeomorphic to a circle, as its parametrization

$$\theta \in [0, 4\pi] \mapsto (\cos \theta, \sin \theta, x^\theta) \in \mathbb{R} \times \mathbb{R} \times S$$

shows. This component contains four trivial solutions: two of them associated with the eigenvalue $\lambda_* = 1$ and the others with $\lambda_* = -1$. They are all simple solutions, and the component D agrees with the statement of Theorem 4.5.

Incidentally, we observe that the projection of D onto the circle \mathcal{C} is a double covering map, and the above parametrization of D is just a lifting of the map $\theta \mapsto (\cos \theta, \sin \theta) \in \mathcal{C}$, $\theta \in [0, 4\pi]$.

Let us consider now the isolated eigenpairs. That is, the ones having $|\lambda| > 1$. In this case (5.4) admits non-zero solutions if and only if $s = 0$ and $\lambda = \pm\sqrt{1+n^2}$, with $n = 1, 2, 3, \dots$. More precisely, these solutions are oscillating and, given any isolated eigenpair $(0, \lambda_*)$, the corresponding solutions of (5.4), plus the zero one, form a two-dimensional subspace of $H^1((0, 2\pi), \mathbb{R}^2)$. This implies that the eigensphere \mathcal{S}_{λ_*} of the unperturbed problem is the geometric circle $\text{Ker}(L - \lambda_*C) \cap S$. Therefore, if x_* is any element of this circle, the connected component in Σ containing the corresponding trivial solution $(0, \lambda_*, x_*)$ does not satisfy the assertion of Theorem 4.5. Thus, the assumption that the eigenpoint (λ_*, x_*) is simple cannot be removed.

5.2. An application. We close by showing how both Theorem 4.5 and the well-known notion of winding number allow us to deduce theoretically, without explicitly solving the differential equation, that the structure of set Σ of solutions (s, λ, x) of the nonlinear boundary value problem

$$\begin{cases} x''(t) + sg(x'(t)) + \lambda x(t) = 0, \\ x(0) = 0 = x(\pi), \quad x \in S \end{cases}$$

is essentially the same as in Example 5.2. Here $g: \mathbb{R} \rightarrow \mathbb{R}$ is an increasing odd C^1 -function, as is the classical air resistance force $g(v) = v|v|$, and the sphere S is as in Example 5.2. The parameter s , when positive, may be regarded as a frictional coefficient.

The problem can be rewritten in the abstract form as follows:

$$(5.6) \quad \begin{cases} Lx + sN(x) = \lambda Cx \\ x \in S. \end{cases}$$

The spaces G and H are the same as in Example 5.2, and so are the operators L and C . The compact map $N: G \rightarrow H$ sends x into the function $N(x): t \mapsto g(x'(t))$. It is not difficult to prove that N is C^1 and its Fréchet differential at $x \in G$ is given by $dN_x(h): t \mapsto g'(x'(t))h'(t)$.

The unperturbed problem is the same as in Example 5.2. Therefore, its eigenvalues are $\lambda_1 = 1, \lambda_2 = 4, \dots, \lambda_n = n^2, \dots$. They are all simple and, consequently, each of them corresponds to a pair of isolated unit eigenpoints.

We will prove that the set Σ of the solutions (s, λ, x) of (5.6) contains infinitely many unbounded components, each of them corresponding to one and only one eigenpoint.

Let S^1 denote the unit circle of \mathbb{C} and let $w: C(S^1) \rightarrow \mathbb{Z}$ stand for the winding number function, defined on the set of the continuous maps from S^1 into itself. Recall that, given $\gamma \in C(S^1)$, $w(\gamma)$ is the same as the Brouwer degree of γ and, speaking loosely, denotes the number of times that γ travels counterclockwise around the origin of \mathbb{C} , and it is negative if the curve travels clockwise.

Call wj the integer valued function that to any non-zero solution x of the parametrized differential equation

$$(5.7) \quad x''(t) + sg(x'(t)) + \lambda x(t) = 0,$$

depending on $s, \lambda \in \mathbb{R}$, assigns the winding number $wj(x)$ of the closed curve $j(x) \in C(S^1)$ defined by

$$z = e^{i\theta} \mapsto \frac{(x'(\theta/2) + ix'(0)x(\theta/2))^2}{x'(\theta/2)^2 + x'(0)^2 x(\theta/2)^2}, \quad \theta \in [0, 2\pi].$$

Notice that, given any non-zero solution x of (5.7), $j(x)$ is well defined, since $x(t)$ and $x'(t)$ cannot be simultaneously zero, due to the uniqueness of the Cauchy problem. Observe also that $j(x)$ is a closed curve, since both the endpoints coincide with $1 \in \mathbb{C}$.

It is convenient to extend the map $x \mapsto j(x)$ to the symmetric set of all the functions $x \in G$ having the property that $x(t)^2 + x'(t)^2 > 0$ for all $t \in [0, \pi]$. We denote this set by \mathcal{X} and we observe that it is open, because of the bounded inclusions $H^2(0, \pi) \hookrightarrow C^1[0, \pi]$ and $H^1(0, \pi) \hookrightarrow C[0, \pi]$.

One can check that, for example, if $x(t) = \sin(nt)$ with $n \in \mathbb{Z}$, then $j(x)$ is the map $z \mapsto z^n$, whose winding number is n .

One can also check that, if a is a positive constant and $x \in \mathcal{X}$, then $j(x)$ and $j(ax)$ are homotopic, therefore they have the same winding number. Moreover, if $x \in \mathcal{X}$, then $j(-x) = 1/\overline{j(x)}$, which is the same as the conjugate map $\overline{j(x)}$ of $j(x)$. Therefore, the winding numbers of $j(x)$ and $j(-x)$ are opposite each other, and this happens for the two unit eigenvectors corresponding to any eigenvalue $\lambda_n = n^2$ of our problem. In fact, this number is n for the unit eigenfunction

$$x_n(t) = \sqrt{\frac{2}{1+n^4}} \sin(nt),$$

and $-n$ for the opposite one.

Observe that, due to the fact that \mathcal{X} is open in G , if two functions of \mathcal{X} are sufficiently close, then the segment joining them lies in \mathcal{X} . Therefore, the corresponding two images under $j: \mathcal{X} \rightarrow C(S^1)$ are homotopic and, consequently, they have the same winding number. Thus, the integer valued function $wj: \mathcal{X} \rightarrow \mathbb{Z}$ is locally constant.

Since the projection map $p: \Sigma \rightarrow \mathcal{X}$ that to any solution (s, λ, x) of (5.6) assigns the function x is continuous, we have the following

Remark 5.4. *The map $\text{wjp}: \Sigma \rightarrow \mathbb{Z}$ that to any solution (s, λ, x) of (5.6) assigns the winding number of the closed curve $j(x)$ is locally constant.*

Let $q_* = (0, \lambda_*, x_*)$ be any trivial solution of (5.6). We want to prove that the connected component D_* of Σ containing q_* is unbounded and does not meet other trivial solutions.

To this purpose, observe first that, since D_* is connected, Remark 5.4 implies $\text{wjp}(q) = \text{wjp}(q_*)$ for all $q \in D_*$. In particular, if $\lambda_* = \lambda_n = n^2$, then $\text{wjp}(q_*)$ is n or $-n$, depending on whether x_* is the above function x_n or its opposite. Thus, D_* does not contain trivial solutions different from q_* , consequence of the fact that the function that to any trivial solution q assigns the integer $\text{wjp}(q)$ is injective.

Finally, from Theorem 4.5 we get that D_* is unbounded, since otherwise D_* would contain a trivial solution different from q_* .

REFERENCES

- [1] Alexander J.C., *A primer on connectivity*, Lecture Notes in Math., Vol. 886, Springer Verlag, Berlin-New York, 1981, 455–483.
- [2] Amann H. - Weiss S.A., *On the uniqueness of the topological degree*, Math. Z., **130** (1973), 39–54.
- [3] Appell J. - De Pascale E. - Vignoli A., *Nonlinear spectral theory*, de Gruyter, Berlin, 2004.
- [4] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *On the persistence of the eigenvalues of a perturbed Fredholm operator of index zero under nonsmooth perturbations*, Z. Anal. Anwend. **36** (2017), no. 1, 99–128.
- [5] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *Global continuation of the eigenvalues of a perturbed linear operator*, Ann. Mat. Pura Appl. **197** (2018), no. 4, 1131–1149.
- [6] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *Global continuation in Euclidean spaces of the perturbed unit eigenvectors corresponding to a simple eigenvalue*, Topol. Methods Nonlinear Anal. **55** (2020), no. 1, 169–184.
- [7] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *Global persistence of the unit eigenvectors of perturbed eigenvalue problems in Hilbert spaces*, Z. Anal. Anwend. **39** (2020), no. 4, 395–419.
- [8] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *The Brouwer degree associated to classical eigenvalue problems and applications to nonlinear spectral theory*, preprint arXiv:1912.03182.
- [9] Benevieri P. - Furi M., *A simple notion of orientability for Fredholm maps of index zero between Banach manifolds and degree theory*, Ann. Sci. Math. Québec, **22** (1998), 131–148.
- [10] Benevieri P. - Furi M., *On the concept of orientability for Fredholm maps between real Banach manifolds*, Topol. Methods Nonlinear Anal. **16** (2000), 279–306.
- [11] Benevieri P. - Furi M., *On the uniqueness of the degree for nonlinear Fredholm maps of index zero between Banach manifolds*, Commun. Appl. Anal. **15** (2011), 203–216.
- [12] Benevieri P. - Furi M. - Pera M.P. - Spadini M., *About the sign of oriented Fredholm operators between Banach spaces*, Z. Anal. Anwendungen **22** (2003), no. 3, 619 – 645.
- [13] Capietto A. - Mawhin J. - Zanolin F., *A continuation approach to superlinear periodic boundary value problems*, J. Differential Equations **88** (1990), no. 2, 347–395.
- [14] Chiappinelli R., *Isolated Connected Eigenvalues in Nonlinear Spectral Theory*, Nonlinear Funct. Anal. Appl. **8** (2003), no. 4, 557–579.
- [15] Chiappinelli R., *Approximation and convergence rate of nonlinear eigenvalues: Lipschitz perturbations of a bounded self-adjoint operator*, J. Math. Anal. Appl. **455** (2017), no. 2, 1720–1732.
- [16] Chiappinelli R., *What do you mean by “nonlinear eigenvalue problems”?*, Axioms **7** (2018), Paper no. 39, 30 pp.
- [17] Chiappinelli R. - Furi M. - Pera M.P., *Normalized Eigenvectors of a Perturbed Linear Operator via General Bifurcation*, Glasg. Math. J. **50** (2008), no. 2, 303–318.
- [18] Chiappinelli R. - Furi M. - Pera M.P., *Topological persistence of the normalized eigenvectors of a perturbed self-adjoint operator*, Appl. Math. Lett. **23** (2010), no. 2, 193–197.
- [19] Chiappinelli R. - Furi M. - Pera M.P., *A new theme in nonlinear analysis: continuation and bifurcation of the unit eigenvectors of a perturbed linear operator*, Communications in Applied Analysis **15** (2011), no. 2, 3 and 4, 299–312.

- [20] Chiappinelli R. - Furi M. - Pera M.P., *Persistence of the normalized eigenvectors of a perturbed operator in the variational case*, Glasg. Math. J. **55** (2013), no. 3, 629–638.
- [21] Chiappinelli R. - Furi M. - Pera M.P., *Topological Persistence of the Unit Eigenvectors of a Perturbed Fredholm Operator of Index Zero*, Z. Anal. Anwend. **33** (2014), no. 3, 347–367.
- [22] Furi M. - Pera M.P., *The forced spherical pendulum does have forced oscillations*, Delay Differential Equations and Dynamical Systems, Busenberg S. and Martelli M. editors, Lecture Notes in Math. 1475, Springer Verlag, Berlin, 1991, 176–183.
- [23] Furi M. - Pera M.P., *On the notion of winding number for closed curves and applications to forced oscillations on even dimensional spheres*, Boll. Un. Mat. Ital. (1993), Vol. 7-A, no. 3, 397–407.
- [24] Furi M. - Pera M.P., *A continuation principle for periodic solutions of forced motion equations on manifolds and applications to bifurcation theory*, Pacific J. Math. **160** (1993), no. 3, 219–244.
- [25] Kato T., *Perturbation Theory for Linear Operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
- [26] Kuratowski C., *Topology*, Vol. 2, Academic Press, New York, 1968.
- [27] Rabinowitz P.H., *Some global results for nonlinear eigenvalue problems*, J. Funct. Anal. **7** (1971), 487–513.
- [28] Smale S., *An infinite dimensional version of Sard’s theorem*, Amer. J. Math. **87** (1965), 861–866.
- [29] Taylor A.E. - Lay D.C., *Introduction to Functional Analysis*, John Wiley & Sons, New York-Chichester-Brisbane, 1980.

PIERLUIGI BENEVIERI - INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, SÃO PAULO - SP - BRASIL - CEP 05508-090 - *E-mail address*: pluigi@ime.usp.br

ALESSANDRO CALAMAI - DIPARTIMENTO DI INGEGNERIA CIVILE, EDILE E ARCHITETTURA, UNIVERSITÀ POLITECNICA DELLE MARCHE, VIA BRECCIE BIANCHE, I-60131 ANCONA, ITALY - *E-mail address*: calamai@dipmat.univpm.it

MASSIMO FURI - DIPARTIMENTO DI MATEMATICA E INFORMATICA “ULISSE DINI”, UNIVERSITÀ DEGLI STUDI DI FIRENZE, VIA S. MARTA 3, I-50139 FLORENCE, ITALY - *E-mail address*: massimo.furi@unifi.it

MARIA PATRIZIA PERA - DIPARTIMENTO DI MATEMATICA E INFORMATICA “ULISSE DINI”, UNIVERSITÀ DEGLI STUDI DI FIRENZE, VIA S. MARTA 3, I-50139 FLORENCE, ITALY - *E-mail address*: mpatrizia.pera@unifi.it