

A fixed point approach for decaying solutions of difference equations

Zuzana Došlá, Mauro Marini and Serena Matucci

Abstract. A boundary value problem associated to the difference equation with advanced argument

$$\Delta(a_n \Phi(\Delta x_n)) + b_n \Phi(x_{n+p}) = 0, \quad n \geq 1 \quad (*)$$

is presented, where $\Phi(u) = |u|^\alpha \operatorname{sgn} u$, $\alpha > 0$, p is a positive integer and the sequences a, b , are positive. We deal with a particular type of decaying solutions of (*), that is the so-called intermediate solutions (see below for the definition) . In particular, we prove the existence of these type of solutions for (*) by reducing it to a suitable boundary value problem associated to a difference equation without deviating argument. Our approach is based on a fixed point result for difference equations, which originates from existing ones stated in the continuous case. Some examples and suggestions for future researches complete the paper.

Keywords. Nonlinear difference equation, functional discrete equations, boundary value problem on the half line, fixed point theorem, decaying solution.

MSC 2010: Primary 39A22 , Secondary 47H10

1 Introduction

Consider the equation

$$\Delta(a_n \Phi(\Delta x_n)) + b_n \Phi(x_{n+p}) = 0, \quad (\text{P})$$

where Δ is the forward difference operator $\Delta x_n = x_{n+1} - x_n$, Φ is the operator $\Phi(u) = |u|^\alpha \operatorname{sgn} u$, $\alpha > 0$, p is a positive integer and the sequences a, b are

positive for $n \geq 1$, and satisfy

$$\sum_{i=1}^{\infty} \Phi^* \left(\frac{1}{a_i} \right) < \infty, \quad \sum_{i=1}^{\infty} b_i = \infty, \quad (1)$$

where Φ^* is the inverse of the map Φ , that is $\Phi^*(u) = |u|^{1/\alpha} \text{sgn } u$.

Equation (P) appears in the discretization process for searching spherically symmetric solutions of certain nonlinear elliptic equations with weighted φ -Laplacian, see, e.g., [9]. A special case of (P) is the discrete half-linear equation

$$\Delta(a_n \Phi(\Delta y_n)) + b_n \Phi(y_{n+1}) = 0, \quad (\text{H})$$

which has been studied extensively from various points of view, especially with regard to the oscillation and the qualitative behavior of nonoscillatory solutions, see [12, Chapter 3] and references therein.

As usual, a nontrivial solution x of (P) is said to be *nonoscillatory* if x_n is either positive or negative for any large n and *oscillatory* otherwise. In virtue of the Sturm separation criterion, see, e.g. [12, Section 8.2.1.], all the solutions of (H) have the same behavior with respect to the oscillation. In other words, either all nontrivial solutions of (H) are nonoscillatory or all the solutions of (H) are oscillatory. Thus, equivalently, we say that (H) is nonoscillatory if (H) has a nonoscillatory solution. Nevertheless, for equation (P) with $p \neq 1$, the situation is different, since in this case oscillatory solutions and nonoscillatory solutions may coexist.

In this paper we deal with the existence of a particular type of nonoscillatory solutions, that is solutions x of (P) such that $x_n > 0$, $\Delta x_n < 0$ for large n and

$$\lim_n x_n = 0, \quad \lim_n x_n^{[1]} = a_n \Phi(\Delta x_n) = -\infty, \quad (2)$$

where $x^{[1]}$ is called the quasidifference of x . Solutions satisfying (2) are called *intermediate solutions*. This terminology originates from the corresponding continuous case and it is due to the fact that, when (1) is satisfied, then any nonoscillatory solution x of (P) satisfies either $\lim_n x_n = \ell_x \neq 0$, or (2), or $\lim_n x_n = 0$, $\lim_n x_n^{[1]} = -\ell_x$, $0 < \ell_x < \infty$, see [5, 7]. The investigation of intermediate solutions is an hard problem, due to difficulties in finding suitable sharp upper and lower bounds for these solutions, see, e.g., [2, page 241] and [14, page 3], in which these facts are pointed out for the continuous case.

In the half-linear case the problem of existence of intermediate solutions has been completely solved by the following.

Theorem 1. *Assume (1).*

(i₁) *Equation (H) does not have intermediate solutions if*

$$\sum_{n=1}^{\infty} b_n \Phi \left(\sum_{k=n+1}^{\infty} \Phi^* \left(\frac{1}{a_k} \right) \right) + \sum_{n=1}^{\infty} \Phi^* \left(\frac{1}{a_{n+1}} \sum_{k=1}^n b_k \right) < \infty.$$

(i₂) *Equation (H) has intermediate solutions if (H) is nonoscillatory and*

$$\sum_{n=1}^{\infty} b_n \Phi \left(\sum_{k=n+1}^{\infty} \Phi^* \left(\frac{1}{a_k} \right) \right) + \sum_{n=1}^{\infty} \Phi^* \left(\frac{1}{a_{n+1}} \sum_{k=1}^n b_k \right) = \infty.$$

Proof. The assertion follows from Theorem 3.1 (b), (c) and Theorem 3.2 in [6], with minor changes. \square

Here, we present a comparison result which allows to solve the boundary value problem [BVP]

$$\begin{cases} \Delta(a_n \Phi(\Delta x_n)) + b_n \Phi(x_{n+p}) = 0, & p > 1, \\ \lim_n x_n = 0, \lim_n x_n^{[1]} = -\infty \end{cases} \quad (3)$$

by reducing it to the existence of intermediate solutions in the half-linear case. The main result is the following.

Theorem 2. *Assume (1) and*

$$\limsup_n b_n < \infty. \quad (4)$$

Then the BVP (3) is solvable, i.e. equation (P) with $p > 1$ has intermediate solutions, if and only if the half-linear equation

$$\Delta(a_{n+p-1} \Phi(\Delta y_n)) + b_n \Phi(y_{n+1}) = 0, \quad (\text{H1})$$

has intermediate solutions.

Theorem 2 will be proved by means of a fixed point result for discrete operators acting in Fréchet spaces, see Theorem 4 below.

Combining Theorem 1 and the comparison Theorem 2, we get necessary and sufficient conditions for existence of intermediate solutions of difference equation with advanced argument.

Corollary 1. *Assume (1) and (4).*

(i₁) *Equation (P) with $p > 1$ has intermediate solutions if (H1) is nonoscillatory and*

$$\sum_{n=1}^{\infty} b_n \Phi \left(\sum_{k=n+1}^{\infty} \Phi^* \left(\frac{1}{a_{k+p-1}} \right) \right) + \sum_{n=1}^{\infty} \Phi^* \left(\frac{1}{a_{n+p}} \sum_{k=1}^n b_k \right) = \infty. \quad (5)$$

(i₂) *Equation (P) with $p > 1$ does not have intermediate solutions if*

$$\sum_{n=1}^{\infty} b_n \Phi \left(\sum_{k=n+1}^{\infty} \Phi^* \left(\frac{1}{a_{k+p-1}} \right) \right) + \sum_{n=1}^{\infty} \Phi^* \left(\frac{1}{a_{n+p}} \sum_{k=1}^n b_k \right) < \infty.$$

2 Fixed point approaches

Boundary value problems for difference equations in \mathbb{R}^n are often solved by reducing the problem to a fixed point equation for a possibly nonlinear operator in a suitable function space. Thus, the existence of a solution is obtained by applying a fixed point theorem, for instance the Tychonoff theorem, the Schauder theorem, the Leray-Schauder continuation principle or Krasnoselkii types fixed point theorems on cones. For a survey on this topic we refer to the papers [1, 3, 15] and the monographies [4, 17]. In particular, in [17, Chapter 2], see also [4, Chapter 5], certain BVPs are studied by means of a nonlinear Leray-Schauder alternative. This approach is based on a very general method given in [13]. In particular, in [13] the authors present a Leray-Schauder continuation principle in locally convex topological vector spaces, which unifies the Leray-Schauder alternative theorem and the Tichonov fixed point theorem. More precisely, let \mathbb{E} be a Hausdorff locally convex topological vector space with a family of seminorms generating the topology. The following holds.

Theorem 3. [13, Theorem 1.1] *Let Q be a convex closed subset of \mathbb{E} and let $T : Q \times [0, 1] \rightarrow \mathbb{E}$ be a continuous map with relatively compact image. Assume that:*

- i₁) $T(x, 0) \in Q$ for any $x \in Q$;*
- i₂) for any $(x, \lambda) \in \partial Q \times [0, 1]$ with $T(x, \lambda) = x$ there exists open neighborhoods U_x of x in \mathbb{E} and I_λ of λ in $[0, 1]$ such that*

$$T\left((U_x \cap \partial Q) \times I_\lambda\right) \subset Q.$$

Then the equation

$$x = T(x, 1)$$

has a solution.

Some of the above quoted results have a discrete counterpart. For instance, a method for solving BVPs associated to difference systems is given in [15, Theorem 2.1]. Due to the peculiarities of the discrete case, it may be applied to functional difference equations, including equations with deviating arguments or sum difference equations.

Now, we present an existence result which generalizes, in the particular case of scalar difference equations, [15, Theorem 2.1].

Denote by \mathbb{N}_n and $\mathbb{N}_{m,n}$, the sets

$$\mathbb{N}_n = \{i \in \mathbb{N} : i \geq n \in \mathbb{N}\}$$

$$\mathbb{N}_{m,n} = \{i \in \mathbb{N}_m : i < n, m, n \in \mathbb{N}, m < n\}$$

and let \mathbb{X} be the space of all real sequences defined on \mathbb{N}_m . Hence \mathbb{X} is a Fréchet space with the topology of pointwise convergence on \mathbb{N}_m . From the discrete Arzelà-Ascoli theorem (see, e.g., [4, Theorem 5.3.1]), any bounded set in \mathbb{X} is relatively compact. We recall that a set $\Omega \subset \mathbb{X}$ is bounded if and only if it consists of sequences which are equibounded on $\mathbb{N}_{m,n}$ for any $n > m$. Clearly, if $\Omega \subset \mathbb{X}$ is bounded, then $\Omega^\Delta = \{\Delta u, u \in \Omega\}$ is bounded, too.

Using, with minor changes, a discrete counterpart of a compactness and continuity result stated in [8, Theorem 1.3] for the continuous case, we have the following.

Theorem 4. *Consider the BVP*

$$\begin{cases} \Delta(a_n \Phi(\Delta x_n)) = g(n, x), & n \in \mathbb{N}_m \\ x \in S, \end{cases} \quad (6)$$

where $g : \mathbb{N}_m \times \mathbb{X} \rightarrow \mathbb{R}$ is a continuous map, and S is a subset of \mathbb{X} .

Let $G : \mathbb{N}_m \times \mathbb{X}^2 \rightarrow \mathbb{R}$ be a continuous map such that $G(k, u, u) = g(k, u)$ for all $(k, u) \in \mathbb{N}_m \times \mathbb{X}$. If there exist a nonempty, closed, convex set $\Omega \subset \mathbb{X}$, and a bounded, closed subset $S_C \subset S \cap \Omega$ such that the problem

$$\begin{cases} \Delta(a_n \Phi(\Delta x_n)) = G(n, x, q), & n \in \mathbb{N}_m \\ x \in S_C, \end{cases} \quad (7)$$

has a unique solution for any $q \in \Omega$ fixed, then (6) has at least a solution.

Proof. Let T be the operator $T : \Omega \rightarrow S_C$ which maps every $q \in \Omega$ into the unique solution $x = T(q)$ of (7). Let us show that the operator T is continuous with relatively compact image. The relatively compactness of $T(\Omega)$ follows immediately since S_C is bounded. To prove the continuity of T in Ω , let $\{q^j\}$ be a sequence in Ω , $q^j \rightarrow q^\infty \in \Omega$, and let $v^j = T(q^j)$. Since $T(\Omega)$ is relatively compact, $\{v^j\}$ admits a subsequence (still indicated with $\{v^j\}$) which converges to $v^\infty \in \mathbb{X}$. As $v^j \in S_C$ and S_C is closed, then $v^\infty \in S_C$. Taking into account the continuity of G , we obtain

$$\Delta(a_n \Phi(\Delta v_k^\infty)) = \lim_j \Delta(a_n \Phi(\Delta v_k^j)) = \lim_j G(k, q^j, v^j) = G(k, q^\infty, v^\infty).$$

The uniqueness of the solution of (7) yields $v^\infty = T(q^\infty)$, and therefore T is continuous on Ω . By the Tychonoff fixed point theorem, T has at least one fixed point in Ω , which is a solution of (6), as it can be easily checked, taking into account that $S_C \subset S$. \square

As follows from the proof of Theorem 4, no explicit form of the fixed point operator is needed for the solvability of (6). A key point for the unique solvability of (7) in Theorem 4 is the choice of the map G . To this aim, in our opinion, the best cases are the following two, namely

$$i_1) \quad G(n, q, x) = \tilde{g}(n, q), \quad i_2) \quad G(n, q, x) = \tilde{g}(n, q)\Phi(x_{n+1}).$$

In the case i_2), the equation in (7) is a half-linear equation and in this situation a very large variety of results is known, see, e.g., [12, Chapter VIII]. An application in this direction is in [15, Section 4].

In the case i_1), that is when the function G does not depend on x and the equation in (7) is affine, Theorem 4 can be particularly useful to solve BVPs associated to difference equations with deviating arguments. Indeed, in this case, it can lead to a BVP associated to a second order difference equation without deviating argument. An application of this fact is in the following section.

3 Proof of Theorem 2

For proving Theorem 2, the following auxiliary result is needed.

Lemma 1. Assume (1) and (4). Let x be an eventually positive decreasing solution of (P) such that $\lim_n x_n = 0$. Then the series

$$\sum_{i=2}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} \sum_{k=1}^{i-1} b_k \Phi(x_{k+p}) \right)$$

converges.

Proof. Without loss of generality, suppose $0 < x_n \leq 1$, $\Delta x_n < 0$ for $n \geq m_0 \geq 1$. We claim that for any $k, j \geq m_0$ we have

$$\left| x_j^{[1]} - x_k^{[1]} \right| \leq B|j - k|, \quad (8)$$

where $B = \sup_{i \geq 1} b_i$. For simplicity, suppose $k \geq j$. Summing (P) we have

$$x_k^{[1]} - x_j^{[1]} = - \sum_{i=j}^{k-1} b_i \Phi(x_{i+p}), \quad (9)$$

where, as usual, $\sum_{i=k_1}^k \gamma_i = 0$ if $k < k_1$. Since $|x_i| \leq 1$ and $b_i \leq B$, the inequality (8) follows. Further, from (9),

$$\begin{aligned} \sum_{i=m_0}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} \sum_{k=m_0}^{i-1} b_k \Phi(x_{k+p}) \right) &= \sum_{i=m_0}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} (x_{m_0}^{[1]} - x_i^{[1]}) \right) = \\ &= \sum_{i=m_0}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} (x_{m_0}^{[1]} - x_i^{[1]} - x_{i+p-1}^{[1]} + x_{i+p-1}^{[1]}) \right). \end{aligned}$$

Since

$$\left| x_i^{[1]} - x_{i+p-1}^{[1]} + x_{i+p-1}^{[1]} - x_{m_0}^{[1]} \right| \leq \left| x_i^{[1]} - x_{i+p-1}^{[1]} \right| + \left| x_{i+p-1}^{[1]} \right| + \left| x_{m_0}^{[1]} \right|,$$

in view of (8) we obtain

$$\begin{aligned} &\left| \sum_{i=m_0}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} (x_i^{[1]} - x_{m_0}^{[1]}) \right) \right| \leq \\ &\leq \sum_{i=m_0}^{\infty} \Phi^* \left(\frac{B(p-1) + |x_{m_0}^{[1]}|}{a_{i+p-1}} + \frac{|x_{i+p-1}^{[1]}|}{a_{i+p-1}} \right). \end{aligned}$$

In virtue of (1), the series

$$\sum_{i=m_0}^{\infty} \Phi^* \left(\frac{B(p-1) + |x_{m_0}^{[1]}|}{a_{i+p-1}} \right)$$

converges. Since

$$\sum_{i=m_0}^{\infty} \Phi^* \left(\frac{|x_{i+p-1}^{[1]}|}{a_{i+p-1}} \right) = - \sum_{i=m_0}^{\infty} \Delta x_{i+p-1} = x_{m_0+p-1},$$

using (1) and the inequality

$$\Phi^*(X + Y) \leq \sigma_{\alpha}(\Phi^*(X) + \Phi^*(Y)),$$

where

$$\sigma_{\alpha} = \begin{cases} 1 & \text{if } \alpha \geq 1 \\ 2^{(1-\alpha)/\alpha} & \text{if } \alpha < 1 \end{cases},$$

we obtain the assertion. \square

Proof of Theorem 2. First, we prove that if (P) has intermediate solutions, then (H1) has intermediate solutions.

Let x be an intermediate solution of (P) and, without loss of generality, assume for $n \geq n_0 \geq 1$

$$0 < x_n < 1, \quad \Delta x_n < 0. \quad (10)$$

In view of (4), there exists $L > 0$ such that for any $n \geq n_0$

$$\sum_{i=n}^{n+p-2} b_i \leq L. \quad (11)$$

Moreover, let M be a positive constant, $M < 1$, such that

$$\Phi(M) \leq \frac{|x_{n_0}^{[1]}|}{L + |x_{n_0}^{[1]}|}. \quad (12)$$

Let \mathbb{X} be the Fréchet space of real sequences defined for $n \geq n_0$, endowed with the topology of convergence on \mathbb{N}_{n_0} , and consider the subset $\Omega \subset \mathbb{X}$ defined by

$$\Omega = \{u \in \mathbb{X} : Mx_{n+p-1} \leq u_n \leq x_{n+p-1}\}.$$

For any $u \in \Omega$ consider the boundary value problem [BVP]

$$\begin{cases} \Delta(a_{n+p-1}\Phi(\Delta z_n)) + b_n\Phi(u_{n+1}) = 0, & n \geq n_0 \\ z_{n_0}^{[1]} = x_{n_0}^{[1]}, \quad \lim_n z_n = 0, \end{cases} \quad (13)$$

where $z^{[1]}$ denotes the quasidifference of z , that is

$$z_n^{[1]} = a_{n+p-1}\Phi(\Delta z_n). \quad (14)$$

For any $u \in \Omega$ we have

$$\sum_{k=n_0}^n b_k\Phi(u_{k+1}) \leq \sum_{k=n_0}^n b_k\Phi(x_{k+p}).$$

Hence, using Lemma 1, we have

$$\lim_n \sum_{i=n}^{\infty} \Phi^* \left(\frac{1}{a_{i+p-1}} \left(\sum_{k=n_0}^{i-1} b_k\Phi(u_{k+1}) \right) \right) = 0.$$

Thus, a standard calculation shows that for any $u \in \Omega$ the BVP (13) has the unique solution z . Let \mathcal{T} be the operator which associates to any $u \in \Omega$ the unique solution z of (13).

Summing the equation in (13) and using (9) we get

$$z_n^{[1]} = x_{n_0}^{[1]} - \sum_{k=n_0}^{n-1} b_k\Phi(u_{k+1}) \geq x_{n_0}^{[1]} - \sum_{k=n_0}^{n-1} b_k\Phi(x_{k+p}) = x_n^{[1]}.$$

Since $x^{[1]}$ is decreasing for $n \geq n_0$ and $p > 1$, we obtain for $n \geq n_0$

$$z_n^{[1]} \geq x_{n+p-1}^{[1]} \quad (15)$$

i.e., in view of (14),

$$a_{n+p-1}\Phi(\Delta z_n) \geq a_{n+p-1}\Phi(\Delta x_{n+p-1}),$$

that is,

$$\Delta z_n \geq \Delta x_{n+p-1}.$$

Since $\lim_i z_i = \lim_i x_i = 0$, we obtain for $n \geq n_0$

$$z_n \leq x_{n+p-1}. \quad (16)$$

Now, let us prove that for $n \geq n_0$

$$z_n \geq Mx_{n+p-1}. \quad (17)$$

Summing the equation in (13) and using (14) we get

$$z_n^{[1]} = x_{n_0}^{[1]} - \sum_{k=n_0}^{n-1} b_k \Phi(u_{k+1}) \leq x_{n_0}^{[1]} - \Phi(M) \sum_{k=n_0}^{n-1} b_k \Phi(x_{k+p}),$$

or, using (9),

$$\begin{aligned} z_n^{[1]} &\leq x_{n_0}^{[1]} + \Phi(M) (x_n^{[1]} - x_{n_0}^{[1]}) = \\ &= \Phi(M)x_{n+p-1}^{[1]} + \Phi(M) (x_n^{[1]} - x_{n+p-1}^{[1]}) + (1 - \Phi(M))x_{n_0}^{[1]}. \end{aligned} \quad (18)$$

From (9), (10) and (11), we also have

$$x_n^{[1]} - x_{n+p-1}^{[1]} = \sum_{i=n}^{n+p-2} b_i \Phi(x_{i+p}) \leq \sum_{i=n}^{n+p-2} b_i \leq L.$$

Thus, from (18) we obtain

$$z_n^{[1]} \leq \Phi(M)x_{n+p-1}^{[1]} + (L + |x_{n_0}^{[1]}|) \Phi(M) + x_{n_0}^{[1]}. \quad (19)$$

In view of (12) we have

$$(L + |x_{n_0}^{[1]}|) \Phi(M) + x_{n_0}^{[1]} \leq 0.$$

Hence, from (19) we get for $n \geq n_0$

$$z_n^{[1]} \leq \Phi(M)x_{n+p-1}^{[1]} \quad (20)$$

or, in view of (14), $\Delta z_n \leq M \Delta x_{n+p-1}$, and (17) follows, since $\lim_i z_i = \lim_i x_i = 0$. Thus, in virtue of (16) and (17), the operator \mathcal{T} maps Ω into itself, that is

$$\mathcal{T}(\Omega) \subset \Omega.$$

Denote by S the boundary conditions in (13), i.e.

$$S = \left\{ v \in \mathbb{X} : a_{n_0+p-1} \Phi(\Delta v_{n_0}) = x_{n_0}^{[1]}, \lim_n v_n = 0 \right\}$$

For any $z \in \mathcal{T}(\Omega)$ we have $z \in S$. Since $\mathcal{T}(\Omega) \subset \Omega$, we get $z \in \Omega \cap S$.

Denote by S_C the subset of \mathbb{X} given by

$$S_C = S \cap \Omega.$$

Since $\lim_n x_n = 0$, it holds

$$S_C = \left\{ v \in \mathbb{X} : a_{n_0+p-1} \Phi(\Delta v_{n_0}) = x_{n_0}^{[1]}, Mx_{n+p-1} \leq v_n \leq x_{n+p-1} \right\}.$$

Thus S_C is a bounded and closed subset of \mathbb{X} . Applying Theorem 4 we obtain that the operator \mathcal{T} has a fixed point $\bar{z} \in S_C$. Clearly the sequence \bar{z} is a solution of (H1) and $\lim_n \bar{z}_n = 0$. Since $\bar{z} \in \mathcal{T}(\Omega)$, from (15) and (20), we get

$$x_{n+p-1}^{[1]} \leq \bar{z}_n^{[1]} \leq \Phi(M)x_{n+p-1}^{[1]}$$

and so $\lim_n \bar{z}_n^{[1]} = -\infty$. Hence \bar{z} is an intermediate solution of (H1).

Now, we prove the vice-versa, that is if (H1) has intermediate solutions, then (P) has intermediate solutions.

The argument is similar to the one above given, with minor changes. Let y be an intermediate solution of (H1) such that for $n \geq n_0 \geq 1$

$$0 < y_n < 1, \quad \Delta y_n < 0,$$

and define

$$n_1 = n_0 + p.$$

In view of (4), there exists $\Lambda > 0$ such that for any $n \geq n_1$

$$\sum_{i=n-p+1}^{n-1} b_i \leq \Lambda. \tag{21}$$

Without loss of generality, we can suppose

$$\Lambda < |y_{n_1}^{[1]}|, \quad (22)$$

where $y_{n_1}^{[1]} = a_{n_1+p-1}\Phi(\Delta y_{n_1})$. Moreover, let $H > 1$ be a positive constant such that

$$\Phi(H) \geq \frac{|y_{n_1}^{[1]}|}{|y_{n_1}^{[1]}| - \Lambda}. \quad (23)$$

Let \mathbb{X}_1 be the Fréchet space of the real sequences defined for $n \geq n_1$, endowed with the topology of convergence on \mathbb{N}_{n_1} . Define the subset Ω_1 of \mathbb{X}_1

$$\Omega_1 = \{u \in \mathbb{X}_1 : y_{n-p+1} \leq u_n \leq Hy_{n-p+1}\}$$

and for any $u \in \Omega_1$ consider the BVP

$$\begin{cases} \Delta(a_n\Phi(\Delta w_n)) + b_n\Phi(u_{n+p}) = 0, & n \geq n_1 \\ w_{n_1}^{[1]} = y_{n_1}^{[1]}, \quad \lim_n w_n = 0, \end{cases} \quad (24)$$

where $w^{[1]}$ and $y^{[1]}$ denote the quasidifferences of w and y , respectively, that is the sequences

$$w_n^{[1]} = a_n\Phi(\Delta w_n), \quad y_n^{[1]} = a_{n+p-1}\Phi(\Delta y_n). \quad (25)$$

As before, for any $u \in \Omega_1$ the BVP (24) has a unique solution w . Thus, let \mathcal{T} be the operator which associates to any $u \in \Omega_1$ the unique solution w of (24). Summing the equation in (24) and using (25) we get

$$w_n^{[1]} = y_{n_1}^{[1]} - \sum_{k=n_1}^{n-1} b_k\Phi(u_{k+p}) \leq y_{n_1}^{[1]} - \sum_{k=n_1}^{n-1} b_k\Phi(y_{k+1}) = y_n^{[1]}. \quad (26)$$

Since $y^{[1]}$ is decreasing for $n \geq n_0$ and $p > 1$, we have $y_n^{[1]} \leq y_{n-p+1}^{[1]}$ for $n \geq n_1$, and from (26) we obtain

$$w_n^{[1]} \leq y_{n-p+1}^{[1]} \quad (27)$$

i.e.

$$\Delta w_n \leq \Delta y_{n-p+1},$$

which implies

$$w_n \geq y_{n-p+1}, \quad (28)$$

since $\lim_i w_i = \lim_i y_i = 0$.

Now, let us prove that

$$w_n \leq H y_{n-p+1} \quad . \quad (29)$$

Summing the equation in (24) we get

$$w_n^{[1]} = y_{n_1}^{[1]} - \sum_{k=n_1}^{n-1} b_k \Phi(u_{k+p}) \geq y_{n_1}^{[1]} - \Phi(H) \sum_{k=n_1}^{n-1} b_k \Phi(y_{k+1}),$$

that is

$$\begin{aligned} w_n^{[1]} &\geq y_{n_1}^{[1]} + \Phi(H) (y_n^{[1]} - y_{n_1}^{[1]}) \\ &= \Phi(H) y_{n-p+1}^{[1]} + \left(y_n^{[1]} - y_{n-p+1}^{[1]} \right) \Phi(H) + (1 - \Phi(H)) y_{n_1}^{[1]}. \end{aligned} \quad (30)$$

From (H1) and (21) we have

$$y_n^{[1]} - y_{n-p+1}^{[1]} = - \sum_{i=n-p+1}^{n-1} b_i \Phi(y_{i+1}) \geq -\Lambda.$$

Thus, from (30) we obtain

$$\begin{aligned} w_n^{[1]} &\geq \Phi(H) y_{n-p+1}^{[1]} - \Lambda \Phi(H) + y_{n_1}^{[1]} - \Phi(H) y_{n_1}^{[1]} \\ &= \Phi(H) y_{n-p+1}^{[1]} + (|y_{n_1}^{[1]}| - \Lambda) \Phi(H) + y_{n_1}^{[1]}, \end{aligned}$$

or, in view of (22) and (23),

$$w_n^{[1]} \geq \Phi(H) y_{n-p+1}^{[1]} \quad (31)$$

i.e.,

$$\Delta w_n \geq H \Delta y_{n-p+1}.$$

Since $\lim_i w_i = \lim_i y_i = 0$, from here we get (29). Hence, in virtue of (28) and (29), the operator \mathcal{T} maps Ω_1 into itself, i.e.

$$\mathcal{T}(\Omega_1) \subset \Omega_1.$$

Using the same argument to the one given in the sufficiency part, denote by S_1 the boundary conditions in (24). Applying Theorem 4 with $S_C = S_1 \cap \Omega_1$, we get that the operator \mathcal{T} has a fixed point $\bar{w} \in S_1 \cap \Omega_1$. Clearly the sequence \bar{w} is a solution of (P) and $\lim_n \bar{w}_n = 0$. Since $\bar{w} \in \mathcal{T}(\Omega_1)$, from (27) and (31), we get

$$Hy_{n-p+1}^{[1]} \leq \bar{w}_n^{[1]} \leq y_{n-p+1}^{[1]}$$

and so $\lim_n \bar{w}_n^{[1]} = -\infty$.

Hence \bar{w} is an intermediate solution of (P) and the proof is complete. \square

4 Suggestions and examples

The following example illustrates Theorem 2 and Corollary 1.

Example 1. Consider the difference equation with advanced argument

$$\Delta((n-p+1)^{1+\alpha}\Phi(\Delta x_n)) + \gamma\Phi(x_{n+p}) = 0, \quad n \geq p \geq 2, \quad (32)$$

where γ is a positive constant. Using Theorem 2 and Corollary 1, it is easy to show that (32) has intermediate solutions if and only if

$$0 < \gamma \leq \left(\frac{1}{1+\alpha}\right)^{\alpha+1}. \quad (33)$$

Indeed, consider the half-linear equation

$$\Delta(n^{1+\alpha}\Phi(\Delta x_n)) + \gamma\Phi(x_{n+1}) = 0. \quad (34)$$

A standard calculation shows that (1) is satisfied. Moreover, using the change of variable

$$y_n = n^{1+\alpha}\Phi(\Delta x_n) \quad (35)$$

the equation (34) is transformed into the generalized discrete Euler equation

$$\Delta(\Phi^*(\Delta y_n)) + \gamma^{1/\alpha} \left(\frac{1}{n+1}\right)^{(1+\alpha)/\alpha} \Phi^*(y_{n+1}) = 0,$$

which is nonoscillatory if γ satisfies (33) and oscillatory if

$$\gamma > \left(\frac{1}{1+\alpha}\right)^{\alpha+1}, \quad (36)$$

see, e.g., [18]. Since the transformation (35) maintains the oscillatory behavior, see, e.g., [11], we get that (34) is nonoscillatory if and only if (33) is satisfied. Moreover, as $a_n = n^{1+\alpha}$, $b_n = \gamma$, we have

$$\sum_{n=p}^{\infty} \Phi^* \left(\frac{1}{a_{n+p}} \sum_{k=p}^n b_k \right) = \gamma^{1/\alpha} \sum_{n=p}^{\infty} \left(\frac{n}{n+p} \right)^{1/\alpha} \frac{1}{n+p} = \infty$$

and so the condition (5) is satisfied. Hence, from Corollary 1– i_1) equation (32) has intermediate solutions if (33) is satisfied. When (36) holds, as claimed, the half-linear equation (34) is oscillatory and it does not admit intermediate solutions. Thus, from Theorem 2, equation (32) does not have intermediate solutions, too.

The existence of intermediate solutions for (P) does not depend on condition (4), as the following example shows.

Example 2. Consider the difference equation with advanced argument

$$\Delta((n+1)! \Delta x_n) + (n+p)! x_{n+p} = 0, \quad p \geq 2. \quad (37)$$

A direct computation shows that $x = \{1/n!\}$ is an intermediate solution of (37). Nevertheless, for (37), assumption (4) is not verified and Theorem 2 cannot be applied. Hence, it is an open problem if Theorem 2 continues to hold when condition (4) failed.

Example 1 suggests the following two comparison results.

Corollary 2. *Assume (1) and (4). Suppose that*

$$a_n \geq n^{1+\alpha}, \quad \text{and} \quad \sum_{n=N}^{\infty} \Phi^* \left(\frac{n}{a_{n+p}} \right) = \infty,$$

where $N \geq p$. Set $L = \sup_{n \geq N} b_n$. If b is bounded away from zero and

$$L < \left(\frac{1}{1+\alpha} \right)^{1+\alpha},$$

then the equation

$$\Delta((n-p+1)^{1+\alpha} \Phi(\Delta x_n)) + b_n \Phi(x_{n+p}) = 0, \quad n \geq p \geq 2, \quad (38)$$

has intermediate solutions.

Proof. Consider the equation

$$\Delta(n^{1+\alpha}\Phi(\Delta x_n)) + L\Phi(x_{n+1}) = 0. \quad (39)$$

Reasoning as in Example 1, we get that (39) is nonoscillatory. Hence, in virtue of the Sturm comparison theorem, see, e.g., [12, Chapter 8.2], the half-linear equation (H) is nonoscillatory. Moreover, since b is bounded away from zero, there exists $\varepsilon > 0$ such that $b_n \geq \varepsilon$ for any $n \geq 1$. Hence

$$\sum_{n=N}^{\infty} \Phi^* \left(\frac{1}{a_{n+1}} \sum_{k=1}^n b_k \right) \geq \Phi^*(\varepsilon) \sum_{n=N}^{\infty} \Phi^* \left(\frac{n}{a_{n+1}} \right) = \infty,$$

which implies (5). Hence, applying Corollary 1– i_1) to the equation (38), we get the assertion. \square

Corollary 3. *Assume (1) and (4). Moreover, suppose that for $n \geq N \geq p$,*

$$a_n \leq n^{1+\alpha}.$$

Set $\ell = \inf_{n \geq N} b_n$. If

$$\ell > \left(\frac{1}{1+\alpha} \right)^{1+\alpha},$$

then the equation (38) does not have intermediate solutions.

Proof. The argument is similar to the one given in Corollary 2. Consider the half-linear equation

$$\Delta(n^{1+\alpha}\Phi(\Delta x_n)) + \ell \Phi(x_{n+1}) = 0. \quad (40)$$

Reasoning as before, we get that (40) is oscillatory. Hence, in virtue of the Sturm comparison theorem, also (H) is oscillatory, and so (H) does not have intermediate solutions. Thus, applying Theorem 2 to the equation (H), we obtain the assertion. \square

Some suggestions for future researches are in order.

1) As claimed, Theorem 3 from [13] gives a very general fixed point result, which is based on a continuation principle in a Hausdorff locally convex space. Further, in [13] the solvability to certain BVPs in the continuous case is also given. It should be interesting to establish corresponding discrete versions of

these existence results, especially for [13, Theorem 2.2], which deals with a scalar equation. It could be useful for studying discrete BVPs when it is hard to find an appropriate bounded closed set Ω which is mapped into itself, as, for instance occurs for intermediate solutions to Emden-Fowler superlinear discrete equation

$$\Delta(a_n|\Delta x_n|^\alpha \operatorname{sgn} \Delta x_n) + b_n|x_{n+1}|^\beta \operatorname{sgn} x_{n+1} = 0, \quad \alpha < \beta,$$

see, e.g., [6, Section V.].

2) The proof of Theorem 2 does not work if $p \leq 0$. Indeed, in this case the half-linear equation (H1) is not defined, due to the shift in the weight coefficient a of the discrete operator

$$(Dx)_n = \Delta(a_n\Phi(\Delta x_n)).$$

Further, when $p < 0$, the sequence $\{u_{n+p}\}$ in BVP (24) has to be defined not only for $n \geq n_1$, but also for any $i \geq n_1 + p$. Consequently, when (P) is an equation with delay, the solvability of (3) requires a different approach, which will be presented in a forthcoming paper [10].

3) Theorem 2 establishes a comparison between the asymptotic decay of intermediate solutions of (P) with the one of an associated half-linear equation. Recently, in some particular cases, a precise asymptotic analysis of intermediate solutions for discrete half-linear equations has been made in the framework of regular variation, see [16]. It should be interesting to apply this approach for obtaining a precise description of the asymptotic behavior also for intermediate solutions of the equations with advanced argument.

References

- [1] Agarwal R.P, O'Regan D., *A fixed-point approach for nonlinear discrete boundary value problems*, Comput. Math. Appl. **36** (1998), 115-121.
- [2] Agarwal R.P, Grace S.R., O'Regan D., *Oscillation Theory for Second Order Linear, Half-Linear, Superlinear and Sublinear Dynamic Equations*. Kluwer Acad. Publ. G, Dordrecht, 2003.
- [3] Agarwal, R.P., Manojlović J., *On the existence and the asymptotic behavior of nonoscillatory solutions of second order quasilinear difference equations*. Funkcial. Ekvac. **56** (2013), 81-109.

- [4] Agarwal R.P., O'Regan D., *Infinite Interval Problems for Differential, Difference and Integral Equations*, Kluwer Academic Publishers, Dordrecht, 2001.
- [5] Cecchi M., Došlá Z., Marini M., *Intermediate solutions for Emden-Fowler type equations: continuous versus discrete*. *Advances Dynam. Systems Appl.* **3** (2008), 161-176.
- [6] Cecchi M., Došlá Z., Marini M., *On oscillation and nonoscillation properties of Emden-Fowler difference equations*, *Cent. Eur. J. Math.*, **7** (2009), 322-334.
- [7] Cecchi M., Došlá Z., Marini M., *Intermediate solutions for nonlinear difference equations with p -Laplacian*. *Advanced Studies in Pure Mathematics* **53** (2009), 33-40.
- [8] Cecchi M., Furi M., Marini M., *On continuity and compactness of some nonlinear operators associated with differential equations in noncompact intervals*, *Nonlinear Anal.*, **9** (1985), 171-180.
- [9] Díaz J.I., *Nonlinear partial differential equations and free boundaries, Vol.I: Elliptic equations*. Pitman Advanced Publ., **106** (1985).
- [10] Došlá Z., Marini M., Matucci S., *Decaying solutions for discrete Emden-Fowler type equations with deviating argument*, preprint.
- [11] Došlý O., Řehák P., *Nonoscillatory criteria for half-linear second order difference equations*, *Computers Math. Appl.* **42** (2001), 453-464.
- [12] Došlý O., Řehák P., *Half-linear Differential Equations*, North-Holland, Mathematics Studies **202**, Elsevier Sci. B.V., Amsterdam, 2005.
- [13] Furi M., Pera M.P., *A continuation method on locally convex spaces and applications to ordinary differential equations on noncompact intervals*, *Ann. Polon. Math.* **47** (1987), 331-346.
- [14] Kamo K., Usami H., *Asymptotic forms of weakly increasing positive solutions for quasilinear ordinary differential equations*. *Electronic J. Differential Equations* **126** (2007), 1-12.

- [15] Marini, M., Matucci, S., Řehák, P., *Boundary value problems for functional difference equations on infinite intervals*, Adv. Difference Equ. **2006**, Art. 31283, 1-14.
- [16] Matucci S., Řehák P., *Rapidly varying decreasing solutions of half-linear difference equations*, Math. Comput. Modelling **49** (2009), 1692-1699.
- [17] O'Regan, D., *Existence theory for nonlinear ordinary differential equations*, Mathematics and its Applications, **398**. Kluwer Academic Publishers Group, Dordrecht, 1997.
- [18] Řehák P., *Comparison theorems and strong oscillation in the half-linear discrete oscillation theory*, Rocky Mt. J. Math. **33** (2003), 333-352.

Authors' addresses:

Zuzana Došlá, Department of Mathematics and Statistics, Masaryk University, Kotlářská 2, CZ-61137 Brno, Czech Republic.

E-mail: `dosla@math.muni.cz`

Mauro Marini, Department of Mathematics and Computer Science "Ulisse Dini", University of Florence, Via di S. Marta 3, I-50139 Florence, Italy.

E-mail: `mauro.marini@unifi.it`

Serena Matucci, Department of Mathematics and Computer Science "Ulisse Dini", University of Florence, Via di S. Marta 3, I-50139 Florence, Italy.

E-mail: `serena.matucci@unifi.it`