

## Supramolecular Forces and Their Interplay in Stabilizing Complexes of Organic Anions: Tuning Binding Selectivity in Water. †

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

Matteo Savastano,<sup>a,c</sup> Carla Bazzicalupi,<sup>a</sup> Celeste García-Gallarín,<sup>b</sup> Maria Dolores López de la Torre,<sup>b</sup> Antonio Bianchi,<sup>\*a</sup> Manuel Melguizo<sup>\*b</sup>

How do different supramolecular forces contribute to the stabilization of complexes of organic anions in water? Oftentimes, when debating such a theme we refer to broad concepts like positive or negative cooperative effects: the focus of the present work is rather on their interplay, i.e. on the way different kinds of stabilizing interactions (salt bridges, H-bonds, anion- $\pi$  interaction,  $\pi$ - $\pi$  stacking, solvent effect, etc.) dialogue among themselves. What happens if we tune the strengths of salt bridges by altering the basicity of the anion? What if we change the geometry of the charged group? How does shifting towards more hydrophilic or hydrophobic anions impact the stability of complexes in water? What happens in the solid state? Will aromatic anions go for a  $\pi$ - $\pi$  stacking or an anion- $\pi$  interaction mode and do they all behave in the same manner? Does the host/guest size make any difference? What if we play with regiochemistry: will one of the isomers be selectively recognized? Here we present a case study featuring the tetrazine-based ligands L1 and L2 and a series of selected organic anions: potentiometric, NMR, XRD data and in silico simulations being employed to render such a complex picture.

### Introduction

Anion coordination in solution revolves around a combination of supramolecular forces, selected among a shortlist by incorporating the required structural features within the receptor: although individually weak, such interactions can collectively furnish enough stabilization to afford polyfunctional ligands capable of strong and selective anion binding.<sup>1,2</sup>

We should, however, keep in mind two facts. First, supramolecular forces are not equally represented in the panorama of anion receptors, hydrogen bond easily taking the lion's share, being among the most stabilizing and directional of intermolecular forces and widely recognized for that, while other forces, among which anion- $\pi$  interaction<sup>3</sup> and halogen bond,<sup>4</sup> are still struggling for the endorsement by part of the scientific community. In second instance, we have to be aware that classifying is an inborn mean of rationalization of the human mind: when we say, for example, that a receptor is hydrogen bond-based, what we mean is that its anion complexes are stabilized by an interplay of supramolecular forces, among which hydrogen bonding plays a major role. This goes beyond semantics, as the importance of such interplay, an

exquisitely supramolecular topic, is receiving an across-the-board growing recognition from researchers working in the most diverse fields of chemistry, spanning from structural biology, with implications for both living beings<sup>5</sup> and new synthetic analogues of biomolecules,<sup>6</sup> up to material chemistry<sup>7</sup> and crystal engineering.<sup>8</sup>

We have recently shown that protonated forms of the tetrazine-based receptors L1 and L2 (Figure 1), decorated with morpholine pendants of variable lengths, bind polyatomic inorganic anions of several different geometries both in aqueous solution and in the solid state.<sup>9</sup> In the case of L2, also binding of spherical halide anions,<sup>10</sup> of some linear polyhalogen derivatives<sup>11</sup> as well as of pseudohalogen anions of the same geometry,<sup>12</sup> has been reported. Crystal structures substantiated an interplay of supramolecular forces, mainly involving strong anion- $\pi$  interactions, salt bridges and CH...anion contacts, while the solution studies highlighted the crucial role of solvent effect in promoting the association phenomena.

The present paper tackles the bigger picture, i.e. the interplay of different supramolecular forces in anion binding in aqueous solution, by inspecting the ligands interactions with an ad hoc series of organic anions.

The new list of studied anionic species was elaborated with the explicit intention to play with the following parameters: i) the basicity of the anion; ii) solvation/capability to establish  $\pi$ - $\pi$  stacking interactions; iii) stereochemistry of the interacting groups; iv) host-guest mutual size.

<sup>a</sup> Department of Chemistry "Ugo Schiff", University of Florence, Via della Lastruccia 3, 50019 Sesto Fiorentino, Italy. E-mail: [antonio.bianchi@unifi.it](mailto:antonio.bianchi@unifi.it)

<sup>b</sup> Department of Inorganic and Organic Chemistry, University of Jaén, 23071 Jaén, Spain. E-mail: [mmelqui@ujaen.es](mailto:mmelqui@ujaen.es)

<sup>c</sup> CSGI, University of Florence, Via della Lastruccia 3, 50019 Sesto Fiorentino, Italy.

† Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x




Figure 1. The tetrazine-based ligands L1 and L2.

The inorganic anions investigated so far, with the obvious exception of  $F^-$  (and the fringe one of  $SCN^-$ ), are all conjugate bases of strong acids. Since the strength of salt bridges and hydrogen bonds in general depends on the relative basicity of the interacting partners, peaking when their  $\Delta pK_a$  approaches 0,<sup>13</sup> no discrimination was possible on this basis in previous cases. Basicity of L1 ( $pK_{a1} = 4.45(3)$ , Table S1) suggested acetate ( $pK_a = 4.51(1)$ , Table S1) as a suitable candidate, together with its essentially non-basic sulfonate analogue, methanesulfonate. Acetate incidentally appears in the Hofmeister series as one of the most hydrated monovalent organic anions. With manifest solvent effects already observed in our preliminary study,<sup>9</sup> and bearing in mind that delocalized aromatic anions are less solvated, we also studied the aromatic analogues benzoate and benzenesulfonate. This offers not only the possibility to tune the strength of solvent effects, but also adds to the mix the possibility of  $\pi$ - $\pi$  stacking interactions, partially moving the complexation phenomena towards the solvent-driven association of organic molecules in polar solvents. In this framework, phthalate and isophthalate, provided the chance of checking the effect of a second anionic group and the influence of the mutual disposition of the divergent binding sites on the anion. This is expected to be a particularly prominent effect for our ligands, as the *s*-tetrazine core in water is documented not just to be poorly solvated, but to be hosted in a clathrate-like water molecule cage without relevant solvent-solute hydrogen bonding.<sup>14</sup> Finally, effect of host-guest size was evaluated varying the length of the aliphatic spacer in the ligand, affecting also flexibility and preorganization,<sup>15</sup> to properly prosecute the evaluation carried out in previous works.

The core of the current paper revolves around the study of anion complexes formation in aqueous solution, investigated mainly through potentiometric titrations. The use of XRD crystal structures of the complexes to identify and assess the role of the key supramolecular interactions in the solid state, consolidated in our previous papers, was once more a precious tool, allowing to draw a parallel with the solution studies. However, due to the coexistence of a large number of complex species (differing both for protonation state and stoichiometry), <sup>1</sup>H NMR experiments and the extensive use of *in silico* simulations have been exploited to bridge the gap between XRD data and potentiometric measurements, allowing for a proper evaluation of the interplay of the different supramolecular forces in aqueous solution.

## Results and Discussion

### Crystal Structures

Among all the possible substrates, only hydrogen phthalate, hydrogen isophthalate and benzenesulfonate afforded anion complexes single crystals of sufficient quality for XRD analysis. Although each of the solved structures is described in detail below, it is interesting to take a synoptic view at them, noticing how all of the expected supramolecular interactions are indeed found in the solid state, but their relevance and overall interplay is different in each case: such synopsis is presented in Table 1. As evidenced by Table 1, all sort of situations are encountered: from an essentially cooperative behaviour between salt bridges formation and  $\pi$ -forces (HPhthalate<sup>-</sup>) to the predominance of charge-charge interactions ( $C_6H_5SO_3^-$  (b)), passing from intermediate cases where salt bridges and  $\pi$  interactions, either anion- $\pi$  ( $C_6H_5SO_3^-$  (a)) or  $\pi$ - $\pi$  stacking (HIsophthalate<sup>-</sup>), establish an active dialogue among themselves. In-depth analysis of the solved crystal structures is provided in the dedicated sections below.

Table 1. Breakdown of the main interactions observed in the crystal structures of anion complexes.

Interaction Type	Anionic Guest in the Crystal Structure			
	HPh <sup>t</sup>	HIPh <sup>t</sup>	$C_6H_5SO_3^-$ (a) <sup>a</sup>	$C_6H_5SO_3^-$ (b) <sup>a</sup>
Anion- $\pi$	+	-	+	-
$\pi$ -Stacking	-	+	-	-
Direct Salt Bridge N-H <sup>+</sup> ...O <sup>b</sup>	+	-	-	+
Bridging Salt Bridge N-H <sup>+</sup> ...O <sup>c</sup>	-	+	+	-
C-H...O Contacts	+	-	+	+

<sup>a</sup> Two non-symmetry related anions (a and b) are found in the crystal structure. Refer to Figure 4 and dedicated discussion section.


<sup>b</sup> Salt bridge was labelled "direct" if it involves the same ligand molecule with which the anion is forming  $\pi$ - $\pi$  or anion- $\pi$  contacts (if any).

<sup>c</sup> Salt bridge was labelled "bridging" if it involves a ligand molecule different from the one with which the anion is forming  $\pi$ - $\pi$  or anion- $\pi$  contacts.

### Crystal Structure of (H<sub>2</sub>L2)(HPhthalate)<sub>2</sub>·2H<sub>2</sub>O

In the phthalate salt, the ligand assumes a chair-type conformation with the morpholine pendant arms placed in trans position with respect to the tetrazine ring (Figure 2). The chair-type conformation was previously observed by XRD analysis of the complexes formed by the ligand with several anions, such as chloride, bromide, nitrate, thiocyanate, perchlorate and hexafluorophosphate.<sup>9,10,12</sup> In all these structures, as well as in the present one, the ligand is centrosymmetric and interacts with two symmetry related anions via  $NH^+ \cdots X$  salt bridges and additional anion- $\pi$  interactions. Actually, the monoprotonated phthalate is almost completely flat, the acidic hydrogen being shared between two oxygens from the two  $-CO_2$  groups. One  $-CO_2$  oxygen is in contact at the same time with the protonated morpholine nitrogen ( $N1 \cdots O3$  distance 2.740(2) Å,  $N1H1 \cdots O3$  distance

1.72(3) Å) and with the tetrazine ring, the geometric parameters defining the anion- $\pi$  interaction being as follows: O3...ring-centroid distance = 3.01 Å, O3...ring-plane distance = 2.97 Å, offset with respect to the normal to plane = 0.49 Å. The external oxygen from the other -CO<sub>2</sub> group is in contact with the aliphatic hydrogen atoms of the morpholine from a symmetry related ligand and contributes to the overall crystal packing stabilization. Interestingly, no mutual  $\pi$ - $\pi$  stacking interactions involve the ligand or the phthalate aromatic ring, while additional contributions to the overall stability are provided by the disordered solvent water molecules.




**Figure 2.** ORTEP drawing of the (H<sub>2</sub>L)(HPhthalate)<sub>2</sub> complex in the (H<sub>2</sub>L)(HPhthalate)<sub>2</sub>·2H<sub>2</sub>O crystal structure. Selected contacts are shown.

### Crystal Structure of (H<sub>2</sub>L)(Hisophthalate)<sub>2</sub>

Differently from the phthalate crystal structure, in the isophthalate salt the ligand assumes a centrosymmetric planar conformation (Figure 3). Both sides of the tetrazine ring are in contact via offset  $\pi$ -stacking with an isophthalate protonated anion. The anions and the tetrazine rings are almost coplanar and placed at 3.19 Å from one another, while the ring centroids are offset by 1.36 Å (Figure 3). Moreover, each anion accepts a salt bridge from the protonated morpholine nitrogen of a symmetry related ligand molecule (N3...O4 distance 2.666(2) Å, N3H1'...O4 distance 1.729(2) Å) and gives head-to-tail O-H...O hydrogen-bonds with the adjacent anions in the crystal lattice (O3...O5' 2.530(2) Å, O3-H2...O5' 1.55(5) Å), forming an infinite zig-zag chain (Figure 3b).

Interestingly, on the basis of the intermolecular potentials evaluated by the Uni force field implemented in Mercury,<sup>16,17</sup> the most stabilizing interactions in the crystal packing are the  $\pi$ - $\pi$  stacking ones (about 54 kJ/mol), while a less extent of stabilization is due to the ligand/anion couples linked by the NH<sup>+</sup>...O salt bridges between morpholine nitrogen and anion (about 30 kJ/mol) and, to an even lower extent, to the head-to-tail H-bonded anions in the anions' zigzag chain (about 25 kJ/mol). Following these data, it could be hypothesized that the  $\pi$ -stacking could be an important stabilizing contribution even in solution.



**Figure 3.** ORTEP drawing of the (H<sub>2</sub>L)(Hisophthalate)<sub>2</sub> crystal structure. (a) lateral view of the complex; (b) details of the crystal packing. Selected contacts shown.


### Crystal Structure of H<sub>2</sub>L(C<sub>6</sub>H<sub>5</sub>SO<sub>3</sub>)<sub>2</sub>·H<sub>2</sub>O

As shown in Figure 4, in this case the ligand assumes a planar conformation. Two symmetry non-equivalent benzenesulfonate anions are present in this structure. Both form a salt bridge with the protonated nitrogen atom of the ligand, however only one exhibits a marked interaction with the tetrazine ring, bringing one of the sulfonate oxygen atoms at 3.440 Å from the ring centroid, the interaction being particularly strong with one of the tetrazine nitrogen atoms (O23...N2 3.066 Å, O-N-centroid angle 93.98°) (Figure 4). It is to be underlined that, as shown in Figure 4, these two interactions, the salt bridge and ring contact, are exerted by the same benzenesulfonate ion with two distinct, symmetry related, ligand molecules. The other two oxygen atoms of the sulfonate group are found either giving rise, as said above, to a bridge-bond interaction with a distinct, symmetry related ligand molecule (N5'...O21 2.700(3) Å, N5'H5'...O21 1.739(2) Å) or multiple CH...O contacts.

The second anion, required to guarantee the electroneutrality of the system, but probably too bulky to be accommodated in proximity of the tetrazine ring, is found mainly interacting through hydrogen bonds with the co-crystallized water molecule (O13...OW1 2.892(3) Å, OW1-H1...O13 167(4)°) and salt bridge with the protonated nitrogen atom of the morpholine moiety (N6...O13 2.733(3) Å, N6H6...O13 1.773(2) Å). Several CH...anion contacts are also recognizable.

As evidenced in Figure 4,  $\pi$ - $\pi$  stacking interaction offers a modest contribution to the overall stability of the structure, the two closest carbon atoms of the benzene ring being in the 3.5-3.8 Å range from the tetrazine. The relative positions of the two rings, probably imposed by their sizes, by the non-planarity of the sulfonate group, by the geometric constraints of the hydrogen bond network and by the preference of the tetrazine for the anionic head rather than for the aromatic portion of the anion, produce a 13.01° angle between the ring planes, limiting

their superimposition. The argument is much more valid for the second benzenesulfonate anion, with an angle in-between the ring planes of  $53.34^\circ$ , which does not give rise to contacts with the tetrazine ring.



**Figure 4.** ORTEP drawing and details of the crystal packing of the  $\text{H}_2\text{L}_2(\text{C}_6\text{H}_5\text{SO}_3)_2 \cdot \text{H}_2\text{O}$  crystal structure. Selected contacts are shown.

### Anion Binding in Solution

As in the case of their inorganic counterparts, the investigated series of organic anions does give rise to detectable interactions with our receptors in water. Stability constants of the anion complexes, obtained through potentiometric titrations performed in 0.10 M  $\text{NMe}_4\text{Cl}$  at  $298.1 \pm 0.1$  K, are reported in Table 2 for L1 and Table 3 for L2.  $^1\text{H}$  NMR experiments were also performed at selected pH values, chosen to be mostly representative of a single complex species in solution: shifting of both anions and ligands signals, demonstrating complex formation, were observed. Complexation induced shift (CIS) values are reported in Table 4 and Table 5 for L1 and L2 respectively.

Equilibrium constants for the formation of anion complexes in solution were obtained through computer-aided analysis of potentiometric titration curves. The HYPERQUAD<sup>18</sup> analysis software furnishes complexes stability constants according to the general equilibria  $j\text{A}^{n-} + k\text{L} + m\text{H}^+ = (\text{A})_j(\text{L})_k(\text{H})_m^{(m-jn)+}$ . Such equilibria are indicative of complex stoichiometry but do not provide any insight about the location of the  $m$  protons within each complex species. Such ambiguity is not due to the data treatment, but inborn to the potentiometric experiment: the glass electrode probes only the point by point free  $\text{H}^+$  concentration in solution, so that, as long as the same  $m$  number of protons are present in a given complex species, their localization cannot affect the shape of the titration curve. Other studies showed that the formation of anion complexes tends not to modify the protonation pattern of ligands, although a modest general shift of NMR signals, corresponding to the increase of ligands basicity brought about by anion complexation, has been observed.<sup>19</sup> Accordingly, the location of protons in the complexes was assumed to be generally regulated by the basicity of the interacting species and the relevant stability constants were calculated following this rule. In those cases where matching of protonation constants between ligand and anion required a closer inspection, in silico simulations were employed: Tables 2 and 3 list complexes

stability constants compiled through the double-criterion of basicity of the isolated components and higher computational stability of the displayed complexes over their possible tautomers. However, possible coexistence in solution of scarce amounts of complex tautomers alongside the most stable form (presented in Table 2 and 3) cannot be completely ruled-out. Since we want to deal with the interplay of different supramolecular forces in solution, whose net sum is mirrored by complexes stability constants, it is important to abstract from the individual data looking for general trends. Conditional stability constants are one of the abstraction tools at our disposal. They can be calculated as a function of pH in the form  $K_{\text{cond}} = \frac{\sum[\text{H}_{i+j}\text{LA}]}{(\sum[\text{H}_i\text{L}] \times \sum[\text{H}_j\text{A}])}$ , where  $i$  and  $j$  are the number of acidic protons on the ligand and on the anion, respectively, allowing direct comparison of the affinity of each ligand for the different anions eluding the problems connected with the different speciation of the systems.<sup>20</sup> As anticipated from the beginning, two main stability trends are expected: salt bridge strength is expected to increase with the basicity of the anion (carboxylates > sulfonates), while stacking forces and solvent effects are anticipated to be more prominent for hydrophobic substrates (aromatic > aliphatic); overall, for the monocharged anion series, this should result in the order of stability benzoate > acetate > benzenesulfonate > methanesulfonate. Examining Figure 5 such trend is indeed found for both ligands. While the difference between sulfonates and carboxylates complexes is

**Table 2.** Equilibrium constants (log  $K$ ) for L1 anion complexes formation determined at  $298.1 \pm 0.1$  K in 0.1 M  $\text{NMe}_4\text{Cl}$  aqueous solution. Figures in parentheses are standard deviations on the last significant figure.

Equilibrium	log $K$
$\text{HL1}^+ + \text{CH}_3\text{SO}_3^- = [\text{HL1}(\text{CH}_3\text{SO}_3)]$	2.21(9)
$\text{H}_2\text{L1}^{2+} + \text{CH}_3\text{SO}_3^- = [\text{H}_2\text{L1}(\text{CH}_3\text{SO}_3)]^+$	1.6(2)
$\text{HL1}^+ + \text{C}_6\text{H}_5\text{SO}_3^- = [\text{HL1}(\text{C}_6\text{H}_5\text{SO}_3)]$	2.49(5)
$\text{H}_2\text{L1}^{2+} + \text{C}_6\text{H}_5\text{SO}_3^- = [\text{H}_2\text{L1}(\text{C}_6\text{H}_5\text{SO}_3)]^+$	2.48(5)
$\text{L1} + \text{CH}_3\text{COO}^- = [\text{L1}(\text{CH}_3\text{COO})]^-$	2.71(8)
$\text{L1} + \text{CH}_3\text{COOH} = [\text{L1}(\text{CH}_3\text{COOH})]$	3.27(5)
$\text{HL1}^+ + \text{CH}_3\text{COOH} = [\text{HL1}(\text{CH}_3\text{COOH})]^+$	3.09(3)
$\text{L1} + \text{C}_6\text{H}_5\text{COO}^- = [\text{L1}(\text{C}_6\text{H}_5\text{COO})]^-$	3.02(1)
$\text{HL1}^+ + \text{C}_6\text{H}_5\text{COO}^- = [\text{HL1}(\text{C}_6\text{H}_5\text{COO})]$	3.16(3)
$\text{H}_2\text{L1}^{2+} + \text{C}_6\text{H}_5\text{COO}^- = [\text{H}_2\text{L1}(\text{C}_6\text{H}_5\text{COO})]^+$	2.95(4)
$\text{L1} + \text{Pht}^{2-} = [\text{L1}(\text{Pht})]^{2-}$	2.88(2)
$\text{L1} + \text{HPht}^- = [\text{L1}(\text{HPht})]^-$	2.87(3)
$\text{HL1}^+ + \text{HPht}^- = [\text{HL1}(\text{HPht})]$	2.56(3)
$\text{H}_2\text{L1}^{2+} + \text{HPht}^- = [\text{H}_2\text{L1}(\text{HPht})]^+$	2.60(3)
$\text{HL1}^+ + \text{IPht}^{2-} = [\text{HL1}(\text{IPht})]^{2-}$	3.40(5)
$\text{HL1}^+ + \text{HIPht}^- = [\text{HL1}(\text{HIPht})]$	2.73(8)
$\text{H}_2\text{L1}^{2+} + \text{HIPht}^- = [\text{H}_2\text{L1}(\text{HIPht})]^+$	3.45(5)

**Table 3.** Equilibrium constants (log *K*) for L2 anion complexes formation determined at 298.1 ± 0.1 K in 0.1 M NMe<sub>4</sub>Cl aqueous solution. Figures in parentheses are standard deviations on the last significant figure.

Equilibrium	log <i>K</i>
HL2 <sup>+</sup> + CH <sub>3</sub> SO <sub>3</sub> <sup>-</sup> = [HL2(CH <sub>3</sub> SO <sub>3</sub> )]	1.41(8)
H <sub>2</sub> L2 <sup>2+</sup> + CH <sub>3</sub> SO <sub>3</sub> <sup>-</sup> = [H <sub>2</sub> L2(CH <sub>3</sub> SO <sub>3</sub> ) <sup>+</sup>	1.8(3)
HL2 <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> SO <sub>3</sub> <sup>-</sup> = [HL2(C <sub>6</sub> H <sub>5</sub> SO <sub>3</sub> )]	1.44(6)
H <sub>2</sub> L2 <sup>2+</sup> + C <sub>6</sub> H <sub>5</sub> SO <sub>3</sub> <sup>-</sup> = [H <sub>2</sub> L2(C <sub>6</sub> H <sub>5</sub> SO <sub>3</sub> ) <sup>+</sup>	1.84(2)
HL2 <sup>+</sup> + CH <sub>3</sub> COO <sup>-</sup> = [HL2(CH <sub>3</sub> COO)]	2.33(9)
H <sub>2</sub> L2 <sup>2+</sup> + CH <sub>3</sub> COO <sup>-</sup> = [H <sub>2</sub> L2(CH <sub>3</sub> COO)] <sup>+</sup>	2.0(1)
H <sub>2</sub> L2 <sup>2+</sup> + CH <sub>3</sub> COOH = [H <sub>2</sub> L2(CH <sub>3</sub> COOH)] <sup>2+</sup>	2.67(6)
[H <sub>2</sub> L2(CH <sub>3</sub> COOH)] <sup>2+</sup> + CH <sub>3</sub> COO <sup>-</sup> = [H <sub>2</sub> L2(CH <sub>3</sub> COOH)(CH <sub>3</sub> COO)] <sup>+</sup>	2.74(6)
L2 + C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup> = [L2(C <sub>6</sub> H <sub>5</sub> COO)] <sup>-</sup>	1.82(9)
HL2 <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup> = [HL2(C <sub>6</sub> H <sub>5</sub> COO)]	2.18(5)
H <sub>2</sub> L2 <sup>2+</sup> + C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup> = [H <sub>2</sub> L2(C <sub>6</sub> H <sub>5</sub> COO)] <sup>+</sup>	2.62(3)
[H <sub>2</sub> L2(C <sub>6</sub> H <sub>5</sub> COO)] <sup>+</sup> + C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup> = [H <sub>2</sub> L2(C <sub>6</sub> H <sub>5</sub> COO) <sub>2</sub> ]	2.69(7)
HL2 <sup>+</sup> + Pht <sup>2-</sup> = [HL2(Pht)] <sup>-</sup>	2.27(2)
H <sub>2</sub> L2 <sup>2+</sup> + Pht <sup>2-</sup> = [H <sub>2</sub> L2(Pht)]	2.6(3)
H <sub>2</sub> L2 <sup>2+</sup> + HPht <sup>-</sup> = [H <sub>2</sub> L2(HPht)] <sup>+</sup>	2.0(7)
H <sub>2</sub> L2 <sup>2+</sup> + H <sub>2</sub> Pht = [H <sub>2</sub> L2(H <sub>2</sub> Pht)] <sup>2+</sup>	1.9(1)
[H <sub>2</sub> L2(Pht)] + Pht <sup>2-</sup> = [H <sub>2</sub> L2(Pht) <sub>2</sub> ] <sup>2-</sup>	2.16(9)
[H <sub>2</sub> L2(Pht)] + HPht <sup>-</sup> = [H <sub>2</sub> L2(HPht)(Pht)] <sup>-</sup>	1.64(3)
[H <sub>2</sub> L2(HPht)] <sup>+</sup> + H <sub>2</sub> Pht = [H <sub>2</sub> L2(H <sub>2</sub> Pht)(HPht)] <sup>+</sup>	2.97(7)
[H <sub>2</sub> L2(H <sub>2</sub> Pht)] <sup>2+</sup> + H <sub>2</sub> Pht = [H <sub>2</sub> L2(H <sub>2</sub> Pht) <sub>2</sub> ] <sup>2+</sup>	2.42(3)
L2 + IPht <sup>2-</sup> = [L2(IPht)] <sup>2-</sup>	1.75(6)
HL2 <sup>+</sup> + IPht <sup>2-</sup> = [HL2(IPht)] <sup>-</sup>	1.79(6)
H <sub>2</sub> L2 <sup>2+</sup> + IPht <sup>2-</sup> = [H <sub>2</sub> L2(IPht)]	2.37(1)

marked, following the expected order, preferential affinity for acetate or benzoate is strongly pH dependent, the latter being invariably favoured in more alkaline media, suggesting a π-π stacking/hydrophobic effect driven association.

Such coincidence between theory and experimental data, although intriguing, can be severely misleading. Accordingly, we will prosecute our search for general trends and re-evaluate selectivity remarks in the light of the whole discussion. The following material is divided into 7 focal points, going as follows: *i*) correlation between complexes stability and charge separation among hosts and guests; *ii*) comparison between carboxylates and sulfonates; *iii*) distinction between aromatic and aliphatic guests; *iv*) differences in complexes stability between the two ligands; *v*) stoichiometries of the formed adducts; *vi*) re-evaluation of selectivity features; *vii*) effect of the regiochemistry of dicarboxylate anions.

In relation to the complexes involving protonated ligands, a manifest poor correlation between complexes stability and substrate-receptor charge separation is observed (point *i*). In

**Table 4.** <sup>1</sup>H NMR CIS observed for anionic complexes of L1. Working pH and most abundant complex species are indicated case by case.

Substrate	CIS <sup>a</sup> Ligand <sup>b</sup>	CIS <sup>a</sup> Anion <sup>c</sup>	pH	Reference Species
Acetate	-0.02	-0.46	3.5	[HA-HL1]
	-0.01			
	-0.02			
Methanesulfonate	-0.97	-0.89	4.0	[A-HL1]
	-0.57			
	-1.02			
Benzoate	0.04	-0.22	8.0	[A-L1]
	0.05	-0.28		
	0.06	-0.22		
Benzenesulfonate	-5.77	-3.63	3.0	[A-H <sub>2</sub> L1]
	-2.08	-3.92		
	-4.86	-3.86		

<sup>a</sup> δ (ppm) / % formation of complex species;

<sup>b</sup> Referring to 1, 3 and 2 groups as designated in Figure 1;

<sup>c</sup> Referring to ortho, para and meta positions respectively for aromatic substrates.

**Table 5.** <sup>1</sup>H NMR CIS observed for anionic complexes of L2. Working pH and most abundant complex species are indicated case by case.

Substrate	CIS <sup>a</sup> Ligand <sup>b</sup>	CIS <sup>a</sup> Anion <sup>c</sup>	pH	Reference Species
Acetate	1.10	-0.09	3.0	[HA-H <sub>2</sub> L2]
	1.10			
	1.09			
	1.10			
Methanesulfonate	1.71	-0.21	3.0	[A-H <sub>2</sub> L2]
	1.72			
	1.71			
	1.72			
Benzoate	1.03	-0.06	4.5	[A-H <sub>2</sub> L2]
	0.93	-0.04		
	0.93	-0.04		
	1.22			
Benzenesulfonate	-0.11	-0.04	3.0	[A-H <sub>2</sub> L2]
	-0.62	-0.05		
	-0.17	-0.03		
	-0.04			

<sup>a</sup> δ (ppm) / % formation of complex species;


<sup>b</sup> Referring to 4, 2, 3 and 1 groups as designated in Figure 1;

<sup>c</sup> Referring to ortho, para and meta positions respectively for aromatic substrates.

particular, for a given anion, increasing the ligands charge does not result at all in the stability gain expected for the formation of a single salt bridge in water (5 ± 1 kJ/mol).<sup>21,22</sup>

This finding is perfectly in line with the evidences from our previous studies, indicating the anion-π interaction, rather than salt bridges, as the main force in play, along with strongly exoentropic solvent effects, in promoting the association of anions with L1 and L2.<sup>9</sup>


Since that was demonstrated for simple inorganic species, reinforcement of this notion is not surprising for the less hydrated, and potentially stronger interacting (due to the



**Figure 5.** Conditional stability constants for complexes of L1 (top) and L2 (bottom) with the series of monovalent anions

possibility of  $\pi$ - $\pi$  stacking), organic anions. Direct involvement of the tetrazine in anion binding is here also suggested by NMR data, as upfield shifts of  $^1\text{H}$  signals are observed, to different extents, for all the anions (Table 4 and Table 5). If in many cases formation of strong salt bridges may be invoked as an explanation, it is noteworthy that shielding is nevertheless observed in simply hydrogen bonded systems, lacking proper  $\pm$  charge separation between the partners (e.g.  $\text{HL1}^+ \cdot \text{HAcetate}$ , Table 4,  $\text{H}_2\text{L2}^{2+} \cdot \text{HAcetate}$ , Table 5), or lacking hydrogen bonds at all (e.g.  $\text{L1} \cdot \text{Benzoate}^-$ , Table 4). If the explanation of the CIS is to be found, at least partially, in aromatic ring currents, then the anion must be located right above the tetrazine ring, as expected for  $\pi$ - $\pi$  or anion- $\pi$  interaction modes: the CIS observed for the  $\text{L1} \cdot \text{Benzoate}^-$  complex (Table 4) being here a good case in point.


Focusing now on complexes involving neutral ligands, they are steadily observed only for carboxylate anions (point *ii*). Contrary to what we could expect, the extra stability provided by the matching of basicity among receptor and substrate protonable sites does not translate into a straightforward extra contribution to complex stability (Table 2 and Table 3). The main reason for this lack of correlation, that becomes evident in the simulated most stable conformations of the complexes (e.g. Figure 6 and Figure 7), is to be ascribed to the different geometries of the anions.



**Figure 6.** Calculated solution conformations of L2 complexes with benzoate ( $\text{A}^-$ ) in solution. Distances in Å. Parameters for anion- $\pi$  contacts given as plane/offset values.

Flat carboxylates are invariably preferred because of their ability to sit on the tetrazine and stick to it, causing extensive desolvation of both host and guest surfaces, while giving rise to salt bridges at the same time, whenever possible (Figure 7 and 8). Things are a little more complicated for the three-dimensional sulfonates, which have to choose between salt bridges and anion- $\pi$  interactions or  $\pi$ - $\pi$  stacking interactions. In the case of neutral ligands, that would mean managing to accommodate their bulkier polar group without getting too close to the electronegative morpholine atoms: as a matter of fact, such complexes are not experimentally observed.

Maintaining the focus on complexes of neutral, non-protonated, ligands, we can also see that they are systematically encountered only for aromatic guests (point *iii*), aliphatic ones being solely represented by the  $\text{L1} \cdot \text{acetate}$  complex. Spontaneous association of non-polar solutes in water is commonplace, and it is well-known to be driven by the hydrophobic effect. The studied aliphatic anions are both missing any contribution from stacking interaction and far less hydrophobic than their aromatic counterparts, resulting in non-detectable, at least by means of our potentiometric method, association equilibria. Interestingly, there is good coincidence between the *in silico* simulation of the benzoate complex of the neutral, not protonated, L1 (Figure 8) and  $^1\text{H}$  NMR data (Table 4), the upfield shift of the anion signals being ascribable to the tetrazine ring current. This reinforces the notion that direct interaction of the  $\pi$  clouds is indeed a factor in play for our complexes. The same can be concluded comparing the stability of L1 complexes with monovalent aliphatic and aromatic anions: the latter being found more stable whenever the speciation of the systems allows for a direct comparison. Similar conclusion cannot be drawn at once for potentiometric data of L2, which,




**Figure 7.** Calculated solution conformations of L1 (top) and L2 (bottom) complexes with benzenesulfonate (A<sup>-</sup>) in solution. Distances in Å. Parameters for anion-π contacts given as plane/offset values.

differently from L1, does not maintain a similar conformation in all of its complexes: its increased flexibility accounts for the more shifting nature of the data, which are less prone to direct comparison (cf. Figure 5 above and its relative discussion at point *vi* for an inclusive analysis based on conditional stability constants).

The scarce or total lack of additivity between π-π stacking interactions and other supramolecular forces for sulfonate anions is manifest in the comparison between the stability of L1 and L2 complexes with methane- and benzene-sulfonate severally (L1: log *K* = 2.21 vs log *K* = 2.49 for monoprotonated complexes, log *K* = 1.6 vs log *K* = 2.48 for diprotonated complexes, respectively; L2: log *K* = 1.41 vs log *K* = 1.41 for monoprotonated complexes, log *K* = 1.8 vs log *K* = 1.84 for diprotonated complexes, respectively; Table 2 and Table 3). The insensitivity of stability constants to the aromatic or aliphatic nature of the guests, log *K* values found close for L1 and equal within the experimental error in the case of L2 complexes of analogous anions, demonstrates the scarce contribution of stacking forces for these tetrahedral anions. It is worth mentioning that the experimental results mirror perfectly the *in silico* studies, which, as shown in Figure 7, demonstrate the marginal involvement of contacts between host and guest aromatic rings in complex stabilization, especially in the case of L2 adducts.

Centring the discussion on the ability of ligands L1 and L2 to form anion complexes, contrary to our previous data for inorganic anions,<sup>9</sup> L1 complexes are here generally more stable than L2 ones (point *iv*). Figure S1 allows for a direct comparison between the two ligands, showing how L1 species are invariably more abundant in a simulated competition setting.

A possible explanation of this trend reversal is provided by the most stable conformations calculated for these complexes in solution. In the case of benzoate, for instance, analogous



**Figure 8.** Calculated solution conformations of L1 complexes with benzoate (A<sup>-</sup>) in solution. Distances in Å. Parameters for anion-π contacts given as plane/offset values.

[L(C<sub>6</sub>H<sub>5</sub>COO)]<sup>-</sup>, [HL(C<sub>6</sub>H<sub>5</sub>COO)] and [H<sub>2</sub>L(C<sub>6</sub>H<sub>5</sub>COO)]<sup>+</sup> (L = L1, L2) species are formed. As shown in Figure 8, L1 is invariably found U-shaped, with all the binding sites converging towards the anion (Figure 8), while this is not always the case for L2 (Figure 6). Furthermore, the shorter methylenic spacer allows for a throughout contribution of the anion arenic portion to the interaction, which is found scarce in all the minimum energy conformations of L2 complexes.

These same observations are also found to be true for another series of analogous complexes, that of benzenesulfonate. As one can see in Figure 7, the more curled-up structure of L1 invariably results in an increased involvement of the benzenic ring in the host-guest interaction.


Accordingly, the higher overall stability observed for L1 complexes appears congruent with what should be expected due to its more rigid structure (higher preorganization/reduced loss of conformational degrees of freedom upon complexation), better convergence of binding sites and higher charge density when fully protonated, compared to its superior homologue L2. The main difference justifying the opposite stability trend observed for inorganic anions (L2 > L1),<sup>9</sup> lies in the geometry and charge distribution of the guest. Trigonal, tetrahedral and octahedral anions offer the more flexible L2 the possibility to wrap around them more easily than the flat aromatic anions of the new series (cf. the case of benzoate, Figure 6), offering several electron-rich binding sites amenable to accepting salt bridges or interacting with the tetrazine ring. On the contrary, L1, whose inability to fold led to the replacement of salt bridges with CH...anion contacts in the case of inorganic anions,<sup>9</sup> now manages to maintain in contact host and guest aromatic portions, while giving rise to salt bridges with the anionic groups protruding from the benzenic ring. This, reinforced by the magnified importance of solvation effects, the hydrophobicity of the anions being greatly increased in comparison to the inorganic ones, explains the discrepancy in relative stability of the complexes among the two series of studied anions.

Coming to the subject of stoichiometry, L2 demonstrates a marked tendency to form 1:2 ligand:anion complexes, which is totally absent in the case of L1 (point *v*). As one can observe for example in Figure 6, monoprotonated complexes are found in the signature chair conformation, stabilized by the concomitant

establishment of both salt bridge and anion- $\pi$  interaction. This interaction mode leaves out the other face of the tetrazine ring and the additional morpholine group for the interaction with a second anion, resembling the centrosymmetric arrangement observed in the crystal structures of inorganic anion complexes ( $\text{NO}_3^-$ ,  $\text{ClO}_4^-$ ,  $\text{PF}_6^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$  and  $\text{SCN}^-$ )<sup>9,10,12</sup> as well as in the phthalate complex (Figure 2). Such conformation is not amenable in solution for the less flexible L1, which, in fact, sticks to the 1:1 stoichiometry.

Interestingly, only carboxylate anions (acetate, benzoate, phthalate, isophthalate) do give rise to 1:2 complexes with L2. This correlates well with the discussion on the influence of the anion geometry. Carboxylate groups lay coplanar to their benzenic ring, allowing the charged groups to form salt bridges while the aromatic moieties can freely engage in parallel displaced  $\pi$ - $\pi$  stacking interactions, which ultimately lead to the chair conformation observed for the ligand in L2 complexes (Figure 6). This is not the case for sulfonate anions, the trigonal shape of the charged  $-\text{SO}_3^-$  group forces the system to choose: if a linear strong salt bridge is formed, imperfect matching in the  $\pi$ - $\pi$  stacking interaction is not avoidable; else, if the stacking interaction is preferred, the contribution of charge-charge interaction is greatly sacrificed. According to our simulations, when the ligands are diprotonated they wrap around the anion as portrayed in Figure 7, two salt bridges taking the upper hand over  $\pi$ - $\pi$  stacking. A point worth mentioning here, reinforcing the notion and showing the good agreement between experimental and *in silico* data, is the fact that diprotonated complexes of benzenesulfonate are found to be equally or marginally more stable than their monoprotinated counterparts for both ligands ( $\log K$  values 2.49 vs 2.48 for L1, 1.84 vs 1.44 for L2, Table 2 and Table 3). This mirrors what observed in the predicted minimum energy conformations (Figure 7), where the second protonation brings not only stronger enthalpic contributions but also entropic losses due to the exposure of the hydrophobic part of the anion to the solvent. Preference of the tetrazine for the polar group rather than the aromatic portion of the anion, as found in the simulations (Figure 7), is also supported, at least in the case of L2, by the arrangement found in the crystals of  $\text{H}_2\text{L2}(\text{C}_6\text{H}_5\text{SO}_3)_2 \cdot \text{H}_2\text{O}$  (Figure 4). The U folding assumed in solution by both diprotonated ligands when forming benzenesulfonate complexes (Figure 7) leaves no possibility of other stoichiometries in solution but 1:1, even for the flexible L2.

A borderline, and thus interesting, case is offered by acetate, which surely maintains the geometrical features of the other carboxylate anions, yet lacks entirely of the aromatic portion, partially invalidating some of the above arguments. In fact, without the stabilizing contribution of  $\pi$ - $\pi$  stacking, the ligand in the  $[\text{H}_2\text{L2}(\text{Acetate})]^+$  complex is predicted to be in a folded conformation by our simulations (Figure 9), closely resembling the situation encountered for sulfonate anions. Differently from them, however, acetate is a base, although of modest strength, thus it can undergo protonation: in the calculated conformation of  $[\text{H}_2\text{L2}(\text{HAcetate})]^{2+}$  the ligand opens up once more in the familiar chair conformation (Figure 9), again resulting in the



**Figure 9.** Calculated solution conformations of selected L2-acetate (A<sup>-</sup>) complexes in solution. Distances in Å. Parameters for anion- $\pi$  contacts given as plane/offset values.

possibility of 1:2 stoichiometry, which is indeed empirically encountered (Table 3).

Overall, L2 tendency to form 1:2 ligand:anion complexes is due to its increased flexibility compared to L1, but it is subject to the geometry of the anions and their basicity. This ultimately leads to discriminate the tetrahedral and essentially non-basic sulfonates from their carboxylate counterparts.

Finally, building on the previous discussion (points *i-v*), we may now re-evaluate properly the selectivity features disclosed by the analysis of conditional stability constants in Figure 5 (point *vi*). Indeed, as commented above, the theoretical order of stability expected on the basis of salt bridge strength and solvation/ $\pi$ -stacking contribution to the overall host-guest interaction is experimentally encountered for the series of monocharged anions; yet, in the light of the close inspection carried out above, we are forced to conclude that other factors subtend to the observed stability trend. If we admit, as demonstrated at point *i*, that we have a poor charge-charge control of the association phenomena, then salt bridge strength may not be invoked to justify the stability gap observed between carboxylate and sulfonate anions complexes. On the contrary, we may argue that looking at the salt bridge contribution only, in the complexes of diprotonated ligands sulfonate anions might even be more stabilized than carboxylate ones: even if in the latter case the intrinsic strength of the salt bridge is higher, carboxylates never form more than 1 salt bridge each in our simulations (Figures 6,8,9), while sulfonates can easily form 2 (Figure 7). The reason for this difference lies in the geometry of the anion, geometry which was also found to be responsible for the different stoichiometries of the formed adducts (point *v*) (affecting selectivity in the case of L2) and for carboxylates overall higher tendency to establish  $\pi$ -stacking interaction (points *ii*). Geometry's regulating function on the cooperativity of the different supramolecular forces is also manifest in the different selectivity observed for aliphatic over aromatic substrates (point *iii*). In the case of carboxylate anions, where cooperativity is favoured, we can clearly see that the aromatic benzoate is highly preferred over the aliphatic acetate starting from alkaline media (where the association resembles that of aromatic molecules due to solvent effect and  $\pi$ - $\pi$  stacking interactions)



until the different speciation of the systems allows so. On the contrary, since the tetrahedral shape of sulfonates does not allow stacking interactions (L2, Figure 7) or allows them poorly (L1, Figure 7), selectivity is indeed found to be scarcely affected (L1), if at all (L2), by the aromatic or aliphatic nature of the substrate.

The overall lesson is that amenability to selected interactions of the single binding sites of a substrate is of no use without an arrangement which favours the cooperativity of different supramolecular forces. In the present case, success of carboxylate guests over sulfonate ones is due to the topology of their binding sites, rather than the net strength of the formed salt bridges.


Lastly, regiochemistry of dicarboxylate anions is taken into account (point *vii*). Despite leading to the very same conclusions, it is instructive to undertake the analysis from a double viewpoint. If we look at the solution data (Table 2 and Table 3) from the ligands' perspective, L1 and L2 present different selectivity for the substrates: L1 forms more stable complexes with isophthalate than with phthalate, while the contrary is true for L2. Conditional stability constants comparisons (Figure S2) reveal that isophthalate binding is preferential in the 2.0-5.0 pH range for L1, while L2 privileges phthalate over its isomer in the 2.0-7.0 pH range. In both cases, selective recognition of one or the other regioisomer is possible in different conditions.

Instead, if we examine the data in terms of the affinity of the same anion for the two different receptors, L1 and L2, we find that both phthalate and isophthalate binds preferentially to L1, rather than L2, over a large pH range (see dedicated selectivity diagrams in Figure S1 and/or compare values of conditional stability constants in Figure S2).

Summarizing, L1 forms more stable complexes than L2 with both anions, which should not surprise according to the general discussion at point *iv*, yet, when the two substrates compete for the same ligand, L1 recognizes preferentially isophthalate over phthalate, while L2 behaves on the contrary, preferring phthalate to its isomer.

A guiding light for rationalizing the data is provided by the *in silico* simulation of the monoprotonated L2 complexes with both anions (Figure 10) and their comparison with the solved crystal structures of complexes featuring these anions (Figure 2 and Figure 3). Isophthalate has an inner tendency to simultaneously interact with both ligand arms, thus reducing the availability of binding sites for a second guest, and to involve its aromatic nucleus in the interaction. It should be noted that such involvement of stacking interaction in anion binding was also the main feature observed in the solved (H<sub>2</sub>L<sub>2</sub>)(Hisophthalate)<sub>2</sub> crystal structure (Figure 3), both visually and according to intermolecular potentials (see **Crystal Structure of (H<sub>2</sub>L)(Hisophthalate)<sub>2</sub>**).

Phthalate, on the contrary, due to the more gathered arrangement of its anionic sites, interacts with the tetrazine ring and just one morpholinic pendant (Figure 10). The L2 ligand assumes an overall chair-like conformation, stabilized by a short hydrogen bond contact (H...O 1.88 Å) and by an anion- $\pi$  interaction involving the same oxygen atom (O...ring-centroid



**Figure 10.** Calculated most stable conformations in solution of HL<sub>2</sub><sup>+</sup> complexes with phthalate (a) and (b) isophthalate dianions, respectively.

distance = 3.14 Å, X...ring-plane distance = 2.99 Å, offset with respect to the normal to plane = 0.94 Å) allowing, as in the case of benzoate (Figure 6), the interaction of a second anion with the residual vacant binding sites of the ligand. Here the resemblance with the (H<sub>2</sub>L)(HPhthalate)<sub>2</sub> crystal structure (Figure 2) is even more striking, strongly supporting a chair arrangement of the ligand which allows for the facile formation of 1:2 ligand:anion complexes in solution. The formation of such complexes necessarily favours phthalate over isophthalate binding by mass action in the case of L2, resulting in the observed selectivity.

On the other hand, the more gathered structure of L1, already shown to account for the across-the-board higher stability of its complexes (point *iv*) but also to prevent complex stoichiometries different from 1:1 (point *v*), joins the tendency of the isophthalate anion to interact simultaneously with the two ligand's arm: 1,3 disposition of the anion binding sites matching far better the para arrangement of the morpholine pendants around the 1,2,4,5 tetrazine core.

It should be mentioned that regiochemistry per se may not account for the detected differences: more subtle factors, better described with the broader concept of stereoelectronic effects, are in play. Geometry itself, beyond the ortho or meta disposition of the anionic sites, is not trivial either. Stereoelectronic reasons allow better planarity for the isophthalate dianion than for phthalate, torn between maximizing electron delocalization on the ring and minimizing charge-charge repulsion between carboxylate groups (Figure 10). The situation is not static either, protonation of phthalate implying its complete flattening due to the formation of a strong intramolecular hydrogen bond, as observed in the solid state for the case of the complexes with the monoprotonated anion (Figure 2). As a consequence, not only the mutual disposition of the divergent binding sites of the two substrates differs beyond

the ortho or meta orientation, pointing in or out of the benzene plane, but it is also subject, at least for phthalate, to noticeable modification due to the protonation state. Solvation effects due to proximal or distal charged sites may also play an active role for these dianions. Basicity of the two regioisomers is also greatly affected: due to the formation of the aforementioned intramolecular hydrogen bond, it is much easier to protonate once phthalate than isophthalate, and, conversely, it is easier to protonate twice isophthalate than phthalate (see Table S1). Interestingly enough, in this game of stereoelectronic effects and basicity constants, phthalate ends up possessing a persistent monoanionic form right where the diprotonated form of L2,  $H_2L2^{2+}$ , starts reaching its maximum (pH 4.0), thus inevitably promoting the formation of 1:2 complexes and favouring selectivity. Isophthalate instead, is left with basicity constants which match almost perfectly those of L1 ( $pK_{a1} = 4.32$  vs  $pK_{a1} = 4.45$  and  $pK_{a2} = 3.28$  vs  $pK_{a2} = 3.45$ , respectively; see Table S1): hence, matching of  $pK_a$ s might be really one of the features contributing to the selectivity observed for the L1-isophthalate system.

## Conclusions

The focus of this work is on the interplay of different supramolecular forces in stabilizing organic anion complexes both in solution and in the solid state. The studied substrates have been selected to possess different but related stereoelectronic properties, allowing the rationalization of a broad sample case of possible interactions. The study has been intentionally undertaken in water, despite it being one of the most challenging solvents for anion binding studies, in an effort to demonstrate how the interplay of different weak forces is able to stabilize anion complexes even in aqueous media, provided that certain criteria of host-guest electronic and structural complementarity are met.

The observed interactions include three main kinds of forces, salt bridge/hydrogen bond, anion- $\pi$  interactions and  $\pi$ - $\pi$  stacking, together with other contributions to complex stability such as the hydrophobic effect or size/shape complementarity with the two homologous ligands L1 and L2. For the series of monoanionic species, remarkable features of selectivity have been observed, as both L1 and L2 complexes show a stability trend which follows the order: benzoate  $\geq$  acetate  $>$  benzenesulfonate  $>$  methanesulfonate, i.e. carboxylate  $>$  sulfonate and aromatic  $>$  aliphatic.

Salt bridge strength, which was anticipated to be higher for carboxylate than for sulfonate anions due to the better matching of ligands' and anions'  $pK_a$ s, could be invoked a priori to justify the data. However, the geometry of the anions has been found to be a controlling factor, so that sulfonates could be even more stabilized by salt-bridges than carboxylates in some cases, due to the fact that, owing to the shape of the charged group, they are able to form 2 salt bridges instead of only 1.

Anion- $\pi$  interactions are also found to play an important role in these complexes. Whether intrinsically chosen by the systems or favoured by the disposition of ligands' hydrogen bond

donors, they are preferred over  $\pi$ - $\pi$  forces in most cases. Their importance for carboxylate anions manifest itself in the stabilization of 1:2 ligand:anion complexes, which are ubiquitous with L2.

Stacking interactions were found to contribute to the overall stability of complexes, but their participation is intermittent. Stacking is found to be the key force promoting association of aromatic carboxylates to neutral non-protonated ligands, yet it contributes almost nothing to the overall stabilization of sulfonate complexes. The high and positive quadrupolar moment of the tetrazine pairs well with the negative ones of all the aromatic substrates, thus enhancing receptor-substrate interactions resembling the association of non-polar molecules in water. Once more, geometry constraints determine whether or not an anion will be able to bring its aromatic portion close to the tetrazine ring.

The same is true also for the hydrophobic effect. One thing is the global hydrophilic/hydrophobic character of an anion, another is effectively receiving a contribution to complex stability due to the solvent effect: for this to happen, it is required that the hydrophobic surfaces are indeed desolvated and brought close to one another in the final complex in solution. This is found to occur in every case for flat carboxylate substrates, while it does not in the case of sulfonates: stability trend mirrors this exact observation.

Geometrical features of the anions appear as the single most important factor in orchestrating the interplay of the different supramolecular forces, thus confirming that utmost stability of supramolecular adducts is much easily achieved with an effective cooperativity of several different forces, rather than by maximizing each one of them as if they were in isolation.

Effect of the size/shape of the ligand accounts for the evidence that L1 more gathered converging disposition of binding sites and rigid structure result in complexes of higher stability compared to L2. The higher flexibility of L2 is instead found to be a governing factor in the formation of complexes with different stoichiometries. Overall, size/shape of the ligands determine the selectivity for the studied anions, including the intriguing one observed for the phthalate regioisomers: L1 preferentially binding to isophthalate, while L2 to phthalate. Although geometry appears once more as a key feature, provided explanation of the experimental data could not overlook global stereoelectronic considerations, exceeding a simplistic view of isomerism.

## Experimental

### Materials

L1 and L2 (3,6-bis(morpholin-4-ylmethyl)-1,2,4,5-tetrazine and 3,6-bis(morpholin-4-ylethyl)-1,2,4,5-tetrazine, respectively) were synthesized as previously described.<sup>9</sup> Pink crystals of  $H_2L2(C_6H_5SO_3)_2 \cdot H_2O$  were obtained upon evaporation at room temperature of an aqueous solution of L2 (0.01M) at pH 4.0 containing an excess of benzenesulfonate. For the phthalate and isophthalate complexes, ethanolic solution of the corresponding dicarboxylic acids were prepared separately and

added dropwise to an ethanolic solution of the ligand without further pH adjustment. Slow evaporation at room temperature led to pink crystals of (H<sub>2</sub>L)(HPhthalate)<sub>2</sub> and (H<sub>2</sub>L)(HIsophthalate)<sub>2</sub> respectively.

#### Potentiometric Measurements

Potentiometric (pH-metric) titrations employed for the determination of equilibrium constants were carried out in 0.1 M NMe<sub>4</sub>Cl degassed aqueous solutions at 298.1 ± 0.1 K by using previously described equipment and procedures.<sup>23</sup> The determined ionic product of water was pK<sub>w</sub> = 13.83(1) (298.1 ± 0.1 K, 0.1 M NMe<sub>4</sub>Cl). High purity commercial reagents were purchased and employed for the potentiometric measurements without further purification. The computer program HYPERQUAD<sup>18</sup> was used to calculate equilibrium constants from potentiometric data deriving from at least three independent titration experiments. Ligands protonation constants were previously determined,<sup>9</sup> while anion protonation constants were re-determined in our experimental conditions: these data are available in the supporting information (Table S1). For complexation studies, ligand concentration was either 5 × 10<sup>-4</sup> M (L1) or 1 × 10<sup>-3</sup> M (L2), while anion concentrations ranged from 2 up to 5 equivalents for each anion. Only in the case of benzenesulfonate measurements were extended up to 10 equivalents of anion. The studied pH range was 3.0–9.0.

#### <sup>1</sup>H NMR

Stock solutions of ligands and anions at the desired pH value were prepared in D<sub>2</sub>O and mixed to obtain the desired anion:ligand ratio: further pH adjustment was performed after mixing whenever needed. pH adjustments were performed by addition of DCl or NaOD solutions, pD and pH were correlated according to the pH = pD – 0.4 equivalence. Spectra were recorded on a Bruker Avance III 400 MHz spectrometer.

#### X-ray Structure Analyses

Pink crystals of (H<sub>2</sub>L<sub>2</sub>)(HPhthalate)<sub>2</sub>·2H<sub>2</sub>O (a), (H<sub>2</sub>L<sub>2</sub>)(HIsophthalate)<sub>2</sub> (b) and H<sub>2</sub>L<sub>2</sub>(C<sub>6</sub>H<sub>5</sub>SO<sub>3</sub>)<sub>2</sub>·H<sub>2</sub>O (c) were used for X-ray diffraction analysis. A summary of the crystallographic data is reported in Table S2. The integrated intensities were corrected for Lorentz and polarization effects and an empirical absorption correction was applied.<sup>24</sup> The structures were solved by direct methods (SIR92).<sup>25</sup> Refinements were performed by means of full-matrix least-squares using SHELXL Version 2014/7.<sup>26</sup> All the non-hydrogen atoms were anisotropically refined. All hydrogen atoms were introduced in calculated position and their coordinates were refined according to the linked atoms, with the exception of those belonging to the water molecule in (a) which were not found in the Fourier difference map and the ammonium hydrogen in (a), the carboxylic hydrogens in (a) and (b) and the water hydrogens in (c) which were instead localized and freely refined with isotropic treatment. CCDC 1835198–1835200 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

#### Molecular Modelling Calculations

Molecular modelling investigations on the following complexes (charges omitted for clarity) in order to take into account possible different protons' distributions, as suggested by potentiometric results:

Acetate: L1/S, L1/HS, L2/HS, HL1/S, HL2/S, HL1/HS, HL2/HS, H<sub>2</sub>L1/S, H<sub>2</sub>L2/S, H<sub>2</sub>L2/HS, HL2/HS/HS, H<sub>2</sub>L2/HS/S

Benzoate: L1/S, L2/S, L1/HS, L2/HS, HL1/S, HL2/S, HL1/HS, HL2/HS, H<sub>2</sub>L1/S, H<sub>2</sub>L2/S, H<sub>2</sub>L2/S/S, HL2/HS/S

Benzenesulfonate: HL1/S, HL2/S, H<sub>2</sub>L1/S, H<sub>2</sub>L2/S

Phthalate and Isophthalate: HL2/S

Calculations were performed by means of the empirical force field method AMBER3 as implemented in the Hyperchem 7.51 package,<sup>27</sup> using an implicit simulation of aqueous environment (ε = 4 r) and atomic charged evaluated at the semiempirical level of theory (PM3)<sup>28</sup>. Potential energy surface of all the systems were explored by means of simulated annealing (T = 600 K, equilibration time = 10 ps, run time = 10 ps and cooling time = 10 ps, time step = 1.0 fs). For each studied system, 80 conformations were sampled.

#### Conflicts of interest

There are no conflicts to declare.

#### Acknowledgements

Financial support from the Italian MIUR (project 2015MP34H3) and from the Spanish MINECO (project MAT2014-60104-C2-2-R) are gratefully acknowledged.

#### References

- 1 K. Bowman-James, A. Bianchi, E. García-España, Eds; *Anion Coordination Chemistry*; Wiley-VHC: New York, **2012**.
- 2 J. L. Sessler, P. A. Gale, W. S. Cho, *Anion Receptor Chemistry*, Monographs in Supramolecular Chemistry, RSC Publishing: Cambridge, U.K., **2006**.
- 3 P. Arranz-Mascarós, C. Bazzicalupi, A. Bianchi, C. Giorgi, M. L. Godino-Salido, M. D. Gutiérrez-Valero, R. Lopez-Garzón, M. Savastano, *J. Am. Chem. Soc.* **2013**, *135*(1), 102–105.
- 4 a) V. Stilinović, G. Horvat, T. Hrenar, V. Nemeč, D. Činčić, *Chem. Eur. J.* **2017**, *23*, 5244; b) Grabowski, S. J., *Chem. Eur. J.* **2013**, *19*, 14600–14611.
- 5 a) X. Lucas, A. Bauzá, A. Frontera, D. Quiñonero, *Chem. Sci.* **2016**, *7*, 1038–1050; b) Y. P. Yurenko, J. Novotný, R. Marek, *Chem. Eur. J.* **2017**, *23*, 5573.
- 6 a) R. V. Nair, S. Kheria, S. Rayavarapu, A. S. Kotmale, B. Jagadeesh, R. G. Gonnade, V. G. Puranik, P. R. Rajamohanam, G. J. Sanjayan, *J. Am. Chem. Soc.* **2013**, *135* (31), 11477–11480; b) G. N. M. Reddy, A. Marsh, J. T. Davis, S. Masiero, S. P. Brown, *Crystal Growth & Design* **2015**, *15* (12), 5945–5954
- 7 a) Y.-L. Wang, A. Laaksonen, M. D. Fayer, *J. Phys. Chem. B* **2017**, *121* (29), 7173–7179; b) M. Lübtow, I. Helmers, V. Stepanenko, R. Q. Albuquerque, T. B. Marder, G. Fernández, *Chem. Eur. J.* **2017**, *23*, 6198.
- 8 a) M. A. Uddin, T. H. Lee, S. Xu, S. Y. Park, T. Kim, S. Song, T. L. Nguyen, S. Ko, S. Hwang, J. Y. Kim, H. Y. Woo, *Chem. Mat.* **2015**, *27* (17), 5997–6007; b) M. Müller, M. Albrecht, V. Gossen, T. Peters, A. Hoffmann, G. Raabe, A. Valkonen, K.

- Rissanen, *Chem. Eur. J.* **2010**, *16*, 12446–12453; c) M. Savastano, C. Bazzicalupi, P. Mariani and A. Bianchi, *Molecules* **2018**, *23*(3), 572.
- 9 M. Savastano, C. Bazzicalupi, C. Giorgi, C. García-Gallarín, M. D. López de la Torre, F. Pichierri, A. Bianchi, M. Melguizo, *Inorg. Chem.* **2016**, *55* (16), 8013–8024.
- 10 M. Savastano, C. Bazzicalupi, C. García-Gallarín, C. Giorgi, M. D. López de la Torre, F. Pichierri, A. Bianchi, M. Melguizo, *Dalton Trans.* **2018**, *47*, 3329–3338.
- 11 M. Savastano, C. Bazzicalupi, C. García, C. Gellini, M. D. López de la Torre, P. Mariani, F. Pichierri, A. Bianchi and M. Melguizo, *Dalton Trans.* **2017**, *46*, 4518–4529.
- 12 M. Savastano, C. García, M. D. López de la Torre, F. Pichierri, C. Bazzicalupi, A. Bianchi and M. Melguizo, *Inorg. Chim. Acta* **2018**, *470*, 133–138.
- 13 a) S. Shan, S. Loh, D. Herschlag, *Science* **1996**, *272*, 97–101; b) G. Gilli, P. Gilli, *The Nature of Hydrogen Bond: Outline of a Comprehensive Hydrogen Bond Theory*; Oxford University Press, New York, **2009**.
- 14 M. Odellius, B. Kirchner, and J. Hutter; *J. Phys. Chem.* **2004**, *108* (11), 2044–2052
- 15 L. K. S. von Krbek, A. J. Achazi, S. Schoder, M. Gaedke, T. Biberger, B. Paulus, C. A. Schalley, *Chem. Eur. J.* **2017**, *23*, 2877.
- 16 A. Gavezzotti, *Acc. Chem. Res.* **1994**, *27* (10), 309–314.
- 17 A. Gavezzotti, G. Filippini, *J. Phys. Chem.* **1994**, *98* (18), 4831–4837
- 18 P. Gans, A. Sabatini, A. Vacca, *Talanta* **1996**, *43*, 1739–1753.
- 19 a) A. Bianchi, M. Micheloni, P. Paoletti, *Inorg. Chim. Acta* **1988**, *151*, 269; b) A. Andrés, J. Aragón, A. Bencini, A. Bianchi, A. Domenech, V. Fusi, E. García-España, P. Paoletti, J.A. Ramírez, *Inorg. Chem.* **1993**, *32*, 3418; c) C. Bazzicalupi, A. Bencini, A. Bianchi, M. Cecchi, B. Escuder, V. Fusi, E. García-España, C. Giorgi, S.V. Luis, G. Maccagni, V. Marcelino, P. Paoletti, B. Valtancoli, *J. Am. Chem. Soc.* **1999**, *121*, 6807.
- 20 C. Bazzicalupi, A. Bianchi, C. Giorgi, M. P. Clares, E. Garcia-España, *Coord. Chem. Rev.* **2011**, *256*, 13–27.
- 21 H.-J. Schneider, *Chem. Soc. Rev.* **1994**, *22*, 227–234.
- 22 H.-J. Schneider, *Angew. Chem. Int. Ed.* **2009**, *48*, 3924–3977.
- 23 C. Bazzicalupi, A. Bianchi, T. Biver, C. Giorgi, S. Santarelli, M. Savastano, *Inorg. Chem.* **2014**, *53*, 12215–12224.
- 24 *CrysAlisPro*, Agilent Technologies, Version 1.171.37.35g (release 09-12-2014 CrysAlis171 .NET); Agilent Technologies: Oxfordshire, England.
- 25 A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori, M. Camalli, *J. Appl. Crystallogr.* **1994**, *27*, 435–436.
- 26 G.M. Sheldrick; *Acta Crystallogr. C* **2015**, *71*, 3–8.
- 27 *Hyperchem*, Release 7.51 for Windows MM System; Hypercube, Inc.: Gainesville, FL, USA, 2002.
- 28 a) J.J.P. Stewart, *J. Comput. Chem.* **1989**, *10*, 209–220; b) J.J.P. Stewart, *J. Comput. Chem.* **1989**, *10*, 221–264.