



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi  
di Firenze

## LIPSCHITZ CONTINUITY FOR ENERGY INTEGRALS WITH VARIABLE EXPONENTS

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

LIPSCHITZ CONTINUITY FOR ENERGY INTEGRALS WITH VARIABLE EXPONENTS / Eleuteri, Michela; Marcellini, Paolo; Mascolo, Elvira. - In: ATTI DELLA ACCADEMIA NAZIONALE DEI LINCEI. RENDICONTI LINCEI. MATEMATICA E APPLICAZIONI. - ISSN 1720-0768. - STAMPA. - 27:(2016), pp. 61-87. [10.4171/RLM/723]

*Availability:*

This version is available at: 2158/1017799 since: 2021-03-17T10:10:59Z

*Published version:*

DOI: 10.4171/RLM/723

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

# LIPSCHITZ CONTINUITY FOR ENERGY INTEGRALS WITH VARIABLE EXPONENTS

MICHELA ELEUTERI – PAOLO MARCELLINI – ELVIRA MASCOLO

ABSTRACT. A regularity result for integrals of the Calculus of Variations with variable exponents is presented. Precisely, we prove that any vector-valued minimizer of an energy integral over an open set  $\Omega \subset \mathbb{R}^n$ , with variable exponent  $p(x)$  in the Sobolev class  $W_{\text{loc}}^{1,r}(\Omega)$  for some  $r > n$ , is locally Lipschitz continuous in  $\Omega$  and an a priori estimate holds.

## 1. INTRODUCTION

In the past few years the study of energy integrals with variable exponent received a large interest. We refer for instance to the integral functional

$$F(u) = \int_{\Omega} a(x)h(|Du|)^{p(x)} dx, \quad (1.1)$$

where  $a(x) > 0$  and  $p(x) > 1$  are continuous functions in  $\Omega$ ,  $h$  is an increasing convex function and  $u$  is a vector-valued map. The variational integral (1.1) exhibits  $p(x)$ -growth, which is a particular case of the so-called nonstandard growth, with an extensive literature on the subject. While existence of minimizers follows from the direct methods of the Calculus of Variations, the regularity problem is not yet completely settled. We stress that in the vector-valued case, as suggested by the well known counterexamples by De Giorgi [10], Giusti-Miranda [18] and more recently by Sverak-Yan [33], Mooney-Savin [30], some structure conditions on the integrand are required for everywhere regularity.

Here we consider an open bounded set  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , a coefficient  $a \in W_{\text{loc}}^{1,r}(\Omega)$  and an exponent  $p \in W_{\text{loc}}^{1,r}(\Omega)$  for some  $r > n$ , an increasing convex function  $h = h(t)$ ,  $h : [0, \infty) \rightarrow [0, \infty)$  and a vector-valued map  $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^N$ , with  $N \geq 1$ . We deal with the local Lipschitz continuity of minimizers, without a prescribed bound on the oscillation of  $p(x)$ , assuming instead the summability of the weak derivatives of  $a$  and  $p$ . Precisely we will prove the following result.

**Theorem 1.1.** *Let  $u \in W_{\text{loc}}^{1,1}(\Omega; \mathbb{R}^N)$  be a local minimizer of the energy integral (1.1). Let  $h : [0, +\infty) \rightarrow [0, +\infty)$  be an increasing convex function, not identically zero, such that for*

---

2000 *Mathematics Subject Classification.* Primary: 35J60, 35B65, 49N60; Secondary: 35J50, 35B45.

*Key words and phrases.* Energy integrals, local minimizers, local Lipschitz continuity,  $p(x)$ -growth, variable exponents.

The authors are supported by GNAMPA (Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni) of INdAM (Istituto Nazionale di Alta Matematica); in particular the first and the third author are supported by the research project GNAMPA-INdAM 2014 “*Materiali speciali e regolarità nel Calcolo delle Variazioni*” (coord. M. Eleuteri).

The authors wish to express their gratitude to the referee for carefully reading the manuscript providing useful comments and remarks.

some  $M_0 > 0$ ,  $t_0 \geq 1$  and  $0 \leq \delta \leq \frac{1}{n} - \frac{1}{r}$

$$h'(t) \leq M_0 t^\delta, \quad h''(t) \leq M_0 \frac{h'(t)}{t}, \quad \forall t \geq t_0, \quad (1.2)$$

with  $a, p \in W_{\text{loc}}^{1,r}(\Omega)$  for  $r > n$ . Then  $u$  is a locally Lipschitz continuous map and there exist constants  $C > 0$  and  $\beta > 1$  such that, for  $0 < \rho < R$ ,

$$\|Du\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq C \left[ \left( \frac{1 + \|a_x\|_{L^r(B_R)} + \|p_x\|_{L^r(B_R)}}{(R - \rho)} \right)^{\frac{1}{n - \frac{1}{r}}} \int_{B_R} \{1 + a(x) h(|Du|)^{p(x)}\} dx \right]^\beta.$$

As usual  $B_\rho$  and  $B_R$  are balls in  $\Omega$  of radii  $\rho$  and  $R$  with the same center. In particular the constants  $C > 0$  and  $\beta > 1$  depend on  $n, r, M_0, t_0$  and on the infimum and the supremum of  $a$  and  $p$  in the ball  $B_R$ .

Examples of  $h$  functions which satisfy the assumptions of Theorem 1.1 are, of course  $h(t) = t$  or

$$h(t) = t \log(1 + t), \quad t \geq 0,$$

more generally  $h(t) = t \log^\beta t$  for large values of  $t$  and for some  $\beta > 0$  and also

$$h(t) = \int_1^t \frac{s^\delta}{\log s} ds, \quad t \geq t_0.$$

As in [9], [16], [23], [34], a different example of a convex function  $h(t)$  satisfying the above conditions is given, for large  $t$ , by

$$h(t) = t^{a+b \sin \log \log t}, \quad (1.3)$$

when  $a > 1$  and  $b > 0$  is sufficiently small (see Section 3). The function  $h$  has  $p, q$ -growth, in the sense that  $t^p \leq h(t) \leq t^q$  for large values of  $t$ , with  $p = a - b$  and  $q = a + b$ .

The proof of Theorem 1.1 will follow through several steps, the main one being a reduction of the given energy functional in (1.1) to the framework considered by the authors in [13].

The energy variable exponent is nowadays a classical topic in the Calculus of Variations, PDEs and Nonlinear Analysis. The large number of papers studying energies involving variable exponents is motivated by the fact that this type of functionals can be considered as a model in the theory of strongly anisotropic materials (see e.g. Zhikov [35] and Zhikov et al. [36]) and in the theory of electrorheological fluids (see e.g. Rajagopal-Růžička [31] and Růžička [32]). More recently, functionals as in (1.1) were considered also in the study of image denoising (see e.g. Chen et al. [4]) and in some models for growth of heterogeneous sandpiles (see e.g. Bocea et al. [2]). The regularity of minimizers has been studied by many authors. For the case  $h(t) = t$  we mention: Chiadò-Piat-Coscia [5], Coscia-Mingione [7], Acerbi-Mingione [1], Esposito-Leonetti-Mingione [15]. The case  $h(t) \sim e^{tm}$ ,  $m > 0$ , was considered by Mascolo-Migliorini [26] and with the  $x, u$  dependence by Eleuteri [12]. A further list of references can be found in Härjulehto-Hästö-Nuortio [20] and Diening-Härjulehto-Hästö-Nuortio [11]. Energies with variable exponents are also studied in the framework of the  $p, q$ -growth; indeed if  $p(x)$  is a continuous function, on a small ball its minimum and its maximum values behave as  $p, q$ , with  $q$  arbitrarily close to  $p$ . We refer to Marcellini [21], [22], [23], [24] and to Mingione [27] for a survey on this subject. Variable exponents were also considered under different aspects in Nonlinear Analysis, for instance with respect to

eigenvalue problems and to the multiplicity of solutions (see e.g. Pucci, Radulescu et al. [3], [28], [29] and, more recently, Colasuonno-Squassina [6]).

## 2. PROOF OF THEOREM 1.1

The proof of Theorem 1.1 follows in several steps:

**2.1. Step 1: localization.** For every  $x_0 \in \Omega$ , there exists  $R_0 > 0$  such that the ball  $B_{R_0}(x_0)$  is contained in  $\Omega$  and, if we set

$$p := \inf\{p(x) : x \in B_{R_0}(x_0)\} > 1 \quad q := \sup\{p(x) : x \in B_{R_0}(x_0)\}(1 + \delta) + \tau \quad (2.1)$$

for some

$$\tau < \frac{1}{1 + \delta} \left( \frac{1}{n} - \frac{1}{r} - \delta \right), \quad (2.2)$$

then

$$\frac{q}{p} < 1 + \frac{1}{n} - \frac{1}{r}. \quad (2.3)$$

Indeed, for any  $\varepsilon_0$  there exists  $R_0 > 0$  such that

$$\sup\{p(x) : x \in B_{R_0}(x_0)\} - \inf\{p(x) : x \in B_{R_0}(x_0)\} < \varepsilon_0.$$

We set

$$\varepsilon_0 := \frac{1}{1 + \delta} \left( \frac{1}{n} - \frac{1}{r} - \delta \right) - \tau.$$

This is possible due to the smallness assumptions on  $\delta$  and  $\tau$  we required. At this point

$$\begin{aligned} \frac{\sup\{p(x) : x \in B_{R_0}(x_0)\}}{\inf\{p(x) : x \in B_{R_0}(x_0)\}} - 1 &< \frac{1}{1 + \delta} \left( \frac{1}{n} - \frac{1}{r} - \delta \right) \frac{1}{\inf\{p(x) : x \in B_{R_0}(x_0)\}} \\ &= \frac{\tau}{\inf\{p(x) : x \in B_{R_0}(x_0)\}} \\ &< \frac{1}{1 + \delta} \left( \frac{1}{n} - \frac{1}{r} - \delta \right) - \frac{\tau}{\inf\{p(x) : x \in B_{R_0}(x_0)\}} \end{aligned}$$

as long as  $p(x) > 1$ . This finally entails, taking into account (2.1)

$$\frac{q}{p} < \frac{[\sup\{p(x) : x \in B_{R_0}(x_0)\} + \tau](1 + \delta)}{\inf\{p(x) : x \in B_{R_0}(x_0)\}} < 1 + \delta + \frac{1}{n} - \frac{1}{r} - \delta = 1 + \frac{1}{n} - \frac{1}{r}.$$

It is obvious that  $\sup\{p(x) : x \in B_{R_0}(x_0)\} < q$ .

**2.2. Step 2: consequences from the assumptions.** Before starting, we need the following elementary properties of convex functions.

Let  $h : [0, \infty) \rightarrow [0, \infty)$  be an increasing convex function fulfilling (1.2). The functions  $h(t)$  and  $h'(t)$  are not identically equal to zero, thus there exists  $t_0 \geq 1$  such that (1.2) holds and

$$h(t_0) > 0 \quad h'(t_0) > 0.$$

Let us set

$$m_0 := \min \left\{ \frac{h(t_0)}{t_0}, h'(t_0) \right\}. \quad (2.4)$$

Then we have

$$h'(t) \geq h'(t_0) \geq m_0 \quad \forall t \geq t_0 \quad (2.5)$$

and moreover by convexity and (2.5)

$$h(t) \geq h(t_0) + h'(t_0)(t - t_0) \geq h(t_0) + m_0(t - t_0) \geq m_0 t_0 + m_0(t - t_0) = m_0 t. \quad (2.6)$$

On the other hand, we can also show that

$$\frac{1}{\tilde{C}_0} h'(t)t \leq h(t) \leq C_0 h'(t)t, \quad (2.7)$$

where

$$\tilde{C}_0 := \max\{M_0 + 1, h'(t_0)t_0\} \quad C_0 = 1 + \frac{h(t_0)}{h'(t_0)}. \quad (2.8)$$

Indeed from the Mean Value Theorem and the fact that  $h$  is increasing

$$h(t) \leq h(0) + h'(t)t \leq h(t_0) + h'(t)t.$$

But  $h$  is convex, so also  $h'$  is increasing; this entails, for all  $t \geq t_0 \geq 1$

$$h(t_0) = h'(t_0) \frac{h(t_0)}{h'(t_0)} \leq \frac{h(t_0)}{h'(t_0)} h'(t_0)t_0 \leq \frac{h(t_0)}{h'(t_0)} h'(t)t, \quad (2.9)$$

thus we get the desired inequality

$$h(t) \leq \left(1 + \frac{h(t_0)}{h'(t_0)}\right) h'(t)t \stackrel{(2.8)}{=} C_0 h'(t)t.$$

We prove now the other inequality in (2.7). Indeed, by the second inequality in (1.2), we have

$$\int_{t_0}^t h''(\tau)\tau \, d\tau \leq M_0 \int_{t_0}^t h'(\tau) \, d\tau$$

and by integration by parts

$$h'(t)t - h'(t_0)t_0 - \int_{t_0}^t h'(\tau) \, d\tau \leq M_0 \int_{t_0}^t h'(\tau) \, d\tau$$

that is

$$h'(t)t \leq (M_0 + 1)(h(t) - h(t_0)) + h'(t_0)t_0 \leq \tilde{C}_0 h(t).$$

Observe that (2.6) and (2.7) imply

$$m_0 t \leq h(t) \leq C_0 M_0 t^{1+\delta} \quad \forall t \geq t_0. \quad (2.10)$$

In the sequel it is not restrictive to assume that

$$m_0 \leq 1 \leq M_0 \quad \inf_{x \in B_{R_0}} a(x) \geq m_0 > 0 \quad \sup_{x \in B_{R_0}} a(x) \leq M_0.$$

**2.3. Step 3: ellipticity and growth assumptions.** Let us denote

$$g(x, |Du|) = a(x)h(|Du|)^{p(x)} \quad g(x, t) = a(x)h(t)^{p(x)}.$$

In this step we show that  $g(x, t)$  satisfies the following growth conditions, for all  $t \geq t_0$  and a.e.  $x \in B_{R_0}$

$$\lambda t^{p(x)-2} \leq \frac{g_t(x, t)}{t} \leq \Lambda t^{(1+\delta)p(x)-2} \quad (2.11)$$

$$\lambda t^{p(x)-2} \leq g_{tt}(x, t) \leq \Lambda t^{(1+\delta)p(x)-2} \quad (2.12)$$

$$|g_{tx}(x, t)| \leq \Lambda \ell(x) t^{(1+\delta)p(x)+\tau-1} \quad (2.13)$$

for  $\lambda, \Lambda$  defined as

$$\lambda := \min \left\{ 1, \frac{p-1}{C_0^2} \right\} p m_0^{q+1}, \quad \Lambda := \frac{q(q-1)}{m_0} C_0^q M_0^{q+2} \tilde{C}_0^2 + \frac{2}{m_0} \max\{\log(C_0 M_0), \delta + 1\}, \quad (2.14)$$

with  $\delta$  and  $\tau$  as in (1.2) and (2.1) respectively and

$$\ell(x) := a_x(x) + p_x(x) \quad \text{for a.e. } x \in B_{R_0}. \quad (2.15)$$

Indeed

$$g_t(x, t) = a(x) p(x) h(t)^{p(x)-1} h'(t)$$

so that

$$\frac{g_t(x, t)}{t} = a(x) p(x) h(t)^{p(x)-1} \frac{h'(t)}{t} \stackrel{(1.2), (2.10)}{\leq} q M_0^{q+1} C_0^{q-1} t^{(1+\delta)p(x)-2} \stackrel{(2.14)}{\leq} \Lambda t^{(1+\delta)p(x)-2}. \quad (2.16)$$

Consider now

$$g_{tt}(x, t) = a(x) p(x) (p(x) - 1) h(t)^{p(x)-2} [h'(t)]^2 + a(x) p(x) h(t)^{p(x)-1} h''(t).$$

We deal first with the second term. We have

$$\begin{aligned} a(x) p(x) h(t)^{p(x)-1} h''(t) &\stackrel{(1.2)}{\leq} M_0 a(x) p(x) h(t)^{p(x)-1} \frac{h'(t)}{t} \\ &\stackrel{(2.16)}{\leq} \Lambda t^{(1+\delta)p(x)-2}. \end{aligned}$$

On the other hand

$$\begin{aligned} a(x) p(x) (p(x) - 1) h(t)^{p(x)-2} [h'(t)]^2 &\stackrel{(2.7)}{\leq} a(x) p(x) (p(x) - 1) \frac{h(t)^{p(x)}}{t^2} \tilde{C}_0^2 \\ &\stackrel{(2.10)}{\leq} C_0^q M_0^{q+1} q (q-1) \tilde{C}_0^2 t^{(1+\delta)p(x)-2} \\ &\stackrel{(2.14)}{\leq} \Lambda t^{(1+\delta)p(x)-2}. \end{aligned}$$

We deal now with the lower bounds. First of all we have

$$\frac{g_t(x, t)}{t} = a(x) p(x) h(t)^{p(x)-1} \frac{h'(t)}{t} \stackrel{(2.5), (2.6)}{\geq} a(x) p(x) [m_0 t]^{p(x)-1} \frac{m_0}{t} \stackrel{(2.14)}{\geq} \lambda t^{p(x)-2}.$$

We also have, using (2.5)

$$\begin{aligned} g_{tt}(x, t) &\geq a(x)p(x)(p(x) - 1)h(t)^{p(x)-2}[h'(t)]^2 \\ &\stackrel{(2.7)}{\geq} \frac{1}{C_0^2} p(p-1) m_0 \frac{h(t)^{p(x)}}{t^2} \stackrel{(2.10)}{\geq} \frac{1}{C_0^2} m_0^{q+1} p(p-1) t^{p(x)-2} \stackrel{(2.14)}{\geq} \lambda t^{p(x)-2}, \end{aligned}$$

where we used the fact that  $m_0 \leq 1$ .

Finally

$$\begin{aligned} g_{tx}(x, t) &= a_x(x)p(x)h(t)^{p(x)-1}h'(t) + a(x)p_x(x)h(t)^{p(x)-1}h'(t) \\ &\quad + a(x)p(x)h(t)^{p(x)-1}h'(t)p_x(x) \log(h(t)) \end{aligned}$$

so that

$$\begin{aligned} g_{tx}(x, t) &\leq \frac{a_x(x)}{m_0} a(x)p(x)h(t)^{p(x)-1}h'(t) + \frac{p_x(x)}{p} a(x)p(x)h(t)^{p(x)-1}h'(t) \\ &\quad + a(x)p(x)h(t)^{p(x)-1}h'(t)p_x(x) \log(h(t)) \\ &\leq \frac{\ell(x)}{m_0} g_t(x, t)(1 + \log(h(t))), \end{aligned}$$

with  $\ell$  as in (2.15). At this point we observe that, for  $t \geq t_0$

$$\log(h(t)) \stackrel{(2.10)}{\leq} \log(C_0 M_0 t^{\delta+1}) = \log(C_0 M_0) + (\delta + 1) \log t \leq C t^\tau$$

where  $\tau$  is as in (2.2) and  $C := \max\{\log(C_0 M_0), \delta + 1\} \geq 1$ . This allows us to conclude that

$$|g_{tx}(x, t)| \leq \frac{\ell(x)}{m_0} g_t(x, t) 2 \max\{\log(C_0 M_0), \delta + 1\} t^\tau \stackrel{(2.14), (2.16)}{\leq} \Lambda \ell(x) t^{(1+\delta)p(x)+\tau-1}.$$

**2.4. Step 4: Approximation.** We construct a sequence of smooth functions  $g^{k\varepsilon}(x, t)$ , related to  $g(x, |Du|) = a(x)h(|Du|)^{p(x)}$ . We will deal with this approximation procedure in two steps. First let us define, for a.e.  $x \in B_{R_0}$

$$g_t^k(x, t) := \begin{cases} a(x)p(x)h(t)^{p(x)-1}h'(t) & 0 \leq t < k \\ a(x)p(x)h(k)^{p(x)-1}h'(k) + \frac{\lambda}{p-1}[t^{p-1} - k^{p-1}] & t \geq k, \end{cases} \quad (2.17)$$

and

$$g^k(x, t) := \int_0^t g_t^k(x, s) ds + g(x, 0). \quad (2.18)$$

Arguing in a similar way as in [13], it is possible to show that the sequence of functions defined by (2.18) satisfies, for  $k$  sufficiently large, the conditions

$$g^k(x, t) \leq g(x, t) \quad (2.19)$$

$$g^k(x, t) \leq g^{k+1}(x, t) \quad (2.20)$$

for all  $t > 0$ , a.e. in  $B_{R_0}$ .

At this point, let us denote with  $p_{\varepsilon_n}$  and  $a_{\varepsilon_n}$  the regularization of the functions  $p$  and  $a$  respectively

$$p_{\varepsilon_n}(x) = \int_B \rho(y)p(x + \varepsilon_n y) dy \quad a_{\varepsilon_n}(x) = \int_B \rho(y)a(x + \varepsilon_n y) dy, \quad (2.21)$$

where  $B$  denotes the unit ball,  $\rho$  is a positive symmetric mollifier such that  $\int \rho = 1$  and where  $\varepsilon_n$  is an infinitesimal sequence of positive numbers. With an abuse of notation in the sequel we denote with  $\varepsilon = \varepsilon_n$  and with  $p_\varepsilon = p_{\varepsilon_n}$ ,  $a_\varepsilon = a_{\varepsilon_n}$ . It is well known that  $p_\varepsilon$  and  $a_\varepsilon$  converge in the strong topology of  $W^{1,r}(B_{R_0})$  and uniformly in  $B_{R_0}$  to the functions  $p$  and  $a$  respectively. Now let us define

$$\tilde{g}_t^{k\varepsilon}(x, t) := \begin{cases} a_\varepsilon(x)p_\varepsilon(x)h(t)^{p_\varepsilon(x)-1}h'(t) & 0 \leq t < k \\ a_\varepsilon(x)p_\varepsilon(x)h(k)^{p_\varepsilon(x)-1}h'(k) + \frac{\lambda}{p-1}[t^{p-1} - k^{p-1}] & t \geq k, \end{cases} \quad (2.22)$$

and

$$\tilde{g}^{k\varepsilon}(x, t) := \int_0^t \tilde{g}_t^{k\varepsilon}(x, s) ds + g(x, 0).$$

Finally consider

$$g^{k\varepsilon}(x, t) := \tilde{g}^{k\varepsilon}(x, t) + \varepsilon (1 + t^2)^{\frac{p}{2}}, \quad \forall t > 0, \text{ a.e. in } B_{R_0}. \quad (2.23)$$

By the properties of function  $g(x, t)$  given in the previous step, since  $p \leq p_\varepsilon(x) \leq q$  in  $B_{R_0}$  and by proceeding as in Lemmas 4.2 and Lemma 4.3 of [13], we have that the sequence  $g^{k\varepsilon}(x, t)$  satisfies the following inequalities a.e. in  $B_{R_0}$  and for all  $t > 0$ , the lower bound in (2.25) and (2.26) ensured by (2.23)

$$g^{k\varepsilon}(x, t) \leq C(k) (1 + t^2)^{\frac{p}{2}} \quad (2.24)$$

$$\varepsilon(1 + t^2)^{\frac{p-2}{2}} \leq \frac{g_t^{k\varepsilon}(x, t)}{t} \leq C(k)(1 + t^2)^{\frac{p-2}{2}} \quad (2.25)$$

$$\min\{p-1, 1\} \varepsilon(1 + t^2)^{\frac{p-2}{2}} \leq g_{tt}^{k\varepsilon}(x, t) \leq C(k)(1 + t^2)^{\frac{p-2}{2}} \quad (2.26)$$

$$|g_{tx}^{k\varepsilon}(x, t)| \leq C(k, \varepsilon, \omega_0) (1 + t^2)^{\frac{p-1}{2}} \quad \forall \omega_0 \subset\subset B_{R_0}. \quad (2.27)$$

Moreover the functions  $g^{k\varepsilon}$  fulfill for a.e.  $x \in B_{R_0}$  and  $t \geq t_0$

$$\lambda t^{p-2} \leq \frac{g_t^{k\varepsilon}(x, t)}{t} \leq \Lambda t^{q-2} \quad (2.28)$$

$$\lambda t^{p-2} \leq g_{tt}^{k\varepsilon}(x, t) \leq \Lambda t^{q-2} \quad (2.29)$$

$$|g_{tx}^{k\varepsilon}(x, t)| \leq \ell_\varepsilon(x)(1 + t^2)^{\frac{q-1}{2}}, \quad (2.30)$$

where  $\ell_\varepsilon \in \mathcal{C}^\infty(B_{R_0})$  is the regularized function of  $\ell$  is defined as in (2.15) and  $\lambda, \Lambda$  are as in (2.11), (2.12).

In the sequel, for simplicity of notations, we assume that  $t_0 = 1$ .

**2.5. Step 5: a priori estimates.** Let  $w \in W_{\text{loc}}^{1,p}(\Omega; \mathbb{R}^N)$  be a local minimizer of the functional (1.1); moreover let us take  $B_R \subset\subset B_{R_0}$  to be a ball of radius  $R$  compactly contained in  $B_{R_0}$ . Consider the following variational problem

$$\inf\{F^{k\varepsilon}(v) : v \in w + W_0^{1,p}(B_R; \mathbb{R}^N)\}. \quad (2.31)$$

where

$$F^{k\varepsilon}(v) = \int_{B_{R_0}} g^{k\varepsilon}(x, |Dv|) dx. \quad (2.32)$$



Since  $F^{k\varepsilon}$  is lower semicontinuous, there exists  $v^{k\varepsilon} \in w + W_0^{1,p}(B_{R_0}; \mathbb{R}^N)$  solution to Problem (2.31). The purpose of this Step is to prove an a priori estimate for the  $L^\infty$ -norm of  $Dv^{k\varepsilon}$  independent of  $k, \varepsilon$ . We claim that, for  $0 < \rho < R < R_0$

$$\|Dv^{k\varepsilon}\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq C \left[ \frac{\left(1 + \|\ell_\varepsilon\|_{L^r(B_{R_0})}^2\right)^{\frac{1}{2}}}{(R - \rho)} \right]^{\frac{\beta}{\frac{1}{n} - \frac{1}{r}}} \left[ \int_{B_R} (1 + g^{k\varepsilon}(x, |Dv^{k\varepsilon}|)) dx \right]^\beta, \quad (2.33)$$

with constants  $C, \beta$  independent of  $k, \varepsilon$ . Once (2.33) is obtained, we observe that, by convolution properties

$$\|\ell_\varepsilon\|_{L^r(B_{R_0})} \leq \|\ell\|_{L^r(B_{R_0})},$$

thus we deduce

$$\|Dv^{k\varepsilon}\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq C \left[ \frac{\left(1 + \|\ell\|_{L^r(B_{R_0})}^2\right)^{\frac{1}{2}}}{(R - \rho)} \right]^{\frac{\beta}{\frac{1}{n} - \frac{1}{r}}} \left[ \int_{B_R} (1 + g^{k\varepsilon}(x, |Dv^{k\varepsilon}|)) dx \right]^\beta. \quad (2.34)$$

The proof of estimate (2.33) turns to be quite similar to the proof of Proposition 3.1 of [13]; here, for sake of clarity, we list the main arguments, referring to [13] for more details. For simplicity of notations, from now on we set

$$f(x, \xi) = f^{k\varepsilon}(x, \xi) = g^{k\varepsilon}(x, |\xi|), \quad v^{k\varepsilon} := u, \quad \ell_\varepsilon := \ell.$$

Observe that, by (2.28), (2.29), (2.30), we have that, for  $|\xi| \geq 1$  and a.e.  $x \in B_{R_0}$

$$\lambda |\xi|^{p-2} |\mu|^2 \leq \sum_{i,j,\alpha,\beta} f_{\xi_i^\alpha \xi_j^\beta}(x, \xi) \mu_i^\alpha \mu_j^\beta, \quad (2.35)$$

$$|f_{\xi_i^\alpha \xi_j^\beta}(x, \xi)| \leq \Lambda |\xi|^{q-2}, \quad (2.36)$$

$$|f_{\xi x}(x, \xi)| \leq \ell(x) |\xi|^{q-1}, \quad (2.37)$$

hold, where  $p$  and  $q$  have been introduced in (2.1). The minimizer  $u$  satisfies the Euler's first variation

$$\int_{B_{R_0}} \sum_{i,\alpha} f_{\xi_i^\alpha}(x, Du) \varphi_{x_i}^\alpha(x) dx = 0 \quad \forall \varphi = (\varphi^\alpha)_{\alpha=1,\dots,N} \in W_0^{1,p}(B_{R_0}; \mathbb{R}^N),$$

and, by using the technique of the different quotients (see for example [14], [17], [19]) we have that

$$u \in W_{\text{loc}}^{2,\min(2,p)}(B_{R_0}; \mathbb{R}^N), \quad (1 + |Du|^2)^{\frac{p-2}{2}} |D^2u|^2 \in L_{\text{loc}}^1(B_{R_0}) \quad (2.38)$$

and the second variation

$$\int_{B_{R_0}} \left\{ \sum_{i,j,\alpha,\beta,s} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) \varphi_{x_i}^\alpha u_{x_s x_j}^\beta + \sum_{i,\alpha,s} f_{\xi_i^\alpha x_s}(x, Du) \varphi_{x_i}^\alpha \right\} dx = 0 \quad (2.39)$$

$$\forall s = 1, \dots, n, \quad \forall \varphi = (\varphi^\alpha)_{\alpha=1,\dots,N} \in W_0^{1,\min(2,p)}(B_{R_0}; \mathbb{R}^N).$$

Let  $\eta \in C_0^1(B_{R_0})$ . For any fixed  $s \in \{1, \dots, n\}$ , we choose

$$\varphi^\alpha = \eta^2 u_{x_s}^\alpha \Phi(|Du| - 1)_+$$

for  $\Phi : [0, +\infty) \rightarrow [0, +\infty)$  an increasing, locally Lipschitz continuous function, with  $\Phi$  and  $\Phi'$  bounded on  $[0, +\infty)$ , such that  $\Phi(0) = 0$  and satisfying

$$\Phi'(s)s \leq c_\Phi \Phi(s) \quad (2.40)$$

for a suitable value of  $c_\Phi$ . In the following  $(a)_+$  denotes the positive part of  $a \in \mathbb{R}$  and we write  $\Phi((|Du| - 1)_+) = \Phi(|Du| - 1)_+$ . Let us compute

$$\varphi_{x_i}^\alpha = 2\eta\eta_{x_i}u_{x_s}^\alpha \Phi(|Du| - 1)_+ + \eta^2u_{x_sx_i}^\alpha \Phi(|Du| - 1)_+ + \eta^2u_{x_s}^\alpha \Phi'(|Du| - 1)_+ [(|Du| - 1)_+]_{x_i}.$$

Here we used the fact that  $u \in W_{\text{loc}}^{1,\infty}(B_{R_0}; \mathbb{R}^N)$ , see Proposition 3.1 of [8]. Plugging this expression in (2.39) we obtain

$$\begin{aligned} 0 &= \int_{B_{R_0}} 2\eta\Phi(|Du| - 1)_+ \sum_{i,j,\alpha,\beta,s} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) \eta_{x_i} u_{x_s}^\alpha u_{x_sx_j}^\beta dx \\ &+ \int_{B_{R_0}} \eta^2\Phi(|Du| - 1)_+ \sum_{i,j,\alpha,\beta,s} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) u_{x_sx_i}^\alpha u_{x_sx_j}^\beta dx \\ &+ \int_{B_{R_0}} \eta^2\Phi'(|Du| - 1)_+ \sum_{i,j,\alpha,\beta,s} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) u_{x_s}^\alpha u_{x_sx_j}^\beta [(|Du| - 1)_+]_{x_i} dx \\ &+ \int_{B_{R_0}} 2\eta\Phi(|Du| - 1)_+ \sum_{i,\alpha,s} f_{\xi_i^\alpha x_s}(x, Du) \eta_{x_i} u_{x_s}^\alpha dx \\ &+ \int_{B_{R_0}} \eta^2\Phi(|Du| - 1)_+ \sum_{i,\alpha,s} f_{\xi_i^\alpha x_s}(x, Du) u_{x_sx_i}^\alpha dx \\ &+ \int_{B_{R_0}} \eta^2\Phi'(|Du| - 1)_+ \sum_{i,\alpha,s} f_{\xi_i^\alpha x_s}(x, |Du|) u_{x_s}^\alpha [(|Du| - 1)_+]_{x_i} dx \\ &=: I_1 + I_2 + I_3 + I_4 + I_5 + I_6. \end{aligned} \quad (2.41)$$

In the following, constants will be denoted by  $C$ , regardless of their actual value. We have that

$$I_1 + I_2 + I_3 = -(I_4 + I_5 + I_6).$$

Consider  $I_1$ ; by the Cauchy-Schwartz inequality, the Young inequality and (2.36), we have

$$\begin{aligned} |I_1| &= \left| \int_{B_{R_0}} 2\eta\Phi(|Du| - 1)_+ \sum_{i,j,s,\alpha,\beta} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) \eta_{x_i} u_{x_s}^\alpha u_{x_sx_j}^\beta dx \right| \quad (2.42) \\ &\leq \int_{B_{R_0}} 2\eta\Phi(|Du| - 1)_+ \left\{ \sum_{i,j,s,\alpha,\beta} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) \eta_{x_i} u_{x_s}^\alpha \eta_{x_j} u_{x_s}^\beta \right\}^{\frac{1}{2}} \\ &\quad \times \left\{ \sum_{i,j,s,\alpha,\beta} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) u_{x_sx_i}^\alpha u_{x_sx_j}^\beta \right\}^{\frac{1}{2}} dx \\ &\leq C \int_{B_{R_0}} |D\eta|^2 \Phi(|Du| - 1)_+ |Du|^q dx \end{aligned}$$

$$+ \frac{1}{2} \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ \sum_{i,j,s,\alpha,\beta} f_{\xi_i^\alpha \xi_j^\beta}(x, Du) u_{x_s x_i}^\alpha u_{x_s x_j}^\beta dx.$$

Therefore

$$\frac{1}{2} I_2 + I_3 \leq C \int_{B_{R_0}} |D\eta|^2 \Phi(|Du| - 1)_+ |Du|^q dx - (I_4 + I_5 + I_6).$$

By proceeding as in [13], (see also Lemma 4.1 of [25]), we have that

$$I_3 \geq \int_{B_{R_0}} \eta^2 \Phi'(|Du| - 1)_+ \frac{g_{tt}(x, |Du|)}{|Du|} \sum_{\alpha} \left( \sum_i u_{x_i}^\alpha [ (|Du| - 1)_+ ]_{x_i} \right)^2 dx \geq 0,$$

and (2.35) implies

$$I_2 \geq \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |Du|^{p-2} |D^2 u|^2 dx.$$

We now deal with  $|I_4|$ . We have, by (2.30)

$$\begin{aligned} |I_4| &= \left| \int_{B_{R_0}} 2\eta \Phi(|Du| - 1)_+ \sum_{i,s,\alpha} f_{\xi_i^\alpha x_s}(x, Du) \eta_{x_i} u_{x_s}^\alpha dx \right| \\ &\leq \int_{B_{R_0}} 2\eta \Phi(|Du| - 1)_+ |g_{tx}(x, |Du|)| \sum_{i,s,\alpha} |\eta_{x_i} u_{x_s}^\alpha| dx \\ &\leq \int_{B_{R_0}} (\eta^2 + |D\eta|^2) \ell(x) \Phi(|Du| - 1)_+ |Du|^q dx. \end{aligned}$$

Consider now  $|I_5|$ . We have, again by (2.30)

$$\begin{aligned} |I_5| &= \left| \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |g_{tx}(x, |Du|)| |D^2 u| dx \right| \\ &\leq \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ \ell(x) |Du|^{q-1} |D^2 u| dx \\ &\leq \int_{B_{R_0}} [\eta^2 \Phi(|Du| - 1)_+ |Du|^{p-2} |D^2 u|^2]^{1/2} [\eta^2 \Phi(|Du| - 1)_+ |\ell(x)|^2 |Du|^{2q-p}]^{1/2} dx \\ &\leq \varepsilon \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |Du|^{p-2} |D^2 u|^2 dx + C_\varepsilon \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |\ell(x)|^2 |Du|^{2q-p} dx, \end{aligned}$$

where in the last line we used the Young inequality. The estimate of  $I_6$  is more delicate. For any  $0 < \delta < 1$  (please, excuse the abuse of notation: the  $\delta$  used here is different from the one in (1.2))

$$\begin{aligned} |I_6| &= \left| \int_{B_{R_0}} \eta^2 |g_{tx}(x, |Du|)| |Du| \Phi'(|Du| - 1)_+ [ (|Du| - 1)_+ ]_{x_i} dx \right| \\ &\leq \int_{B_{R_0}} \eta^2 \Phi'(|Du| - 1)_+ \ell(x) |Du|^q |D^2 u| dx \end{aligned}$$

$$\begin{aligned}
&= \int_{B_{R_0}} \eta^2 \Phi'(|Du| - 1)_+ \ell(x) [ (|Du| - 1)_+ + \delta ] [ (|Du| - 1)_+ + \delta ]^{-1} |Du|^q |D^2u| dx \\
&\leq \int_{B_{R_0}} \eta^2 \left\{ \frac{1}{c_\Phi} \Phi'(|Du| - 1)_+ [ (|Du| - 1)_+ + \delta ] |Du|^{p-2} |D^2u|^2 \right\}^{1/2} \\
&\quad \times \left\{ c_\Phi \Phi'(|Du| - 1)_+ |\ell(x)|^2 |Du|^{2q-p+2} [ (|Du| - 1)_+ + \delta ]^{-1} \right\}^{1/2} dx \\
&\leq \frac{\varepsilon}{c_\Phi} \int_{B_{R_0}} \eta^2 \Phi'(|Du| - 1)_+ [ (|Du| - 1)_+ + \delta ] |Du|^{p-2} |D^2u|^2 dx \\
&\quad + C_\varepsilon c_\Phi \int_{B_{R_0}} \eta^2 \Phi'(|Du| - 1)_+ |\ell(x)|^2 |Du|^{2q-p+2} [ (|Du| - 1)_+ + \delta ]^{-1} dx.
\end{aligned}$$

To estimate the first term in the last inequality we split  $B_{R_0} = \{x : |Du(x)| \geq 2\} \cup \{x : |Du(x)| < 2\}$  and we observe that in the set  $\{x : |Du(x)| \geq 2\}$  we have  $(|Du| - 1)_+ \geq 1$ , since  $\delta < 1$

$$(|Du| - 1)_+ + \delta \leq 2(|Du| - 1)_+. \quad (2.43)$$

Therefore we have, using (2.40)

$$\begin{aligned}
&\int_{|Du| \geq 2} \eta^2 \Phi'(|Du| - 1)_+ [ (|Du| - 1)_+ + \delta ] |Du|^{p-2} |D^2u|^2 dx \\
&+ \int_{1 < |Du| < 2} \eta^2 \Phi'(|Du| - 1)_+ [ (|Du| - 1)_+ + \delta ] |Du|^{p-2} |D^2u|^2 dx \\
&\leq 2 \int_{|Du| \geq 2} \eta^2 \Phi'(|Du| - 1)_+ (|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx \\
&+ \int_{1 < |Du| < 2} \eta^2 \Phi'(|Du| - 1)_+ (|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx \\
&+ \delta \int_{1 < |Du| < 2} \eta^2 \Phi'(|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx \\
&\leq 2 c_\Phi \int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx \\
&+ \delta \int_{1 < |Du| < 2} \eta^2 \Phi'(|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx.
\end{aligned}$$

Now, for  $\varepsilon$  sufficiently small, by collecting the previous estimates, we deduce

$$\begin{aligned}
&\int_{B_{R_0}} \eta^2 \Phi(|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx \quad (2.44) \\
&\leq C c_\Phi \int_{B_{R_0}} (\eta^2 + |D\eta|^2) (1 + \ell^2(x)) |Du|^{2q-p} \\
&\quad \times [\Phi(|Du| - 1)_+ |Du|^{2q-p} + \Phi'(|Du| - 1)_+ |Du|^2 [ (|Du| - 1)_+ + \delta ]^{-1}] dx \\
&+ \delta \int_{1 < |Du| < 2} \eta^2 \Phi'(|Du| - 1)_+ |Du|^{p-2} |D^2u|^2 dx,
\end{aligned}$$

with a constant  $C$  depending on  $n, r, p, q$ .

Let define the function

$$\Phi(s) := (1 + s)^{\gamma-2} s^2 \quad \gamma \geq 0; \quad (2.45)$$

which satisfies (2.40) with  $c_\Phi = 2(1 + \gamma)$ . We can approximate  $\Phi$  in (2.45) by a sequence of functions  $\Phi_r$ , each of them being equal to  $\Phi$  in the interval  $[0, r]$ , and then extended to  $[r, +\infty)$  with the constant value  $\Phi(r)$ . Then we insert  $\Phi_r$  in (2.44) and, passing to the limit as  $r \rightarrow +\infty$  by the Monotone Convergence Theorem, we obtain for every  $0 < \delta < 1$

$$\begin{aligned} & \int_{B_{R_0}} \eta^2 (1 + (|Du| - 1)_+)^{\gamma-2} (|Du| - 1)_+^2 |Du|^{p-2} |D^2u|^2 dx \\ & \leq C (1 + \gamma)^2 \int_{B_{R_0}} (\eta^2 + |D\eta|^2) (1 + \ell(x)^2) (1 + (|Du| - 1)_+)^{\gamma+2q-p} dx \\ & \quad + \delta c(\gamma) \int_{1 < |Du| < 2} \eta^2 |Du|^{p-2} |D^2u|^2 dx, \end{aligned} \quad (2.46)$$

as

$$\frac{(|Du| - 1)_+}{(|Du| - 1)_+ + \delta} \leq 1 \quad \forall \delta > 0.$$

By the elementary inequality (for the proof see [13])

$$C_1 (1 + |\xi|^2)^{\frac{p-2}{2}} \leq |\xi|^{p-2} \leq C_2 (1 + |\xi|^2)^{\frac{p-2}{2}}, \quad |\xi| \geq t_0,$$

with  $C_1, C_2$  depending on  $t_0$  (here  $t_0 = 1$ ), we have

$$\int_{1 < |Du| < 2} \eta^2 |Du|^{p-2} |D^2u|^2 dx \leq C \int_{1 < |Du| < 2} \eta^2 (1 + |Du|^2)^{\frac{p-2}{2}} |D^2u|^2 dx < +\infty$$

and, due to (2.38), the last integral is finite. As  $\delta \rightarrow 0$ , the last term in (2.46) vanishes.

Now, let us define

$$m := \left(\frac{r}{2}\right)' = \frac{r}{r-2}. \quad (2.47)$$

Since  $\ell \in L^r(B_{R_0})$ , by the Hölder inequality

$$\begin{aligned} & \int_{B_{R_0}} \eta^2 (1 + (|Du| - 1)_+)^{\gamma-2} (|Du| - 1)_+^2 |Du|^{p-2} |D((|Du| - 1)_+)|^2 dx \\ & \leq C (1 + \gamma)^2 H \left[ \int_{B_{R_0}} (\eta^2 + |D\eta|^2)^m (1 + (|Du| - 1)_+)^{(\gamma+2q-p)m} dx \right]^{\frac{1}{m}}, \end{aligned} \quad (2.48)$$

where  $C$  depends also on  $r$  and  $|B_{R_0}|$  (and so on  $n$ ) and

$$H := (1 + \|\ell\|_{L^r(B_{R_0})}^2). \quad (2.49)$$

Let us introduce

$$G(t) = 1 + \int_0^t \sqrt{\Phi(s)} (1 + s)^{\frac{p-2}{2}} ds = 1 + \int_0^t (1 + s)^{\frac{\gamma}{2} + \frac{p}{2} - 2} s ds; \quad (2.50)$$

since  $p \leq q \leq 2q - p$  we get

$$[G(t)]^2 \leq 4(1 + t)^{\gamma+p} \leq 4(1 + t)^{\gamma+2q-p}. \quad (2.51)$$

Moreover

$$G_t(t) = \sqrt{\Phi(t)} (1+t)^{\frac{p-2}{2}} \stackrel{(2.45)}{=} (1+t)^{\frac{\gamma}{2} + \frac{p}{2} - 2} t. \quad (2.52)$$

Set  $w = \eta G((|Du| - 1)_+)$ , we have

$$\begin{aligned} & \int_{B_{R_0}} |D(\eta G((|Du| - 1)_+))|^2 dx \\ & \leq 2 \int_{B_{R_0}} |D\eta|^2 |G((|Du| - 1)_+)|^2 dx \\ & \quad + 2 \int_{B_{R_0}} \eta^2 [G_t((|Du| - 1)_+)]^2 [D((|Du| - 1)_+)]^2 dx \\ & \leq C (1+\gamma)^2 H \left[ \int_{B_{R_0}} (\eta^2 + |D\eta|^2)^m [1 + (|Du| - 1)_+^{(\gamma+2q-p)m}] dx \right]^{\frac{1}{m}}. \end{aligned} \quad (2.53)$$

Now, let  $2^* = \frac{2n}{n-2}$  for  $n > 2$ , while  $2^*$  equal to any fixed real number larger than 2, if  $n = 2$ . By Sobolev's inequality there exists a constant  $C$  such that

$$\left\{ \int_{B_{R_0}} [\eta G((|Du| - 1)_+)]^{2^*} dx \right\}^{\frac{2}{2^*}} \leq C \int_{B_{R_0}} |D(\eta G((|Du| - 1)_+))|^2 dx. \quad (2.54)$$

Moreover, since  $r > n$ , we have

$$1 \leq m := \frac{r}{r-2} < \frac{n}{n-2} = \frac{2^*}{2}. \quad (2.55)$$

Observe that

$$(2q-p)m = 2(q-p)m + pm; \quad (2.56)$$

and in view of the strict inequality in (2.3), we infer the existence of  $0 < \epsilon < 1$  such that

$$(q-p) + \epsilon \left( \frac{1}{n} - \frac{1}{r} \right) \leq p \left( \frac{1}{n} - \frac{1}{r} \right). \quad (2.57)$$

Set now

$$0 < \tilde{M} := 2(q-p)m + p(m-1) + \epsilon \quad 0 < \tilde{N} := p - \epsilon. \quad (2.58)$$

We have

$$\tilde{M} + \tilde{N} = (2q-p)m \quad \text{and} \quad \tilde{M} > (2q-p)m - p. \quad (2.59)$$

By the assumptions on  $p, q$  and the definition of  $\tilde{M}$  and  $\tilde{N}$ , we claim that

$$\frac{1}{(\gamma+p)^2} \left[ \int_{B_{R_0}} \eta^{2^*} [1 + (|Du| - 1)_+]^{(\gamma + \frac{\tilde{M}}{m}) \frac{2^*}{2} + \tilde{N}} dx \right]^{\frac{2}{2^*}} \leq 4 \left( \int_{B_{R_0}} [\eta G((|Du| - 1)_+)]^{2^*} dx \right)^{\frac{2}{2^*}}. \quad (2.60)$$

By the definition of  $G$ , if we set  $t := (|Du| - 1)_+$ , (2.60) is proved once we get

$$\frac{1}{\gamma+p} (1+t)^{\left(\frac{\gamma}{2} + \frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*}\right)} \leq 2G(t) = 2 \left( 1 + \int_0^t (1+s)^{\frac{\gamma}{2} + \frac{p}{2} - 2} s ds \right). \quad (2.61)$$

If  $t \leq 1$ ; (2.61) holds, in fact

$$\frac{1}{\gamma + p} (1 + t)^{\left(\frac{\gamma}{2} + \frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*}\right)} \leq \frac{2}{\gamma + p} \leq 2 \leq 2 \left(1 + \int_0^t (1 + s)^{\frac{\gamma}{2} + \frac{p}{2} - 2} s \, ds\right).$$

Let now  $t \geq 1$ ; then (2.61) becomes, after differentiation

$$\frac{\frac{\gamma}{2} + \frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*}}{\gamma + p} (1 + t)^{\frac{\gamma}{2} + \frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*} - 1} \leq 2(1 + t)^{\frac{\gamma}{2} + \frac{p}{2} - 2} t. \quad (2.62)$$

Since

$$\frac{1}{2m} - \frac{1}{2^*} = \frac{1}{n} - \frac{1}{r},$$

we have that

$$(q - p) + \frac{p}{2} - \frac{p}{2m} + \frac{\epsilon}{2m} + \frac{p}{2^*} - \frac{\epsilon}{2^*} \leq \frac{p}{2}$$

which implies, by the definition of  $\tilde{M}$  and  $\tilde{N}$

$$\frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*} \leq \frac{p}{2}. \quad (2.63)$$

We then have

$$\frac{\frac{\gamma}{2} + \frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*}}{\gamma + p} (1 + t)^{\frac{\tilde{M}}{2m} + \frac{\tilde{N}}{2^*}} \leq \frac{1}{2} (1 + t)^{\frac{p}{2}} = \frac{1}{2} (1 + t)^{\frac{p-2}{2}} (1 + t)^{t \geq 1} \leq (1 + t)^{\frac{p-2}{2}} t$$

and so (2.62) is satisfied.

By collecting (2.53), (2.54) and (2.60), we obtain

$$\begin{aligned} & \left[ \int_{B_{R_0}} \eta^{2^*} [1 + (|Du| - 1)_+]^{\left(\gamma + \frac{\tilde{M}}{m}\right) \frac{2^*}{2}} [1 + (|Du| - 1)_+]^{\tilde{N}} dx \right]^{\frac{2}{2^*}} \\ & \leq C H (\gamma + 2q - p)^4 \left[ \int_{B_{R_0}} (\eta^2 + |D\eta|^2)^m [1 + (|Du| - 1)_+]^{(\gamma + 2q - p)m} dx \right]^{\frac{1}{m}}, \end{aligned} \quad (2.64)$$

where the constant  $C$  only depends on  $n, r, p, q, \lambda, \Lambda$ .

Now, let us choose  $0 < \rho < R < R_0$  and let  $\eta$  to be equal to 1 in  $B_\rho$ , with  $\text{supp } \eta \subset B_R$  and such that  $|D\eta| \leq \frac{1}{(R-\rho)}$ . Let us denote by

$$\kappa := \gamma m + \tilde{M} = (\gamma + 2q - p)m - \tilde{N}.$$

From (2.64) it follows

$$\begin{aligned} & \left\{ \int_{B_\rho} [1 + (|Du| - 1)_+]^{\kappa \frac{2^*}{2m}} [1 + (|Du| - 1)_+]^{\tilde{N}} dx \right\}^{\frac{2m}{2^*}} \\ & \leq C H^m \left( \frac{(\kappa + \tilde{N})^2}{R - \rho} \right)^{2m} \int_{B_R} [1 + (|Du| - 1)_+]^\kappa [1 + (|Du| - 1)_+]^{\tilde{N}} dx. \end{aligned} \quad (2.65)$$

Fixed  $\bar{R}$  and  $\bar{\rho}$ , with  $\bar{R} > \bar{\rho}$ , we define the decreasing sequence of radii  $\{\rho_i\}_{i \geq 0}$

$$\rho_i = \bar{\rho} + \frac{\bar{R} - \bar{\rho}}{2^i}, \quad \forall i \geq 0, \quad R - \rho := \rho_i - \rho_{i+1} = \frac{\bar{R} - \bar{\rho}}{2^{i+1}}$$

with  $\rho_0 = \bar{R} > \rho_i > \rho_{i+1} > \bar{\rho}$  and the increasing sequence  $\{\kappa_i\}_{i \geq 0}$

$$\kappa_0 := \tilde{M} \quad \kappa_{i+1} = \kappa_i \frac{2^*}{2m} \quad i \geq 0.$$

We rewrite (2.65) and we obtain for every  $i \geq 0$

$$A_{i+1} \leq C_i A_i, \quad (2.66)$$

where

$$A_i := \left( \int_{B_{\rho_i}} [1 + (|Du| - 1)_+]^{\kappa_i} [1 + (|Du| - 1)_+]^{\tilde{N}} dx \right)^{\frac{1}{\kappa_i}}$$

$$C_i := \left[ C H^m \left( \frac{(\kappa_i + \tilde{N})^{\frac{3}{2}} 2^{i+1}}{\bar{R} - \bar{\rho}} \right)^{2m} \right]^{\frac{1}{\kappa_i}}.$$

By iteration of (2.66), we deduce

$$\left\{ \int_{B_{\bar{\rho}}} [1 + (|Du| - 1)_+]^{\kappa_0 \left(\frac{2^*}{2m}\right)^{i+1}} [1 + (|Du| - 1)_+]^{\tilde{N}} dx \right\}^{\left(\frac{2m}{2^*}\right)^{i+1}}$$

$$\leq \tilde{C} \int_{B_{\bar{R}}} [1 + (|Du| - 1)_+]^{(2q-p)m} dx, \quad (2.67)$$

where

$$\tilde{C} \leq \prod_{k=0}^{\infty} \left[ C H^m \left( \frac{(\kappa_k + \tilde{N})^{\frac{3}{2}} 2^{k+1}}{\bar{R} - \bar{\rho}} \right)^{2m} \right]^{\left(\frac{2m}{2^*}\right)^k} \leq \frac{C H^{2\left(\frac{1}{n} - \frac{1}{r}\right)}}{(\bar{R} - \bar{\rho})^{\frac{2 \cdot 2^* m}{2^* - 2m}}}, \quad (2.68)$$

with a constant  $C = C(n, r, p, q)$ . Let us denote

$$\tau := \frac{2 \cdot 2^* m}{2^* - 2m} = \frac{1}{\frac{1}{n} - \frac{1}{r}}; \quad (2.69)$$

thus (2.67) implies

$$\left\{ \int_{B_{\bar{\rho}}} [1 + (|Du| - 1)_+]^{\kappa_0 \left(\frac{2^*}{2m}\right)^{i+1}} dx \right\}^{\left(\frac{2m}{2^*}\right)^{i+1}} \leq C \left[ \frac{\sqrt{H}}{(\bar{R} - \bar{\rho})} \right]^{\tau} \int_{B_{\bar{R}}} [1 + (|Du| - 1)_+]^{(2q-p)m} dx. \quad (2.70)$$

At this point we pass to the limit as  $i \rightarrow +\infty$ , obtaining

$$\sup_{x \in B_{\bar{\rho}}} [1 + (|Du|(x) - 1)_+]^{\tilde{M}} = \lim_{i \rightarrow +\infty} \left\{ \int_{B_{\bar{\rho}}} [1 + (|Du| - 1)_+]^{\tilde{M} \left(\frac{2^*}{2m}\right)^{i+1}} dx \right\}^{\left(\frac{2m}{2^*}\right)^{i+1}}$$

$$\leq C \left[ \frac{\sqrt{H}}{(\bar{R} - \bar{\rho})} \right]^{\tau} \int_{B_{\bar{R}}} [1 + (|Du| - 1)_+]^{(2q-p)m} dx. \quad (2.71)$$



Set

$$V(x) := 1 + (|Du|(x) - 1)_+ \quad \text{and} \quad s := (2q - p)m; \quad (2.72)$$

the estimate (2.71) becomes, for any  $\rho < R < R_0$

$$\sup_{x \in B_\rho} |V(x)| \leq C \left( \left[ \frac{\sqrt{H}}{(R - \rho)} \right]^{\frac{\tau}{s}} \|V\|_{L^s(B_R)} \right)^{\frac{s}{M}}. \quad (2.73)$$

Our aim is to estimate the essential supremum of  $|Du|$  in terms of its  $L^p$ -norm. By classical interpolation inequalities

$$\|V\|_{L^s(B_\rho)} \leq \|V\|_{L^p(B_\rho)}^{\frac{p}{s}} \|V\|_{L^\infty(B_\rho)}^{1-\frac{p}{s}}, \quad (2.74)$$

and (2.73) and (2.74) give

$$\|V\|_{L^s(B_\rho)} \leq C^{1-\frac{p}{s}} \|V\|_{L^p(B_\rho)}^{\frac{p}{s}} \left( \left[ \frac{\sqrt{H}}{(\bar{R} - \bar{\rho})} \right]^{\frac{\tau}{s}} \|V\|_{L^s(B_R)} \right)^\theta \quad (2.75)$$

where

$$\theta := \frac{s}{M} \left( 1 - \frac{p}{s} \right) = \frac{1}{M} (s - p) = \frac{1}{M} [(2q - p)m - p] < 1. \quad (2.76)$$

For  $0 < \bar{\rho} < \bar{R}$  and for every  $k \geq 0$ , let us define

$$\rho_k := \bar{R} - (\bar{R} - \bar{\rho})2^{-k} \quad B_k := \|V\|_{L^s(B_{\rho_k})}.$$

By inserting in (2.75)  $\rho = \rho_k$  and  $R = \rho_{k+1}$ , (so that  $R - \rho = (\bar{R} - \bar{\rho})2^{-(k+1)}$ ), we have for every  $k \geq 0$

$$B_k \leq C^{1-\frac{p}{s}} \|V\|_{L^p(B_{\bar{R}})}^{\frac{p}{s}} \left( 2^{\frac{\tau}{s}(k+1)} \left[ \frac{\sqrt{H}}{(\bar{R} - \bar{\rho})} \right]^{\frac{\tau}{s}} B_{k+1} \right)^\theta. \quad (2.77)$$

By iteration of (2.77), we deduce for  $k \geq 0$

$$B_0 \leq \left( C^{1-\frac{p}{s}} \left[ \frac{\sqrt{H}}{(\bar{R} - \bar{\rho})} \right]^{\frac{\tau}{s}\theta} \|V\|_{L^p(B_{\bar{R}})}^{\frac{p}{s}} \right)^{\sum_{i=0}^k \theta^i} 2^{\frac{\tau}{s} \sum_{i=0}^{k+1} i\theta^i} (B_{k+1})^{\theta^{k+1}}. \quad (2.78)$$

By (2.76), the series appearing in (2.78) are convergent. Since  $B_k$  is bounded independently of  $k$ , i.e.

$$B_{k+1} \leq \|V\|_{L^s(B_{\bar{R}})},$$

we can pass to the limit as  $k \rightarrow +\infty$  and we obtain for every  $0 < \rho < R$  with a constant  $C = C(n, r, p, q)$  independent of  $k$

$$\|V\|_{L^s(B_\rho)} \leq C \left( \left[ \frac{\sqrt{H}}{(R - \rho)} \right]^{\frac{\tau}{s}\theta} \|V\|_{L^p(B_R)}^{\frac{p}{s}} \right)^{\frac{1}{1-\theta}}. \quad (2.79)$$

Combining (2.73) and (2.79), by setting  $\rho' = \frac{(R+\rho)}{2}$  we have

$$\|V\|_{L^\infty(B_\rho)} \leq C \left( \left[ \frac{\sqrt{H}}{(\rho' - \rho)} \right]^{\frac{\tau}{s}(1-\theta)} \left[ \frac{\sqrt{H}}{(R - \rho')} \right]^{\frac{\tau}{s}\theta} \|V\|_{L^p(B_R)}^{\frac{p}{s}} \right)^{\frac{s}{M} \frac{1}{1-\theta}}, \quad (2.80)$$

which implies

$$\|Du\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq \|V\|_{L^\infty(B_\rho)} \leq C \left[ \frac{\sqrt{H}}{(R - \rho)} \right]^{\tilde{\beta}} \left( \int_{B_R} (1 + |Du|^p) dx \right)^\beta,$$

with

$$\beta = \frac{1}{\tilde{M} - s + p} = \frac{1}{\epsilon} > 1, \quad \tilde{\beta} = \frac{1}{\frac{1}{n} - \frac{1}{r}} \frac{1}{\epsilon} = \frac{1}{\frac{1}{n} - \frac{1}{r}} \beta. \quad (2.81)$$

Since

$$\frac{\lambda}{p-1} t^p \leq g^{k\varepsilon}(x, t) = a_\varepsilon(x) h(t)^{p_\varepsilon(x)} \quad \forall t \geq 1 \quad \text{a.e. in } B_{R_0},$$

the estimate (2.33) follows.

**2.6. Step 6: comparison and conclusion.** We go back to Problem (2.31) and we observe that, since

$$t^p \leq C(1 + g^{k\varepsilon}(x, t)) \quad \forall t > 0, \quad \text{for a.e. } x \in B_{R_0},$$

by the minimality of  $v^{k\varepsilon}$ , we have

$$\int_{B_R} |Dv^{k\varepsilon}|^p dx \leq C \int_{B_R} [1 + g^{k\varepsilon}(x, |Dv^{k\varepsilon}|)] dx \leq C \int_{B_R} [1 + g^{k\varepsilon}(x, |Dw|)] dx. \quad (2.82)$$

Moreover, by the convolution properties, as  $\varepsilon \rightarrow 0$

$$g^{k\varepsilon}(x, |Dw|) \rightarrow g^k(x, |Dw|) \quad \text{a.e. in } B_{R_0},$$

and

$$g^{k\varepsilon}(x, |Dw|) \leq C(k)(1 + |Dw|^2)^{\frac{p}{2}} \in L^1(B_{R_0}).$$

The Lebesgue Dominated Convergence Theorem and (2.19) imply

$$\lim_{\varepsilon} \int_{B_R} g^{k\varepsilon}(x, |Dw|) dx = \int_{B_R} g^k(x, |Dw|) dx \stackrel{(2.19)}{\leq} \int_{B_R} a(x) h(|Dw|)^{p(x)} dx. \quad (2.83)$$

By collecting (2.82) and (2.83)

$$\sup_{\varepsilon} \int_{B_R} |Dv^{k\varepsilon}|^p dx \leq C \int_{B_R} [1 + a(x) h(|Dw|)^{p(x)}] dx. \quad (2.84)$$

Therefore there exists  $v^k \in w + W_0^{1,p}(B_R; \mathbb{R}^N)$  such that

$$v^{k\varepsilon} \rightharpoonup v^k \quad \text{weakly in } W^{1,p}(B_R; \mathbb{R}^N).$$

Moreover, by (2.82) and (2.83) we get, for all  $B_\rho \subset\subset B_R$

$$v^{k\varepsilon} \overset{*}{\rightharpoonup} v^k \quad \text{weakly star in } W^{1,\infty}(B_\rho; \mathbb{R}^N).$$

By the semicontinuity of the norm and (2.84), we obtain

$$\int_{B_R} |Dv^k|^p dx \leq \liminf_{\varepsilon} \int_{B_R} |Dv^{k\varepsilon}|^p dx \leq C \int_{B_R} [1 + a(x)h(|Dw|)^{p(x)}] dx. \quad (2.85)$$

On the other hand, (2.34) and (2.83) imply

$$\|Dv^k\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq \liminf_{\varepsilon} \|Dv^{k\varepsilon}\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq \hat{C} \left[ \int_{B_R} [1 + a(x)h(|Dw|)^{p(x)}] dx \right]^\beta =: M, \quad (2.86)$$

where we set for brevity

$$\hat{C} := C \left[ \frac{\left(1 + \|\ell\|_{L^r(B_{R_0})}^2\right)^{\frac{1}{2}}}{(R - \rho)} \right]^{\frac{\beta}{\frac{1}{n} - \frac{1}{r}}}.$$

Thus we can deduce that, up to subsequences, there exist  $v \in w + W_0^{1,p}(B_R; \mathbb{R}^N)$  such that

$$\begin{aligned} v^k &\rightharpoonup v && \text{weakly in } W^{1,p}(B_R; \mathbb{R}^N) \\ v^k &\overset{*}{\rightharpoonup} v && \text{weakly star in } W^{1,\infty}(B_\rho; \mathbb{R}^N) \quad \text{for all } B_\rho \subset\subset B_R. \end{aligned}$$

Let us show that  $v$  is a solution to the problem

$$\inf \left\{ \int_{B_R} a(x)h(|Dv|)^{p(x)} dx : v \in w + W_0^{1,p}(B_R, \mathbb{R}^N) \right\}. \quad (2.87)$$

To this end, using the semicontinuity of the functional  $\int_{B_\rho} g^{k_0}(x, |Dw|) dx$  and (see (2.20))

$$g^{k_0}(x, t) \leq g^k(x, t) \quad \forall k \geq k_0,$$

we get

$$\int_{B_\rho} g^{k_0}(x, |Dv^k|) dx \leq \liminf_{\varepsilon} \int_{B_\rho} g^{k_0}(x, |Dv^{k\varepsilon}|) dx \leq \liminf_{\varepsilon} \int_{B_\rho} g^k(x, |Dv^{k\varepsilon}|) dx. \quad (2.88)$$

Since, up to subsequences,  $g^{k\varepsilon}(x, t)$  converges as  $\varepsilon \rightarrow 0$ , a.e. in  $B_{R_0} \times [0, +\infty)$  to  $g^k(x, t)$ , by Egorov theorem, fixed  $K = \{\xi \in \mathbb{R}^{Nn} : |\xi| \leq M + 1\}$ , for every  $\delta > 0$  there exists  $A_\delta$  with  $|A_\delta| < \delta$  such that  $g^{k\varepsilon}$  converges to  $g^k$  uniformly in  $(B_\rho \setminus A_\delta) \times K$ . Thus

$$\limsup_{\varepsilon} \int_{B_\rho \setminus A_\delta} g^k(x, |Dv^{k\varepsilon}|) dx = \limsup_{\varepsilon} \int_{B_\rho \setminus A_\delta} g^{k\varepsilon}(x, |Dv^{k\varepsilon}|) dx$$

and due to (2.86)

$$\limsup_{\varepsilon} \int_{B_\rho \cap A_\delta} g^k(x, |Dv^{k\varepsilon}|) dx \leq C(k) |A_\delta| (1 + M^q),$$

with  $C(k)$  independent of  $\delta$ . Thus, putting together the previous inequalities, (2.88) gives

$$\int_{B_\rho} g^{k_0}(x, |Dv^k|) dx \leq \limsup_{\varepsilon} \int_{B_R} g^{k\varepsilon}(x, |Dv^{k\varepsilon}|) dx + C(k) |A_\delta| (1 + M^q)$$

so that, letting  $\delta \rightarrow 0$ , by (2.83)

$$\int_{B_\rho} g^{k_0}(x, |Dv^k|) dx \leq \limsup_\varepsilon \int_{B_R} g^{k\varepsilon}(x, |Dw|) dx = \int_{B_R} g^k(x, |Dw|) dx.$$

At this point, by the lower semicontinuity of the functional  $\int_{B_\rho} g^{k_0}(x, |Dw|) dx$ , and the Lebesgue Dominated Convergence Theorem applied to the sequence of functions  $g^k(x, |Dw|)$ , we obtain

$$\int_{B_\rho} g^{k_0}(x, |Dv|) dx \leq \liminf_k \int_{B_\rho} g^{k_0}(x, |Dv^k|) dx \leq \int_{B_R} g(x, |Dw|) dx.$$

Finally, letting  $k_0 \rightarrow +\infty$  and  $\rho \rightarrow R$

$$\int_{B_R} a(x)h(|Dv|)^{p(x)} dx \leq \int_{B_R} a(x)h(|Dw|)^{p(x)} dx, \quad (2.89)$$

and passing to the limit in (2.86), we get

$$\|Dv\|_{L^\infty(B_\rho; \mathbb{R}^{Nn})} \leq \hat{C} \left[ \int_{B_R} [1 + a(x)h(|Dw|)^{p(x)}] dx \right]^\beta. \quad (2.90)$$

Then  $w$  and  $v$  are two solutions to Problem (2.87), but since  $g$  is not strictly convex for all  $t > 0$ , we may not conclude that  $w = v$  in  $B_R$ . Let us define

$$E_0 := \left\{ x \in B_R : \left| \frac{Dw(x) + Dv(x)}{2} \right| > 1 \right\} \quad \text{and} \quad \bar{w} := \frac{w + v}{2}.$$

If  $E_0$  has positive measure, then from the convexity of  $g(x, \cdot)$  we have

$$\int_{B_R \setminus E_0} g(x, |D\bar{w}|) dx \leq \frac{1}{2} \int_{B_R \setminus E_0} g(x, |Dw|) dx + \frac{1}{2} \int_{B_R \setminus E_0} g(x, |Dv|) dx. \quad (2.91)$$

Now, by the strict convexity of  $g(x, t)$  for  $t \geq 1$ , we have

$$\int_{B_R \cap E_0} g(x, |D\bar{w}|) dx < \frac{1}{2} \int_{B_R \cap E_0} g(x, |Dw|) dx + \frac{1}{2} \int_{B_R \cap E_0} g(x, |Dv|) dx. \quad (2.92)$$

Adding (2.91) and (2.92), we get a contradiction with the minimality of  $w$  and  $v$ . Therefore the set  $E_0$  has zero measure, which implies that

$$\sup_{B_\rho} |Dw(x)| \leq \sup_{B_\rho} |Dw(x) + Dv(x)| + \sup_{B_\rho} |Dv(x)| \leq 2 + \sup_{B_\rho} |Dv(x)|$$

and the main estimate follows by (2.90). □

### 3. AN EXAMPLE

In this section we show that  $h(t)$ , defined in (1.3) with  $t_0$  to be chosen later,

$$h(t) = t^{a+b \sin \log \log t} = e^{(a+b \sin \log \log t) \log t}, \quad \forall t \geq t_0, \quad (3.1)$$

satisfies the assumptions of Theorem 1.1. First let us notice that  $h$  satisfies, for large  $t$ , the growth condition

$$t^p \leq h(t) \leq t^q, \quad t \geq t_0$$

with  $p = a - b$  and  $q = a + b$ .

Moreover, the first derivative, when  $t \geq t_0$ , is

$$\begin{aligned} h'(t) &= t^{a+b \sin \log \log t} \left[ b \cos \log \log t \cdot \frac{1}{\log t} \cdot \frac{1}{t} \cdot \log t + (a + b \sin \log \log t) \cdot \frac{1}{t} \right] \\ &= t^{a-1+b \sin \log \log t} [b \cos \log \log t + (a + b \sin \log \log t)]. \end{aligned}$$

Then  $h'(t) \geq 0$  for  $t \geq t_0$  if  $a \geq \sqrt{2}b$  (since  $\sin \alpha + \cos \alpha \geq -\sqrt{2}$  for all  $\alpha \in \mathbb{R}$ ).

The second derivative, when  $t \geq t_0$ , similarly is

$$\begin{aligned} h''(t) &= t^{a-2+b \sin \log \log t} \cdot [b \cos \log \log t + (a - 1 + b \sin \log \log t)] \cdot [b \cos \log \log t + (a + b \sin \log \log t)] \\ &\quad + t^{a-1+b \sin \log \log t} \left[ b (\cos \log \log t - \sin \log \log t) \cdot \frac{1}{\log t} \cdot \frac{1}{t} \right] \\ &= t^{a-2+b \sin \log \log t} \cdot \left\{ [b \cos \log \log t + (a - 1 + b \sin \log \log t)] \cdot [b \cos \log \log t + (a + b \sin \log \log t)] \right. \\ &\quad \left. + b (\cos \log \log t - \sin \log \log t) \cdot \frac{1}{\log t} \right\}; \end{aligned}$$

since the last addendum converges to zero at  $t \rightarrow +\infty$ , if  $a - 1 > \sqrt{2}b$ , for large values of  $t$   $h''(t) > 0$  and we also have

$$0 < \frac{h''(t) \cdot t}{h'(t)} \leq \frac{a - 1 + \sqrt{2}b}{a - \sqrt{2}b}.$$

Therefore the function  $h(t)$  satisfies (1.2) for a given  $\delta > 0$ , with  $a, b > 0$  such that

$$1 + \sqrt{2}b < a < a + b < 1 + \delta,$$

with  $t_0 > 0$  large in dependence of  $a, b$ . This is possible if  $1 < a < 1 + \delta$  and  $b$  is sufficiently small, say  $b < \min \left\{ \frac{a-1}{\sqrt{2}}, 1 + \delta - a \right\}$ .

## REFERENCES

- [1] E. ACERBI, G. MINGIONE: Regularity results for a class of functionals with nonstandard growth, *Arch. Rat. Mech. Anal.*, **156** (2001), 121-140.
- [2] M. BOCEA, M. MIHĂILESCU, M. PEREZ-LLANOS, J. D. ROSSI: Models for growth of heterogeneous sandpiles via Mosco convergence, *Asymptotic Analysis*, to appear.
- [3] M.M. BOUREANU, P. PUCCI, V. RĂDULESCU: Multiplicity of solutions for a class of anisotropic elliptic equations with variable exponent, *Complex Var. Elliptic Equ.*, **56** (2011), 755-767.
- [4] Y. CHEN, S. LEVINE, M. RAO: Variable Exponent, Linear Growth Functionals in Image Restoration, *SIAM J. Appl. Math.*, **66** (4) (2006), 1383-1406.
- [5] V. CHIADÒ-PIAT, A. COSCIA: Hölder continuity of minimizers of functionals with variable growth exponent, *Manuscripta Math.*, **93** (1997), 283-299.
- [6] F. COLASUONNO, M. SQUASSINA: Eigenvalues for double phase variational integrals, (2015), preprint.
- [7] A. COSCIA, G. MINGIONE: Hölder continuity of the gradient of  $p(x)$ -harmonic mappings, *C.R. Acad. Sci. Paris*, **328** (1999), 363-368.

- [8] G. CUPINI, M. GUIDORZI, E. MASCOLO: Regularity of minimizers of vectorial integrand with  $p - q$  growth, *Non Linear Analysis TMA* **4** (2003) 591-616.
- [9] A. DALL'AGLIO, E. MASCOLO, G. PAPI: Local boundedness for minima of functionals with nonstandard growth conditions, *Rend. Mat., Serie VII*, **18** (1998), 305-326.
- [10] E. DE GIORGI: Un esempio di estremali discontinue per un problema variazionale di tipo ellittico, *Boll. Un. Mat. Ital.*, (4) **1** (1968), 135-137.
- [11] L. DIENING, P. HÄRJULEHTO, P. HÄSTO, M. RŮŽIČKA: Lebesgue and Sobolev spaces with variable exponents, *Lecture Notes in Mathematics*, vol. 2017, 2011.
- [12] M. ELEUTERI: Hölder continuity results for a class of functionals with non standard growth, *Boll. Un. Mat. Ital.*, (8) **7 - B** (2004), 129-157.
- [13] M. ELEUTERI, P. MARCELLINI, E. MASCOLO: Lipschitz estimates for systems with ellipticity conditions at infinity, *Ann. Mat. Pura Appl.*, (2015), to appear.
- [14] L. ESPOSITO, F. LEONETTI, G. MINGIONE: Regularity results for minimizers of irregular integrals with  $(p, q)$  growth, *Forum Mathematicum* **14** (2002) 245-272.
- [15] L. ESPOSITO, F. LEONETTI, G. MINGIONE: Sharp regularity for functionals with  $(p, q)$  growth, *J. Differential Equations*, **204** (2004), 5-55.
- [16] N. FUSCO, C. SBORDONE: Higher integrability of the gradient of minimizers of functionals with non-standard growth conditions, *Comm. Pure Appl. Math.*, **43** (1990), no. 5, 673-683.
- [17] M. GIAQUINTA: *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, Annals of Mathematics Studies, 105. Princeton University Press, (1983).
- [18] E. GIUSTI, M. MIRANDA: Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni, *Boll. Un. Mat. Ital.* (4) **1** (1968), 219-226.
- [19] E. GIUSTI: *Direct methods in the calculus of variations*, World Scientific Publishing Co., Inc., River Edge, NJ (2003).
- [20] P. HÄRJULEHTO, P. HÄSTO, U. V. LE, M. NUORTIO: Overview of differential equations with non-standard growth, *Nonlinear Anal.*, **72** (2010), 4551-4574.
- [21] P. MARCELLINI: Regularity of minimizers of integrals in the calculus of variations with non standard growth conditions, *Arch. Rational Mech. Anal.*, **105** (1989), 267-284.
- [22] P. MARCELLINI: Regularity and existence of solutions of elliptic equations with  $p - q$ -growth conditions, *J. Differential Equations*, **90** (1991), 1-30.
- [23] P. MARCELLINI: Regularity for elliptic equations with general growth conditions, *J. Differential Equations*, **105** (1993), 296-333.
- [24] P. MARCELLINI: Regularity for some scalar variational problems under general growth conditions, *J. Optim. Theory Appl.*, **90** no. 1 (1996), 161-181.
- [25] P. MARCELLINI, G. PAPI: Nonlinear elliptic systems with general growth, *J. Differential Equations*, **221** no. 1 (2006), 412-443.
- [26] E. MASCOLO, A.P. MIGLIORINI: Everywhere regularity for vectorial functionals under general growth, *ESAIM: Control Optim. Calc. Var.* **9** (2003), 399-418.
- [27] G. MINGIONE: Regularity of minima: an invitation to the dark side of the calculus of variations, *Appl. Math.*, **51** (2006), 355-426.
- [28] M. MIHĂILESCU, P. PUCCI, V. RADULESCU: Nonhomogeneous boundary value problems in anisotropic Sobolev spaces, *C. R. Math. Acad. Sci. Paris* **345** (2007), 561-566.
- [29] M. MIHĂILESCU, P. PUCCI, V. RADULESCU: Eigenvalue problems for anisotropic quasilinear elliptic equations with variable exponent, *J. Math. Anal. Appl.*, **340** (2008), 687-698 .
- [30] C. MOONEY, O. SAVIN: Some singular minimizers in low dimensions in the Calculus of Variations, preprint arXiv:1503.00671 (2015), submitted.
- [31] K.R. RAJAGOPAL, M. RŮŽIČKA: On the modeling of electrorheological materials, *Mech. Res. Commun.*, **23** (1996), 401-407.
- [32] M. RŮŽIČKA: Flow of shear dependent electrorheological fluids, *C. R. Acad. Sci. Paris*, **329** (1999), 393-398.
- [33] V. SVERÁK, X. YAN: A singular minimizer of a smooth strongly convex functional in three dimensions, *Calc. Var. Partial Differential Equations*, **10** (2000), 213-221.

- [34] G. TALENTI: Boundedness of minimizers, *Hokkaido Math. J.*, **19** (1990), no. 2, 259-279.
- [35] V.V. ZHIKOV: Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.*, **50** (1986), 675-710.
- [36] V.V. ZHIKOV, S.M. KOZLOV, O.A. OLEINIK: Homogenization of differential operators and integral functionals, *Springer-Verlag, Berlin* (1994).

*E-mail address:* `eleuteri@math.unifi.it`

*E-mail address:* `marcellini@math.unifi.it`

*E-mail address:* `mascolo@math.unifi.it`

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, 50134 - FIRENZE, ITALY