



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Degree of non-Kählerianity for 6-dimensional nilmanifolds

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Degree of non-Kählerianity for 6-dimensional nilmanifolds / Angella, Daniele; Franzini, Maria Giovanna; Rossi, Federico Alberto. - In: MANUSCRIPTA MATHEMATICA. - ISSN 0025-2611. - STAMPA. - 148:(2015), pp. 177-211. [10.1007/s00229-015-0734-x]

Availability:

This version is available at: 2158/1009017 since: 2021-03-09T09:25:55Z

Published version:

DOI: 10.1007/s00229-015-0734-x

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

DEGREE OF NON-KÄHLERIANITY FOR 6-DIMENSIONAL NILMANIFOLDS

DANIELE ANGELLA, MARIA GIOVANNA FRANZINI, AND FEDERICO ALBERTO ROSSI

ABSTRACT. We use Bott-Chern cohomology to measure the non-Kählerianity of 6-dimensional nilmanifolds endowed with the invariant complex structures in M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa’s classification, [14]. We investigate the existence of pluriclosed metric in connection with such a classification.

INTRODUCTION

The *Bott-Chern cohomology* and the *Aeppli cohomology* provide further tools to investigate the complex geometry of non-Kähler manifolds: they are, in a sense, a bridge between the Dolbeault cohomology and the de Rham cohomology of a complex manifold X . Introduced by A. Aeppli in [1], the cohomology groups

$$H_{BC}^{\bullet,\bullet}(X) := \frac{\ker \partial \cap \ker \bar{\partial}}{\text{im } \partial \bar{\partial}} \quad \text{and} \quad H_A^{\bullet,\bullet}(X) := \frac{\ker \partial \bar{\partial}}{\text{im } \partial + \text{im } \bar{\partial}}$$

have raised much interest in several areas of Mathematics and Physics: see, e.g., B. Bigolin [8, 9], J. Varouchas [41], L. Alessandrini and G. Bassanelli [2], and, more recently, M. Schweitzer [37] and R. Kooistra [26] (in the context of cohomology theories), J.-M. Bismut [11, 12] (in the context of Chern characters), and L.-S. Tseng and S.-T. Yau [38] (in the framework of generalized geometry and type II string theory).

The Bott-Chern cohomology turns out to be particularly interesting in the non-Kähler case. In fact, for compact Kähler manifolds, or, more in general, for compact complex manifolds satisfying the $\partial\bar{\partial}$ -*Lemma* (that is, the very special cohomological property that every ∂ -closed $\bar{\partial}$ -closed d-exact form is $\partial\bar{\partial}$ -exact), the Bott-Chern cohomology coincides with the Dolbeault cohomology.

In [6], the first author and A. Tomassini proved that, on a compact complex manifold X , a Frölicher-type inequality relates the dimension of $H_{BC}^{\bullet,\bullet}(X)$ and the

2010 *Mathematics Subject Classification.* 57T15; 32Q57; 53C558.

Key words and phrases. nilmanifold; cohomology; Bott-Chern; non-Kähler; pluriclosed; SKT.

This work was supported by GNSAGA of INdAM. During the preparation of this work, the first author has been granted with a research fellowship by Istituto Nazionale di Alta Matematica INdAM, and has been supported by the Project PRIN “Varietà reali e complesse: geometria, topologia e analisi armonica”, by the Project FIRB “Geometria Differenziale e Teoria Geometrica delle Funzioni”, and by GNSAGA of INdAM.

To appear in *manuscripta mathematica*. The final publication is available at Springer via <http://dx.doi.org/10.1007/s00229-015-0734-x>.

dimension of $H_{dR}^\bullet(X; \mathbb{C})$, namely, [6, Theorem A],

$$(1) \quad \text{for every } k \in \mathbb{N}, \quad \sum_{p+q=k} (\dim_{\mathbb{C}} H_{BC}^{p,q}(X) + \dim_{\mathbb{C}} H_A^{p,q}(X)) \geq 2 \dim_{\mathbb{C}} H_{dR}^k(X; \mathbb{C}),$$

and that the equality in (1) holds if and only if X satisfies the $\partial\bar{\partial}$ -Lemma, [6, Theorem B].

A special class of manifolds, which turn out to provide useful examples in studying geometric and cohomological properties of (almost-)complex manifolds, is given by the class of *nilmanifolds*, namely, compact quotients $X = \Gamma \backslash G$ of a connected simply-connected nilpotent Lie group G by a discrete co-compact subgroup Γ .

By K. Nomizu's theorem [31, Theorem 1], the finite-dimensional vector space $\wedge^\bullet \mathfrak{g}^*$ of linear forms on the dual of the Lie algebra \mathfrak{g} naturally associated to G yields a minimal model for the de Rham complex $(\wedge^\bullet X, d)$ of X ; this means, in particular, that the de Rham cohomology of X can be computed as the cohomology of a finite-dimensional subcomplex of $(\wedge^\bullet X, d)$.

On the other hand, nilmanifolds cannot be formal, [23, Theorem 1], and hence, in particular, neither they satisfy the $\partial\bar{\partial}$ -Lemma nor they admit Kähler metrics.

The 6-dimensional nilmanifolds can be classified in terms of their Lie algebra, up to isomorphisms, in 34 classes, according to V. V. Morozov's classification, [30], see also [28]. As regards their complex geometry, the works by S. M. Salamon [36], L. Ugarte and R. Villacampa [40], A. Andrada, M. L. Barberis, and I. Dotti [4], and M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa [14] give a complete classification, up to equivalence, of the invariant complex structures on 6-dimensional nilmanifolds.

Even if they admit no Kähler metric, nilmanifolds can be endowed with other special Hermitian metrics, whose associated $(1, 1)$ -form satisfies conditions weaker than the Kähler condition, see, e.g., [29, 10, 39, 21, 27]. A special class of Hermitian metrics that recently have arisen interest in several areas of Mathematics and Physics is the class of *pluriclosed* (also called *strong-Kähler with torsion*, shortly, SKT) metrics, namely, the Hermitian metrics whose associated $(1, 1)$ -form is $\partial\bar{\partial}$ -closed, see, e.g., [10, 21, 34, 20, 13].

In this note, we are concerned especially in studying cohomological properties of the 6-dimensional nilmanifolds endowed with invariant complex structures; furthermore, we investigate cohomological properties of special classes of nilmanifolds admitting pluriclosed metrics. More precisely, we compute explicitly the dimensions of the Bott-Chern cohomology groups for each of the complex structures in M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa's classification, [14]: in view of the Frölicher-type inequality (1) and of K. Hasegawa's theorem [23, Theorem 1], such dimensions, together with the Betti numbers, measure, in some sense, the non-Kählerianity of nilmanifolds. We provide some sufficient conditions assuring the validity of the $\partial\bar{\partial}$ -Lemma on a compact complex manifold. Then, we study invariant pluriclosed structures on 6-dimensional nilmanifolds, proving that only seven of the classes in the cohomological classification by means of the Bott-Chern cohomology contain invariant complex structures admitting invariant pluriclosed metrics, see Theorem 4.1. On a 6-dimensional nilmanifold endowed with an invariant complex structure, the existence of pluriclosed metrics and the existence of

balanced metrics are complementary properties: indeed, by [21, Proposition 1.4], or also [3, Remark 1], any Hermitian metric being both pluriclosed and balanced is in fact Kähler, and by [21, Theorem 1.2] the pluriclosed property is satisfied by either all invariant Hermitian metrics or by none; we refer to [27] for the complementary study of balanced structures on 6-dimensional nilmanifolds: there, A. Latorre, L. Ugarte, and R. Villacampa study the behaviour of the cohomology in relation to the existence of balanced Hermitian metrics on 6-dimensional nilmanifolds endowed with invariant complex structures. In [34, Theorem 14], the third author and A. Tomassini provided a classification of 8-dimensional nilmanifolds endowed with a invariant complex structure such that every invariant Hermitian metric is pluriclosed; such classification consists of two classes, the first of which contains manifolds of the type $M^6 \times \mathbb{T}^2$, where M^6 is a 6-dimensional nilmanifold admitting pluriclosed structures and \mathbb{T}^2 is the standard 2-dimensional torus endowed with the standard complex structure. Hence, we investigate cohomological properties of manifolds of such type $M^6 \times \mathbb{T}^2$.

During the preparation of the final draft, we have been informed by L. Ugarte that A. Latorre, L. Ugarte, and R. Villacampa performed similar computations in [27], with the aim to further investigate the behaviour of the Bott-Chern cohomology under deformations of the complex structure, and the possible relation between cohomological properties and the existence of balanced or strongly-Gauduchon metrics; we refer to [27] for further details.

Acknowledgments. The authors would like to warmly thank Adriano Tomassini for his constant support and encouragement and for many interesting conversations and useful suggestions. We wish to warmly thank Luis Ugarte, Adela Latorre and Raquel Villacampa for pointing out the preprint [27] and for their several suggestions and remarks, which highly improved this paper. Thanks are also due to Nicola Enrietti for helpful suggestions concerning the use of Maple for Complex Geometry. Many thanks to the anonymous Referee, whose suggestions improved the presentation of the paper.

1. PRELIMINARIES AND NOTATION

1.1. Bott-Chern cohomology of non-Kähler manifolds. Let X be a compact complex manifold. Other than Dolbeault cohomology, another important tool to study the complex geometry of X is provided by the *Bott-Chern* and *Aeppli cohomologies*:

$$H_{BC}^{\bullet,\bullet}(X) := \frac{\ker \partial \cap \ker \bar{\partial}}{\text{im } \partial \bar{\partial}}, \quad H_A^{\bullet,\bullet}(X) := \frac{\ker \partial \bar{\partial}}{\text{im } \partial + \text{im } \bar{\partial}}.$$

Note that, while the Dolbeault cohomology groups are not symmetric, one has that, for every $p, q \in \mathbb{N}$, the complex conjugation induces the isomorphisms $H_{BC}^{p,q}(X) \simeq H_{BC}^{q,p}(X)$ and $H_A^{p,q}(X) \simeq H_A^{q,p}(X)$.

One has the natural maps of (bi-)graded \mathbb{C} -vector spaces

$$\begin{array}{ccccc}
 & & H_{BC}^{\bullet,\bullet}(X) & & \\
 & \swarrow & \downarrow & \searrow & \\
 H_{\bar{\partial}}^{\bullet,\bullet}(X) & & H_{dR}^{\bullet,\bullet}(X; \mathbb{C}) & & H_{\partial}^{\bullet,\bullet}(X) \\
 & \swarrow & \downarrow & \searrow & \\
 & & H_A^{\bullet,\bullet}(X) & &
 \end{array}$$

which are, in general, neither injective nor surjective. A compact complex manifold is said to satisfy the $\partial\bar{\partial}$ -Lemma if every ∂ -closed $\bar{\partial}$ -closed d-exact form is $\partial\bar{\partial}$ -exact, namely, if the map $H_{BC}^{\bullet,\bullet}(X) \rightarrow H_{dR}^{\bullet,\bullet}(X; \mathbb{C})$ is injective: this turns out to be equivalent to all the above maps being isomorphisms, [18, Lemma 5.15, Remark 5.16, 5.21]. The compact complex manifolds admitting a Kähler metric, or, more in general, belonging to *class C of Fujiki*, [22], satisfy the $\partial\bar{\partial}$ -Lemma, [18, Lemma 5.11, Corollary 5.23].

Fixed a J -Hermitian metric g on X , consider the 4th order self-adjoint elliptic differential operators

$$\tilde{\Delta}_{BC} := (\partial\bar{\partial})(\partial\bar{\partial})^* + (\partial\bar{\partial})^*(\partial\bar{\partial}) + (\bar{\partial}^*\partial)(\bar{\partial}^*\partial)^* + (\bar{\partial}^*\partial)^*(\bar{\partial}^*\partial) + \bar{\partial}^*\bar{\partial} + \partial^*\partial$$

and

$$\tilde{\Delta}_A := \partial\partial^* + \bar{\partial}\bar{\partial}^* + (\partial\bar{\partial})^*(\partial\bar{\partial}) + (\partial\bar{\partial})(\partial\bar{\partial})^* + (\bar{\partial}\partial^*)^*(\bar{\partial}\partial^*) + (\bar{\partial}\partial^*)(\bar{\partial}\partial^*)^* ,$$

see [25, Proposition 5], see also [37, §2.b, §2.c], [8, §5.1]; by means of them, one gets a Hodge theory for the Bott-Chern, respectively the Aeppli, cohomology: indeed, one has the following isomorphisms:

$$H_{BC}^{\bullet,\bullet}(X) \simeq \ker \tilde{\Delta}_{BC} = \ker \partial \cap \ker \bar{\partial} \cap \ker (\partial\bar{\partial})^* ,$$

[37, Théorème 2.2], and

$$H_A^{\bullet,\bullet}(X) \simeq \ker \tilde{\Delta}_A = \ker (\partial\bar{\partial}) \cap \ker \partial^* \cap \ker \bar{\partial}^* ,$$

[37, §2.c]. In particular, one gets, [37, Corollaire 2.3, §2.c], that

$$\dim_{\mathbb{C}} H_{BC}^{\bullet,\bullet}(X) < +\infty \quad \text{and} \quad \dim_{\mathbb{C}} H_A^{\bullet,\bullet}(X) < +\infty ,$$

and that, for every $p, q \in \mathbb{N}$,

$$*: H_{BC}^{p,q}(X) \xrightarrow{\simeq} H_A^{\dim_{\mathbb{C}} X - q, \dim_{\mathbb{C}} X - p}(X) .$$

For any $k \in \mathbb{N}$, set

$$\Delta^k(X) := \sum_{p+q=k} (\dim_{\mathbb{C}} H_{BC}^{p,q}(X) + \dim_{\mathbb{C}} H_A^{p,q}(X)) - 2b_k ,$$

where $b_k := \dim_{\mathbb{R}} H_{dR}^k(X; \mathbb{R})$ is the k^{th} Betti number of X . Note that, if X satisfies the $\partial\bar{\partial}$ -Lemma, then $\Delta^k = 0$ for every $k \in \mathbb{N}$.

The first author and A. Tomassini proved that, [6, Theorem A],

$$\text{for every } k \in \mathbb{N}, \quad \Delta^k(X) \geq 0 ,$$

and that, [6, Theorem B], if $\Delta^k(X) = 0$ for every $k \in \mathbb{N}$, then X satisfies the $\partial\bar{\partial}$ -Lemma.

Note that $\Delta^k(X) = \Delta^{\dim_{\mathbb{C}} X - k}(X)$ for every $k \in \mathbb{N}$; hence, on a 6-dimensional manifold endowed with a complex structure, just $\Delta^1(X)$, $\Delta^2(X)$, and $\Delta^3(X)$ have to be computed.

1.2. Invariant complex structures on 6-dimensional nilmanifolds. Let $X = \Gamma \backslash G$ be a nilmanifold, that is, a compact quotient of a connected simply-connected nilpotent Lie group G by a discrete co-compact subgroup Γ of G ; denote the Lie algebra associated to G by \mathfrak{g} , and let $\mathfrak{g}_{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$.

Dealing with (*G-left-invariant*) objects on X , we mean objects on X induced by objects on G which are invariant under the action of G on itself given by left-translations; equivalently, through left-translations, any invariant object on X is uniquely determined by an object on spaces constructed starting with the Lie algebra \mathfrak{g} .

V. V. Morozov classified in [30] the 6-dimensional nilpotent Lie algebras, up to isomorphism, in 34 different classes, see also [28]. In [36], S. M. Salamon identified the 18 classes of 6-dimensional nilpotent Lie algebra admitting a linear integrable complex structure: up to equivalence¹, the linear integrable non-nilpotent complex structures² have been classified by L. Ugarte and R. Villacampa in [40], the linear integrable Abelian complex structures³ have been classified by A. Andrada, M. L. Barberis, and I. Dotti in [4], and lastly the linear integrable nilpotent non-Abelian complex structures have been classified by M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa in [14].

We recall in Table 1 the classification in [14]. As a matter of notation, we identify a Lie algebra \mathfrak{g} (or the associated connected simply-connected Lie group G , or the associated nilmanifold $X = \Gamma \backslash G$, where Γ is a discrete co-compact subgroup of G) by its structure equations, namely, writing

$$\mathfrak{h}_5 := (0^4, 13 + 42, 14 + 23) := (0, 0, 0, 0, 13 + 42, 14 + 23) ,$$

we mean that there exists a basis $\{e_j\}_{j \in \{1, \dots, 6\}}$ of \mathfrak{g} such that, with respect to the dual basis $\{e^j\}_{j \in \{1, \dots, 6\}}$ of \mathfrak{g}^* , the structure equations are

$$\begin{cases} de^1 = de^2 = de^3 = de^4 = 0 \\ de^5 = e^1 \wedge e^3 + e^4 \wedge e^2 \\ de^6 = e^1 \wedge e^4 + e^2 \wedge e^3 \end{cases} ,$$

where usually we also shorten, e.g., $e^{AB} := e^A \wedge e^B$. Similarly, we identify a complex structure on a nilmanifold by its structure equations in terms of a coframe of the holomorphic cotangent bundle, namely, writing

$$J_2 := (0, 0, \omega^{12}) \quad \text{on } \mathfrak{h}_5 ,$$

¹We recall that two linear integrable complex structures J and J' on a Lie algebra \mathfrak{g} are said to be *equivalent* if there exists an automorphism $F: \mathfrak{g} \rightarrow \mathfrak{g}$ of the Lie algebra such that $J = F^{-1} \circ J' \circ F$.

²We recall that a linear integrable complex structure J on a $2n$ -dimensional Lie algebra \mathfrak{g} is called *nilpotent* if there is a basis $\{\omega^1, \dots, \omega^n\}$ for $(\mathfrak{g}^{1,0})^*$ with respect to which the structure equations are of the form $d\omega^j = \sum_{h < k < j} A_{hk}^j \omega^h \wedge \omega^k + \sum_{h, k < j} B_{hk}^j \omega^h \wedge \bar{\omega}^k$ with $A_{hk}^j, B_{hk}^j \in \mathbb{C}$.

³We recall that a linear integrable complex structure J on a Lie algebra \mathfrak{g} is called *Abelian* if $[Jx, Jy] = [x, y]$ for any $x, y \in \mathfrak{g}$.

we mean that there exists an invariant coframe $\{\omega^1, \omega^2, \omega^3\}$ of the $\mathcal{C}^\infty(X; \mathbb{C})$ -module $T^{1,0}X$, where $X = \Gamma \backslash G$ is a nilmanifold with Lie algebra \mathfrak{h}_5 , such that the structure equations read

$$\begin{cases} d\omega^1 = d\omega^2 = 0 \\ d\omega^3 = \omega^1 \wedge \omega^2 \end{cases},$$

where usually we shorten, e.g., $\omega^{A\bar{B}} := \omega^A \wedge \bar{\omega}^B$.

Other than the conditions listed in Table 1, as in [14] we assume that the parameters satisfy $\lambda \geq 0$, $c \geq 0$, $B \in \mathbb{C}$, and $D \in \mathbb{C}$.

1.3. Cohomologies of nilmanifolds. The space $\wedge^\bullet \mathfrak{g}^*$, equivalently, the space of invariant differential forms on X , yields a sub-complex $(\wedge^\bullet \mathfrak{g}^*, d)$ of the de Rham complex, $(\wedge^\bullet X, d)$, where $d: \wedge^\bullet \mathfrak{g}^* \rightarrow \wedge^{\bullet+1} \mathfrak{g}^*$ is induced by $d|_{\wedge^1 \mathfrak{g}^*}: \wedge^1 \mathfrak{g}^* \ni \alpha \mapsto d\alpha := -\alpha([\cdot, \cdot]) \in \wedge^2 \mathfrak{g}^*$. By K. Nomizu's theorem [31, Theorem 1], the map $(\wedge^\bullet \mathfrak{g}^*, d) \rightarrow (\wedge^\bullet X, d)$ is a quasi-isomorphism, that is, the de Rham cohomology of X can be computed using just invariant forms. (The same result holds true, more in general, for completely-solvable solvmanifolds, as proved by A. Hattori, [24, Corollary 4.2]; a counter-example for general solvmanifolds has been provided in [17, Corollary 4.2, Remark 4.3] by P. de Bartolomeis and A. Tomassini.)

Analogously, an invariant complex structure J on X induces a bi-graduation on the space of invariant forms, equivalently $\wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^*$, and a sub-double-complex $(\wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^*, \partial, \bar{\partial}) \hookrightarrow (\wedge^{\bullet, \bullet} X, \partial, \bar{\partial})$; this inclusion induces an injective morphism in cohomology, [15, Lemma 9], namely,

$$(2) \quad \frac{\ker(\bar{\partial}: \wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^* \rightarrow \wedge^{\bullet, \bullet+1} \mathfrak{g}_{\mathbb{C}}^*)}{\text{im}(\bar{\partial}: \wedge^{\bullet, \bullet-1} \mathfrak{g}_{\mathbb{C}}^* \rightarrow \wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^*)} \hookrightarrow H_{\bar{\partial}}^{\bullet, \bullet}(X).$$

Similarly, one has an injective morphism into Bott-Chern cohomology, [5, Lemma 3.6],

$$(3) \quad \frac{\ker(d: \wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^* \rightarrow \wedge^{\bullet+1, \bullet} \mathfrak{g}_{\mathbb{C}}^* \oplus \wedge^{\bullet, \bullet+1} \mathfrak{g}_{\mathbb{C}}^*)}{\text{im}(\partial\bar{\partial}: \wedge^{\bullet-1, \bullet-1} \mathfrak{g}_{\mathbb{C}}^* \rightarrow \wedge^{\bullet, \bullet} \mathfrak{g}_{\mathbb{C}}^*)} \hookrightarrow H_{BC}^{\bullet, \bullet}(X).$$

By saying that the Dolbeault, respectively the Bott-Chern, cohomology of X is (G -left-)invariant, we mean that the inclusion (2), respectively (3), is actually an isomorphism. The invariance of the Bott-Chern cohomology can be seen as a consequence of the invariance of the Dolbeault cohomology, as proved by the first author, [5, Theorem 3.7].

It is not known whether, for any invariant complex structure on a nilmanifold, the Dolbeault cohomology is invariant, [33, Conjecture 1], see also [16, page 5406], [15, page 112]. However, this turns out to be true for a large class of invariant complex structures: more precisely, the following result holds.

Theorem 1.1 ([35, Theorem 1], [16, Main Theorem], [15, Theorem 2, Remark 4], [32, Theorem 1.10], [5, Theorem 3.8]). *Let $X = \Gamma \backslash G$ be a nilmanifold endowed with an invariant complex structure J . The Dolbeault cohomology, and the Bott-Chern cohomology of X are invariant, provided one of the following conditions holds:*

- X is holomorphically parallelizable, [35, Theorem 1];
- J is an Abelian complex structure (i.e., $[Jx, Jy] = [x, y]$ for any $x, y \in \mathfrak{g}$), [15, Remark 4];

- J is a nilpotent complex structure (i.e., there is an invariant coframe $\{\omega^1, \dots, \omega^n\}$ for $(T^{1,0}X)^*$ with respect to which the structure equations of X are of the form

$$d\omega^j = \sum_{h < k < j} A_{hk}^j \omega^h \wedge \omega^k + \sum_{h, k < j} B_{hk}^j \omega^h \wedge \bar{\omega}^k$$

with $\{A_{hk}^j, B_{hk}^j\}_{j,h,k} \subset \mathbb{C}$, [16, Main Theorem];

- J is a rational complex structure (i.e., $J(\mathfrak{g}_{\mathbb{Q}}) \subseteq \mathfrak{g}_{\mathbb{Q}}$ where $\mathfrak{g}_{\mathbb{Q}}$ is the rational structure for \mathfrak{g} induced by Γ —where a rational structure for \mathfrak{g} is a \mathbb{Q} -vector space such that $\mathfrak{g} = \mathfrak{g}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R}$), [15, Theorem 2];
- \mathfrak{g} admits a torus-bundle series compatible with J and with the rational structure induced by Γ , [32, Theorem 1.10].

We recall that, since every 6-dimensional nilpotent Lie algebra, except \mathfrak{h}_7 , admits a stable torus-bundle series compatible with any linear complex structure and any rational structure, [32, Theorem B], one gets the following result.

Corollary 1.2 ([33, Corollary 3.10]). *For any 6-dimensional nilmanifold endowed with an invariant complex structure and with Lie algebra non-isomorphic to $\mathfrak{h}_7 = (0, 0, 0, 12, 13, 23)$, both the Dolbeault cohomology and the Bott-Chern cohomology are invariant.*

Lastly, we recall that, if X satisfies the $\partial\bar{\partial}$ -Lemma, then the differential graded algebra (\wedge^*X, d) is formal, [18, Main Theorem]. By K. Hasegawa's theorem [23, Theorem 1], no nilmanifold is formal except for the torus. In particular, the $\partial\bar{\partial}$ -Lemma does not hold for any complex structure on a non-torus nilmanifold.

2. BOTT-CHERN COHOMOLOGY OF 6-DIMENSIONAL NILMANIFOLDS

In this section, we provide the results of the computation of the Bott-Chern cohomology for each of the complex structure in M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa's classification, [14]. In view of [33, Corollary 3.10], and noting that the unique (up to equivalence) complex structure on the nilmanifold corresponding to \mathfrak{h}_7 is a rational complex structure, one is reduced to study invariant Δ_{BC} -harmonic forms. Note that the Dolbeault, respectively Bott-Chern, cohomology being invariant is not an up-to-equivalence property: that is, the Bott-Chern numbers $h_{BC}^{p,q} := \dim_{\mathbb{C}} H_{BC}^{p,q}(X)$ in Table ?? provide a complete picture *except for* nilmanifolds with Lie algebra isomorphic to \mathfrak{h}_7 . Such computations have been performed with the aid of a symbolic calculations software. In Figure 1, we provide a graphical visualization of the non-Kählerianity of such 6-dimensional nilmanifolds by means of the degrees Δ^1 , Δ^2 , and Δ^3 .

Theorem 2.1. *Consider a 6-dimensional nilpotent Lie algebra endowed with a linear integrable complex structure. Equivalently, consider a 6-dimensional nilmanifold with Lie algebra non-isomorphic to $\mathfrak{h}_7 = (0, 0, 0, 12, 13, 23)$ and endowed with any invariant complex structure, or a nilmanifold with Lie algebra isomorphic to \mathfrak{h}_7 and endowed with the invariant complex structure in M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa's classification, [14], see Table 1. Then the dimensions of the Bott-Chern cohomology groups are provided in Table ??.*

Remark 2.2. During the preparation of the final version of this note, L. Ugarte informed us on a similar paper in preparation by A. Latorre, L. Ugarte, and R. Villacampa, [27]: they computed the dimensions of the Bott-Chern cohomology groups for any complex structure on a 6-dimensional nilpotent Lie algebra using the classification of [14], with the aim to study variations of the cohomology under deformations of the complex structure, and the behaviour of the cohomology in relation to the existence of balanced Hermitian metrics or strongly-Gauduchon metrics. Since the main purpose (other than some results and, possibly, the techniques of computation) of our paper is partly different from [27], we believe that no crucial overlapping could distract the reader.

Remark 2.3. As a consequence of the computations, we notice that the degrees $\Delta^1 \in \{0, 2\}$ and $\Delta^3 \in \{0, 4, 8, 12\}$ do not depend on the chosen linear integrable complex structure on a fixed 6-dimensional nilpotent Lie algebra, while $\Delta^2 \in \{0, \dots, 9\}$ does depend on it; more precisely, the numbers $h_{BC}^{p,q}$ depend just on the Lie algebra whenever $p+q$ odd (compare also [5, Remark 5.2]). Furthermore, on a 6-dimensional Lie algebra endowed with a linear integrable complex structure, one has that the Abelianity of the Lie algebra is equivalent to the condition $\Delta^3 = 0$.

3. DETECTING THE VALIDITY OF THE $\partial\bar{\partial}$ -LEMMA

In this section, we prove some straightforward formulas concerning

$$\left\{ \Delta^k := \Delta^k(X) := \sum_{p+q=k} (\dim_{\mathbb{C}} H_{BC}^{p,q}(X) + \dim_{\mathbb{C}} H_A^{p,q}(X)) - 2 \dim_{\mathbb{C}} H_{dR}^k(X; \mathbb{C}) \right\}_{k \in \mathbb{N}},$$

in order to detect the validity of the $\partial\bar{\partial}$ -Lemma on a compact complex manifold X just in terms of few numbers.

Let X be a compact complex manifold of complex dimension n . As a matter of notation, denote, for $k \in \mathbb{N}$,

$$b_k := \dim_{\mathbb{C}} H_{dR}^k(X; \mathbb{C})$$

and, for $\sharp \in \{\partial, \bar{\partial}, BC, A\}$ and $(p, q) \in \mathbb{N}^2$ and $k \in \mathbb{N}$,

$$h_{\sharp}^{p,q} := \dim_{\mathbb{C}} H_{\sharp}^{p,q}(X) \quad \text{and} \quad h_{\sharp}^k := \sum_{p+q=k} \dim_{\mathbb{C}} H_{\sharp}^{p,q}(X);$$

we recall that the following equalities hold: for any $(p, q) \in \mathbb{N}^2$,

$$h_{BC}^{p,q} = h_{BC}^{q,p} = h_A^{n-p, n-q} = h_A^{n-q, n-p} \quad \text{and} \quad h_{\bar{\partial}}^{p,q} = h_{\partial}^{q,p} = h_{\bar{\partial}}^{n-p, n-q} = h_{\partial}^{n-q, n-p}$$

and, for any $k \in \mathbb{N}$,

$$b_k = b_{2n-k}, \quad h_{BC}^k = h_A^{2n-k}, \quad \text{and} \quad h_{\bar{\partial}}^k = h_{\partial}^k = h_{\bar{\partial}}^{2n-k} = h_{\partial}^{2n-k}.$$

Denote the topological Euler-Poincaré characteristic of X by

$$\chi_{\text{top}} := \sum_{k=0}^{2n} (-1)^k b_k = \sum_{k=0}^{2n} \sum_{p+q=k} (-1)^k h_{\bar{\partial}}^{p,q},$$

see, e.g., [19, Theorem VI.11.7].

One has the equalities

$$b_n = (-1)^n \chi_{\text{top}} (\# \mathcal{P} \sum_{k=0}^{n-1} (-1)^{n-k-1} b_k) \quad \text{and} \quad h_{\bar{\partial}}^n = (-1)^n \chi_{\text{top}} + 2 \sum_{k=0}^{n-1} (-1)^{n-k-1} h_{\bar{\partial}}^k.$$

Proposition 3.1. *Let X be a compact complex manifold. For any $k \in \mathbb{N}$, it holds that $\Delta^{2k+1} \equiv 0 \pmod{2}$.*

Proof. Since

$$\begin{aligned} \Delta^{2k+1} &= h_{BC}^{2k+1} + h_A^{2k+1} - 2b_{2k+1} \\ &= 2 \left(\sum_{\substack{p+q=2k+1 \\ p < q}} (h_{BC}^{p,q} + h_A^{p,q}) - b_{2k+1} \right) \equiv 0 \pmod{2}, \end{aligned}$$

the proposition follows. \square

Proposition 3.2. *Let X be a nilmanifold endowed with a complex structure, and suppose that the complex dimension n of X is odd. Then, $\Delta^n \equiv 0 \pmod{4}$.*

Proof. Since the Euler-Poincaré characteristic χ_{top} of a nilmanifold vanishes, one has

$$\begin{aligned} \Delta^n &= h_{BC}^n + h_A^n - 2b_n = 2h_{BC}^n - 2b_n \\ &= 4 \left(\sum_{\substack{p+q=n \\ p < q}} h_{BC}^{p,q} - \sum_{k=0}^{n-1} (-1)^{n-k-1} b_k \right) \equiv 0 \pmod{4}, \end{aligned}$$

from which the proposition follows. \square

Proposition 3.3. *Let X be a compact complex manifold of complex dimension n . If either*

- $\Delta^k = 0$ for every $k \equiv n \pmod{2}$, and
- for every $k \in \mathbb{N}$, it holds that $\sum_{p+q=2k+1} \dim_{\mathbb{C}} \frac{\text{im } \bar{\partial} \cap \text{im } \partial \cap \wedge^{p,q} X}{\text{im } \partial \bar{\partial}} = 0$,

or

- $\Delta^k = 0$ for every $k \equiv n-1 \pmod{2}$, and
- for every $k \in \mathbb{N}$, it holds that $\sum_{p+q=2k} \dim_{\mathbb{C}} \frac{\text{im } \bar{\partial} \cap \text{im } \partial \cap \wedge^{p,q} X}{\text{im } \partial \bar{\partial}} = 0$;

then X satisfies the $\partial \bar{\partial}$ -Lemma.

Proof. Consider the bi-graded finite-dimensional \mathbb{C} -vector spaces

$$A^{\bullet,\bullet}(X) := \frac{\text{im } \bar{\partial} \cap \text{im } \partial}{\text{im } \partial \bar{\partial}} \quad \text{and} \quad F^{\bullet,\bullet}(X) := \frac{\ker \partial \bar{\partial}}{\ker \bar{\partial} + \ker \partial},$$

and denote, for every $(p, q) \in \mathbb{N}^2$ and $k \in \mathbb{N}$,

$$a^{p,q} := \dim_{\mathbb{C}} A^{p,q}(X) \quad \text{and} \quad a^k := \sum_{p+q=k} \dim_{\mathbb{C}} A^{p,q}(X)$$

and

$$f^{p,q} := \dim_{\mathbb{C}} F^{p,q}(X) \quad \text{and} \quad f^k := \sum_{p+q=k} \dim_{\mathbb{C}} F^{p,q}(X).$$

Following the same argument used in [37] to prove the duality between Bott-Chern and Aeppli cohomology groups, we prove the duality between $A^{\bullet,\bullet}$ and $F^{\bullet,\bullet}$; it follows that, for any $(p, q) \in \mathbb{N}^2$ and $k \in \mathbb{N}$, the equalities

$$a^{p,q} = a^{q,p} = f^{n-p,n-q} = f^{n-q,n-p} \quad \text{and} \quad a^k = f^{2n-k}$$

hold.

Indeed, note that the pairing

$$A^{\bullet,\bullet} \times F^{\bullet,\bullet} \rightarrow \mathbb{C}, \quad ([\alpha], [\beta]) \mapsto \int_X \alpha \wedge \bar{\beta},$$

is non-degenerate: choose a Hermitian metric g on X ; if $[\alpha] \in A^{\bullet,\bullet} \subseteq H_{BC}^{\bullet,\bullet}(X)$, then there exists a $\tilde{\Delta}_{BC}$ -harmonic representative $\tilde{\alpha}$ in $[\alpha] \in A^{\bullet,\bullet}$, by [37, Corollaire 2.3], that is, $\partial\tilde{\alpha} = \bar{\partial}\tilde{\alpha} = \partial\bar{\partial}\tilde{\alpha} = 0$; hence, $[\tilde{\alpha}] \in F^{\bullet,\bullet}$, and $([\tilde{\alpha}], [\tilde{\alpha}]) = \int_X \tilde{\alpha} \wedge \bar{\tilde{\alpha}}$ is zero if and only if $\tilde{\alpha}$ is zero if and only if $[\alpha] \in A^{\bullet,\bullet}$ is zero.

We further note that, by J. Varouchas' exact sequences, [41, §3.1], one has the following equality, see [6, §3], for any $k \in \mathbb{N}$:

$$(5) \quad h_{BC}^k + h_A^k = 2h_{\bar{\partial}}^k + a^k + f^k.$$

By using (5) and (4), one gets

$$\begin{aligned} \Delta^n &= h_{BC}^n + h_A^n - 2b_n = 2h_{\bar{\partial}}^n + a^n + f^n - 2b_n \\ &= 2 \sum_{k=0}^{n-1} (-1)^{n-k-1} (2h_{\bar{\partial}}^k - 2b_k) + a^n + f^n \\ &= 2 \sum_{k=0}^{n-1} (-1)^{n-k-1} (h_{BC}^k + h_A^k - 2b_k) + 2 \sum_{k=0}^{n-1} (-1)^{n-k} (a^k + f^k) + (a^n + f^n) \\ &= 2 \sum_{k=0}^{n-1} (-1)^{n-k-1} \Delta^k + 2 \sum_{k=0}^{2n} (-1)^{n-k} a^k; \end{aligned}$$

in particular, it follows that

$$\sum_{k \equiv n \pmod{2}} \Delta^k + 2 \sum_{k=1}^n a^{2k-1} = \sum_{k \equiv n-1 \pmod{2}} \Delta^k + 2 \sum_{k=0}^n a^{2k}.$$

Hence, the proposition follows by using [6, Theorem B]. \square

In particular, by using the K. Hasegawa theorem [23, Theorem 1, Corollary], one gets straightforwardly the following corollary.

Corollary 3.4. *Let $X = \Gamma \backslash G$ be a 6-dimensional nilmanifold endowed with a complex structure. If either*

$$\Delta^1 = \Delta^3 = 0 \quad \text{and} \quad \dim_{\mathbb{C}} \frac{\text{im } \bar{\partial} \cap \text{im } \partial \cap \wedge^{2,1} X}{\text{im } \partial \bar{\partial}} = 0,$$

or

$$\Delta^2 = 0 \quad \text{and} \quad \dim_{\mathbb{C}} \frac{\text{im } \bar{\partial} \cap \text{im } \partial \cap \wedge^{1,1} X}{\text{im } \partial \bar{\partial}} = 0,$$

then X is diffeomorphic to a torus.

4. PLURICLOSED METRICS ON 6-DIMENSIONAL NILMANIFOLDS

A very special class of metrics in Complex Geometry is provided by Kähler metrics, namely, by J -Hermitian metrics g on a complex manifold (X, J) such that the associated $(1, 1)$ -form $\omega := g(J \cdot, \cdot)$ is d-closed. By Ch. Benson and C. S. Gordon's theorem [7, Theorem A], or by K. Hasegawa's theorem [23, Theorem 1, Corollary], if a nilmanifold admits a Kähler structure, then it is diffeomorphic to a torus. Hence, a natural question is to investigate the problem of the existence of special Hermitian metrics, other than the Kähler metrics, on nilmanifolds endowed with complex structures, and the possible connection with cohomological properties; for instance, given a complex manifold (X, J) of complex dimension n , a Hermitian metric g on X , with associated $(1, 1)$ -form ω , is called *balanced* if $d\omega^{n-1} = 0$, [29], and it is called *pluriclosed* (or *strong Kähler with torsion*, shortly SKT) if $\partial\bar{\partial}\omega = 0$, [10]. In particular, we study here pluriclosed metrics.

The following result points out which of the complex manifolds in Table ?? admit pluriclosed metrics.

Theorem 4.1. *Let X be a 6-dimensional nilmanifold and J be an invariant complex structure on X . Then (X, J) admits an invariant pluriclosed metric if and only if*

$$[\mathbf{00}] (X, J) = (\mathfrak{h}_1, J);$$

$$[\mathbf{01b}] (X, J) = (\mathfrak{h}_2, J_1^D) \text{ with } D \in \{i, -i\};$$

$$[\mathbf{02b}] (X, J) = (\mathfrak{h}_2, J_2^D) \text{ with } D \in \mathbb{C} \text{ such that } \operatorname{Re}D = 1 \text{ and } \operatorname{Im}D > 0;$$

$$[\mathbf{06b}] (X, J) = (\mathfrak{h}_4, J_2^D) \text{ with } D = 1;$$

$$[\mathbf{09b}^*] (X, J) = (\mathfrak{h}_5, J_3^{(\lambda, D)}) \text{ with } \lambda = 0 \text{ and } D = \frac{1}{2} + iy \text{ such that } 0 < 4y^2 < 3;$$

$$[\mathbf{09c}] (X, J) = (\mathfrak{h}_5, J_3^{(\lambda, D)}) \text{ with } \lambda = 0 \text{ and } D = \frac{1}{2};$$

$$[\mathbf{12}] (X, J) = (\mathfrak{h}_8, J),$$

where we use the notation in §1.2 and §2, and in [14].

Proof. It is well known that, if a 6-dimensional nilmanifold endowed with an invariant complex structure admits an invariant pluriclosed metric, then its Lie algebra is isomorphic to $\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_4, \mathfrak{h}_5$, or \mathfrak{h}_8 , see [21, Theorem 3.2], or [39, Theorem 3.3]. Moreover, one knows that, on a 6-dimensional nilmanifold endowed with an invariant complex structure, the pluriclosed condition is satisfied by either all the invariant Hermitian metrics or by none, [21, Theorem 1.2]. Hence we consider only the complex structures on $\mathfrak{h}_1, \mathfrak{h}_2, \mathfrak{h}_4, \mathfrak{h}_5, \mathfrak{h}_8$ and test the pluriclosed condition on the standard invariant Hermitian metric g with fundamental form $\sum_{j=1}^3 \omega^j \wedge \bar{\omega}^j$, where $\{\omega^j\}_{j \in \{1,2,3\}}$ is a given invariant coframe for $T^{1,0}X$, according to the notation in [14]. Since the first case, \mathfrak{h}_1 , is the Kählerian one, we focus our attention on non-toric nilmanifolds, and, after a simple computation, we get the results summarized in Table 3. Imposing the pluriclosed condition on the standard metric and matching with the conditions in Table ??, we get the proposition. \square

Remark 4.2. As a consequence of the computations summarized in Table ?? and of Theorem 4.1, we note that, restricting to the class of 6-dimensional nilmanifolds endowed with invariant pluriclosed structures, the number $h_{BC}^{p,q}$ depends just on the Lie algebra whenever $(p, q) \notin \{(3, 1), (2, 2), (1, 3)\}$, see also Remark 2.3.

With the aim of studying cohomological properties in relation to the existence of special metrics, we provide the following example, which shows a curve of complex structures admitting pluriclosed metrics and with jumping Bott-Chern cohomology numbers.

Example 4.3 (*A curve of complex structures on a 6-dimensional nilmanifold such that each complex structure admits a pluriclosed metric and the dimensions of the Bott-Chern cohomology groups are not constant*). Let X be a nilmanifold with associated Lie algebra

$$\mathfrak{h}_2 := (0^4, 12, 34) .$$

Consider the family $\{J_t\}_{t \in \mathbb{R}}$ of invariant complex structures on X such that an invariant coframe for the $\mathcal{C}^\infty(X)$ -module of the $(1,0)$ -forms on (X, J_t) is given by $\{\psi_t^1, \psi_t^2, \psi_t^3\}$ satisfying

$$\begin{cases} d\psi_t^1 = 0 \\ d\psi_t^2 = 0 \\ d\psi_t^3 = t\psi_t^1 \wedge \psi_t^2 + \psi_t^1 \wedge \bar{\psi}_t^1 + t\psi_t^1 \wedge \bar{\psi}_t^2 + (t^2 + i)\psi_t^2 \wedge \bar{\psi}_t^2 \end{cases} .$$

Note that

- for $t = 0$, the invariant complex structure J_t is equivalent to the invariant Abelian complex structure J_1^D with $D = i$ (case **01b** in Table ??);
- for $t \neq 0$, the invariant complex structure J_t is equivalent to the invariant non-Abelian complex structure J_2^D with $D = 1 + \frac{1}{t^2}i$ (case **02b** in Table ??),

where we use the notation in §1.2 and §2, and [14].

In particular, by Theorem 4.1, (X, J_t) admits an invariant pluriclosed metric for every $t \in \mathbb{R}$. Notwithstanding, note that

$$\dim_{\mathbb{C}} H_{BC}^{3,1}(X, J_0) = \dim_{\mathbb{C}} H_{BC}^{1,3}(X, J_0) = 3 ,$$

and

$$\dim_{\mathbb{C}} H_{BC}^{3,1}(X, J_t) = \dim_{\mathbb{C}} H_{BC}^{1,3}(X, J_t) = 2 \quad \text{for } t \neq 0 ,$$

while, for $(p, q) \notin \{(3, 1), (1, 3)\}$ one has $\dim_{\mathbb{C}} H_{BC}^{p,q}(X, J_0) = \dim_{\mathbb{C}} H_{BC}^{p,q}(X, J_t)$ for every $t \in \mathbb{R}$.

Remark 4.4. An analogous example is provided by a nilmanifold X with associated Lie algebra \mathfrak{h}_5 and endowed with the family $\left\{ J_t := J_3^{(0, \frac{1}{2} + i t)} \right\}_{t \geq 0, 4t^2 < 3}$ of invariant complex structures on X such that, with respect to an invariant coframe $\{\psi_t^1, \psi_t^2, \psi_t^3\}$ of $T^{1,0}X$, one has

$$\begin{cases} d\psi_t^1 = 0 \\ d\psi_t^2 = 0 \\ d\psi_t^3 = \psi_t^1 \wedge \psi_t^2 + \psi_t^1 \wedge \bar{\psi}_t^1 + \left(\frac{1}{2} + i t\right) \psi_t^2 \wedge \bar{\psi}_t^2, \end{cases}$$

(case **09** in Table 1); in fact, there exists a pluriclosed metric for every t such that $t \geq 0$ and $4t^2 < 3$, and we have the jumping of

$$\dim_{\mathbb{C}} H_{BC}^{2,2}(X, J_0) = 8 \neq 7 = \dim_{\mathbb{C}} H_{BC}^{2,2}(X, J_t) \quad \text{for } 0 < t < \frac{\sqrt{3}}{2}$$

(cases **09c** and **09b'** in Table ??).

Remark 4.5. Note that cases **09b'** and **09b''** provide examples of linear integrable non-equivalent complex structures on \mathfrak{h}_5 having the same dimensions of the Bott-Chern cohomology groups but admitting, respectively not admitting pluriclosed metrics.

For the sake of completeness, we provide the following example, compare also [14, Theorem 5.9]: it shows a curve of complex structures on a 6-dimensional nilmanifold admitting both pluriclosed and balanced metrics.

Example 4.6 (*A curve of compact complex manifolds endowed with special metrics*). Let X be a nilmanifold with associated Lie algebra

$$\mathfrak{h}_4 := (0^4, 12, 14 + 23) ,$$

and consider on it the family $\{J_2^D\}_{D \in \mathbb{R} \setminus \{0\}}$ of invariant non-Abelian complex structures whose $(1, 0)$ -forms are generated by $\{\omega_D^1, \omega_D^2, \omega_D^3\}$ as a $C^\infty(X; \mathbb{C})$ -module and such that

$$\begin{cases} d\omega_D^1 & := 0 \\ d\omega_D^2 & := 0 \\ d\omega_D^3 & := \omega_D^{12} + \omega_D^{1\bar{1}} + \omega_D^{1\bar{2}} + D\omega_D^{2\bar{2}} \end{cases} .$$

By [21, Theorem 3.2] and [39, Theorem 26], the Lie algebra \mathfrak{h}_4 admits both pluriclosed and balanced invariant Hermitian structures.

Let

$$\Omega_D^{r,s,t,u,v,z} := i \left(r^2 \omega_D^{1\bar{1}} + s^2 \omega_D^{2\bar{2}} + t^2 \omega_D^{3\bar{3}} \right) + u \omega_D^{1\bar{2}} - \bar{u} \omega_D^{2\bar{1}} + v \omega_D^{2\bar{3}} - \bar{v} \omega_D^{3\bar{2}} + z \omega_D^{1\bar{3}} - \bar{z} \omega_D^{3\bar{1}}$$

the generic J_2^D -Hermitian structure, where

$$r, s, t \in \mathbb{R} \setminus \{0\} \quad \text{and } u, v, z \in \mathbb{C}$$

satisfy

$$\begin{cases} r^2 s^2 > |u|^2 \\ s^2 t^2 > |v|^2 \\ r^2 t^2 > |z|^2 \\ r^2 s^2 t^2 + 2 \operatorname{Re}(i \bar{u} \bar{v} z) > t^2 |u|^2 + r^2 |v|^2 + s^2 |z|^2 \end{cases} ,$$

see, e.g., [40, pages 3-4].

By computing

$$\partial\bar{\partial}\Omega_D^{r,s,t,u,v,z} = i t^2 (1 - D) \omega_D^{12\bar{1}\bar{2}} ,$$

we have that $\Omega_D^{r,s,t,u,v,z}$ is pluriclosed if and only if $D = 1$, compare also [21, Theorem 1.2].

Computing

$$d(\Omega_D^{r,s,t,u,v,z})^2 = \frac{1}{2} \rho_D^{r,s,t,u,v,z} \omega_D^{123\bar{1}\bar{2}} - \frac{1}{2} \overline{\rho_D^{r,s,t,u,v,z}} \omega_D^{12\bar{1}\bar{2}\bar{3}}$$

where

$$\rho_D^{r,s,t,u,v,z} := D t^2 r^2 - D |v|^2 + t^2 s^2 - |z|^2 + i t^2 u + v \bar{z} ,$$

we have that $\Omega_D^{r,s,t,u,v,z}$ is balanced if and only if D, r, s, t, u, v, z are such that $\rho_D^{r,s,t,u,v,z} = 0$.

Firstly, suppose that $0 < D < \frac{1}{4}$. Let $s, t \in \mathbb{R} \setminus \{0\}$ be such that $s^2 > (D + s^2)^2$ and consider the J_D^D -Hermitian metric

$$\Omega_D^{s,t} := \Omega_D^{1,s,t,i(D+s^2),0,0} = i \left(\omega^{1\bar{1}} + s^2 \omega^{2\bar{2}} + t^2 \omega^{3\bar{3}} \right) + i (D + s^2) \cdot \left(\omega^{1\bar{2}} + \omega^{2\bar{1}} \right);$$

a straightforward computation shows that $\Omega_D^{s,t}$ is balanced, see also [14, §5].

In the case $D = \frac{1}{4}$, it turns out that J_D^D admits no balanced metric, see also [14, Theorem 5.9]. Indeed, for the sake of completeness, we note that, if $\tilde{\Omega}$ were a balanced metric, then one should have

$$0 = \int_X \tilde{\Omega}^2 \wedge d \left(\omega_{\frac{1}{4}}^3 - \overline{\omega_{\frac{1}{4}}^3} \right) = \int_X \tilde{\Omega}^2 \wedge \left(\sqrt{2} \omega_{\frac{1}{4}}^1 + \frac{\sqrt{2}}{2} \omega_{\frac{1}{4}}^2 \right) \wedge \overline{\left(\sqrt{2} \omega_{\frac{1}{4}}^1 + \frac{\sqrt{2}}{2} \omega_{\frac{1}{4}}^2 \right)} > 0,$$

which yields an absurd.

Summarizing, we have a curve such that:

- for $0 < D < \frac{1}{4}$, there exists a balanced metric on (X, J_D^D) ;
- for $D = \frac{1}{4}$, the complex manifold $(X, J_{\frac{1}{4}}^{\frac{1}{4}})$ admits no balanced metric and no pluriclosed metric;
- for $D = 1$, there exists a pluriclosed metric on (X, J_2^1) .

Compare also [14, Theorem 5.9], where the same example is studied in order to prove that the existence of balanced metrics and the existence of strongly-Gauduchon metrics are not closed under deformations of the complex structure.

Remark 4.7. For the study of the behaviour of the cohomology in relation to the existence of Hermitian balanced metrics or strongly-Gauduchon metrics, we refer to [27] by A. Latorre, L. Ugarte, and R. Villacampa, where, for any balanced structure, they also determine L.-S. Tseng and S.-T. Yau's spaces parametrizing the deformations in type IIB supergravity, see [38].

5. 8-DIMENSIONAL NILMANIFOLDS AND PLURICLOSED METRICS

While, for 6-dimensional nilmanifolds endowed with an invariant complex structure, the condition that a given invariant metric is pluriclosed depends just on the associated Lie algebra and not on the chosen invariant Hermitian metric, [21, Theorem 1.2], this holds no more true in dimension greater than 6, see [34, §2]. The 8-dimensional nilmanifolds endowed with an invariant complex structure such that every invariant Hermitian metric is pluriclosed have been classified by the third author and A. Tomassini in [34, Theorem 13] into two families, compare also [20, Theorem 4.1] (as a consequence, it turns out that an invariant complex structure on a 8-dimensional nilmanifold is pluriclosed for every invariant Hermitian metric if and only if it is *astheno-Kähler*⁴ for every invariant Hermitian metric, [34, Theorem 15]); more precisely, the first family consists of nilmanifolds of the type $M^6 \times \mathbb{T}^2$, where M^6 is a 6-dimensional nilmanifold endowed with an invariant complex structure such that any invariant Hermitian metric on it is pluriclosed, [21, Theorem 1.2], and \mathbb{T}^2 is the standard 2-dimensional torus endowed with the standard complex structure. In this section, we study the Bott-Chern cohomology of such 8-dimensional nilmanifolds $M^6 \times \mathbb{T}^2$. Recall that the possible M^6 have been

⁴We recall that a Hermitian metric g on a complex manifold (X, J) of complex dimension n is said to be *astheno-Kähler* (according to the terminology by J. Jost and S.-T. Yau) if the associated $(1, 1)$ -form $\omega := g(J \cdot, \cdot)$ satisfies $\partial \bar{\partial} \omega^{n-2} = 0$.

classified in [21, Theorem 3.2] by A. Fino, M. Parton, and S. Salamon in terms of their associated Lie algebra.

More precisely, in Table ?? we summarize the nilmanifolds of such type $M^6 \times \mathbb{T}^2$, that is, the nilmanifolds possibly endowed with invariant pluriclosed structures and belonging to the first class in [34, Theorem 13]. Then, in Table ??, we summarize the results of the computations of the dimensions of the Bott-Chern cohomology groups and the degree of non-Kählerianity for such 8-dimensional nilmanifolds, pointing out the existence of pluriclosed structures. (As regards notation, we follow the same conventions as in §1.2 and §2, and [14].)

TABLES

\sharp	algebra	b^1	b^2	b^3	complex structure	conditions ($\lambda \geq 0, c \geq 0, B \in \mathbb{C}, D \in \mathbb{C}$) $(S(B, c) := c^4 - 2(B ^2 + 1)c^2 + (B ^2 - 1)^2)$
00	$\mathfrak{h}_1 := (0, 0, 0, 0, 0, 0)$	6	15	20	$J := (0, 0, 0)$	
01	$\mathfrak{h}_2 := (0, 0, 0, 0, 12, 34)$	4	8	10	$J_1^D := (0, 0, \omega^{11} + D\omega^{22})$, $J_2^D := (0, 0, \omega^{12} + \omega^{11} + \omega^{12} + D\omega^{22})$	$\text{Im}D \in \{1, -1\}$ $\text{Im}D > 0$
03	$\mathfrak{h}_3 := (0, 0, 0, 0, 0, 12 + 34)$	5	9	10	$J_1 := (0, 0, \omega^{11} + \omega^{22})$ $J_2 := (0, 0, \omega^{11} - \omega^{22})$	
04						
05	$\mathfrak{h}_4 := (0, 0, 0, 0, 12, 14 + 23)$	4	8	10	$J_1 := (0, 0, \omega^{11} + \omega^{12} + \frac{1}{4}\omega^{22})$ $J_2^D := (0, 0, \omega^{12} + \omega^{11} + \omega^{12} + D\omega^{22})$	$D \in \mathbb{R} \setminus \{0\}$
06						
07	$\mathfrak{h}_5 := (0, 0, 0, 0, 13 + 42, 14 + 23)$	4	8	10	$J_1^D := (0, 0, \omega^{11} + \omega^{12} + D\omega^{22})$, $J_2 := (0, 0, \omega^{12})$ $J_3^{(\lambda, D)} := (0, 0, \omega^{12} + \omega^{11} + \lambda\omega^{12} + D\omega^{22})$	$D \in [0, \frac{1}{4}]$ $(\lambda, D) \in \{(0, x + iy) \in \mathbb{R} \times \mathbb{C} : y \geq 0, 4y^2 < 1 + 4x\}$ $\cup \{(\lambda, iy) \in \mathbb{R} \times \mathbb{C} : 0 < \lambda^2 < \frac{1}{2}, 0 \leq y < \frac{\lambda^2}{2}\}$ $\cup \{(\lambda, iy) \in \mathbb{R} \times \mathbb{C} : \frac{1}{2} \leq \lambda^2 < 1, 0 \leq y < \frac{1-\lambda^2}{2}\}$ $\cup \{(\lambda, iy) \in \mathbb{R} \times \mathbb{C} : \lambda^2 > 1, 0 \leq y < \frac{\lambda^2-1}{2}\}$
08						
09						
10	$\mathfrak{h}_6 := (0, 0, 0, 0, 12, 13)$	4	9	12	$J := (0, 0, \omega^{12} + \omega^{11} + \omega^{13})$	
11	$\mathfrak{h}_7 := (0, 0, 0, 0, 12, 13, 23)$	3	8	12	$J := (0, \omega^{11}, \omega^{12} + \omega^{12})$	
12	$\mathfrak{h}_8 := (0, 0, 0, 0, 0, 12)$	5	11	14	$J := (0, 0, \omega^{11})$	
13	$\mathfrak{h}_9 := (0, 0, 0, 0, 12, 14 + 25)$	4	7	8	$J := (0, \omega^{11}, \omega^{12} + \omega^{21})$	
14	$\mathfrak{h}_{10} := (0, 0, 0, 0, 12, 13, 14)$	3	6	8	$J := (0, \omega^{11}, \omega^{12} + \omega^{21})$	
15	$\mathfrak{h}_{11} := (0, 0, 0, 0, 12, 13, 14 + 23)$	3	6	8	$J^B := (0, \omega^{11}, \omega^{12} + B\omega^{12} + B-1 \omega^{21})$	$B \in \mathbb{R} \setminus \{0, 1\}$
16	$\mathfrak{h}_{12} := (0, 0, 0, 0, 12, 13, 24)$	3	6	8	$J^B := (0, \omega^{11}, \omega^{12} + B\omega^{12} + B-1 \omega^{21})$	$\text{Im}B \neq 0$
17	$\mathfrak{h}_{13} := (0, 0, 0, 0, 12, 13 + 14, 24)$	3	5	6	$J^{(B, c)} := (0, \omega^{11}, \omega^{12} + B\omega^{12} + c\omega^{21})$	$c \neq B-1 , (c, B) \neq (0, 1), S(B, c) < 0$
18	$\mathfrak{h}_{14} := (0, 0, 0, 0, 12, 14, 13 + 42)$	3	5	6	$J^{(B, c)} := (0, \omega^{11}, \omega^{12} + B\omega^{12} + c\omega^{21})$	$c \neq B-1 , (c, B) \neq (0, 1), S(B, c) = 0$
19	$\mathfrak{h}_{15} := (0, 0, 0, 0, 12, 13 + 42, 14 + 23)$	3	5	6	$J_1 := (0, \omega^{11}, \omega^{21})$ $J_2^c := (0, \omega^{11}, \omega^{12} + c\omega^{21})$ $J_3^{(B, c)} := (0, \omega^{11}, \omega^{12} + B\omega^{12} + c\omega^{21})$	$c \neq 1$ $c \neq B-1 , (c, B) \neq (0, 1), S(B, c) > 0$
20						
21						
22	$\mathfrak{h}_{16} := (0, 0, 0, 0, 12, 14, 24)$	3	5	6	$J^B := (0, \omega^{11}, \omega^{12} + B\omega^{12})$	$ B = 1, B \neq 1$
23	$\mathfrak{h}_{19} := (0, 0, 0, 0, 12, 23, 14 - 35)$	3	5	6	$J_1 := (0, \omega^{13} + \omega^{13}, i(\omega^{12} - \omega^{21}))$ $J_2 := (0, \omega^{13} + \omega^{13}, -i(\omega^{12} - \omega^{21}))$	
24						
25	$\mathfrak{h}_{26}^+ := (0, 0, 12, 13, 23, 14 + 25)$	2	4	6	$J_1 := (0, \omega^{13} + \omega^{13}, i\omega^{11} + i(\omega^{12} - \omega^{21}))$ $J_2 := (0, \omega^{13} + \omega^{13}, i\omega^{11} - i(\omega^{12} - \omega^{21}))$	
26						

TABLE 1. M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa's classification of linear integrable complex structures on 6-dimensional nilpotent Lie algebras up to equivalence, [14].

#	algebra	complex structure	$\partial\bar{\partial}\left(\sum_{j=1}^3\omega^j\wedge\bar{\omega}^j\right)$
01	\mathfrak{h}_2	J_1^D	$(D + \bar{D})\omega^{12\bar{1}\bar{2}}$
02		J_2^D	$(-2 + D + \bar{D})\omega^{12\bar{1}\bar{2}}$
05	\mathfrak{h}_4	J_1	$-\frac{1}{2}\omega^{12\bar{1}\bar{2}}$
06		J_2^D	$2(D - 1)\omega^{12\bar{1}\bar{2}}$
07	\mathfrak{h}_5	J_1^D	$(2D - 1)\omega^{12\bar{1}\bar{2}}$
08		J_2	$-\omega^{12\bar{1}\bar{2}}$
09		$J_3^{(\lambda, D)}$	$(-1 - \lambda^2 + D + \bar{D})\omega^{12\bar{1}\bar{2}}$
12	\mathfrak{h}_8	J	0

TABLE 3. Non-toric 6-dimensional nilmanifolds endowed with invariant complex structures and possibly with invariant pluriclosed metrics.

”””

”””

”””

FIGURES




FIGURE 1. Non-Kählerianity of 6-dimensional nilmanifolds endowed with the invariant complex structures in M. Ceballos, A. Otal, L. Ugarte, and R. Villacampa’s classification, [14].

REFERENCES

- [1] A. Aeppli, On the cohomology structure of Stein manifolds, *Proc. Conf. Complex Analysis (Minneapolis, Minn., 1964)*, Springer, 1965, 58–70.
- [2] L. Alessandrini, G. Bassanelli, The class of compact balanced manifolds is invariant under modifications, *Complex analysis and geometry (Trento, 1993)*, Lecture Notes in Pure and Appl. Math. **173**, Dekker, 1996, 1–17.
- [3] B. Alexandrov, S. Ivanov, Vanishing theorems on Hermitian manifolds, *Differ. Geom. Appl.* **14** (2001), no. 3, 251–265.
- [4] A. Andrada, M. L. Barberis, I. Dotti, Classification of abelian complex structures on 6-dimensional Lie algebras, *J. Lond. Math. Soc. (2)* **83** (2011), no. 1, 232–255. Corrigendum: Classification of abelian complex structures on six-dimensional Lie algebras, *J. Lond. Math. Soc. (2)* **87** (2013), no. 1, 319–320.
- [5] D. Angella, The cohomologies of the Iwasawa manifold and of its small deformations, *J. Geom. Anal.* **23** (2013), no. 3, 1355–1378.
- [6] D. Angella, A. Tomassini, On the $\partial\bar{\partial}$ -Lemma and Bott-Chern cohomology, *Invent. Math.* **192** (2013), no. 1, 71–81.
- [7] Ch. Benson, C. S. Gordon, Kähler and symplectic structures on nilmanifolds, *Topology* **27** (1988), no. 4, 513–518.
- [8] B. Bigolin, Gruppi di Aeppli, *Ann. Scuola Norm. Sup. Pisa (3)* **23** (1969), no. 2, 259–287.
- [9] B. Bigolin, Osservazioni sulla coomologia del $\partial\bar{\partial}$, *Ann. Scuola Norm. Sup. Pisa (3)*, **24**, (1970), no. 3, 571–583.
- [10] J.-M. Bismut, A local index theorem for non-Kähler manifolds, *Math. Ann.* **284** (1989), no. 4, 681–699.
- [11] J.-M. Bismut, Laplacien hypoelliptique et cohomologie de Bott-Chern, *C. R. Math. Acad. Sci. Paris* **349** (2011), no. 1-2, 75–80.
- [12] J.-M. Bismut, *Hypoelliptic Laplacian and Bott-Chern cohomology. A theorem of Riemann-Roch-Grothendieck in complex geometry*, Progress in Mathematics, **305**, Birkhäuser/Springer, Cham, 2013.
- [13] G. R. Cavalcanti, Hodge theory and deformations of SKT manifolds”, [arXiv:1203.0493v5](https://arxiv.org/abs/1203.0493v5) [math.DG].
- [14] M. Ceballos, A. Otal, L. Ugarte, R. Villacampa, Invariant Complex Structures on 6-Nilmanifolds: Classification, Frölicher Spectral Sequence and Special Hermitian Metrics, *J. Geom. Anal.* (2014), DOI: 10.1007/s12220-014-9548-4.
- [15] S. Console, A. Fino, Dolbeault cohomology of compact nilmanifolds, *Transform. Groups* **6** (2001), no. 2, 111–124.
- [16] L. A. Cordero, M. Fernández, A. Gray, L. Ugarte, Compact nilmanifolds with nilpotent complex structures: Dolbeault cohomology, *Trans. Amer. Math. Soc.* **352** (2000), no. 12, 5405–5433.
- [17] P. de Bartolomeis, A. Tomassini, On solvable generalized Calabi-Yau manifolds, *Ann. Inst. Fourier* **56** (2006), no. 5, 1281–1296.
- [18] P. Deligne, Ph. Griffiths, J. Morgan, D. Sullivan, Real homotopy theory of Kähler manifolds, *Invent. Math.* **29** (1975), no. 3, 245–274.
- [19] J.-P. Demailly, *Complex Analytic and Differential Geometry*, <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>, 2009.
- [20] N. Enrietti, A. Fino, L. Vezzoni, Tamed Symplectic forms and Strong Kahler with torsion metrics, *J. Symplectic Geom.* **10** (2012), no. 2, 203–224.
- [21] A. Fino, M. Parton, S. Salamon, Families of strong KT structures in six dimensions, *Comment. Math. Helv.* **79** (2004), no. 2, 317–340.
- [22] A. Fujiki, On automorphism groups of compact Kähler manifolds, *Invent. Math.* **44** (1978), no. 3, 225–258.
- [23] K. Hasegawa, Minimal models of nilmanifolds, *Proc. Amer. Math. Soc.* **106** (1989), no. 1, 65–71.
- [24] A. Hattori, Spectral sequence in the de Rham cohomology of fibre bundles, *J. Fac. Sci. Univ. Tokyo Sect. I* **8** (1960), no. 1960, 289–331.
- [25] K. Kodaira, D. C. Spencer, On deformations of complex analytic structures, III, Stability theorems for complex structures, *Ann. of Math. (2)*, **71** (1960), no. 1, 43–76.

- [26] R. Kooistra, *Regulator currents on compact complex manifolds*, Ph.D. thesis, University of Alberta.
- [27] A. Latorre, L. Ugarte, R. Villacampa, On the Bott-Chern cohomology and balanced Hermitian nilmanifolds, *Internat. J. Math.* **25** (2014), no. 6, 1450057, 24 pp.
- [28] L. Magnin, Sur les algèbres de Lie nilpotentes de dimension ≤ 7 , *J. Geom. Phys.* **3** (1986), no. 1, 119–144.
- [29] M. L. Michelsohn, On the existence of special metrics in complex geometry, *Acta Math.* **143** (1983), no. 3-q4, 261–295.
- [30] V. V. Morozov, Classification of nilpotent Lie algebras of sixth order, *Izv. Vysš. Učebn. Zaved. Matematika* **1958** (1958), no. 4 (5), 161–171.
- [31] K. Nomizu, On the cohomology of compact homogeneous spaces of nilpotent Lie groups, *Ann. of Math. (2)* **59** (1954), no. 3, 531–538.
- [32] S. Rollenske, Geometry of nilmanifolds with left-invariant complex structure and deformations in the large, *Proc. London Math. Soc.* **99** (2009), no. 2, 425–460.
- [33] S. Rollenske, Dolbeault cohomology of nilmanifolds with left-invariant complex structure, in W. Ebeling, K. Hulek, K. Smoczyk (eds.), *Complex and Differential Geometry: Conference held at Leibniz Universität Hannover, September 14 – 18, 2009*, Springer Proceedings in Mathematics **8**, Springer, 2011, 369–392.
- [34] F. A. Rossi, A. Tomassini, On strong Kähler and astheno-Kähler metrics on nilmanifolds, *Adv. Geom.* **12** (2012), no. 3, 431–446.
- [35] Y. Sakane, On compact complex parallelisable solvmanifolds, *Osaka J. Math.* **13** (1976), no. 1, 187–212.
- [36] S. M. Salamon, Complex structures on nilpotent Lie algebras, *J. Pure Appl. Algebra* **157** (2001), no. 2-3, 311–333.
- [37] M. Schweitzer, Autour de la cohomologie de Bott-Chern, [arXiv:0709.3528v1](https://arxiv.org/abs/0709.3528v1) [math.AG].
- [38] L.-S. Tseng, S.-T. Yau, Generalized cohomologies and supersymmetry, *Comm. Math. Phys.* **326** (2014), no. 3, 875–885.
- [39] L. Ugarte, Hermitian structures on six-dimensional nilmanifolds, *Transform. Groups* **12** (2007), no. 1, 175–202.
- [40] L. Ugarte, R. Villacampa, Non-nilpotent complex geometry of nilmanifolds and heterotic supersymmetry, *Asian J. Math.* **18** (2014), no. 2, 229–246.
- [41] J. Varouchas, Propriétés cohomologiques d’une classe de variétés analytiques complexes compactes, *Séminaire d’analyse P. Lelong-P. Dolbeault-H. Skoda, années 1983/1984*, 233–243, Lecture Notes in Math., **1198**, Springer, Berlin, 1986.

(Daniele Angella) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127, PISA, ITALY

Email address: angella@mail.dm.unipi.it

Current address: Centro di Ricerca Matematica “Ennio de Giorgi”, Collegio Puteano, Scuola Normale Superiore, Piazza dei Cavalieri 3, 56126 Pisa, Italy

Email address: daniele.angella@sns.it

Email address: daniele.angella@gmail.com

(Maria Giovanna Franzini) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE 53/A, 43124, PARMA, ITALY

Email address: mariagiovanna.franzini@studenti.unipr.it

(Federico Alberto Rossi) DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DI MILANO BICOCCA, VIA COZZI 53, 20125, MILANO, ITALY

Email address: f.rossi46@campus.unimib.it