



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Possible hybrid speciation for two Malagasy species of Piper L. (Piperaceae)**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Possible hybrid speciation for two Malagasy species of Piper L. (Piperaceae) / Enrico Palchetti,  
Massimo Gori,  
Stefano Biricolti,  
Alberto Masoni,

*Availability:*

The webpage <https://hdl.handle.net/2158/1213147> of the repository was last updated on 2021-07-19T10:20:50Z

*Published version:*

DOI: 10.13128/caryologia-1133

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)



**Citation:** E. Palchetti, M. Gori, S. Biricolti, A. Masoni, L. Bini, C. Tani, S. Falsini, E. Corti, A. Papini (2020) Possible hybrid speciation for two Malagasy species of *Piper* L. (*Piperaceae*). *Caryologia* 73(4): 27-38. doi: 10.13128/caryologia-1133

**Received:** November 12, 2020

**Accepted:** November 15, 2020

**Published:** May 19, 2021

**Copyright:** © 2020 E. Palchetti, M. Gori, S. Biricolti, A. Masoni, L. Bini, C. Tani, S. Falsini, E. Corti, A. Papini. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/caryologia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Competing Interests:** The Author(s) declare(s) no conflict of interest.

**Author Contributions:** Conceptualization, Palchetti, Biricolti and Gori; methodology, Gori, Papini; software, Papini; validation, Palchetti, Gori, Papini, Biricolti, Calamai; formal analysis, Papini, Gori, Biricolti, Bini; investigation, Gori, Bini, Falsini, Corti, Calamai; resources, Palchetti; data curation, Gori; writing—original draft preparation, all authors; project administration, Palchetti; funding acquisition, Palchetti. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Company TozziGreen, La Tour - 26 Etage Rue Ravoninahitriniarivo – Ankorondrano, Antananarivo 101 Madagascar.

## Possible hybrid speciation for two Malagasy species of *Piper* L. (*Piperaceae*)

ENRICO PALCHETTI<sup>1</sup>, MASSIMO GORI<sup>1,4</sup>, STEFANO BIRICOLTI<sup>1,\*</sup>, ALBERTO MASONI<sup>1</sup>, LORENZO BINI<sup>4</sup>, CORRADO TANI<sup>2</sup>, SARA FALSINI<sup>2</sup>, EMILIO CORTI<sup>2</sup>, ALESSIO PAPINI<sup>2,3</sup>

<sup>1</sup> Department of Agriculture, Food, Environment and Forestry (DAGRI) University of Florence, Piazzale delle Cascine, 18, 50144 Florence (FI), Italy; E-mail: [enrico.palchetti@unifi.it](mailto:enrico.palchetti@unifi.it)

<sup>2</sup> Department of Biology, University of Florence, Via Micheli, 3, Firenze, Italy

<sup>3</sup> CSET Tropical Herbarium University of Florence, Italy; E-mail: [alpapini@unifi.it](mailto:alpapini@unifi.it)

<sup>4</sup> Interdepartmental Service Centre for Agricultural, Chemical and Industrial Biotechnologies (CIBIACI), University of Florence, Via Romana, 21, 50125 Florence, Italy; E-mail: [massimo.gori@unifi.it](mailto:massimo.gori@unifi.it)

\*Corresponding author. E-mail: [stefano.biricolti@unifi.it](mailto:stefano.biricolti@unifi.it)

**Abstract.** Two new species of genus *Piper* L. from Madagascar: *Piper malgassicum* Papini, Palchetti, M. Gori & Rota Nodari and *Piper tsarasotrae* Papini, Palchetti, M. Gori & Rota Nodari, were analyzed to investigate their phylogenetic position and evolutionary history. Both plastidial and nuclear markers were used for sequencing. The plastidial markers (*ndhF* and *trnL* intron) showed a close relationship between the two species with respect to the other species of *Piper*. Both species appeared phylogenetically related to the African *P. guineense* and the Malagasy/Mascarenhas endemic *P. borbonense*. The nuclear marker (*G3pdh*) amplification produced two separate sets of sequences: “long” sequences and “short” sequences, characterized by some long deletions. Analyzing together the nuclear sequences, we observed that the “long” sequence of *P. tsarasotrae* had a stricter relationship to the African accessions of *P. guineense*, while the accession of *P. malgassicum* was more strictly related to *P. borbonense*. On the contrary both “short” sequences of *P. malgassicum* and *P. tsarasotrae* resulted phylogenetically related to Asian accessions and more distantly related to the formerly cited species. This unexpected result was tentatively explained with a more ancient hybridization event between an ancestor of *P. malgassicum* and *P. tsarasotrae* (and possibly *P. borbonense*) and an Asian species of *Piper*. The Asian contribution would have produced the ancestors carrying the “short” sequences. A more recent hybridization event would have led to the separation of *P. malgassicum* from *P. tsarasotrae* with an African pollen-derived genome contribution from *P. guineense* or, more probably, an ancestor thereof, to an ancestor of *P. tsarasotrae*. The chromosome numbers of *P. tsarasotrae* ( $2n =$  about 38) and *P. malgassicum* ( $2n =$  about 46), were more similar to the Asian species than to the American species. Unfortunately, no chromosome number of the African species *P. guineense* is currently available, to compare the chromosomal numbers.

**Keywords:** *Piper malgassicum*, *Piper tsarasotrae*, *Piperaceae*, chromosomes, hybridization, DNA sequences, *G3pdh*, *trnL*, *ndhF*, Malagasy biodiversity.

## 1. INTRODUCTION

Genus *Piper* L. (Piperaceae) is one of the largest genera of Angiosperms, with more than 2000 species (Quijano-Abril et al. 2006) and were considered belonging to a basal group of angiosperms, the so called “paleoherbs” (Loconte et al. 1991).

*Piper* is a pantropical genus developing highly variable growth forms (Isnard et al. 2012), with the highest biodiversity in the American continent with a number of species ranging from 500 (Burger 1972; Tebbs 1993), to 1100 (Jaramillo and Manos 2001), later increased to more than 1800 (Ulloa Ulloa et al. 2017), many of them with a small distribution area (Quijano-Abril et al. 2006).

The separation of species is often tricky, due to the small size of the floral parts and hence the number of synonyms may be high (Suwanphakdee et al. 2016), while other species tend to get naturalized (Smith et al. 2008). While only two species are known as native of the African continent, *P. guineense* Thonn. and *P. capense* L. f., more species are known of Madagascar, even if some of them are known only for a single or few herbarium samples. The currently recognized species in Madagascar are *P. heimii* C. DC., *P. pachyphyllum* Baker and possibly *P. borbonense* (Miq.) C. DC., described for the Bourbon island, nowadays La Reunion (Weil et al. 2017), belonging to the Mascarenhas Islands. However, its presence in Madagascar was affirmed by De Candolle (1869; 1923). The fact that *P. borbonense* is cultivated makes more complex to understand its real distribution area (Palchetti et al. 2018).

*Piper malgassicum* Papini, Palchetti, M. Gori, Rota Nodari and *Piper tsarasotrae* Papini, Palchetti, M. Gori, Rota Nodari, were recently described as new Malagasy species (Palchetti et al. 2018) and are of economic interest, since their dried fruits are often mixed with *P. borbonense* to produce the typical Malagasy spice called in local language “voatsiperifery” pepper.

The aim of the investigation was to understand how the malagasy species might have been originated and their relationships with the African and the Asian species. This knowledge will help to understand how the Malagasy species used as spices may be related to *P. nigrum* with possible future biotechnological implications. The chosen method for answering the research goal was the analysis of DNA sequences both of nuclear and plastidial origin and the chromosome numbers of *P. malgassicum* and *P. tsarasotrae*.

## 2. MATERIALS AND METHODS


A first round of sample collection within the internal area of Madagascar was conducted in 2016 and the

samples have been submitted to analyses. The results have been reported in Palchetti et al. (2018) but, in order to get a deeper knowledge about the genetic asset of the two species and to confirm the obtained results a second round of sample collection has been carried on in 2019. 4 new plants were collected in two different areas of the Ambositra region in Madagascar. The first 2 plants, belonging to the *P. malgassicum* type, were collected in the tropical rainy forest of Vohiday and the second 2 plants, belonging to the *P. tsarasotrae* type, in the semi-dry area of the Tsaratsotra village. These plants were compared with the samples of *P. tsarasotrae* and *P. malgassicum* which have been used for a previous investigation that included the description of the species (Palchetti et al. 2018). Samples were conserved either in ethanol 96% either as herbarium sample by the ET (Tropical Herbarium of Florence, CSET, <https://www.bio.unifi.it>). Some seeds were also germinated in Florence for karyotyping. The DNA used for this work was extracted from tissue conserved in ethanol 96% (Murray et al. 1996; Bressan et al. 2014).

DNA was extracted from 40 mg of the ethanol preserved leaves after drying under vacuum. The starting material was inserted in 2 ml tube, together with tungsten carbide beads, frozen in liquid nitrogen and finely ground in a tissue homogenizer (Tissue Lyser<sup>®</sup>, Qiagen). DNA was extracted using Invisorb Spin Plant Mini kit (Stratec molecular<sup>®</sup>) according to the manufacture’s guidelines.

Amplification of the trnL (UAA) intron (*trnL*) and the low copy nuclear gene glyceraldehyde 3-phosphate dehydrogenase (*G3pdh*) followed respectively the protocols by Taberlet et al. (1991) and Strand et al. (1997).

Two new primer pairs were designed using the chloroplast genome sequence of *P. kadsura* (GenBank<sup>®</sup>: KT223569.1) as template to cover the entire NADH dehydrogenase F (*ndhF*) plastid gene: *ndhF*-F3\_forward 5'-AGGTTCTTATCGAGCCGCTT-3' and *ndhF*-F3\_reverse 5'-GTAAGAAGAAATGCGCCCC-3' and *ndhF*-F10\_forward 5'-CTTCGCCGTATGTGGGCTTT-3' and *ndhF*-F10\_reverse 5'-TCGACCAAAGCAAGCAAGAG-3'. The amplicons have been directly and bi-directionally sequenced by using the corresponding primers for each amplified sequence. Since direct sequencing of *G3pdh* showed fragments of extra peaked sequencing data, we proceeded with cloning with InsTAclone PCR Cloning Kit (Thermo Scientific<sup>®</sup>) of the *G3pdh* amplification products. Several colonies for each cloned sample were amplified using T7 and SP6 primers whose sites are located at the boundaries of the cloning region. PCR products were purified using the QIAquick PCR Purification Kit (Qiagen) and sent to the University of



**Figure 1.** Alignment of the long and short *G3pdh* sequences isolated from *P. malgassicum* and *P. tsarasotrae* using BioEdit software (Hall 1999). Shaded fragments represent the primers used for amplification.

Florence internal sequencing service CIBIACI ([www.cibiaci.unifi.it](http://www.cibiaci.unifi.it)). Manual correction and assembly of the sequences was performed using the software Multalin (Corpet 1988) and MEGA7 (Kumar et al. 2016). Unexpectedly, two DNA sequences were obtained, after

removing the cloning vector fragments, showing a different size: 965bp and 1058bp which were named “short” and “long” sequences respectively (Figure 1).

At a first sight only the long sequences of *G3pdh* have been considered as the right ones because, as

observed by Smith et al. (2008), no paralogs have been detected in a great deal of other *Piper* species and therefore other results have been discarded as PCR artifacts. In the present work a second thorough revision of the sequencing output has been carried on and showed the presence of overlapping peaks in all the samples and an additional round of analysis of the colonies confirmed the presence of the short sequences. To rule out any doubt, two additional specimen for each species has been collected and submitted to amplification and cloning in order to confirm the presence of the short sequence. As all the samples showed the same pattern, we decide to use also this “short” sequence to study the phylogeny of these two *piper* species, by comparing with all the accession present online.

The sequences used during our investigation are available in GenBank®: *Piper tsarasotrae* *G3pdh* long sequence (MH234634), *G3pdh* short sequence (MT793801), *trnL* (MH234638), *ndhF* (MH234636) and *Piper malgassicum*: *G3pdh* long sequence (MH234633), *G3pdh* short sequence (MT793800), *trnL* (MH234637), *ndhF* (MH234635).

### 2.1. Phylogenetic analysis

The DNA sequences were aligned with CLUSTALX 2.0 and checked by eye for manual adjustment. The plastidial and the nuclear sequences were aligned separately to produce matrices that were later combined with the software combinex2\_0.py (Python version 2.6.4; Biopython 1.57), by A. Papini, released under GPL license and available at [www.unifi.it/caryologia/PapiniPrograms.html](http://www.unifi.it/caryologia/PapiniPrograms.html) as implemented in Bandara et al. (2013) and in Simeone et al. (2016).

The phylogenetic analysis was executed on both cpDNA (*ndhF* and *trnL*) and nuclear sequences (*G3pdh*). Maximum parsimony analysis was performed with PAUP\* 4.0b1 (Swofford 1998, 2001). The genbank sequences of *P. humistratum* Görts & K. U. were used as outgroups both in the nuclear and the plastid genes matrix, following the previous phylogenetic analysis by Smith et al. (2008). This sequences used as outgroup resulted belonging to the sister clade with respect to the clade containing the African species and the other related clades in Smith et al. (2008). References of the other species used in the analysis are summarized (with GenBank® codes) in Table 1 in Smith et al. (2008). All characters had equal weight and unordered state transitions. Gaps were coded with the “simple indel coding” model (Simmons and Ochoterena 2000), with the software Gap-coder (Young and Healy 2003) and added to the final matrix after the DNA sequences as in Papini et al. (2004).

The evolutionary model implemented in MrBayes for treating gaps was the same as that proposed by Lewis et al. (2001) for treating morphological data, the Mk model, justified as simple absence/presence of the character without *a priori* assignment of different weights.

We used MrMODELTEST 2.0 (Nylander 2004) to choose the best evolutionary model of DNA sequences on the basis of the Akaike information criterion (Akaike 1974). The best model was used as settings with MrBayes 3.2.7 (Ronquist et al. 2012) for Bayesian Inference. A maximum likelihood (ML) phylogenetic analysis was carried out with RaxML (Stamatakis 2014) and the resulting trees were edited with Figtree (Rambaut and Drummond 2010). We mapped the support on the tree branches with the results of the Bayesian phylogenetic analysis after removing the first trees with low likelihood values as “burn-in”, as in Papini et al. (Papini et al. 2007; Papini et al. 2011). The remaining trees were used to produce a 50% majority-rule consensus tree in which the percentage indicated on branches was used as a measure of the Bayesian posterior probability.

### 2.2. Karyological analysis

Chromosomes images were obtained from somatic mitoses recorded from root tips of only one plant living in a pot. The procedure was the same as in Mosti et al. (2011) and Mousavi et al. (2013), with a pretreatment in 8-hydroxyquinoline and fixation in Carnoy. Then the material was hydrolyzed in HCl and then stained with Lacto-propionic-orceine.

We observed metaphase plates of meristematic cells, with the technique of fresh squashes of root tips. Chromosome counts were made during direct observations with the microscope, and later recounted on enlarged digital images. Images were recorded with a microscope Leica DM RB Fluo.

## 3. RESULTS


Amplification of two plastid fragments named *ndhF* and *trnL* intron was carried on and the amplicons correctly sequenced producing reads of 1860 bp and 920 bp, respectively. Cloning of the amplicon of the nuclear gene *G3pdh* of *P. malgassicum* and *P. tsarasotrae* allowed to isolate two haplotypes, which were named “long” (1060bp for *P. malgassicum* and 1045bp for *P. tsarasotrae*) and “short” 965 bp (for both species) after their size. We used a total of 71 sequences, considering separately the short and long sequences of *P. malgassicum* and *P. tsarasotrae* for *G3pdh* and the plastid sequences




**Figure 2.** Maximum likelihood tree produced by RAXML with nuclear sequences. The supports above or below the branches are, respectively, the bootstrap resampling support with maximum likelihood criterion produced by RAXML, and the bayesian support calculated including the information derived from indels. In case the bayesian support is lower than 50, it is not indicated on the figure.

matrix. The total alignment of the *G3pdh* region was 1127 nucleotides long including gaps. The final parts of the sequences were very variable and hence the align-

ment was ambiguous. For this reason, we excluded the characters from position 957 to 1127. The rest of the alignment was used for indels (gap) coding (with the



**Figure 3.** Maximum likelihood tree produced by RAXML with chloroplast sequences. The support indexes indicated on the tree are the same as in Figure 2 (maximum likelihood bootstrap and bayesian support).



**Figure 4.** Chromosomes. A) *P. malgassicum* number of chromosomes:  $2n = \text{about } 46$ . Bar =  $5 \mu\text{m}$ ; B) *P. tsarasotrae*:  $2n = \text{about } 38$ . Bar =  $5 \mu\text{m}$ .

software gapcoder), resulting in further 99 characters that were inserted after the nucleotide sequences. The plastid genes *ndhF* and *trnL* were inserted one after the other in the sequence, producing an aligned matrix of 2016 characters. The coding of indels resulted in further 115 characters. RAxML applied on the nuclear *G3pdh* matrix (indels coding excluded) produced a maximum likelihood tree with bootstrap support obtained with 1000 replicates (Figure 2).

The support on branches corresponds to maximum likelihood bootstrap support (left) and Bayesian support with gaps (on the right). The same method was using for the plastid matrix (Figure 3).

Comparing the two maximum likelihood trees, the one based on nuclear DNA data (*G3pdh* sequences) and that obtained with plastid markers, we could observe that in the first case *P. malgassicum*, clustered together and as sister group of *P. borbonense* (Figure 2), another species from an island, La Reunion, which lies relatively close to Madagascar. This relationship is corroborated by 100% maximum likelihood bootstrap (MLS) and bayesian (BS) support. The other Malagasy species, *P. tsarasotrae*, typical of arid forest, was more strictly related to the entries of the African species *P. guineense*, with 100% MLS and 100% BS. All these species formed a well characterized clade with 89% MLS and 100% BS and their closest species appeared to be Asian species *P. caninum*, (Figure 2). The BS without considering gaps coding gave the same support in this clade.

The “short” sequences of *G3pdh* of both *P. tsarasotrae* and *P. malgassicum* clustered together within a group of Asian species, mainly originating from Malaysia and Australia with 98% MLS and 100% BS (Figure 2).

The (phylogenetic) story told by the data obtained from chloroplast genome sequences was quite different: the Malagasy species *P. tsarasotrae* and *P. malgassicum* clustered together with the phylogeographically close *P. borbonense* with 90% MLS and 100% BS, while the 5 accessions of the African *P. guineense* were in a more external condition with respect to the former group and separated in two groups, one from Cameroon (NW Africa) and one from Uganda/Kenya (Central-East Africa). All these species together formed a monophyletic group with 64% MLS and 96% BS (95% bayesian support in the analysis without gaps). Also in this case *P. caninum*, together with *P. rothianum*, was the outgroup to the African + Malagasy species (Figure 3) with 80% MLS and 98% BS (99% without gaps). Adding indels data to the matrix did not appear to increase the support value of nodes in the plastidial genes tree.

The counted chromosome numbers varied from  $2n=46\pm 2$  in *P. malgassicum* (Figure 4A) to  $2n=36\pm 2$  in *P. tsarasotrae* (Figure 4B). The uncertainty in the counts, that should be taken only as preliminary result, derived from the small size of the chromosomes (many of them less than  $1 \mu\text{m}$  of length), the low amount of metaphases in the root tips of the plants cultivated in Florence and the apparently small size of the mitotic spindle, leading



to partial overlapping of many of the small chromosomes.


The two currently known areal of the two species (two new localities discovered here) is shown in Figure 5.

#### 4. DISCUSSION

The fact that the phylogenetic history based on the chloroplast markers told a different tale with respect to the tree produced with nuclear markers may be explained with a possible ancient hybridization/introgression event with pollen coming from an ancestor of the African *P. guineense* and reaching the ancestor of *P. tsarasotrae*, that would hence share some part of the nuclear genome with the African species. The only species of *Piperaceae* analyzed under the point of view of the type of plastid inheritance was a species of *Peperomia*, which resulted to have only maternal plastidial inheritance (Corriveau and Coleman 1988). The presence of the “short” *G3pdh* nuclear sequences may be related to a still more ancient hybridization event involving the ancestor of the Malagasy species and some ancestor of Asian origin. Also in this case probably, with Asian pollen entering in contact with the ancestor’s stigma of the Malagasy species. As a matter of fact the closest relatives to the African species *sensu lato* (including the Malagasy and the Reunion species) are Asian, with the closest species (among those here sampled) apparently from Malaysia (Figure 3 and Figure 4). Apparently interspecific hybrids can be obtained in genus *Piper* also experimentally (Vanaja et al. 2008), while the hybrid origin of several Andean species was already proposed by Quijano-Abril et al. (2006).

The presence of paralogs of *G3pdh* in angiosperms may represent a problem in several phylogenetic analysis (Hurteau and Spivack 2002; Liu et al. 2009; Sun et al 2012). However, here most of the indels were found in the introns of the gene and hence we are not able to assess the functionality of the short sequences.

The preliminary results about the chromosome numbers scored about  $2n=46+2$  in *P. malgassicum* and  $2n=36+2$  in *P. tsarasotrae*. The uncertainty in the counts was due to the small dimensions of the chromosomes that were observed in most of the species of the genus, together with stickiness (Samuel 1987; Samuel and Morawetz 1989), the low amount of metaphases in the root tips of the plants cultivated *in vitro* and the apparently small size of the mitotic fuse, leading to partial overlapping of many of the small chromosomes. The mitotic spindle can reach dimensions up to 60  $\mu\text{m}$  (Wühr et al. 2008; Petry 2016), while in *P. malgassicum* and *P. tsarasotrae* it was about 15-20  $\mu\text{m}$  (see Figure 4).



**Figure 5.** Geographical localization of the sampling area for *P. tsarasotrae* (yellow) and *P. malgassicum* (white). Area of the sampling campaign of 2018 (a and b) and 2019 (A and B) for *P. tsarasotrae* (a/A) and *P. malgassicum* (b/B) respectively.

The chromosome numbers in genus *Piper* are very variable, ranging from  $2n=26$  to  $2n=104$ , with some species apparently able to possess several possible chromosome numbers (Samuel 1987). Most new world species show a karyotype of  $2n=26$  and  $x=13$  (Samuel and Morawetz 1989), with some exceptions having  $2n=28$  chromosomes (Maugini 1953). In Asia tetraploids  $2n=52$  would prevail (Samuel 1987). No data was available for African and Malagasy species up to the here presented results. However, the clear difference in karyotype between *P. tsarasotrae* and *P. malgassicum*, two species otherwise strictly phylogenetically related, may confirm a possible hybridization/introgression event with a species with a different chromosome number with respect to the ancestor of the Malagasy species. As a matter of fact, also Nair et al. (1993) explained the observation of a triploid plant of *P. nigrum* ( $2n=78$ ) as the result of a natural crossing between  $2n=52$  and  $2n=104$  plants.

The progeny showed a range of variation from  $2n=52$  to  $2n=104$  and production of aneuploid viable pollen (Nair et al. 1993). Hybridization may influence diversity, including gene flow from one taxon to another (introgression) and the formation of new, stable hybrid taxa and, possibly, speciation (Mallet 2007; Vallejo-Marín and Hiscock 2016).

As preliminary guess, the two different chromosome numbers of the Malagasy species may have arisen as a consequence of hybridization of a  $2n=52$  species with a  $2n=26$  (*P. tsarasotrae*) and with another  $2n=52$  species (*P. malgassicum*), respectively, with following aneuploid reductions. The two species have close areals (they are almost sympatric), even if they tend to occupy different habitats, more arid *P. tsarasotrae* (not lianous habit) and more humid (lianus habit) *P. malgassicum*. Such proximity of two closely related species may be considered another possible indication of a relatively fast speciation event.

A discordance between plastid and nuclear inheritance inferred through DNA sequencing has been often related to reticulate evolution and species of hybrid origin (García et al. 2014; Stefanović et al. 2007; Aubriot et al. 2018), even if the karyological data may be a decisive evidence, it has been rarely used in relationship to the DNA sequence evidence as, for instance in Selvi et al. (2002). Here the hypothesis of an hybrid origin for the two investigated species may explain the presence of a double *G3pdh* sequence in both of them.

The relationship between the Malagasy species and *P. guineense* with the asian species as members of *Piper* s. s. was already proposed by Jaramillo and Callejas (2004) and Jaramillo et al. (2008) as a result of a dispersal event, and our results do not disagree with this position. Apparently, in Africa and Madagascar the conditions leading to the wide diversification observed in South-american *Piper* (Martines et al. 2015) are lacking or less capable of influencing the speciation process.

## 5. CONCLUSIONS

The surprising discrepancy between the nuclear and the plastid phylogeny could be explained with an ancestral introgression event due probably to pollen contribution from an ancestor of the African mainland *P. guineense* towards the ancestor of *P. tsarasotrae*. The presence of possible paralogs of the nuclear gene *G3pdh*, clustering together with more distantly related Asian species lead to the hypothesis that a second more ancient hybridation/introgression event would have occurred between south Asian species and the ancestor of the Malagasy species. The chromosome numbers observed

in the Malagasy species would confirm different evolutionary history.

Further studies about the karyotypes of the Malagasy species, the african *P. guineense* and *P. borbonense* will be necessary together with the investigation of the possible presence of short paralog sequences of *G3pdh* in *P. borbonense*.

## ACKNOWLEDGMENTS

the Authors would like to acknowledge Dr. Nicola Gandolfi from the NGO TsiryParma, Ambositra, Madagascar, for the support given in the germplasm collection in field.

## REFERENCES

- Akaike H (1974) A new look at the statistical model identification. IEEE Trans Autom Control 19: 716–723. <https://doi.org/10.1109/TAC.1974.1100705>
- Aubriot X, Knapp S, Syfert MM, Poczai P and Buerki S (2018) Shedding new light on the origin and spread of the brinjal eggplant (*Solanum melongena* L.) and its wild relatives. Am J Bot 105: 1175–1187. <https://doi.org/10.1002/ajb2.1133>
- Bandara N, Papini A, Mosti S, Brown T, Smith L (2013) A phylogeny of *Onobrychis* and its relationships with allied genera of Hedysareae. Turk Journ of Bot. 37(6): 981–992. <https://doi.org/10.3906/bot-1210-32>
- Bressan E A, Rossi ML, Lee TSG, Figueira A (2014) Extraction of high-quality DNA from ethanol-preserved tropical plant tissues.” BMC research notes 7: 268. <https://doi.org/10.1186/1756-0500-7-268>
- Burger W (1972) Evolutionary trends in the Central American Species of *Piper* (Piperaceae). Brittonia 24: 356–362. <https://doi.org/10.2307/2805498>
- Corpet F (1988) Multiple sequence alignment with hierarchical clustering. Nucleic Acids Res 16:10881–10890. <https://doi.org/10.1093/nar/16.22.10881>
- Corriveau JL, Coleman AW (1988) Rapid screening method to detect potential biparental inheritance of plastid DNA and results for over 200 angiosperm species. Am J Bot 75: 1443–1458. <https://doi.org/10.2307/2444695>
- De Candolle C (1869) Piperaceae. In: De Candolle C, editor. Prodrromus Systematis Naturalis Regni Vegetabilis, Vol. 16. Part 1. Paris, France: Masson, pp. 235–471 (in Latin).
- De Candolle C (1923) Piperacearum clavis analytica. Candollea 1: 65–415 (in Latin).

- García MA, Costea M, Kuzmina M, Stefanović S (2014) Phylogeny, character evolution, and biogeography of *Cuscuta* (dodders; Convolvulaceae) inferred from coding plastid and nuclear sequences. *Am J Bot* 101:670–690. doi:10.3732/ajb.1300449
- Hall TA (1999) BioEdit: A User-Friendly Biological Sequence Alignment Editor and Analysis Program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41: 95–98. https://doi.org/10.14601/Phytopathol\_Mediterr-14998u1.29
- Hurteau GJ, Spivack SD (2002) mRNA-specific reverse transcription-polymerase chain reaction from human tissue extracts. *Anal Biochem* 307:304–315. https://doi.org/10.1016/S0003-2697(02)00058-1
- Isnard S, Prosperi J, Wanke S, Wagner ST, Samain M-S, Trueba S, Frenze L, Neinhuis C, Rowe NP (2012) Growth form evolution in Piperales and its relevance for understanding Angiosperm diversification: an integrative approach combining plant architecture, anatomy, and biomechanics. *Int J Plant Sci* 173: 610–639. https://doi.org/10.1086/665821
- Jaramillo MA and Callejas R (2004) Current perspectives on the classification and phylogenetics of the genus *Piper* L. Pp. 179–198. in *Piper: A model genus for studies of chemistry, ecology, and evolution*, eds. L. A. Dyer and A. N. Palmer. Boston: Kluwer Academic.
- Jaramillo MA, Callejas R, Davidson C, Smith JF, Stevens AC, and Tepe EJ (2008) A Phylogeny of the Tropical Genus *Piper* Using ITS and the Chloroplast Intron psbJ–petA. *Systematic Botany* 33(4): 647–60. doi:10.1600/036364408786500244.
- Jaramillo MA, Manos PS (2001) Phylogeny and patterns of floral diversity in the genus *Piper* (Piperaceae). *Am J Bot* 88:706–716. https://doi.org/10.2307/2657072
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33: 1870–1874. https://doi.org/10.1093/molbev/msw054
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23: 2947–2948. https://doi.org/10.1093/bioinformatics/btm404
- Lewis PO (2001) A likelihood approach to estimating phylogeny from discrete morphological character data. *Syst Biol* 50: 913–925. https://doi.org/10.1080/106351501753462876
- Liu Y-J, Zheng D, Balasubramanian S, Carriero N, Khurana E, Robilotto R, Gerstein MB (2009) Comprehensive analysis of the pseudogenes of glycolytic enzymes in vertebrates: the anomalously high number of GAPDH pseudogenes highlights a recent burst of retrotranspositional activity. *BMC Genomics* 10: 480. https://doi.org/10.1186/1471-2164-10-480
- Loconte H, Stevenson DW (1991) Cladistics of the Magnoliidae. *Cladistics* 7:267–296. https://doi.org/10.1111/j.1096-0031.1991.tb00038.x
- Mallet J (2007) Hybrid speciation. *Nature* 446: 279–283.
- Martines C, Carvalho MR, Marinan S, Jaramillo CA (2015) A Late Cretaceous *Piper* (Piperaceae) from Colombia and diversification patterns for the genus. *Am J Bot* 102(2): 273–289. https://doi.org/10.3732/ajb.1400427
- Maugini E. (1953) Ricerche Cito-Embriologiche su *Piper Medium* Jacq. var. *ceanothifolium* (HBK) Trel. et Yun: (con 14 figure nel testo). *Caryologia* 5(3): 282–287.
- Mosti S., Fiorini G. and Papini A. (2011) Karyological investigations on several species of genus *Rebutia* Sect. *Digitorebutia* (Cactaceae). *Caryologia* 64(3):350–359 https://doi.org/10.1080/00087114.2011.10589802
- Mousavi S H, Hassandokht M R, Choukan R, Sepahvand N, Khosrowchahli M, Papini A. (2013) Cytological study of chromosome and genome composition of Iranian lettuce (*Lactuca sativa* L.) accessions. *Caryologia* 66(1): 41–49. *Digitorebutia* (Cactaceae). *Caryologia* 64(3): 302–308. https://doi.org/10.1080/00087114.2013.780440
- Murray MG and Pitas JW (1996) “Plant DNA from alcohol-preserved samples.” *Plant molecular biology reporter* 14: 261–265. https://doi.org/10.1007/BF02671661
- Nair RR, Sasikumar B, Ravindran PN (1993) Polyploidy in a cultivar of black pepper (*Piper nigrum* L.) and its open pollinated progenies. *Cytologia* 58: 27–31.
- Nylander JAA, (2004) Mr Modeltest, version 1.0b. Department of Systematic Zoology, EBC, Uppsala University, Uppsala, Sweden.
- Palchetti E, Biricolti S, Gori M., Rota Nodari G, Gandolfi N, Papini A (2018) Two new Malagasy species of genus *Piper* L. (Piperaceae): *Piper malgassicum* and *Piper tsarasotrae* and their phylogenetic position. *Turk J of Bot* 42(5): 622–610. doi:10.3906/bot-1712-2
- Papini A, Banci F, Nardi E (2007) Molecular evidence of polyphyletism in the plant genus *Carum* L. (Apiaceae). *Genet Mol Biol* 30: 475–482. https://doi.org/10.1590/S1415-47522007000300029
- Papini A, Simeone MC, Bellarosa R, Spada F, Schirone B (2011) *Quercus macranthera* Fisch. & Mey. Ex Hohen. and *Quercus iberica* M. Bieb.: taxonomic definition and systematic relationships with European oaks inferred from nuclear Internal Transcribed Spacer (ITS) data. *Plant Biosyst* 145: 37–49. https://

- doi.org/10.1080/11263504.2010.502684
- Papini A, Trippanera GB, Maggini F, Filigheddu R, Biondi E (2004) New insights in *Salicornia* L. and allied genera (Chenopodiaceae) inferred from nrDNA sequence data. *Plant Biosystems* 138(3): 215–223 <https://doi.org/10.1080/11263500400006977>
- Petry S (2016) Mechanisms of Mitotic Spindle Assembly. *Annu Rev Biochem* 85:659–683. <https://doi.org/10.1146/annurev-biochem-060815-014528>
- Quijano-Abril MA, Callejas-Posada R, Miranda-Esquivel DR (2006) Areas of endemism and distribution patterns for Neotropical *Piper* species (Piperaceae). *J Biogeogr* 33:1266–1278. <https://doi.org/10.1111/j.1365-2699.2006.01501.x>
- Rambaut A, Drummond A (2010) FigTree v1.3.1. <http://tree.bio.ed.ac.uk/software/figtree>. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, United Kingdom.
- Ronquist F, Teslenko M, Van Der Mark P, Ayacceres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* 61: 539–42. <https://doi.org/10.1093/sysbio/sys029>
- Samuel R (1987) Chromosome Numbers in *Piper*. *Kew Bull* 42: 465. <https://doi.org/10.2307/4109710>
- Samuel R, Morawetz W (1989) Chromosomal evolution within Piperaceae. *Pl. Syst. Evol.* 166: 105–117.
- Selvi F, Papini A, Bigazzi M (2002) Systematics of *Nonea* (Boraginaceae-Boragineae): new insights from phenetic and cladistic analyses. *Taxon* 51: 719–730. <https://doi.org/10.2307/1555025>
- Simeone MC, Grimm GW, Papini A, Vessella F, Cardoni S, Tordoni E, Piredda R, Franc A, Denk T (2016) Plastome data reveal multiple geographic origins of *Quercus* Group Ilex. *PeerJ* 4:e1897 <https://doi.org/10.7717/peerj.1897>
- Simmons MP, Ochoterena H (2000) Gaps as Characters in Sequence-Based Phylogenetic Analyses. *Syst Biol* 49: 369–381. <https://doi.org/10.1093/sysbio/49.2.369>
- Smith JF, Stevens AC, Tepe EJ, Davidson C (2008) Placing the origin of two species-rich genera in the late cretaceous with later species divergence in the tertiary: a phylogenetic, biogeographic and molecular dating analysis of *Piper* and *Peperomia* (Piperaceae). *Plant Syst Evol* 275:9–30. <https://doi.org/10.1007/s00606-008-0056-5>
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Stefanović S, Kuzmina M, and Costea M (2007) Delimitation of major lineages within *Cuscuta* subgenus *Grammica* (dodders; Convolvulaceae) using plastid and nuclear DNA sequences. *Am J Bot* 94: 568–589. <https://doi.org/10.3732/ajb.94.4.568>
- Strand AE, Leebens-Mack J, Milligan BG (1997) Nuclear DNA-based markers for plant evolutionary biology. *Mol Ecol* 6: 113–118. <https://doi.org/10.1046/j.1365-294X.1997.00153.x>
- Sun Y, Li Y, Luo D, Liao DJ (2012) Pseudogenes as weaknesses of ACTB (Actb) and GAPDH (Gapdh) used as reference genes in reverse transcription and polymerase chain reaction. *PLoS ONE* 7:e41659. <https://doi.org/10.1371/journal.pone.0041659>
- Suwanphakdee C, Simpson DA, Hodkinson TR, Chantaranonthai P (2016) Taxonomic notes on the genus *Piper* (Piperaceae). *Nord J Bot* 34:605–618. <https://doi.org/10.1111/njb.01114>
- Swofford DL (1998) PAUP\* 4.1. Phylogenetic Analysis Using Parsimony. Test version. Sunderland (MA): Sinauer Associates.
- Swofford DL (2001) Laboratory of Molecular Systematics Smithsonian Institution. 143
- Taberlet P, Gielly L, Pautou G, Bouvet J (1991) Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Mol Biol* 17: 1105–1109. <https://doi.org/10.1007/BF00037152>
- Tebbs MC (1993) Piperaceae. In: Kubitzki K, Rohwer JG, Bittrich V, editors. Flowering Plants Dicotyledons. The Families and Genera of Vascular Plants, Vol. 2. Berlin, Germany: Springer, pp. 516–520.
- Ulloa Ulloa C, Acevedo-Rodríguez P, Beck S, Belgrano MJ, Bernal R, Berry PE, Brako L, Celis M, Davidse G, Forzza RC, Gradstein SR, Hokche O, Leon B, Leon-Yanez S, Magill RE, Neill DA, Nee M, Raven PH, Stimmel H, Strong MT, Villaseñor JL, Zarucchi JL, Zuloaga FO, Jørgensen PM (2017) An integrated assessment of the vascular plant species of the Americas. *Science* 358:1614–1617. <https://doi.org/10.1126/science.aao0398>
- Vallejo-Marín M, Hiscock SJ (2016) Hybridization and hybrid speciation under global change. *New Phytologist* 211(4): 1170–1187.
- Vanaja T, Neema VP, Mammooty KP, Rajeshkumar R (2008) Development of a promising interspecific hybrid in black pepper (*Piper nigrum* L.) for *Phytophthora* foot rot resistance. *Euphytica* 161:437–445. <https://doi.org/10.1007/s10681-007-9602-4>
- Weil M, Shum Cheong Sing A, Méot JM, Boulanger R, Bohuon P (2017) Impact of blanching, sweating and drying operations on pungency, aroma and color of *Piper borbonense*. *Food Chem* 219:274–281. <https://doi.org/10.1016/j.foodchem.2016.09.144>

- Wühr M, Chen Y, Dumont S, Groen AC, Needleman DJ, Salic A, Mitchison TJ (2008) Evidence for an Upper Limit to Mitotic Spindle Length. *Curr Biol* 18:1256–1261. <https://doi.org/10.1016/j.cub.2008.07.092>
- Young ND, Healy J (2003) GapCoder automates the use of indel characters in phylogenetic analysis. *BMC Bioinformatics* 4: 6 <https://doi.org/10.1186/1471-2105-4-6>