



Activation of carbonic anhydrase isoforms involved in modulation of emotional memory and cognitive disorders with histamine agonists, antagonists and derivatives

Gustavo Provensi, Alessio Nocentini, Maria Beatrice Passani, Patrizio Blandina & Claudiu T. Supuran

To cite this article: Gustavo Provensi, Alessio Nocentini, Maria Beatrice Passani, Patrizio Blandina & Claudiu T. Supuran (2021) Activation of carbonic anhydrase isoforms involved in modulation of emotional memory and cognitive disorders with histamine agonists, antagonists and derivatives, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 36:1, 719-726, DOI: [10.1080/14756366.2021.1891051](https://doi.org/10.1080/14756366.2021.1891051)

To link to this article: <https://doi.org/10.1080/14756366.2021.1891051>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 02 Mar 2021.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

RESEARCH PAPER



Activation of carbonic anhydrase isoforms involved in modulation of emotional memory and cognitive disorders with histamine agonists, antagonists and derivatives

Gustavo Provensi^a , Alessio Nocentini^a , Maria Beatrice Passani^b, Patrizio Blandina^a and Claudiu T. Supuran^a 

^aDepartment of NEUROFARBA, University of Florence, Section of Pharmacology and Toxicology, Firenze, Italy; ^bDepartment of Health Science, University of Florence, Section of Clinical Pharmacology and Oncology, Firenze, Italy

ABSTRACT

Carbonic anhydrases (CAs, EC 4.2.1.1) activators were shown to be involved in memory enhancement and learning in animal models of cognition. Here we investigated the CA activating effects of a large series of histamine based compounds, including histamine receptors (H₁R – H₄R) agonists, antagonists and other derivatives of this autacoid. CA activators may be thus useful for improving cognition as well as in diverse therapeutic areas (phobias, obsessive-compulsive disorder, generalised anxiety, post-traumatic stress disorders), for which activation of this enzyme was recently shown to be involved.

ARTICLE HISTORY


Received 30 December 2020
Revised 8 February 2021
Accepted 10 February 2021

KEYWORDS

Carbonic anhydrase; activators; cognition-related disorders; histamine receptors; agonists/antagonists

1. Introduction

CO₂ is generated in most metabolic processes, being one of the simplest molecules involved in crucial physiologic processes in all life kingdoms^{1,2}. The carbonic anhydrases (CAs, EC 4.2.1.1) are metalloenzymes which catalyse its interconversion to bicarbonate (Equation (1))^{3–6}, generating also a proton, and thus a pH disequilibrium, which is used in most biological systems as a readily available buffering system⁷.



The reaction also occurs without a catalyst, but at physiological pH values it is exceedingly slow for meeting metabolic needs, as CO₂ is a poorly water-soluble gas, which can also damage cellular components (e.g. membranes, mitochondria, etc.)^{6,7}. On the other side, its conversion to water-soluble ions (bicarbonate and protons) counteracts this effect, and although interfering with the pH balance, is used to control homeostasis and metabolism, making CAs crucial enzymes in many physiological and pathological conditions^{3–7}. In fact, in vertebrates at least 16 CA isoforms belonging to the α -CA genetic family are known, whereas in other organisms all over the phylogenetic tree at least seven other CA families were described so far, the β -, γ -, δ -, ζ -, η -, θ - and ι -CAs^{8–14}. In humans 15 CAs are expressed, 12 of which are catalytically active: the cytosolic CA I-III, VII and XIII, the membrane-bound CA IV, the mitochondrial CA VA and VB, the secreted (in saliva and tears) CA VI, and the transmembrane CA IX, XII and XIV (the acatalytic forms are CA VIII, X and XI)^{4,15–19}. Many of these enzymes are drug targets, as their inhibitors show pharmacological applications for drugs treating edoema, glaucoma, obesity, epilepsy and tumours^{4–6}.

The human central nervous system (CNS), as well as the choroid plexus, contains a multitude of CA isoforms, although their particular functions are not yet completely understood¹⁷. We will consider here mainly the CAs present in CNS, as the compounds investigated here for modulating their activity (i.e. the CA activators – CAAs) may also have interesting applications in therapy, which started to be considered only recently^{20–23}. The nervous system CA isoforms comprise: the cytosolic CA I (expressed in the motor neurons in the spinal cord), CA II (present in the choroid plexus, oligodendrocytes, myelinated tracts, astrocytes and myelin sheaths); CA III (in the choroid plexus), the membrane-associated CA IV (located on the luminal surface of cerebral capillaries and associated with the blood-brain barrier, being present also in the cortex, hippocampus and thalamus). The mitochondrial CA VA is expressed in astrocytes and in neurons, whereas CA VB seems to be absent in the SNC¹⁷. CA VII and VIII are present in high levels throughout the cortex, hippocampus and thalamus, although CA VIII is acatalytic, whereas CA VII shows a good enzymatic activity with CO₂/bicarbonate as substrates⁴. The acatalytic CA X is expressed in the myelin sheath, whereas CA XI (also acatalytic) is present in the neural cell body and astrocytes¹⁷. CA IX and CA XII are transmembrane proteins overexpressed in many neurologic cancers^{18,19}, whereas CA XIII seems not to be present in the brain. CA XIV is expressed in nuclei and nerve tracts associated with pontine, medullary and hippocampal functions being also located on the plasma membrane of some neurons and on axons of mammalian brain¹⁷.


The most investigated CAAs are the amino acids, the biogenic amines (histamine, serotonin, catecholamines and their derivatives), and to some extent also the oligopeptides or small

CONTACT Alessio Nocentini  alessio.nocentini@unifi.it; Claudiu T. Supuran  claudiu.supuran@unifi.it  Department of NEUROFARBA, University of Florence, Section of Pharmaceutical and Nutraceutical Sciences, Polo Scientifico, Via U. Schiff 6, Firenze, Sesto Fiorentino 50019, Italy

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

proteins, although these activators were less investigated²⁰. The CAAs were demonstrated to participate in the catalytic cycle of the enzyme, forming enzyme-activator complexes, as described in Equation (2):²⁰



The activator molecule forms a complex with the enzyme, binding in an active site region distinct of that of the classical CA inhibitors^{24,25}, which generally bind to the metal ion⁴⁻⁶. The activator molecule must incorporate proton shuttling moieties, which take part to the rate-determining step of the catalytic cycle, i.e. the transfer of protons from the zinc-coordinated water molecule to the external reaction medium, with formation of the nucleophilic, zinc hydroxide species of the enzyme²⁰. In the wild type enzyme, this proton shuttling is achieved by residue His64 (in many CA isoforms), found within the middle of the active site cleft, and which possess the imidazole moiety able to transfer protons in the pH range of 6–8²⁰⁻²⁴. His64 was shown to possess two conformations: the *in* one, orientated towards the bottom of the active site, and the *out* one, orientated towards the external part of the active site, favouring thus the proton wiring^{20,24}. In such processes, within the enzyme-activator complexes, the proton transfer becomes intramolecular, being more efficient compared to the intermolecular transfer to buffer molecules (which are not bound within the enzyme cavity)²⁰. X-ray crystallography has been performed on several other hCA I/II – activator complexes, among which those with histamine, L- and D-His, L- and D-Phe, D-Trp, L-adrenaline as well as pyridinium derivatives of histamine^{20,23-26}. A schematic representation of the activators bound to CA is shown in Figure 1.

Thus, histamine was the main compound used to obtain new CAAs²⁷, but many of its rather simple derivatives as well as drugs belonging to the histamine receptors (H₁R, H₂R, H₃R and H₄R) agonists/antagonists, were not yet been investigated for their potential activating effects. Here we report the first such study, including in our investigations 28 such derivatives which have been assayed as activators of four pharmacologically significant isoforms, hCA I, II and VII (cytosolic isoforms) and hCA IV (membrane-anchored enzyme).

2. Materials and methods

2.1. Chemistry

Histamine **1** and compounds **2–30** were commercially available, highest purity reagents from Sigma-Aldrich, Milan Italy.

2.2. Carbonic anhydrase activation

A stopped-flow method²⁸ has been used for assaying the CA catalysed CO₂ hydration activity with Phenol red as indicator, working at the absorbance maximum of 557 nm, following the initial rates of the CA-catalysed CO₂ hydration reaction for 10–100 s. For each activator, at least six traces of the initial 5–10% of the reaction have been used for determining the initial velocity. The uncatalyzed rates were determined in the same manner and subtracted from the total observed rates. Stock solutions of activator (0.1 mM) were prepared in distilled-deionized water and dilutions up to 0.1 nM were done thereafter with the assay buffer. The activation constant (K_A), defined similarly with the inhibition constant (K_i), was obtained by considering the classical Michaelis–Menten




Figure 1. CA activation mechanisms. Activators bind in the middle of the active site and contain a proton shuttle moiety (PSM) of the amine, imidazole or carboxylate type with an appropriate pK_a for the proton transfer processes, usually in the range of 6–8.

Equation (3), which has been fitted by nonlinear least squares by using PRISM 3:

$$v = v_{\max} / \{ 1 + K_M / [S] (1 + [A]_f / K_A) \} \quad (3)$$

where [A]_f is the free concentration of activator.

Working at substrate concentrations considerably lower than K_M ([S] ≪ K_M), and considering that [A]_f can be represented in the form of the total concentration of the enzyme ([E]_t) and activator ([A]_t), the obtained competitive steady-state equation for determining the activation constant is given by Equation (4):

$$v = v_0 K_A / \left\{ K_A + \left([A]_t - 0.5 \{ ([A]_t + [E]_t + K_A) - \sqrt{([A]_t + [E]_t + K_A)^2 - 4[A]_t[E]_t} \} \right) \right\} \quad (4)$$

where v₀ represents the initial velocity of the enzyme-catalysed reaction in the absence of an activator²⁹⁻³². Enzyme concentrations in the assay system were in the range of 6.5–12.0 nM.

3. Results and discussion

As mentioned above, histamine **1** (Figure 2) was one of the first CAAs to be investigated in detail²⁴, but except for histidine (L- and D-enantiomers), other histamine derivatives were not yet assayed for their potential CA activating effects.

Considering the relatively large number of histamine receptors (H₁R–H₄R) as well as the huge number of agonists/antagonists developed for the management of various disorders, among which allergies, gastritis and gastric ulcers, narcolepsy, acute unilateral vestibulopathy, and atopic dermatitis^{33,34}, there is a large number of compounds incorporating fragments of the histamine chemotype as well as a wealth of structural modifications which mimic this autacoid.

Some of these compounds, possessing structures **2–30** (Figure 1), were included in our study for investigating their possible CA activating effects against four pharmacologically relevant human isoforms, hCA I, II, IV and VII. The compounds were numbered according to their similarity to the lead histamine **1** and are: the H₁R agonist 2-(2-aminoethyl)thiazole **6**; the H₂R agonists impromidine **16** and nordimaprit **19**; the H₃R agonists Nπ-methylhistamine **3**, α-methylhistamine **4**, methimipip **8**, proxyfan **9**, imetit **14**, VUF16839 **23**; the H₁R antagonists pyrillamine **24**, loratadine **29**;




Figure 2. Histamine 1 and derivatives 2–30 acting as histamine receptors agonists/antagonists (for reviews see refs.^{33,34}) investigated here as CAAs.

the H₂R antagonists metiamide **12**, cimetidine **13**, ranitidine **17**, tiotidine **18**, zolantidine **20**; the H₃R antagonists ciproxifan **10**, clobenpropit **15**, ABT239 **22**, GSK189254A **28**, GSK334429B **30**; the H₄R antagonists JNJ39758979 **25**, JNJ7777120 **26**, A940894 **27**; the mixed modulators of the histaminergic system N τ -methylhistamine **2**, 4-methylhistamine **5**, 1-methylhistidine **7**, burimamide **11**, beta-histidine **21**.

In some of these compounds, such as the methyl-histamines **2–5**, the thiazolyl derivative **6** or τ -methyl-His **7**, the histamine chemotype is readily observable, whereas the remaining compounds incorporate more drastic changes of the basic structure,

but all of them possess moieties which can in principle shuttle protons in the pH range of 6–8 which, as mentioned earlier²⁰, lead to CA activation.

The following structure-activity relationship (SAR) can be worked out from the data reported in [Table 1](#) for activation of the four isoforms hCA I, II, IV and VII:

- Compounds **17**, **20**, **22**, **25–30** did not induce any activation of the tested CA isoforms (K_As >100 μ M). Consistently, these derivatives do not possess the histamine chemotype in their structures and/or other moieties that clearly make CA

Table 1. hCA I, II, IV and VII activation with compounds 2–30 (Figure 2) by a stopped-flow CO₂ hydrase assay.²⁸ Histamine 1 used as standard.

Compound	K _A (μM) ^a				Data		Ref
	hCA I	hCA II	hCA IV	hCA VII	BBB crossing	Central action	
1	2.1	125	25.3	37.5	–	+	35
2	0.11	8.91	3.21	2.07			
3	3.1	0.43	7.6	0.23			
4	0.12	0.082	2.91	1.25		+	36
5	0.36	5.4	5.13	0.39		+	37
6	0.87	7.45	1.02	0.7			
7	0.052	0.57	13.9	0.19			
8	3.16	5.24	4.66	0.12		+	38
9	3.15	7.66	8.01	0.52	+	+	39
10	4.29	9.9	8.12	0.11	+	+	40
11	0.88	8.39	9.07	0.43		+	41
12	0.98	8.75	9.62	1.01		+	42
13	8.79	6.3	8.54	0.59		+	43
14	4.25	8.31	8.05	1.00		+	44
15	>100	2.1	6.59	5.31	+	+	45
16	0.72	2.14	3.3	0.10			46
17	>100	>100	>100	>100		+	47
18	>100	>100	32.3	45.5		+	48
19	1.36	6.93	9.08	5.21			
20	>100	>100	>100	>100	+	+	49
21	>100	13.5	9.9	7.05	+	+	50
22	>100	>100	>100	>100	+	+	51
23	>100	9.82	15.9	>100		+	52
24	5.23	9.62	6.78	2.05	+	+	53
25	>100	>100	>100	>100			
26	>100	>100	>100	>100	+	+	54
27	>100	>100	>100	>100			
28	>100	>100	>100	>100	+	+	55
29	>100	>100	>100	>100	+	+	56
30	>100	>100	>100	5.06	+	+	57

^aFrom three different assays (errors within ± 10% of the reported values). – means no BBB crossing; + means that there are evidences of BBB crossing and central action; no sign means that no literature data are available.

activation possible. As a unique exception, compound 30 reported a 5.06 μM selective activation of hCA VII. It should be stressed that other derivatives, such as 18, 19, 21, 23 and 24, do not directly include imidazole-like scaffolds in their chemical structure, but showed however to possess significant CA activation profiles in a low micromolar range (KAs between 1.36 and 45.5 μM) and are thus included in the SAR discussion. These compounds possess however protonatable moieties of the secondary amine or guanidine type in their molecule, which like the imidazole may shuttle protons and thus act as CAAs.

- ii. The cytosolic and ubiquitous hCA isoforms I and II were quite efficiently activated by most active derivatives (that are 2–16, 18, 19, 21, 23 and 24), in a low micromolar to submicromolar range. Intriguingly, derivatives 15, 21 and 23 did not produce any hCA I activation up to 100 μM, whereas 18 did not activate neither hCA I nor hCA II. The methylation of histamine 1 at position N τ (2), α (4), 4 (5) and the imidazole/thiazole swap (6) increased up to one order of magnitude the hCA I activation profile (from 2.1 to 0.11 μM). The presence of an extra proton transfer group (COOH) as in 7 (N τ -methyl-histidine) further improved 2-fold the KA (52 vs 110 nM) against hCA I compared to compound 2. The N τ -methylation of histamine (3, KA of 3.1 μM) slightly decreased the activation efficacy of the molecule, situations also encountered for the inclusion of the aliphatic amine into a cycle (8, KA of 3.16 μM) or the amine/ether swap (as in 9 and 10, KAs of 3.15 and 4.29 μM). Substituting the amine group with N-linked thioureas as in 11 and 12 improved 2-fold the KAs towards hCA I with respect to histamine.

Among the remaining derivatives, only the bis-imidazole 16 reported an improved KA over the lead towards this isoform (KA of 0.72 μM). In fact, the presence of S-linked thioureas worsened the activation efficacy 2- (14, KA of 4.25 μM) or 4-fold (13, KA of 8.79 μM) with respect to the lead histamine. Oddly, the S-linked thiourea 19 showed a 2-fold improved KA compared to histamine, although bearing a dimethylamino group in place of the imidazole ring. As an exception, the N-rich compound 24 interestingly activated hCA I just two times less than histamine, in spite of a completely diverse structure. On the contrary, all histamine derivatives here reported showed a superior hCA II activation efficacy with respect to the lead (KA > 100 μM). Among mostly low micromolar CAAs (KAs in the range 2.1–13.5 μM), derivatives 3, 4 and 7 stood out as the most potent hCA II modulators of the study (KAs in the range 82 nM–0.57 μM). In particular, the α -methylation of histamine, as in 4, induced the largest increase of efficacy, up to a KA of 82 nM, when compared to the ineffective (as CAA) lead molecule.

- iii. No submicromolar KA values was measured for 1–30 as hCA IV activators. Indeed, all KAs are in a rather flat low micromolar range (KAs in the range 1.02–13.9 μM), making this membrane-associated isozyme the less activated one by the compounds tested in this work. Interestingly, with the exception of compound 18, all derivatives were more efficient CAAs than the lead histamine, which showed a KA of 25.3 μM. Of note, the imidazole/thiazole swap led to the most effective activation increase with respect to the lead, with a KA of 1.02 μM in case of derivative 6.
- iv. The other cytosolic isoform investigated here, hCA VII, was the most effectively activated one by the compounds investigated in this study. Indeed, a wide subset of KAs were detected in a submicromolar range (from 0.10 to 45.5 μM) for some of these derivatives. All of them showed much better activation profile than the reference compound histamine (KA of 37.5 μM) towards hCA VII. Contrariwise to hCA I and II, the most efficient CAAs were not detected among the methylhistamine derivatives 2–5: N τ -methylhistidine 7, N-methylpiperidine 8 and the aryl ether 10, showed KAs ranging between 110 and 190 nM. The bis-imidazole 16 stood out as the most effective hCA VII activator of the study with a KA of 100 nM. Intriguingly, compound 23 did not activate hCA VII below 100 μM, whereas derivative 30, previously classified among the inactive compounds for the other CA isoforms, weakly activated this CNS-associated CA (KA of 5.06 μM). In fact, this isoform is one of the most widely spread in the brain, probably being involved in crucial metabolic/pH regulation processes, while it is not expressed in other tissues. It is thus relevant that a rather wide set of compounds was detected here (8, 9, 10, 13, 16), which showed a promising isoform selectivity towards hCA VII over the ubiquitous CAAs (up to 100-fold over hCA II).

Table 2 also include the literature references regarding the compounds ability to cross the BBB, which presumably should also lead to brain CA activating effects, as well as evidences for their action at central level.

4. Conclusions and future perspectives

In the present study, we investigated the CA activating effects of a series of histamine receptors agonists/antagonists (compounds 2–30 in Figure 2) towards four hCA isoforms expressed in human

brain, that are CA I, II, IV and VII. Though all derivatives possess moieties which can in principle shuttle protons in the pH range of 6–8, a consistent subset of them (**17**, **20**, **22**, **25–30**), not having the histamine chemotype in their structure, did not report any activation effect for the tested CA isoforms ($K_{As} > 100 \mu\text{M}$). hCA I and II were effectively activated by methylhistamine derivatives (**2–7**), whereas more intriguing SAR were observed for hCA VII with more lipophilic groups (as in **6**, **8** or **10**) promoting greater and more selective isoform activation. Of note, a subset of selective hCA VII activators was identified, that could serve to drive the identification and optimisation of new brain specific CAAs. We are currently witnessing a second youth period for CAAs, because of innovative pharmacological studies spurring researchers to take into account these lately neglected agents for their potential clinical relevance in the treatment of emotional memory disorders, including the improvement of the clinical efficacy of exposure-based treatments of obsessive-compulsive disorders, phobias, generalised anxiety, and post-traumatic stress disorder. The here gathered data might also provide more insights on the pharmacodynamics of therapeutically used histamine modulators, whose therapeutic action and/or side effects could be related to polypharmacology. Overall, this work might bring new lights on the intricate relationship between CA activation and brain physiology.

Acknowledgements

This paper is dedicated to the 90th anniversary of Professor Alexandru T. Balaban, mentor and friend to some of us.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by the Italian Ministry for University and Research (MIUR), grant PRIN: prot. 2017XYBP2R (to CTS).

ORCID

Gustavo Provensi  <http://orcid.org/0000-0002-6182-6848>
Alessio Nocentini  <http://orcid.org/0000-0003-3342-702X>
Claudiu T. Supuran  <http://orcid.org/0000-0003-4262-0323>

References

- (a) Arias-Hidalgo M, Yuan Q, Carta F, et al. CO_2 permeability of rat hepatocytes and relation of CO_2 permeability to CO_2 production. *Cell Physiol Biochem* 2018;46:1198–208.(b) Arias-Hidalgo M, Hegemann J, Tsiavalariis G, et al. CO_2 and HCO_3^- permeability of the rat liver mitochondrial membrane. *Cell Physiol Biochem* 2016;39:2014–24.
- (a) Iteel F, Al-Samir S, Öberg F, et al. CO_2 permeability of cell membranes is regulated by membrane cholesterol and protein gas channels. *Faseb J* 2012;26:5182–91. (b) Endeward V, Musa-Aziz R, Cooper GJ, et al. Evidence that aquaporin 1 is a major pathway for CO_2 transport across the human erythrocyte membrane. *Faseb J* 2006;20:1974–81.
- (a) Supuran CT. Structure and function of carbonic anhydrases. *Biochem J* 2016;473:2023–32. (b) Supuran CT. Advances in structure-based drug discovery of carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2017;12:61–88.(c) Supuran CT. How many carbonic anhydrase inhibition mechanisms exist? *J Enzyme Inhib Med Chem* 2016;31:345–60.(d) Supuran CT. Exploring the multiple binding modes of inhibitors to carbonic anhydrases for novel drug discovery. *Expert Opin Drug Discov* 2020;15:671–86.
- (a) Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nat Rev Drug Discov* 2008;7:168–81.(b) Alterio V, Di Fiore A, D'Ambrosio K, et al. Multiple binding modes of inhibitors to carbonic anhydrases: how to design specific drugs targeting 15 different isoforms? *Chem Rev* 2012;112:4421–68.(c) De Simone G, Alterio V, Supuran CT. Exploiting the hydrophobic and hydrophilic binding sites for designing carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2013;8:793–810.
- (a) Scozzafava A, Supuran CT, Carta F. Antiobesity carbonic anhydrase inhibitors: a literature and patent review. *Expert Opin Ther Pat* 2013;23:725–35. (b) Supuran CT. Applications of carbonic anhydrases inhibitors in renal and central nervous system diseases. *Expert Opin Ther Pat* 2018;28:713–21.(c) Supuran CT. The management of glaucoma and macular degeneration. *Expert Opin Ther Pat* 2019;29:745–7.(d) Fabrizi F, Mincione F, Somma T, et al. A new approach to antiglaucoma drugs: carbonic anhydrase inhibitors with or without NO donating moieties. Mechanism of action and preliminary pharmacology. *J Enzyme Inhib Med Chem* 2012;27:138–47.
- (a) Masini E, Carta F, Scozzafava A, Supuran CT. Antiglaucoma carbonic anhydrase inhibitors: a patent review. *Expert Opin Ther Pat* 2013;23:705–16.(b) Supuran CT, Altamimi ASA, Carta F. Carbonic anhydrase inhibition and the management of glaucoma: a literature and patent review 2013–2019. *Expert Opin Ther Pat* 2019;29:781–92.(c) De Simone G, Supuran CT. (In)organic anions as carbonic anhydrase inhibitors. *J Inorg Biochem* 2012;111:117–29.(d) Supuran CT. Carbon- versus sulphur-based zinc binding groups for carbonic anhydrase inhibitors? *J Enzyme Inhib Med Chem* 2018;33:485–95.
- (a) Lee SH, McIntyre D, Honess D, et al. Carbonic anhydrase IX is a pH-stat that sets an acidic tumour extracellular pH in vivo. *Br J Cancer* 2018;119:622–30.(b) Ruusuvaari E, Kaila K. Carbonic anhydrases and brain pH in the control of neuronal excitability. *Subcell Biochem* 2014;75:271–90. (c) Chesler M, Kaila K. Modulation of pH by neuronal activity. *Trends Neurosci* 1992;15:396–402.(d) Occhipinti R, Boron WF. Role of carbonic anhydrases and inhibitors in acid-base physiology: insights from mathematical modeling. *Int J Mol Sci* 2019;20:3841.
- (a) Mishra CB, Tiwari M, Supuran CT. Progress in the development of human carbonic anhydrase inhibitors and their pharmacological applications: where are we today? *Med Res Rev* 2020;40:2485–565.(b) Capasso C, Supuran CT. An overview of the alpha-, beta- and gamma-carbonic anhydrases from Bacteria: can bacterial carbonic anhydrases shed new light on evolution of bacteria? *J Enzyme Inhib Med Chem* 2015;30:325–32.(c) Nocentini A, Supuran CT. Advances in the structural annotation of human carbonic anhydrases and impact on future drug discovery. *Expert Opin Drug Discov* 2019;14:1175–97.
- (a) Capasso C, Supuran CT. Inhibition of bacterial carbonic anhydrases as a novel approach to escape drug resistance. *Curr Top Med Chem* 2017;17:1237–48. (b) Capasso C, Supuran CT. Bacterial, fungal and protozoan carbonic

- anhydrases as drug targets. *Expert Opin Ther Targets* 2015; 19:1689–704.(c) Ozensoy Guler O, Supuran CT, Capasso C. Carbonic anhydrase IX as a novel candidate in liquid biopsy. *J Enzyme Inhib Med Chem* 2020;35:255–60.(d) Supuran CT, Capasso C. Biomedical applications of prokaryotic carbonic anhydrases. *Expert Opin Ther Pat* 2018;28:745–54.
10. (a) Kaur J, Cao X, Abutaleb NS, et al. Optimization of acetazolamide-based scaffold as potent inhibitors of vancomycin-resistant enterococcus. *J Med Chem* 2020;63:9540–62.(b) Supuran CT, Capasso C. Antibacterial carbonic anhydrase inhibitors: an update on the recent literature. *Expert Opin Ther Pat* 2020;30:963–82.
 11. (a) Del Prete S, Bua S, Supuran CT, Capasso C. Escherichia coli γ -carbonic anhydrase: characterisation and effects of simple aromatic/heterocyclic sulphonamide inhibitors. *J Enzyme Inhib Med Chem* 2020;35:1545–54.(b) Del Prete S, De Luca V, Bua S, et al. The effect of substituted benzene-sulfonamides and clinically licensed drugs on the catalytic activity of CynT2, a carbonic anhydrase crucial for escherichia coli life cycle. *Int J Mol Sci* 2020;21:4175.(c) Angeli A, Ferraroni M, Pinteala M, et al. Crystal structure of a tetrameric type II β -carbonic anhydrase from the pathogenic bacterium *Burkholderia pseudomallei*. *Molecules* 2020;25:2269.
 12. (a) Del Prete S, Nocentini A, Supuran CT, Capasso C. Bacterial ι -carbonic anhydrase: a new active class of carbonic anhydrase identified in the genome of the Gram-negative bacterium *Burkholderia territorii*. *J Enzyme Inhib Med Chem* 2020;35:1060–8.(b) Angeli A, Pinteala M, Maier SS, et al. Inhibition of α -, β -, γ -, δ -, ζ - and η -class carbonic anhydrases from bacteria, fungi, algae, diatoms and protozoans with famotidine. *J Enzyme Inhib Med Chem* 2019;34:644–50.
 13. (a) Angeli A, Etxebeste-Mitxelorena M, Sanmartín C, et al. Tellurides bearing sulfonamides as novel inhibitors of leishmanial carbonic anhydrase with potent antileishmanial activity. *J Med Chem* 2020;63:4306–14.(b) Nocentini A, Osman SM, Almeida IA, et al. Appraisal of anti-protozoan activity of nitroaromatic benzenesulfonamides inhibiting carbonic anhydrases from *Trypanosoma cruzi* and *Leishmania donovani*. *J Enzyme Inhib Med Chem* 2019;34:1164–71. (c) da Silva Cardoso V, Vermelho AB, Ricci Junior E, et al. Antileishmanial activity of sulphonamide nanoemulsions targeting the β -carbonic anhydrase from *Leishmania* species. *J Enzyme Inhib Med Chem* 2018;33:850–7.
 14. (a) Vermelho AB, da Silva Cardoso V, Ricci Junior E, et al. Nanoemulsions of sulfonamide carbonic anhydrase inhibitors strongly inhibit the growth of *Trypanosoma cruzi*. *J Enzyme Inhib Med Chem* 2018;33:139–46.(b) Vermelho AB, Capaci GR, Rodrigues IA, et al. Carbonic anhydrases from *Trypanosoma* and *Leishmania* as anti-protozoan drug targets. *Bioorg Med Chem* 2017;25:1543–55.(c) Nocentini A, Cadoni R, Dumy P, et al. Carbonic anhydrases from *Trypanosoma cruzi* and *Leishmania donovani* chagasi are inhibited by benzoxaboroles. *J Enzyme Inhib Med Chem* 2018;33:286–9.
 15. (a) Ozensoy Guler O, Capasso C, Supuran CT. A magnificent enzyme superfamily: carbonic anhydrases, their purification and characterization. *J Enzyme Inhib Med Chem* 2016;31:689–94. (b) Supuran CT. Carbonic anhydrases and metabolism. *Metabolites* 2018;8:25.(c) Supuran CT, Nicolae A, Popescu A. Carbonic anhydrase inhibitors. Part 35. Synthesis of Schiff bases derived from sulfanilamide and aromatic aldehydes: the first inhibitors with equally high affinity towards cytosolic and membrane-bound isozymes. *Eur J Med Chem* 1996;31:431–8.
 16. (a) Supuran CT. Carbonic anhydrase inhibition and the management of neuropathic pain. *Expert Rev Neurother* 2016; 16:961–8. (b) Supuran CT. Acetazolamide for the treatment of idiopathic intracranial hypertension. *Expert Rev Neurother* 2015;15:851–6.
 17. (a) Halmi P, Parkkila S, Honkaniemi J. Expression of carbonic anhydrases II, IV, VII, VIII and XII in rat brain after kainic acid induced status epilepticus. *Neurochem Int* 2006;48:24–30. (b) Thiry A, Dogne JM, Supuran CT, Masereel B. Carbonic anhydrase inhibitors as anticonvulsant agents. *Curr Top Med Chem* 2007;7:855–64.(c) Patrikainen MS, Tolvanen MEE, Aspatwar A, et al. Identification and characterization of a novel zebrafish (*Danio rerio*) pentraxin-carbonic anhydrase. *PeerJ* 2017;5:e4128.
 18. (a) Supuran CT, Alterio V, Di Fiore A, et al. Inhibition of carbonic anhydrase IX targets primary tumors, metastases, and cancer stem cells: three for the price of one. *Med Res Rev* 2018;38:1799–836. (b) Supuran CT. Carbonic anhydrase inhibitors as emerging agents for the treatment and imaging of hypoxic tumors. *Expert Opin Investig Drugs* 2018;27:963–70.(c) Nocentini A, Supuran CT. Carbonic anhydrase inhibitors as antitumor/antimetastatic agents: a patent review (2008–2018). *Expert Opin Ther Pat* 2018;28:729–40.
 19. (a) Angeli A, Carta F, Nocentini A, et al. Carbonic anhydrase inhibitors targeting metabolism and tumor microenvironment. *Metabolites* 2020;10:412.(b) McDonald PC, Chia S, Bedard PL, et al. A phase 1 study of SLC-0111, a novel inhibitor of carbonic anhydrase IX, in patients with advanced solid tumors. *Am J Clin Oncol* 2020;43:484–90.(c) Neri D, Supuran CT. Interfering with pH regulation in tumours as a therapeutic strategy. *Nat Rev Drug Discov* 2011;10:767–77.
 20. (a) Supuran CT. Carbonic anhydrase activators. *Future Med Chem* 2018;10:561–73.(b) Blandina P, Provensi G, Passsani MB, et al. Carbonic anhydrase modulation of emotional memory. Implications for the treatment of cognitive disorders. *J Enzyme Inhib Med Chem* 2020;35:1206–14.
 21. Canto de Souza L, Provensi G, Vullo D, et al. Carbonic anhydrase activation enhances object recognition memory in mice through phosphorylation of the extracellular signal-regulated kinase in the cortex and the hippocampus. *Neuropharmacology* 2017;118:148–56.
 22. Schmidt SD, Costa A, Rani B, et al. The role of carbonic anhydrases in extinction of contextual fear memory. *Proc Natl Acad Sci USA* 2020;117:16000–8.
 23. (a) Izquierdo I, Furini CR, Myskiw JC. Fear memory. *Physiol Rev* 2016;96:695–750.(b) Provensi G, Costa A, Izquierdo I, et al. Brain histamine modulates recognition memory: possible implications in major cognitive disorders. *Br J Pharmacol* 2020;177:539–56.(c) Provensi G, Passani MB, Costa A, et al. Neuronal histamine and the memory of emotionally salient events. *Br J Pharmacol* 2020;177:557–69.(d) Benetti F, Furini CR, de Carvalho Myskiw J, et al. Histamine in the basolateral amygdala promotes inhibitory avoidance learning independently of hippocampus. *Proc Natl Acad Sci USA* 2015;112:E2536–42.
 24. (a) Briganti F, Mangani S, Orioli P, et al. Carbonic anhydrase activators: X-ray crystallographic and spectroscopic investigations for the interaction of isozymes I and II with histamine. *Biochemistry* 1997;36:10384–92.(b) Temperini C,

- Scozzafava A, Vullo D, Supuran CT. Carbonic anhydrase activators. Activation of isozymes I, II, IV, VA, VII, and XIV with L- and D-histidine and crystallographic analysis of their adducts with isoform II: engineering proton-transfer processes within the active site of an enzyme. *Chemistry* 2006;12:7057–66.(c) Temperini C, Scozzafava A, Supuran CT. Carbonic anhydrase activators: the first X-ray crystallographic study of an adduct of isoform I. *Bioorg Med Chem Lett* 2006;16:5152–6.
25. (a) Temperini C, Scozzafava A, Supuran CT. Carbonic anhydrase activation and the drug design. *Curr Pharm Des* 2008;14:708–15.(b) Temperini C, Innocenti A, Scozzafava A, et al. Carbonic anhydrase activators: L-Adrenaline plugs the active site entrance of isozyme II, activating better isoforms I, IV, VA, VII, and XIV. *Bioorg Med Chem Lett* 2007;17:628–35.(c) Temperini C, Innocenti A, Scozzafava A, Supuran CT. Carbonic anhydrase activators: kinetic and X-ray crystallographic study for the interaction of D- and L-tryptophan with the mammalian isoforms I–XIV. *Bioorg Med Chem* 2008;16:8373–8.(d) Temperini C, Scozzafava A, Vullo D, Supuran CT. Carbonic anhydrase activators. Activation of isoforms I, II, IV, VA, VII, and XIV with L- and D-phenylalanine and crystallographic analysis of their adducts with isozyme II: stereospecific recognition within the active site of an enzyme and its consequences for the drug design. *J Med Chem* 2006;49:3019–27.
 26. (a) Dave K, Ilies MA, Scozzafava A, et al. An inhibitor-like binding mode of a carbonic anhydrase activator within the active site of isoform II. *Bioorg Med Chem Lett* 2011;21:2764–8. (b) Bhatt A, Mondal UK, Supuran CT, et al. Crystal structure of carbonic anhydrase II in complex with an activating ligand: implications in neuronal function. *Mol Neurobiol* 2018;55:7431–7.
 27. (a) Saada MC, Montero JL, Vullo D, et al. Carbonic anhydrase activators: gold nanoparticles coated with derivatized histamine, histidine, and carnosine show enhanced activatory effects on several mammalian isoforms. *J Med Chem* 2011;54:1170–7. (b) Scozzafava A, Supuran CT. Carbonic anhydrase activators – part 21. Novel activators of isozymes I, II and IV incorporating carboxamido and ureido histamine moieties. *Eur J Med Chem* 2000;35:31–9.(c) Scozzafava A, Supuran CT. Carbonic anhydrase activators: high affinity isozymes I, II, and IV activators, incorporating a beta-alanyl-histidine scaffold. *J Med Chem* 2002;45:284–91.(d) Ilies M, Banciu MD, Ilies MA, et al. Carbonic anhydrase activators: design of high affinity isozymes I, II, and IV activators, incorporating tri-/tetrasubstituted-pyridinium-azole moieties. *J Med Chem* 2002;45:504–10.
 28. Khalifah RG. The carbon dioxide hydration activity of carbonic anhydrase. I. Stop-flow kinetic studies on the native human isoenzymes B and C. *J Biol Chem* 1971;246:2561–73.
 29. Vullo D, Innocenti A, Nishimori I, et al. Carbonic anhydrase activators: nocentini of the human isoforms VII (cytosolic) and XIV (transmembrane) with amino acids and amines. *Bioorg Med Chem Lett* 2007;17:4107–12.
 30. (a) Pastorekova S, Vullo D, Nishimori I, et al. Carbonic anhydrase activators: activation of the human tumor-associated isozymes IX and XII with amino acids and amines. *Bioorg Med Chem* 2008;16:3530–6. (b) Vullo D, Nishimori I, Innocenti A, et al. Carbonic anhydrase activators: an activation study of the human mitochondrial isoforms VA and VB with amino acids and amines. *Bioorg Med Chem Lett* 2007;17:1336–40.
 31. (a) Saada MC, Vullo D, Montero JL, et al. Mono- and di-halogenated histamine, histidine and carnosine derivatives are potent carbonic anhydrase I, II, VII, XII and XIV activators. *Bioorg Med Chem* 2014;22:4752–8. (b) Saada MC, Vullo D, Montero JL, et al. Carbonic anhydrase I and II activation with mono- and dihalogenated histamine derivatives. *Bioorg Med Chem Lett* 2001;21:4884–7.
 32. (a) Supuran CT, Barboiu M, Luca C, et al. Carbonic anhydrase activators. Part 14. Synthesis of mono- and bis- pyridinium salt derivatives of 2-amino-5-(2-aminoethyl)- and 2-amino-5-(3-aminopropyl)-1,3,4-thiadiazole, and their interaction with isozyme II. *Eur J Med Chem* 1996;31:597–606. (b) Scozzafava A, Supuran CT. Carbonic anhydrase activators. Part 24. High affinity isozymes I, II and IV activators, derivatives of 4-(4-chlorophenylsulfonylureido-amino acyl)ethyl-1H-imidazole. *Eur J Pharm Sci* 2000;10:29–41.
 33. Tiligada E, Ennis M. Histamine pharmacology: from Sir Henry Dale to the 21st century. *Br J Pharmacol* 2020;177:469–89.
 34. (a) Passani MB, Benetti F, Blandina P, et al. Histamine regulates memory consolidation. *Neurobiol Learn Mem* 2017;145:1–6. (b) Provensi G, Costa A, Passani MB, Blandina P. Donepezil, an acetylcholine esterase inhibitor, and ABT-239, a histamine H3 receptor antagonist/inverse agonist, require the integrity of brain histamine system to exert biochemical and procognitive effects in the mouse. *Neuropharmacology* 2016;109:139–47.
 35. Abbott NJ. Inflammatory mediators and modulation of blood-brain barrier permeability. *Cell Mol Neurobiol* 2000;20:131–47.
 36. Kitanaka J, Kitanaka N, Hall SF, et al. Histamine H3 receptor agonists decrease hypothalamic histamine levels and increase stereotypical biting in mice challenged with methamphetamine. *Neurochem Res* 2011;36:1824–33.
 37. Giovannini MG, Bartolini L, Bacciottini L, et al. Effects of histamine H3 receptor agonists and antagonists on cognitive performance and scopolamine-induced amnesia. *Behav Brain Res* 1999;104:147–55.
 38. Kitbunnadaj R, Zuiderveld OP, Christophe B, et al. Identification of 4-(1H-imidazol-4(5)-ylmethyl)pyridine (Immethridine) as a Novel, Potent, and Highly Selective Histamine H3 Receptor Agonist. *J Med Chem* 2004;47:2414–7.
 39. Henry MB, Zheng S, Duan C, et al. Hwa antidiabetic properties of the histamine H3 receptor protean agonist proxyfan. *Endocrinology* 2011;152:828–35.
 40. Ligneau X, Lin J, Vanni-Mercier G, et al. Neurochemical and behavioral effects of ciproxifan, a potent histamine H3-receptor antagonist. *J Pharmacol Exp Ther* 1998;287:658–66.
 41. Hough LB, Nalwalk JW, Barnes WG, et al. A third life for burimamide. Discovery and characterization of a novel class of non-opioid analgesics derived from histamine antagonists. *Ann N Y Acad Sci* 2000;909:25–40.
 42. Nowak JZ. Effects of histamine H1- and H2-receptor antagonists on dopamine, noradrenaline and serotonin systems in rat brain. *Pol J Pharmacol Pharm* 1980;32:451–61.
 43. Jonsson KA, Eriksson SE, Kagevi I, et al. Cimetidine, but not oxmetidine, penetrates into the cerebrospinal fluid after a single intravenous dose. *Br J Clin. Pharmacol* 1982;14:815–9.
 44. Clapp RH, Luckman SM. Proxyfan acts as a neutral antagonist of histamine H3 receptors in the feeding-related hypothalamic ventromedial nucleus. *Br J Pharmacol* 2012;167:1099–110.

45. Mochizuki T, Jansen FP, Leurs R, et al. Brain penetration of the histamine H3 receptor antagonists thioperamide and clobenpropit in rat and mouse, determined with ex vivo [¹²⁵I]iodophenpropit binding. *Brain Res* 1996;743:178–83.
46. Boertje SB, Le Beau D, Ward S. Impromidine-induced changes in the permeability of the blood-brain barrier of normotensive and spontaneously hypertensive rats. *Res Commun Chem Pathol Pharmacol* 1990;69:249–52.
47. Slugg PH, Haug MT, Pippenger CE. Ranitidine pharmacokinetics and adverse central nervous system reactions. *Arch Intern Med* 1992;152:2325–9.
48. Hough LB, Nalwalk JW. Inhibition of morphine antinociception by centrally administered histamine H2 receptor antagonists. *Eur J Pharmacol* 1992;215:69–74.
49. Calcutt CR, Ganellin CR, Griffiths R, et al. Zolantidine (SK&F 95282) is a potent selective brain-penetrating histamine H2-receptor antagonist. *Br J Pharmacol* 1988;93:69–78.
50. Tighilet B, Trottier S, Lacour M. Dose- and duration-dependent effects of betahistine dihydrochloride treatment on histamine turnover in the cat. *Eur J Pharmacol* 2005;523:54–63.
51. Esbenshade TA, Fox GB, Krueger KM, et al. Pharmacological properties of ABT-239 [4-(2-{2-[(2R)-2-methylpyrrolidinyl]ethyl}-benzofuran-5-yl)benzonitrile]: I. Potent and selective histamine H3 receptor antagonist with drug-like properties. *J Pharmacol Exp Ther* 2005;313:165–75.
52. Wágner G, Mocking TAM, Arimont M, et al. 4-(3-Aminoazetid-1-yl)pyrimidin-2-amines as High-Affinity Non-imidazole Histamine H3 Receptor Agonists with in Vivo Central Nervous System Activity. *J Med Chem* 2019;62:10848–66.
53. Yamazaki M, Terasaki T, Yoshioka K, et al. Carrier-mediated transport of H1-antagonist at the blood-brain barrier: Mepyramine uptake into bovine brain capillary endothelial cells in primary monolayer cultures. *Pharm Res* 1994;11:975–8.
54. Dettori I, Gaviano L, Melani A, et al. A selective histamine h4 receptor antagonist, JNJ7777120, is protective in a rat model of Transient Cerebral Ischemia. *Front Pharmacol*. 2018;9:e01231.
55. Plisson C, Gunn RN, Cunningham VJ, et al. 11C-GSK189254: a selective radioligand for *in vivo* central nervous system imaging of histamine H3 receptors by PET. *J Nucl Med* 2009;50:2064–72.
56. Nakamura T, Hiraoka K, Harada R, et al. Brain histamine H1 receptor occupancy after oral administration of desloratadine and loratadine. *Pharmacol Res Perspect* 2019;7:e00499.
57. Medhurst AD, Briggs MA, Bruton G, et al. Structurally novel histamine H3 receptor antagonists GSK207040 and GSK334429 improve scopolamine-induced memory impairment and capsaicin-induced secondary allodynia in rats. *Biochem Pharmacol* 2007;73:1182–94.