



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Role of adiponectin in the metabolism of skeletal muscles in collagen VI-related myopathies

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Role of adiponectin in the metabolism of skeletal muscles in collagen VI-related myopathies / Gamberi, Tania; Magherini, Francesca; Mannelli, Michele; Chrisam, Martina; Cescon, Matilde; Castagnaro, Silvia; Modesti, Alessandra; Braghetta, Paola; Fiaschi, Tania. - In: JOURNAL OF MOLECULAR MEDICINE. - ISSN 0946-2716. - ELETTRONICO. - 97:(2019), pp. 0-0. [10.1007/s00109-019-01766-0]

Availability:

This version is available at: 2158/1152913 since: 2019-07-02T20:32:59Z

Published version:

DOI: 10.1007/s00109-019-01766-0

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

(Article begins on next page)

[Click here to view linked References](#)

Role of adiponectin in the metabolism of skeletal muscles in collagen VI-related myopathies

Tania Gamberi, Francesca Magherini, Michele Mannelli, Martina Chrisam¹, Matilde Cescon¹,
Silvia Castagnaro¹, Alessandra Modesti, Paola Braghetta¹ and Tania Fiaschi*

*Dipartimento di Scienze Biomediche, Sperimentali e Cliniche “Mario Serio”, Università degli Studi
di Firenze, Italia; ¹Dipartimento di Medicina Molecolare, Università degli Studi di Padova, Italia*

*Corresponding author: Tania Fiaschi, Dipartimento di Scienze Biomediche, Sperimentali e
Cliniche “Mario Serio”, Università degli Studi di Firenze, viale Morgagni 50, 50134 Firenze, Italia.

e-mail: tania.fiaschi@unifi.it

1
2 **Abstract**
3

4 The role of adiponectin has been particularly deepened in diabetic muscles while the study of
5 adiponectin in hereditary myopathies has been marginal investigated. Here, we report the study
6
7 about adiponectin effects in *Col6a1*^{-/-} (collagen VI-null) mice. *Col6a1*^{-/-} mice show myopathic
8
9 phenotype closer to that of patients with Bethlem myopathy, thus representing an excellent animal
10
11 model for the study of this hereditary disease. Our findings demonstrate that *Col6a1*^{-/-} mice have
12
13 decreased plasma adiponectin content and diseased myoblasts have an impaired autocrine secretion
14
15 of the hormone. Moreover, *Col6a1*^{-/-} myoblasts show decreased glucose up-take and mitochondria
16
17 with depolarized membrane potential and impaired functionality, as supported by decreased oxygen
18
19 consumption. Exogenous addition of globular adiponectin modifies the features of *Col6a1*^{-/-}
20
21 myoblasts, becoming closer to that of the healthy myoblasts. Indeed, globular adiponectin enhances
22
23 glucose up-take in *Col6a1*^{-/-} myoblasts, modifies mitochondrial membrane potential and restores
24
25 oxygen consumption, turning closer to those of wild-type myoblasts. Finally, increase of plasma
26
27 adiponectin level in *Col6a1*^{-/-} mice is induced by fasting, a condition that has been previously
28
29 shown to lead to the amelioration of the dystrophic phenotype. Collectively, our results demonstrate
30
31 that exogenous replenishment of adiponectin reverses metabolic abnormalities observed in
32
33 *Col6a1*^{-/-} myoblasts.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Congenital muscular dystrophies represent a large and heterogeneous group of inherited muscle disorders with a severe and progressive clinical course. In particular, mutations in any of the genes encoding for the three main chains ($\alpha 1$, $\alpha 2$, $\alpha 3$) composing collagen VI (COL6), an extracellular matrix protein remarkably abundant in the endomysium of skeletal muscles, cause multiple muscle diseases, including Bethlem myopathy (BM) and Ullrich congenital muscular dystrophy (UCMD) [1]. COL6 null (*Col6a1*^{-/-}) mice display a myopathic phenotype resembling that of BM patients, and showed to be an excellent animal model for this pathology [2]. Muscles lacking COL6 are characterized by the presence of dilated sarcoplasmic reticulum and dysfunctional mitochondria, leading to muscle wasting [3]. These muscles exhibit accumulation of abnormal organelles due to an autophagy impairment. Amelioration of the dystrophic phenotype can be obtained through the reactivation of the autophagic flux, by either nutritional approaches or by pharmacological and genetic tools, leading to the removal of dysfunctional organelles [3, 4].

The action of adiponectin in skeletal muscle plays key roles ranging from metabolic to differentiating ones. Regarding the metabolic function, adiponectin affects glucose up-take, glycogen synthesis, glycolysis activation and triglyceride degradation [5-7]. In addition, adiponectin promotes myogenesis both in satellite cells [8] and myoblasts [9] and the differentiating role in myoblasts is linked to activation of autophagy [10]. In muscle cells, the binding of adiponectin to its specific AdipoR1 muscle receptor leads to the activation of a signalling cascade in which APMK plays a central role [7]. Adiponectin circulating in the plasma is mainly secreted from the adipose tissue, however, autocrine adiponectin production occurs in several tissues, including skeletal muscle which is one of the main site of local secretion [11]. Plasma adiponectin level is greatly affected by obesity. Obese individuals show significantly decreased adiponectin level in the blood, thus predisposing them to the onset of the metabolic syndrome [12]. Although the role of

1 adiponectin in healthy skeletal muscle is well documented, its effects on dystrophic muscles are still
2 unclear.
3
4

5 Here, we report that *Col6a1*^{-/-} mice display decreased circulating adiponectin level and that
6
7 *Col6a1*^{-/-} myoblasts-enriched cultures (hereafter referred to as “myoblasts”) have impaired
8
9 autocrine adiponectin secretion. In addition, *Col6a1*^{-/-} myoblasts display decreased glucose up-take,
10
11 higher glutamine dependence and decreased glucose consumption. Interestingly, exogenous
12
13 addition of adiponectin improves *Col6a1*^{-/-} metabolic behaviour, bringing it closer to that of wild-
14
15 type (WT) myoblasts. Indeed, adiponectin stimulation of *Col6a1*^{-/-} myoblasts increases glucose up-
16
17 take and oxygen consumption and induces glutamine dependence close to that of WT myoblasts. In
18
19 addition, we show that fasting, previously demonstrated to ameliorate the dystrophic phenotype of
20
21 *Col6a1*^{-/-} mice, induces an up-regulation of plasma adiponectin level in *Col6a1*^{-/-} mice. Overall,
22
23 these findings show that the modulation of adiponectin levels both *in vitro* and *in vivo* could
24
25 improve the **metabolic** defects due to the absence of COL6.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Materials and methods

Materials. Unless differently specified, all reagents were obtained from Sigma-Aldrich, Inc.; PVDF membrane was from Millipore; anti-adiponectin, anti-AdipoR1, anti-GAPDH, anti-GLUT4 antibodies and test ELISA for murine adiponectin assay were from Abcam; Alexa 488 fluorescent secondary antibodies were from Pierce. Adiponectin was from Alexis. JC-1 probe was from Molecular Probe; K-LATE kit for lactate assay was from Megazyme; [³H] 2-deoxy-glucose was from Perkin Elmer; ECL detection reagents was from GE Healthcare.

Mice. Diaphragm and tibialis anterior were isolated from 8 week-old male *Col6a1*^{-/-} (collagen VI-null) mice and compared age-matched male *Col6a1*^{+/+} (WT) mice. We housed mice in individual cages in an environmentally controlled room (23 °C, 12-h light-dark cycle) and provided food and water *ad libitum*. For fasted mice, we removed chow in the morning and maintained mice for 24 hours with no food but free access to water. Mouse procedures were approved by the Ethics Committee of the University of Padua and authorized by the Italian Ministry of Health according to D. Lgs. 26/2014 implementing Directive 2010/63 / EU.

Cell Culture. Myoblast-enriched cultures (hereafter referred to as “myoblasts”) were isolated from 8-week-old of WT and *Col6a1*^{-/-} muscles (diaphragm and tibialis anterior) by enzymatic and mechanical dissociation, following the protocol from Rando and Blau [13]. Isolated cells were transferred to gelatin-coated dishes and then cultured in DMEM supplemented with 10% FBS in 5% CO₂ humidified atmosphere and used until passage 2 as maximum.

Immunoblot analysis. Muscle protein extracts were prepared as previously described [14] and immunoblot analysis was performed as previously reported [10].

1 **Immunofluorescence.** Myoblasts were grown on glass coverslips and for confocal analysis, sub-
2 confluent myoblasts were treated as previously reported [10].
3

4 **Adiponectin assay.** Plasma and local adiponectin were assayed by ELISA test following
5 manufacturer's instruction. Plasma from WT and *Col6a1*^{-/-} mice was diluted 1:400 and adiponectin
6 value obtained by ELISA test was normalized on plasma total protein content. Autocrine production
7 of adiponectin was assayed in culture medium of WT and *Col6a1*^{-/-} myoblasts. In particular, WT
8 and *Col6a1*^{-/-} myoblasts were cultured for a day in serum-free medium before collecting the
9 medium. Total protein content of myoblasts was used for normalization of the value obtained by
10 Elisa test.
11
12
13
14
15
16
17
18
19
20
21

22 **Glucose up-take.** Glucose up-take was performed using [³H] 2-deoxy-glucose (0.5 mCi/mL, final
23 concentration) diluted in a buffered solution (140 mmol/L NaCl, 20 mmol/L Hepes/Na, 2.5 mmol/L
24 MgSO₄, 1 mmol/L CaCl₂, and 5 mmol/L KCl, pH7.4) for 15 minutes at 37 °C. Cells were
25 subsequently washed with cold PBS and lysed with 0.1 mol/L NaOH. Incorporated radioactive
26 glucose was assayed by scintillation counter and the obtained value was then normalized on total
27 protein content.
28
29
30
31
32
33
34
35
36
37

38 **Lactate assay.** Lactate amount was assayed in cell medium using K-LATE kit according to
39 manufacturer's instructions. Lactate amount was then normalized on protein content of the same
40 sample.
41
42
43
44
45

46 **MTT assay.** 0.5 mg/ml of 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2 H tetrazolium bromide
47 (MTT) was added to the cells. After 1 h at 37 °C, cells were extensively washed with PBS and then
48 1 ml of DMSO was added to the culture. The absorption was measured at 570 nm.
49
50
51
52
53

54 **Measurement of oxygen consumption.** Myoblasts were trypsinized, washed with PBS and
55 suspended in complete DMEM medium at the concentration of 10⁶ cells/ml. One ml of cell
56
57
58
59

1 suspension was transferred to an airtight chamber maintained at 37 °C. Oxygen consumption was
2 measured using a Clark-type O₂ electrode (Oxygraph Hansatech). Oxygen content was monitored
3
4 for at least 10 min. The rate of decrease in oxygen content, related to protein amount was taken as
5
6 index of the respiratory ability.
7
8
9

10 **Statistical analysis.** Data are presented as mean ± S.D. from at least three independent experiments.

11
12 Statistical analysis of the data was performed by Student's t-test or by one-way ANOVA using

13
14 Graph Pad Prism 4.0. *p*-values <0.05 were considered statistically significant.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

1
2
3 **1. Analysis of circulating and local adiponectin content in *Col6a1*^{-/-} mice.** We planned to
4 study adiponectin in *Col6a1*^{-/-} mice, a good animal model for UCMD and BM [2]. For this aim, we
5 chose the muscles diaphragm and tibialis anterior. Diaphragm is characterized by red fibres, a great
6 amount of mitochondria and oxidative metabolism. Conversely, tibialis anterior has fermentative
7 metabolism and is composed of white fibres containing few mitochondria. We checked adiponectin
8 and AdipoR1 levels in the two muscles isolated from **male** WT and *Col6a1*^{-/-} mice. No difference
9 in the expression level of both adiponectin and AdipoR1 has been observed in diaphragm and
10 tibialis anterior from *Col6a1*^{-/-} mice in comparison to WT (Fig. 1A). As well, our analysis of
11 adiponectin expression in myoblasts isolated from WT and *Col6a1*^{-/-}, diaphragm and tibialis
12 anterior showed that adiponectin expression is quite similar in *Col6a1*^{-/-} and WT myoblasts (Fig.
13 1B).

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 Hence, we analyzed both circulating adiponectin in plasma of WT and *Col6a1*^{-/-} mice and
31 autocrine adiponectin secretion in WT and *Col6a1*^{-/-} myoblast culture medium. Our findings
32 showed decreased circulating adiponectin (of about 30%) in *Col6a1*^{-/-} plasma in comparison to WT
33 (Fig. 2A). Remarkably, the analysis of autocrine production of adiponectin from diaphragm-isolated
34 myoblasts led to the same result. Indeed, *Col6a1*^{-/-} cells showed a 30% reduction of adiponectin
35 secretion when compared to WT myoblasts (Fig. 2B). **Body weight of WT and *Col6a1*^{-/-} mice**
36 **showed no appreciable differences as shown in Fig. 2C.**

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

diaphragm and tibialis anterior of *Col6a1*^{-/-} females are similar to WT females (Suppl. Fig. 1B), as observed in male *Col6a1*^{-/-} mice (Fig. 1A).

Collectively, these results show that *Col6a1*^{-/-} mice have decreased circulating adiponectin and that autocrine production of adiponectin by *Col6a1*^{-/-} myoblasts is defective, thus suggesting a decreased extracellular adiponectin availability for skeletal muscles in diseased mice.

2. Role of adiponectin in glucose metabolism of *Col6a1*^{-/-} myoblasts. Proteomic analysis in *Col6a1*^{-/-} muscles suggested an impairment of glycolysis [15]. To verify whether decreased glycolysis could be due to an impairment of glucose entry, we analyzed glucose up-take in myoblasts isolated from WT and *Col6a1*^{-/-} diaphragm and tibialis anterior. This analysis showed that *Col6a1*^{-/-} myoblasts have decreased glucose assumption in comparison to WT cells isolated from both muscles (Fig. 3A and 3B). Globular adiponectin (gAd) exerts in skeletal muscle several physiological actions, ranging from metabolic to differentiating functions [10, 11]. In particular, gAd has high affinity for the binding with AdipoR1 and greatly affects glucose up-take in skeletal muscle [16]. Hence, we planned to stimulate WT and *Col6a1*^{-/-} myoblasts with gAd. Stimulation with gAd enhanced glucose up-take in *Col6a1*^{-/-} myoblasts (Fig. 3A and 3B) with higher increase in *Col6a1*^{-/-} myoblasts isolated from tibialis anterior (Fig. 3B). We tested whether the impairment of glucose up-take in *Col6a1*^{-/-} myoblasts was due to a lower expression of GLUT4, the primary glucose transporter in skeletal muscle. As shown in Fig. 4, the expression level of GLUT4 was similar in WT and *Col6a1*^{-/-} myoblasts. However, confocal analysis suggests that gAd stimulation could alter intracellular distribution of GLUT4 in *Col6a1*^{-/-} myoblasts (Suppl. fig. 1).

The impaired glucose up-take observed in myoblasts from diaphragm and tibialis anterior of *Col6a1*^{-/-}, suggested that *Col6a1*^{-/-} myoblasts could import the amino acid glutamine that could be driven towards Krebs cycle. To test this possibility, we performed glucose up-take in the absence of glutamine, thus forcing *Col6a1*^{-/-} myoblasts to use glucose. Glutamine depletion induced *Col6a1*^{-/-}

1 myoblasts to introduce glucose instead of glutamine (Fig. 5A and 5B), thus enhancing the glucose
2 up-take. Our observation suggested that *Col6a1*^{-/-} myoblasts preferentially use glutamine instead of
3 glucose for their metabolism. Indeed, in the presence of both glucose and glutamine *Col6a1*^{-/-}
4 myoblasts showed enhanced metabolic activity in comparison to WT cells. Conversely, when
5 glutamine is removed, *Col6a1*^{-/-} myoblasts showed decreased metabolic activity, becoming similar
6 to that of WT cells (Fig. 5C). In order to deeper characterize the energetic metabolism of *Col6a1*^{-/-}
7 myoblasts, we also measured the production of lactate. This analysis evidenced that *Col6a1*^{-/-}
8 myoblasts from diaphragm and tibialis anterior secreted quite similar amount of lactate in
9 comparison to healthy cells (Fig. 6A and 6B). Interestingly, lactate production was greatly increased
10 in *Col6a1*^{-/-} myoblasts after gAd stimulation in both muscles examined (Fig. 6A and 6B).

11
12 Overall, these findings suggest that *Col6a1*^{-/-} myoblasts introduce less glucose in
13 comparison to WT cells, preferring glutamine for their viability. In the presence of both glucose and
14 glutamine, gAd stimulation enhances glucose up-take and increases lactate production in *Col6a1*^{-/-}
15 myoblasts in both muscles. On the other hand, glutamine withdrawal forced myoblasts to increase
16 glucose up-take and gAd stimulation raises glucose intake.

17
18 **3. Adiponectin stimulation restores mitochondrial membrane potential and increases oxygen**
19 **consumption in *Col6a1*^{-/-} myoblasts.** *Col6a1*^{-/-} muscles exhibit altered mitochondria characterized
20 by high membrane depolarization that is greatly attenuated after prolonged fasting [3]. To detect the
21 presence of altered mitochondria also in *Col6a1*^{-/-} myoblasts, we used the mitochondrial probe JC-
22 1. Fig. 7A shows that *Col6a1*^{-/-} myoblasts contained mitochondria with depolarized membranes
23 (evidenced by green fluorescence) in comparison to WT cells (showing red fluorescent, typical of
24 normal mitochondrial membrane potential). gAd induces mitochondria biogenesis through the
25 regulation of PGC-1 α and genetic removal of AdipoR1 in muscle resulted in decreased expression
26 and deacetylation of PGC-1 α and reduced mitochondrial enzymes [17]. To test whether adiponectin
27

1 could ameliorate mitochondrial membrane depolarization, *Col6a1*^{-/-} myoblasts have been
2 stimulated with gAd. This treatment led to the shift from green to red fluorescence in *Col6a1*^{-/-}
3 mitochondria, which appeared similar to WT cells (Fig. 7A). This finding demonstrates that gAd
4 ameliorates altered mitochondrial features in *Col6a1*^{-/-} myoblasts, thus restoring normal membrane
5 potential.
6
7
8
9
10

11
12 To detect whether the depolarization observed in *Col6a1*^{-/-} mitochondria is associated with
13 altered functionality, we analyzed oxygen consumption. We observed that *Col6a1*^{-/-} myoblasts
14 consume lesser amount of oxygen (about 30%) in comparison to WT cells (Fig.7B). Interestingly,
15 [18] gAd stimulation restored oxygen consumption similar to that of WT cells (Fig. 7B).
16
17
18
19
20
21
22

23 Overall, these findings show that gAd induces beneficial effects on mitochondria in *Col6a1*^{-/-}
24 myoblasts, thus ameliorating membrane potential and the defective oxygen consumption. The
25 benefits following gAd treatment are not due to enhanced expression of mitochondrial complexes,
26 since no differences in the expression of these proteins were observed between WT and *Col6a1*^{-/-}
27 myoblasts (data not shown).
28
29
30
31
32
33
34
35

36 **4. Fasting in *Col6a1*^{-/-} mice increases circulating adiponectin.** Fasting in *Col6a1*^{-/-} mice
37 improves the dystrophic phenotype [3]. Moreover, fasting raises both serum and cerebrospinal fluid
38 levels of adiponectin [18]. This led us to hypothesise that fasting in *Col6a1*^{-/-} mice could enhance
39 both circulating and local muscular production of adiponectin. To test this hypothesis, we detected
40 the expression level of adiponectin in muscles of fasting *Col6a1*^{-/-} mice. Results showed that only
41 tibialis anterior of fasted *Col6a1*^{-/-} mice exhibit higher amount of adiponectin in comparison to WT
42 fasted mice (Fig. 8A). Conversely, adiponectin level remains quite similar in the diaphragm of
43 fasted *Col6a1*^{-/-} and WT mice (Fig. 8A). Interestingly, fasting affects the amount of circulating
44 adiponectin in plasma. Indeed, while normally fed *Col6a1*^{-/-} mice showed decreased circulating
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 adiponectin (Fig. 2A), fasting increased the amount of circulating hormone that become similar to
2 WT mice (Fig. 8B).
3
4

5 These findings demonstrate that fasting promotes adiponectin expression and secretion, thus
6 suggesting that adiponectin could be involved in the improvement of dystrophic phenotype
7
8 observed in *Col6a1*^{-/-} mice following fasting.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discussion

1
2
3 The metabolic and differentiating roles of adiponectin in skeletal muscle have been well-
4 established and deeply studied [7]. However, the involvement of adiponectin in inherited muscle
5 diseases is just at the beginning. It has been reported that DMD patients show a significant
6 reduction of High Molecular Weight complexes of adiponectin [19], but the consequence of this
7 decrease has not been investigated yet. Here, we report the study on adiponectin in *Col6a1*^{-/-} mice,
8 having a targeted inactivation of the α 1 chain of collagen VI (COL6), and show a myopathic
9 phenotype resembling that of BM patients [2]. Whether the local and endocrine adiponectin levels
10 are changed and whether the exogenous addition of adiponectin undergoes beneficial effects for the
11 diseased phenotype has not considered yet in this pathology. This is an important point to be
12 addressed, since adiponectin could be considered a new tool for the improvement of the diseased
13 muscles. An interesting finding is that *Col6a1*^{-/-} mice have decreased amount of circulating
14 adiponectin in comparison to healthy mice. In addition to endocrine adiponectin production, several
15 tissues, including skeletal muscle, have autocrine adiponectin production [9, 11]. We found that
16 muscles and myoblasts from *Col6a1*^{-/-} have intracellular adiponectin content similar to healthy
17 mice, thus showing that myopathy does not affect adiponectin transcription and translation.
18 However, secretion of adiponectin is greatly decreased in *Col6a1*^{-/-} myoblasts in comparison to
19 WT, thus demonstrating that myopathy affects both endocrine and skeletal muscle autocrine
20 adiponectin secretion in *Col6a1*^{-/-} mice. **Particularly, we found that both males and females of**
21 ***Col6a1*^{-/-} mice show reduced circulating adiponectin in comparison to WT, thus suggesting that the**
22 **decrease of plasma adiponectin is not related to gender but rather to the pathological condition.** The
23 mechanisms driving adiponectin secretion are well elucidated [20-22]. However, some factors may
24 compromise the secretion of this adipokine. Among these, oxidative stress has been reported to
25 inhibit adiponectin secretion by adipose tissue [23, 24]. Several studies recognised the key role of
26 oxidative stress and abnormal production of reactive oxygen species in the pathophysiology of
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 muscular dystrophies [25-27]. In particular, the presence of oxidative stress, due to an accumulation
2 of reactive oxygen species by monoamine oxidase activity, has been reported in muscles of *Col6a1*⁻
3
4 ⁻ mice [28]. Hence, oxidative stress observed in *Col6a1*⁻ muscles may be the cause leading to the
5
6 impairment of adiponectin secretion. In addition, *Col6a1*⁻ mice show dilated sarcoplasmic
7
8 reticulum in muscle cells that may affect the translation of those proteins destined to secretion as
9
10 adiponectin [3]. For what adipose tissue is concerned, *Col6a1*⁻ mice show increased adipocyte cell
11
12 size due to a more flexibility of the adipocyte matrix, thus allowing the increase of adipocytes size
13
14 in the absence of associated necrosis and inflammation. This leads to metabolic dysfunction and
15
16 fibrosis formation in *Col6a1*⁻ adipose tissue in a diabetic state [29].
17
18
19
20

21
22 Hence, our results show that *Col6a1*⁻ mice have impaired both circulating and local (at
23
24 least at skeletal muscle level) adiponectin production, thus generating a situation in which the
25
26 hormone is not sufficiently available for target tissues as skeletal muscle.
27
28

29
30 Decreased amount of adiponectin could be involved in the altered metabolic behaviour
31
32 observed in myoblasts of *Col6a1*⁻ mice in comparison to those from healthy animals. We found
33
34 that myoblasts from diaphragm and tibialis anterior of *Col6a1*⁻ mice show impaired glucose up-
35
36 take. Furthermore, our results highlight another altered behaviour of *Col6a1*⁻ myoblasts.
37
38 Particularly myoblasts isolated from diaphragm of *Col6a1*⁻ mice show greater glutamine-
39
40 dependence which appears indispensable for myoblast metabolism. When cells take less glucose,
41
42 glutamine can be used for cell metabolism instead of glucose. In skeletal muscle, glutamine can
43
44 meet different fates since it can be used as nitrogen donor in the synthesis of amino acids and
45
46 nucleosides, converted to α -ketoglutarate or citrate, thus supporting the tricarboxylic acid cycle,
47
48 [30, 31]. Proteomic analysis performed on *Col6a1*⁻ muscles suggested that diaphragm of *Col6a1*⁻
49
50 mice is characterized by blunted glycolysis and impaired tricarboxylic acid cycle [15]. In
51
52 agreement, we observe decreased glucose uptake, likely responsible for the decreased glycolysis
53
54 observed by De Palma *et al.* [15]. In skeletal muscle, adiponectin has essential and pleiotropic roles,
55
56
57
58
59

1 ranging from metabolic to differentiating function [8-11, 32]. Concerning the metabolic role,
2 adiponectin promotes glucose up-take via GLUT4 translocation [16] and regulates fatty acid
3
4 metabolism, increasing fatty acid uptake and oxidation, and suppressing fatty acid synthesis through
5
6 the activation of AMPK, p38 MAPK and PPAR α signalling [5, 6, 33]. We found that treatment of
7
8 *Col6a1*^{-/-} myoblasts with exogenous adiponectin induces several beneficial effects at the metabolic
9
10 level. Adiponectin enhances glucose up-take, promotes lactate production and decreases glutamine
11
12 dependence, approaching that of WT cells. In addition, adiponectin enhances glucose up-take in
13
14 *Col6a1*^{-/-} myoblasts after glutamine withdrawal. Hence, exogenous addition of adiponectin could
15
16 restore the proper metabolic signalling in *Col6a1*^{-/-} myoblasts.
17
18
19
20
21

22 As already reported in intact fibres [3], *Col6a1*^{-/-} myoblasts contain depolarized and less
23
24 functional mitochondria with decreased capacity to consume oxygen. In this scenario, addition of
25
26 exogenous adiponectin can play a key role in restoring proper metabolism in *Col6a1*^{-/-} muscles.
27
28 Indeed, we found that adiponectin promotes beneficial effects also at the mitochondrial level in
29
30 *Col6a1*^{-/-} myoblasts. The role of adiponectin in mitochondria of muscle cells has been clearly
31
32 defined. By AdipoR1, adiponectin promotes the activation of Ca²⁺/calmodulin-dependent protein
33
34 kinase β (CaMKK β), AMPK and SIRT1 signalling pathways, increased the expression and
35
36 decreased acetylation of peroxisome proliferator-activated receptor c coactivator-1 α (PGC-1 α), thus
37
38 leading to increased mitochondria in myotubes [17]. Although the mechanism needs further
39
40 investigations, adiponectin treatment improves mitochondrial potential and increased oxygen
41
42 consumption in *Col6a1*^{-/-} mitochondria similar to that of WT cells.
43
44
45
46
47

48 Skeletal muscles of *Col6a1*^{-/-} mice have impaired autophagy that leads to dysfunctional
49
50 mitochondria and spontaneous apoptosis [3]. The myopathic phenotype is greatly ameliorated by
51
52 forcing activation of autophagy by different approaches. In particular, fasting in *Col6a1*^{-/-} mice
53
54 restores normal autophagy associated with a general amelioration of the dystrophic features [3].
55
56 Adiponectin is involved in autophagy activation in skeletal muscle since adiponectin-activated
57
58
59
60
61
62
63
64
65

1 autophagy is essential for the myogenic role of the hormone thus preventing the formation of the
2 myopathic phenotype [10]. Since a correlation between fasting and adiponectin has been previously
3 reported [34], we hypothesized that fasting could increase circulating adiponectin. Interestingly, we
4 found that fasted *Col6a1*^{-/-} mice show circulating adiponectin levels similar to that of healthy
5 animals. This observation could suggest a role of adiponectin in the induction of the beneficial
6 effects observed in fasted *Col6a1*^{-/-} mice. In fasting condition, only tibialis anterior of *Col6a1*^{-/-}
7 mice shows increased adiponectin production in comparison to tibialis anterior of WT mice, while
8 no difference has been observed in diaphragm of *Col6a1*^{-/-} and WT mice. Since the two muscles
9 have distinct metabolism (i.e. tibialis anterior has fermentative metabolism while diaphragm is an
10 oxidative muscle), the higher adiponectin content in tibialis anterior in fasted *Col6a1*^{-/-} mice could
11 help the muscle to sustain fermentation and the right production of ATP. Thus, a greater amount of
12 adiponectin in the tibialis anterior of *Col6a1*^{-/-} mice could help to overcome the observed metabolic
13 defects.

14 Overall, our results show several abnormalities regarding adiponectin in *Col6a1*^{-/-} mice.
15 Among these, the decrease of local and circulating levels of adiponectin plays a key role since this
16 decrease could be involved in the metabolic alterations observed in *Col6a1*^{-/-} mice. These
17 metabolic defects are compensated by the treatment with exogenous adiponectin that leads *Col6a1*^{-/-}
18 myoblast behaviour similar to that of WT cells. Interestingly, myopathy does not influence the
19 expression level of the muscle-specific receptor for adiponectin, AdipoR1, which remains
20 unchanged in all examined muscle, thus making possible the treatment of the diseased muscles with
21 exogenous adiponectin.

1
2
3
4
5
6 **Figure legends**
7
8
9

10 **Fig. 1. Expression level of adiponectin in muscles and myoblasts from WT and *Col6a1*^{-/-} mice.**

11
12 A) Anti-adiponectin and anti-AdipoR1 immunoblots in diaphragm and tibialis anterior of muscles
13 from WT and *Col6a1*^{-/-} mice. B) Analysis of adiponectin expression in myoblasts isolated from
14 diaphragm and tibialis anterior of WT and *Col6a1*^{-/-} mice. GAPDH immunoblots demonstrate that
15 the same amount of total proteins was loaded in each lane. Bar graphs show the expression level of
16 adiponectin and AdipoR1 in each muscle reported as arbitrary units (a.u.). **p*<0.01. acrp30:
17 adiponectin. n=5 mice/group.
18
19
20
21
22
23
24
25
26
27

28 **Fig. 2. Analysis of plasma and autocrine adiponectin level.** A) Plasma adiponectin level in WT
29 and *Col6a1*^{-/-} mice and B) Autocrine secretion of adiponectin from myoblasts isolated from
30 diaphragm of WT and *Col6a1*^{-/-} mice, obtained using ELISA test. The values are reported as a
31 percentage of decrease of *Col6a1*^{-/-} in comparison to WT. n=5 mice/group. C) **Body weight of WT**
32 **and *Col6a1*^{-/-} mice.** **p*<0.01. acrp30: adiponectin.
33
34
35
36
37
38
39
40
41

42 **Fig. 3. Glucose up-take in myoblasts from WT and *Col6a1*^{-/-} mice.** Glucose up-take was
43 analyzed in myoblasts isolated from diaphragm (A) and tibialis anterior (B) of WT and *Col6a1*^{-/-}
44 mice. Sub-confluent myoblasts were treated with low glucose medium (5mM final) overnight.
45 Where indicated, adiponectin (1 µg/ml) was added to the cells for 30 minutes before the addition of
46 [3H] 2-deoxy-glucose. Radioactive counts were normalized with total protein content and reported
47 as arbitrary units (a.u.). **p*< 0.001. gAd: globular adiponectin.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 4. Analysis of the glucose transporter GLUT4 in myoblasts from WT and *Col6a1*^{-/-} mice.

Myoblasts isolated from diaphragm of WT and *Col6a1*^{-/-} mice were used to analysed GLUT4 expression level by immunoblot. GLUT4 level has been normalized using anti-GAPDH immunoblot and reported in the bar graph as arbitrary unit (a.u.).

Fig. 5. Glucose up-take and MTT assay in WT and *Col6a1*^{-/-} myoblasts after glutamine withdrawal.

Myoblasts, isolated from diaphragm (panel A) and tibialis anterior (panel B) of WT and *Col6a1*^{-/-} mice, were cultured in complete medium without glutamine (gln) for 24 hours. Cells were then stimulated with gAd (1µg/ml) for 30 minutes before the addition of [³H] 2-deoxy-glucose. Radioactive counts were normalized with total protein content and reported as arbitrary units (a.u.). C) MTT assay in myoblasts isolated form diaphragm of WT and *Col6a1*^{-/-} mice. Cells were cultured in complete medium with or without gln (2 mM final) for 48 hours before performing MTT assay. **p*<0.01; ***p*<0.001. gAd: globular adiponectin.

Fig. 6. Lactate production in WT and *Col6a1*^{-/-} myoblasts.

Myoblasts isolated from diaphragm (A) and tibialis anterior (B) were serum deprived overnight in high glucose medium (25mM final) with or without gAd (1µg/ml). The values of lactate were normalized on protein content and reported as arbitrary unit (a.u.). **p*< 0.01. gAd: globular adiponectin.

Fig. 7. Role of adiponectin in mitochondria of *Col6a1*^{-/-} myoblasts.

Myoblasts were isolated from diaphragm of WT and *Col6a1*^{-/-} mice and mitochondrial membrane potential were assayed using JC-1 probe as shown in A). Cells were seeded on coverslips, serum-depleted overnight and then gAd (1 µg/ml) was added for 24 h. JC-1 probe (5 µM) was added to the cells for 15 min at 37 °C and then immediately observed using confocal microscope. Images are representative of four independent experiments. B) Oxygen consumption assay was performed as reported in Material and Methods. The values were reported as percent increase or decrease considering WT myoblasts as 100%. **p*<0.01; gAd: globular adiponectin.

1
2 **Fig. 8. Analysis of plasma and expression levels of adiponectin in fasted WT and *Col6a1*^{-/-}**
3 **mice.** WT and *Col6a1*^{-/-} mice were fasted for 24 hours. A) Acrp30 and AdipoR1 expression levels
4 in diaphragm and tibialis anterior muscles. The values are reported in the bar graph as arbitrary
5 units (a.u.) obtained by the ratio between adiponectin or AdipoR1 and GAPDH. B) Analysis of
6 adiponectin in plasma of WT and *Col6a1*^{-/-} mice performed by ELISA test. The values obtained by
7 ELISA test were normalized on plasmatic protein content. **p*<0.01; acrp30: adiponectin. n=5
8 mice/group.
9
10
11
12
13
14
15
16
17

18 **Supplementary figure 1. Analysis of adiponectin in females of *Col6a1*^{-/-} mice.** A) Plasma
19 adiponectin level obtained using ELISA test and reported as percentage of decrease of *Col6a1*^{-/-} in
20 comparison to WT. n=5 mice/group. **p*<0.01; acrp30: adiponectin. B) Intracellular level of
21 adiponectin in diaphragm and tibialis anterior by immunoblot. GAPDH immunoblot has been used
22 for normalization. Bar graphs show the expression level of adiponectin reported as arbitrary unit
23 (a.u.).
24
25
26
27
28
29
30
31
32

33 **Supplementary figure 2.** Confocal images of myoblasts isolated from diaphragm of WT and
34 *Col6a1*^{-/-} mice incubated with (right panel), or without (left panel) gAd (1 µg/ml) for 30 minutes.
35 The intracellular location of GLUT4 is shown as green signal obtained using a specific primary
36 anti-GLUT4 antibody (ThermoFisher Scientific) and a secondary antibody conjugated to Alexa-
37 488. Nuclei have been labelled with DAPI. The images are representative of three independent
38 experiments. gAd: globular adiponectin.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgments.

This work was supported by the Italian Ministry of University and Research (MIUR).

Conflict of interest: the authors declare no conflict of interest

Bibliography

1. Lampe AK, Bushby KM (2005) Collagen VI related muscle disorders. *J Med Genet* 42: 673-685. DOI 10.1136/jmg.2002.002311
2. Bonaldo P, Braghetta P, Zanetti M, Piccolo S, Volpin D, Bressan GM (1998) Collagen VI deficiency induces early onset myopathy in the mouse: an animal model for Bethlem myopathy. *Hum Mol Genet* 7: 2135-2140
3. Grumati P, Coletto L, Sabatelli P, Cescon M, Angelin A, Bertaglia E, Blaauw B, Urciuolo A, Tiepolo T, Merlini L, et al. (2010) Autophagy is defective in collagen VI muscular dystrophies, and its reactivation rescues myofiber degeneration. *Nat Med* 16: 1313-1320. DOI 10.1038/nm.2247
4. Chrisam M, Pirozzi M, Castagnaro S, Blaauw B, Polishchuck R, Cecconi F, Grumati P, Bonaldo P (2015) Reactivation of autophagy by spermidine ameliorates the myopathic defects of collagen VI-null mice. *Autophagy* 11: 2142-2152. DOI 10.1080/15548627.2015.1108508
5. Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, et al. (2002) Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 8: 1288-1295. DOI 10.1038/nm788
6. Fruebis J, Tsao TS, Javorschi S, Ebbets-Reed D, Erickson MR, Yen FT, Bihain BE, Lodish HF (2001) Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci U S A* 98: 2005-2010. DOI 10.1073/pnas.041591798
7. Yamauchi T, Iwabu M, Okada-Iwabu M, Kadowaki T (2014) Adiponectin receptors: a review of their structure, function and how they work. *Best Pract Res Clin Endocrinol Metab* 28: 15-23. DOI 10.1016/j.beem.2013.09.003
8. Fiaschi T, Giannoni E, Taddei ML, Chiarugi P (2012) Globular adiponectin activates motility and regenerative traits of muscle satellite cells. *PLoS One* 7: e34782. DOI 10.1371/journal.pone.0034782
9. Fiaschi T, Cirelli D, Comito G, Gelmini S, Ramponi G, Serio M, Chiarugi P (2009) Globular adiponectin induces differentiation and fusion of skeletal muscle cells. *Cell Res* 19: 584-597. DOI 10.1038/cr.2009.39
10. Gamberi T, Modesti A, Magherini F, D'Souza DM, Hawke T, Fiaschi T (2016) Activation of autophagy by globular adiponectin is required for muscle differentiation. *Biochim Biophys Acta* 1863: 694-702. DOI 10.1016/j.bbamcr.2016.01.016
11. Fiaschi T, Magherini F, Gamberi T, Modesti PA, Modesti A (2014) Adiponectin as a tissue regenerating hormone: more than a metabolic function. *Cell Mol Life Sci* 71: 1917-1925. DOI 10.1007/s00018-013-1537-4
12. Wang ZV, Scherer PE (2016) Adiponectin, the past two decades. *J Mol Cell Biol* 8: 93-100. DOI 10.1093/jmcb/mjw011
13. Rando TA, Blau HM (1994) Primary mouse myoblast purification, characterization, and transplantation for cell-mediated gene therapy. *J Cell Biol* 125: 1275-1287
14. Gamberi T, Fiaschi T, Valocchia E, Modesti A, Mantuano P, Rolland JF, Sanarica F, De Luca A, Magherini F (2018) Proteome analysis in dystrophic mdx mouse muscle reveals a drastic alteration of key metabolic and contractile proteins after chronic exercise and the potential modulation by anti-oxidant compounds. *J Proteomics* 170: 43-58. DOI 10.1016/j.jprot.2017.09.009
15. De Palma S, Leone R, Grumati P, Vasso M, Polishchuk R, Capitanio D, Braghetta P, Bernardi P, Bonaldo P, Gelfi C (2013) Changes in muscle cell metabolism and mechanotransduction are associated with myopathic phenotype in a mouse model of collagen VI deficiency. *PLoS One* 8: e56716. DOI 10.1371/journal.pone.0056716
16. Ceddia RB, Somwar R, Maida A, Fang X, Bikopoulos G, Sweeney G (2005) Globular adiponectin increases GLUT4 translocation and glucose uptake but reduces glycogen synthesis in rat skeletal muscle cells. *Diabetologia* 48: 132-139. DOI 10.1007/s00125-004-1609-y

17. Iwabu M, Yamauchi T, Okada-Iwabu M, Sato K, Nakagawa T, Funata M, Yamaguchi M, Namiki S, Nakayama R, Tabata M, et al. (2010) Adiponectin and AdipoR1 regulate PGC-1alpha and mitochondria by Ca(2+) and AMPK/SIRT1. *Nature* 464: 1313-1319. DOI 10.1038/nature08991
18. Kadowaki T, Yamauchi T, Kubota N (2008) The physiological and pathophysiological role of adiponectin and adiponectin receptors in the peripheral tissues and CNS. *FEBS Lett* 582: 74-80. DOI 10.1016/j.febslet.2007.11.070
19. Hathout Y, Marathi RL, Rayavarapu S, Zhang A, Brown KJ, Seol H, Gordish-Dressman H, Cirak S, Bello L, Nagaraju K, et al. (2014) Discovery of serum protein biomarkers in the mdx mouse model and cross-species comparison to Duchenne muscular dystrophy patients. *Hum Mol Genet* 23: 6458-6469. DOI 10.1093/hmg/ddu366
20. Jin D, Sun J, Huang J, Yu X, Yu A, He Y, Li Q, Yang Z (2015) Peroxisome proliferator-activated receptor γ enhances adiponectin secretion via up-regulating DsbA-L expression. *Mol Cell Endocrinol* 411: 97-104. DOI 10.1016/j.mce.2015.04.015
21. He Y, Lu L, Wei X, Jin D, Qian T, Yu A, Sun J, Cui J, Yang Z (2016) The multimerization and secretion of adiponectin are regulated by TNF-alpha. *Endocrine* 51: 456-468. DOI 10.1007/s12020-015-0741-4
22. DeClercq V, d'Eon B, McLeod RS (2015) Fatty acids increase adiponectin secretion through both classical and exosome pathways. *Biochim Biophys Acta* 1851: 1123-1133. DOI 10.1016/j.bbali.2015.04.005
23. Tang H, Sebastian BM, Axhemi A, Chen X, Hillian AD, Jacobsen DW, Nagy LE (2012) Ethanol-induced oxidative stress via the CYP2E1 pathway disrupts adiponectin secretion from adipocytes. *Alcohol Clin Exp Res* 36: 214-222. DOI 10.1111/j.1530-0277.2011.01607.x
24. Parola M, Marra F (2011) Adipokines and redox signaling: impact on fatty liver disease. *Antioxid Redox Signal* 15: 461-483. DOI 10.1089/ars.2010.3848
25. Rando TA, Disatnik MH, Yu Y, Franco A (1998) Muscle cells from mdx mice have an increased susceptibility to oxidative stress. *Neuromuscul Disord* 8: 14-21
26. Whitehead NP, Yeung EW, Allen DG (2006) Muscle damage in mdx (dystrophic) mice: role of calcium and reactive oxygen species. *Clin Exp Pharmacol Physiol* 33: 657-662. DOI 10.1111/j.1440-1681.2006.04394.x
27. Tidball JG, Wehling-Henricks M (2007) The role of free radicals in the pathophysiology of muscular dystrophy. *J Appl Physiol* (1985) 102: 1677-1686. DOI 10.1152/jappphysiol.01145.2006
28. Menazza S, Blaauw B, Tiepolo T, Toniolo L, Braghetta P, Spolaore B, Reggiani C, Di Lisa F, Bonaldo P, Canton M (2010) Oxidative stress by monoamine oxidases is causally involved in myofiber damage in muscular dystrophy. *Hum Mol Genet* 19: 4207-4215. DOI 10.1093/hmg/ddq339
29. Khan T, Muise ES, Iyengar P, Wang ZV, Chandalia M, Abate N, Zhang BB, Bonaldo P, Chua S, Scherer PE (2009) Metabolic dysregulation and adipose tissue fibrosis: role of collagen VI. *Mol Cell Biol* 29: 1575-1591. DOI 10.1128/MCB.01300-08
30. Le A, Lane AN, Hamaker M, Bose S, Gouw A, Barbi J, Tsukamoto T, Rojas CJ, Slusher BS, Zhang H, et al. (2012) Glucose-independent glutamine metabolism via TCA cycling for proliferation and survival in B cells. *Cell Metab* 15: 110-121. DOI 10.1016/j.cmet.2011.12.009
31. Wise DR, Ward PS, Shay JE, Cross JR, Gruber JJ, Sachdeva UM, Platt JM, DeMatteo RG, Simon MC, Thompson CB (2011) Hypoxia promotes isocitrate dehydrogenase-dependent carboxylation of α -ketoglutarate to citrate to support cell growth and viability. *Proc Natl Acad Sci U S A* 108: 19611-19616. DOI 10.1073/pnas.1117773108
32. Fiaschi T, Tedesco FS, Giannoni E, Diaz-Manera J, Parri M, Cossu G, Chiarugi P (2010) Globular adiponectin as a complete mesoangioblast regulator: role in proliferation, survival, motility, and skeletal muscle differentiation. *Mol Biol Cell* 21: 848-859. DOI 10.1091/mbc.E09-04-0310
33. Yoon MJ, Lee GY, Chung JJ, Ahn YH, Hong SH, Kim JB (2006) Adiponectin increases fatty acid oxidation in skeletal muscle cells by sequential activation of AMP-activated protein kinase, p38 mitogen-activated protein kinase, and peroxisome proliferator-activated receptor alpha. *Diabetes* 55: 2562-2570. DOI 10.2337/db05-1322

34. Kubota N, Yano W, Kubota T, Yamauchi T, Itoh S, Kumagai H, Kozono H, Takamoto I, Okamoto S, Shiuchi T, et al. (2007) Adiponectin stimulates AMP-activated protein kinase in the hypothalamus and increases food intake. *Cell Metab* 6: 55-68. DOI 10.1016/j.cmet.2007.06.003

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1



A**B**

Figure 2




A**B****C**

Figure 3




Figure 4




Figure 5




Figure 6



A**B**

Figure 7







Figure 8

A



B





Click here to access/download
Supplementary Material
Suppl figure 1.tif





Click here to access/download
Supplementary Material
Suppl figure 2.tif

