



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

On the generating functions of pattern-avoiding Motzkin paths

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

On the generating functions of pattern-avoiding Motzkin paths / Bean C.; Bernini A.; Cervetti M.; Ferrari L..
- In: JOURNAL OF SYMBOLIC COMPUTATION. - ISSN 0747-7171. - STAMPA. - 113:(2022), pp. 126-138.
[10.1016/j.jsc.2022.02.006]

Availability:

This version is available at: 2158/1262218 since: 2022-03-24T15:41:52Z

Published version:

DOI: 10.1016/j.jsc.2022.02.006

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze
(<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

(Article begins on next page)

On the generating functions of pattern-avoiding Motzkin paths^{*}

Christian Bean^{*}

School of Computer Science, Reykjavik University, Reykjavik, Iceland

Antonio Bernini, Luca Ferrari

Dipartimento di Matematica e Informatica, Università di Firenze, Firenze, Italy

Matteo Cervetti

LIB, Université de Bourgogne Franche-Comté, Dijon, France

Abstract

Using a recursive approach, we show that the generating function $F(x)$ for sets of Motzkin paths avoiding a single (not necessarily consecutive) pattern is rational over x and the Catalan generating function $C(x) = \frac{1-\sqrt{1-4x^2}}{2x^2}$, where x keeps track of the length of the path. Moreover, an algorithm is provided for finding the generating function in the more general case of an arbitrary set of patterns. In addition, this algorithm allows us to find a combinatorial specification for pattern-avoiding Motzkin paths, which can be used not only for enumeration, but also for exhaustive and random generation.

Keywords: lattice path, generating function, Motzkin path, pattern, combinatorial specification

2020 MSC: 05A10, 05A15

^{*}Research partially supported by grant 207178-052 from the Icelandic Research Fund. A.B. and L.F. are members of the INdAM research group GNCS. They are partially supported by the 2020-2021 INdAM-GNCS project: “Combinatoria delle permutazioni, delle parole e dei grafi: algoritmi e applicazioni”

^{*}Corresponding author

Email addresses: `christianbean@ru.is` (Christian Bean), `antonio.bernini@unifi.it` (Antonio Bernini), `luca.ferrari@unifi.it` (Luca Ferrari), `matteocervetti27@gmail.com` (Matteo Cervetti)

1. Introduction

A *Motzkin path* of length n is a lattice path starting at $(0,0)$ and ending at $(n,0)$ consisting of *up steps* ($U = (1,1)$), *down steps* ($D = (1,-1)$) and *horizontal steps* ($H = (1,0)$) that never goes below the x -axis. We represent
5 Motzkin paths as words over the alphabet $\{U, D, H\}$. In Figure 1, the Motzkin path $UHUUDHDD$ is shown.




Figure 1: The Motzkin path $UHUUDHDD$.

Let \mathcal{M} be the set of all Motzkin paths, \mathcal{M}_H be the set of Motzkin paths that start with a horizontal step, and \mathcal{M}_U be the set of Motzkin paths starting with an up step. With this we have

$$\mathcal{M} = \{\epsilon\} \sqcup \mathcal{M}_H \sqcup \mathcal{M}_U, \quad (1)$$

where ϵ represents the empty path and we use \sqcup to denote disjoint union. The “folklore” result (see for example Donaghey and Shapiro [1]) on Motzkin paths says that every Motzkin path in \mathcal{M}_H can be written as Hw for some w in \mathcal{M} ,
10 and every Motzkin path in \mathcal{M}_U can be written $UxDy$ for some x, y in \mathcal{M} as shown in Figure 1.




Figure 2: A pictorial representation of the structural decomposition of Motzkin paths.

If we let m_n be the number of length n Motzkin paths then it follows that

m_n satisfies the recurrence relation

$$m_0 = m_1 = 1, \quad m_n = m_{n-1} + \sum_{i=0}^{n-2} m_i m_{n-i-2}$$

and moreover if we let $M(x) = \sum_{n \geq 0} m_n x^n$ be the generating function, then $M(x)$ satisfies the minimal polynomial

$$1 + (x - 1)M(x) + x^2 M(x)^2 = 0.$$

Akin to investigations for other combinatorial structures (eminently for permutations, but also for graphs), there has been interest in studying properties related to the notion of *patterns* in the context of lattice paths. For paths it has
15 been common to consider a pattern as a sequence of *contiguous* letters, see for instance Asinowski et al. [2] and Sapounakis et al. [3], to cite just a couple of references. This may be due to the fact that, as we have also remarked above, the set of lattice paths of a certain type can be conveniently seen as a formal language, and it is common in theoretical computer science to study contiguous
20 patterns, or *factors*, of the words of a formal language (e.g. *pattern matching* and related problems). In this work, however, we will deal with a notion of pattern that is closer to the one usually studied for permutations, namely we will consider a pattern as a *subword* (whose letters are not necessarily contiguous) of a given word. Below we give the necessary notations and definitions in the
25 specific case of Motzkin paths.

A Motzkin path p *contains* a *pattern* q in $\{U, H, D\}^*$, written $q \preceq p$, if q occurs as a subword in p . If p does not contain q we say p *avoids* q and write $q \not\preceq p$. For a set P of patterns, we say a path *avoids* P if it avoids all $q \in P$ and define the set of Motzkin paths avoiding P as

$$\text{Av}(P) = \{p \in \mathcal{M} \mid p \text{ avoids } P\}.$$

If a path does not avoid P we say it *contains* P and define the set of Motzkin paths containing P as

$$\text{Co}(P) = \{p \in \mathcal{M} \mid p \text{ contains } P\},$$

i.e. the set of Motzkin paths containing at least one pattern in P . For instance, consider again the Motzkin path in Figure 1. It is easy to check that it contains the pattern $UHDH$, but it avoids the pattern $UDHH$. Notice that, according to our definition of containment for sets of patterns, such a path also contains
30 the set $P = \{UHDH, UDHH\}$.

The set $\text{Av}(H)^1$ is the set of Dyck paths which are counted by the Catalan numbers. Recall that Dyck paths are defined like Motzkin paths, except that they do not use horizontal steps. In Bacher et al. [4], it was shown that any set of Dyck paths avoiding a single pattern has a rational generating function.
35 In this paper, we show a similar statement holds for the set of Motzkin paths, alongside an algorithm for effectively computing the generating functions even in the more general case of a set of patterns.

Theorem 1.1. Let q be a fixed pattern and let a_n be the number of q -avoiding Motzkin paths of length n . Then the generating function $\Delta_q(x) = \sum_{n \geq 0} a_n x^n$ is
40 rational over x and $C(x) = \sum_{n \geq 0} C_n x^{2n}$, where C_n is the n -th Catalan number.

The paper is organised as follows. In Section 2, we outline an algorithm for computing a *combinatorial specification*, in the sense of Flajolet and Sedgewick [5], for sets of Motzkin paths avoiding an arbitrary set of patterns. Such a specification then gives a method for computing the generating function but also
45 the ability to sample uniformly from these sets. In Section 3, we give a proof of Theorem 1.1. The strategy of the proof consists of describing a recursive procedure to compute the generating function $\Delta_q(x)$; such a procedure also depends on certain auxiliary generating functions, which are in turn described in a recursive fashion. Finally, in Section 4 some suggestions for further research
50 are given.

In closing this Introduction, we remark that, whenever we will consider Dyck paths, we will usually enumerate them according to the length, rather than (as it is usual) the semilength. As a consequence, our version of the Catalan generating

¹We will not include the braces in our notation, i.e. we write $\text{Av}(H)$ rather than $\text{Av}(\{H\})$

function is $C(x) = \frac{1-\sqrt{1-4x^2}}{2x^2}$, hence, for all $n \in \mathbb{N}$, $[x^{2n}]C(x) = C_n = \frac{1}{n+1} \binom{2n}{n}$
 55 (the n -th Catalan number) and $[x^{2n+1}]C(x) = 0$ (where $[x^n]F(x)$ denotes the
 n -th coefficient of the generating function $F(x)$).

2. The algorithm

For a set P of patterns, we define $\text{Av}_H(P)$ to be the Motzkin paths avoiding
 P and beginning with an H step,

$$\text{Av}_H(P) = \text{Av}(P) \cap \mathcal{M}_H,$$

and $\text{Av}_U(P)$ to be the Motzkin paths avoiding P and beginning with a U step,

$$\text{Av}_U(P) = \text{Av}(P) \cap \mathcal{M}_U.$$

The set of Motzkin paths avoiding P can be partitioned in the same manner as
 the set of all Motzkin paths,

$$\text{Av}(P) = \{\epsilon\} \sqcup \text{Av}_H(P) \sqcup \text{Av}_U(P). \quad (2)$$

2.1. Starting with H

Using the following, we can enumerate the set $\text{Av}_H(P)$.

Theorem 2.1. For a set of patterns P , let P_U , P_D , and P_H be the sets of
 patterns in P beginning with U , D and H , respectively, and

$$P' = P_U \cup P_D \cup \{p \mid Hp \in P_H\}$$

then

$$\text{Av}_H(P) = \{Hp \mid p \in \text{Av}(P')\}.$$

60 *Proof.* Let Hq be a path in $\text{Av}_H(P)$. It follows that q must avoid every pattern
 in P_U and P_D . If q contains an occurrence of some p for $Hp \in P_H$, then Hq
 must contain an occurrence of Hp . Therefore, q must avoid p .

On the contrary, let $Hq \in \{Hp \mid p \in \text{Av}(P')\}$, then as q avoids P_U and
 P_D , it follows that Hq also avoids P_U and P_D . As q avoids all p coming from
 65 P_H , the path Hq must avoid Hp . Hence we have shown that Hq avoids
 P . □

If a_n counts the number of length n paths in $\text{Av}_H(P)$ and b_n counts the number of length n paths in $\text{Av}(P')$ then by the above Theorem we get $a_k = b_{k-1}$ for all $k \geq 1$.

70 2.2. Starting with U

Every Motzkin path in $\text{Av}_U(P)$ can be written as $UxDy$ for some x, y in $\text{Av}(P)$, however not each choice of x and y from $\text{Av}(P)$ yields a Motzkin path in $\text{Av}_U(P)$. For example, if $P = \{HH\}$, and $x = H$ and $y = H$, then we get the path $UHDH$ which contains an occurrence of HH . In order to capture this, we
75 introduce the notion of *crossing patterns*.

A Motzkin path $p \in \mathcal{M}_U$ contains the crossing pattern $\ell - r$, where ℓ and r are words over the alphabet $\{U, D, H\}$, if p can be written $UxDy$ where UxD contains ℓ and y contains r . Otherwise, we say it *avoids* $\ell - r$. If either ℓ or r is ϵ we write $-r$ and $\ell-$, respectively, and call these patterns *local*. We use the
80 notation of $\text{Av}(P)$ and $\text{Co}(P)$ as before for crossing patterns.

With our new definition, for the case of $P = \{HH\}$ we have

$$\text{Av}_U(P) = \text{Av}_U(-HH, H-H, HH-).$$

For a path $UxDy$ in $\text{Av}_U(P)$, if x avoids H then $UxDy$ avoids $H-H$ and $HH-$. However, if x contains H then $UxDy$ avoids $-H$ and $HH-$. That is

$$\text{Av}_U(-HH, H-H, HH-) = \text{Av}_U(-HH, H-) \sqcup (\text{Av}_U(-H, HH-) \cap \text{Co}(H-)). \quad (3)$$

Every path in $\text{Av}_U(-HH, H-)$ can be written $UxDy$ where $x \in \text{Av}(H)$ and $y \in \text{Av}(HH)$. Similarly, every path in $\text{Av}_U(-H, HH-) \cap \text{Co}(H-)$ can be written $UxDy$ where $x \in \text{Av}(HH) \cap \text{Co}(H)$ and $y \in \text{Av}(H)$. This argument is shown pictorially in Figure 3.

Theorem 2.1 says that

$$\text{Av}_H(HH) = \{Hp | p \in \text{Av}(H)\}. \quad (4)$$

Let $\Delta_{HH}(x)$ be the generating function for $\text{Av}(HH)$ and $C(x) = \frac{1-\sqrt{1-4x^2}}{2x^2}$ be the generating function for $\text{Av}(H)$, then it follows from Equations (2), (3),




Figure 3: A pictorial representation of Equation (3) for $\text{Av}_U(HH)$.

and (4) that $\Delta_{HH}(x)$ satisfies the equation

$$\Delta_{HH}(x) = 1 + xC(x) + x^2C(x)\Delta_{HH}(x) + x^2(\Delta_{HH}(x) - C(x))C(x). \quad (5)$$

From (5), replacing $C(x)$ with $\frac{1-\sqrt{1-4x^2}}{2x^2}$ and squaring after suitable manipulations, we find that $\Delta_{HH}(x)$ satisfies the minimal polynomial

$$(4x^4 - x^2)\Delta_{HH}(x)^2 + (4x^3 - 4x^2 - x + 1)\Delta_{HH}(x) + 5x^2 - 1.$$

85 In passing, we observe that the generating function $\Delta_{HH}(x)$ is interesting in itself. In fact, the coefficients of even index are Catalan numbers, whereas the coefficients of odd index (which count, by the way, Motzkin paths having exactly one horizontal step with respect to the length) are the binomial coefficients $\binom{2n+1}{n+1}$, i.e., sequence A001700 in the Online Encyclopedia of Integer Sequences
90 (OEIS) [6].

We generalize this idea in the following theorem.

Theorem 2.2. For any finite sets P and Q of patterns there exist sets of local crossing patterns P_1, P_2, \dots, P_k and Q_1, Q_2, \dots, Q_k such that

$$\text{Av}_U(P) \cap \bigcap_{q \in Q} \text{Co}(q) = \bigsqcup_{i=1}^k \left(\text{Av}_U(P_i) \cap \bigcap_{q \in Q_i} \text{Co}_U(q) \right).$$

Proof. For a pattern p we define the set $s(p)$ consisting of all crossing patterns $\ell - r$ such that ℓr is p ,

$$s(p) = \{\ell - r \mid \ell, r \in \{U, D, H\}^* \text{ and } \ell r = p\}.$$

It follows that every Motzkin path in $\text{Av}_U(P)$ must avoid $\bigsqcup_{p \in P} s(p)$ and every Motzkin path in $\bigcap_{q \in Q} \text{Co}(q)$ must contain some pattern in $s(q)$ for each q in Q ,

i.e.

$$\text{Av}_U(P) \cap \bigcap_{q \in Q} \text{Co}(q) = \text{Av}_U \left(\bigsqcup_{p \in P} s(p) \right) \cap \bigcap_{q \in Q} \text{Co}_U(s(q)).$$

For a set of patterns Q we can partition the set of Motzkin paths containing Q into those that avoid a pattern q in Q and those that contain q . In our case, if we set $Q = \{q_1, q_2, \dots, q_h\}$, we are avoiding a set of crossing patterns $\tilde{P} = \bigsqcup_{p \in P} s(p)$ and must contain the sets of crossing patterns $\tilde{Q}_1, \tilde{Q}_2, \dots, \tilde{Q}_h$, where $\tilde{Q}_i = s(q_i)$, for all $i = 1, 2, \dots, h$. Therefore, if $q \in \tilde{Q}_1$, we have

$$\begin{aligned} \text{Av}_U(\tilde{P}) \cap \bigcap_{i=1}^h \text{Co}_U(\tilde{Q}_i) &= \left(\text{Av}_U(\tilde{P} \cup \{q\}) \cap \text{Co}_U(\tilde{Q}_1 \setminus \{q\}) \cap \bigcap_{i=2}^h \text{Co}_U(\tilde{Q}_i) \right) \\ &\sqcup \left(\text{Av}_U(\tilde{P}) \cap \text{Co}_U(\{q\}) \cap \bigcap_{i=2}^h \text{Co}_U(\tilde{Q}_i) \right). \end{aligned}$$

By iterating this process, and perhaps rearranging the order of the \tilde{Q}_i , this will result in writing

$$\text{Av}_U(P) \cap \bigcap_{q \in Q} \text{Co}(q) = \bigsqcup_{i=1}^h \left(\text{Av}_U(\tilde{P}_i) \cap \bigcap_{q \in \tilde{Q}_i} \text{Co}_U(q) \right).$$

where the \tilde{P}_i and \tilde{Q}_i are sets of (not necessarily local) crossing patterns.

If $\ell - r$ in \tilde{Q}_1 then it follows that the paths must contain both $\ell -$ and $-r$, i.e. they contain every pattern in the set $\tilde{Q}'_1 = (\tilde{Q}_1 \setminus \{\ell - r\}) \sqcup \{\ell -, -r\}$ and we have

$$\text{Av}_U(\tilde{P}_1) \cap \bigcap_{q \in \tilde{Q}_1} \text{Co}_U(q) = \text{Av}_U(\tilde{P}_1) \cap \bigcap_{q \in \tilde{Q}'_1} \text{Co}_U(q). \quad (6)$$

If $\ell - r$ in \tilde{P}_1 then we can partition the paths to those that avoid $\ell -$ and those that contain $\ell -$ giving

$$\begin{aligned} \text{Av}_U(\tilde{P}_1) \cap \bigcap_{q \in \tilde{Q}_1} \text{Co}_U(q) &= \left(\text{Av}_U(\tilde{P}_1 \sqcup \{\ell -\}) \cap \bigcap_{q \in \tilde{Q}_1} \text{Co}_U(q) \right) \\ &\sqcup \left(\text{Av}_U(\tilde{P}_1 \sqcup \{-r\}) \cap \bigcap_{q \in \tilde{Q}_1 \sqcup \{\ell -\}} \text{Co}_U(q) \right). \end{aligned} \quad (7)$$

By repeated application of Equations (6) and (7), simplifying the avoidance and containment sets, and possibly reordering the \tilde{P}_i and \tilde{Q}_i , we get the desired disjoint union where all of the crossing patterns in the resulting sets P_i and Q_i are local. \square

The case analysis of Theorem 2.2 will result in a disjoint union with sets of the form $\text{Av}_U(P) \cap \bigcap_{q \in Q} \text{Co}_U(q)$ for some sets of local crossing patterns P and Q . As the pattern containment conditions for these sets of Motzkin paths become local we get the following theorem.

Theorem 2.3. Let P and Q be sets of local crossing patterns. Let P_r (Q_r) be the right local patterns in P (Q). Let P_ℓ (Q_ℓ) be the patterns obtained by taking the left local patterns in P (Q) and removing a single U from the left and single D from the right if such exists. Then,

$$\text{Av}_U(P) \cap \bigcap_{q \in Q} \text{Co}_U(q) = \{UxDy \mid x \in \text{Av}_U(P_\ell) \cap \bigcap_{q \in Q_\ell} \text{Co}_U(q), y \in \text{Av}_U(P_r) \cap \bigcap_{q \in Q_r} \text{Co}_U(q)\}.$$

Let a_n be the number of length n Motzkin paths in $\text{Av}(P) \cap \bigcap_{q \in Q} \text{Co}(q)$, b_n be the number of length n Motzkin paths in $\text{Av}(P_\ell) \cap \bigcap_{q \in Q_\ell} \text{Co}(q)$, and c_n be the number of length n Motzkin paths in $\text{Av}(P_r) \cap \bigcap_{q \in Q_r} \text{Co}(q)$. Theorem 2.3 implies $a_n = \sum_{i=0}^{n-2} b_i c_{n-i-2}$.

2.3. Finding a specification

Combinatorial exploration, introduced in Bean [7], is an automatic method for finding (*combinatorial*) *specifications*. It consists of a systematic application of *strategies* to create (*combinatorial*) *rules* about a (*combinatorial*) *class* of interest. Each rule describes how to build a class from other classes using well-defined *constructors*. In this paper, we only use the disjoint union and Cartesian product constructors. Using these rules, the method then finds a specification which can be used, for example, to count the number of objects of each size, generate objects, and sample uniformly at random. This entire procedure has been implemented as the `comb_spec_searcher` Python package by Bean et al. [8].

Theorems 2.1, 2.2, and 2.3 encode strategies for finding specifications for pattern-avoiding Motzkin paths. Moreover, as the recursive application of these theorems result in either shortening of the patterns being avoided and contained or reducing the size of the sets being avoided and contained, they give a
120 finite process that will always result in a specification. Our Python implementation, which uses the `comb_spec_searcher` package, can be found on GitHub by Bean [9].

We do one final example to illustrate the recursive nature of the theorems.

2.4. Enumerating Motzkin paths avoiding $UHHD$

125 In this section, we outline how Theorems 2.1, 2.2, and 2.3 are used to enumerate $\text{Av}(UHHD)$.

We first apply Equation (2) to get

$$\text{Av}(UHHD) = \{\epsilon\} \sqcup \text{Av}_H(UHHD) \sqcup \text{Av}_U(UHHD). \quad (8)$$

Theorem 2.1 tells us that

$$\text{Av}_H(UHHD) = \{H\} \times \text{Av}(UHHD). \quad (9)$$

Theorem 2.2 tells us that we can find sets of local crossing patterns to describe $\text{Av}_U(UHHD)$. We follow the algorithm outlined in the proof of the theorem to get this description. The set of crossing patterns coming from $UHHD$ is

$$s(UHHD) = \{-UHHD, U - HHD, UH - HD, UHH - D, UHHD-\}.$$

Therefore, $\text{Av}_U(UHHD) = \text{Av}_U(s(UHHD))$. As we have no sets of patterns Q to contain as in Theorem 2.2, we apply Equation 7 to get

$$\begin{aligned} \text{Av}_U(s(UHHD)) &= \text{Av}_U(-UHHD, -HHD, UH-) \cap \text{Co}_U(U-) \\ &\quad \sqcup (\text{Av}_U(-HD, UHH - D, UHHD-) \cap \text{Co}_U(UH-)) \\ &= \text{Av}_U(-HHD, UH-) \cap \text{Co}_U(U-) \\ &\quad \sqcup \text{Av}_U(-HD, UHH-) \cap \text{Co}_U(UH-) \\ &\quad \sqcup \text{Av}_U(-D, UHHD-) \cap \text{Co}_U(UHH-) \end{aligned} \quad (10)$$

Note, the case where you are avoiding $U-$ is precisely the empty set since all of the paths contain $U-$, and therefore we have not included this in our equations.

Each of the disjoint sets on the right are defined by local crossing patterns and so we apply Theorem 2.3 to each of these sets. This gives the equations

$$\begin{aligned} \text{Av}_U(-HHD, UH-) \cap \text{Co}_U(U-) &= \{UD\} \times (\text{Av}(H) \cap \text{Co}(\epsilon)) \times \text{Av}(HHD) \\ \text{Av}_U(-HD, UHH-) \cap \text{Co}_U(UH-) &= \{UD\} \times (\text{Av}(HH) \cap \text{Co}(H)) \times \text{Av}(HD) \\ \text{Av}_U(-D, UHHD-) \cap \text{Co}_U(UHH-) &= \{UD\} \times (\text{Av}(HH) \cap \text{Co}(HH)) \times \text{Av}(D). \end{aligned} \quad (11)$$

As a path can not both avoid and contain HH , the latter set is in fact empty.

130 The algorithm would actually spot this sooner as it is not hard to argue that a path cannot simultaneously avoid $UHHD-$ and contain $UHH-$ since the final letter before the split in a path is D .

In the previous section we enumerated $\text{Av}(HH) \cap \text{Co}(H)$ by utilising the set difference

$$\text{Av}(HH) \cap \text{Co}(H) = \text{Av}(HH) \setminus \text{Av}(H)$$

which implies its generating function is $\Delta_{HH}(x) - C(x)$. Although this is a legitimate method for enumeration, due to the set difference operation above,

135 the sampling of the Motzkin paths will not be efficient. Therefore our algorithm instead continues to apply the theorems in order to only use disjoint unions and Cartesian products.

In Equations (11), we need to expand further the sets $\text{Av}(HH) \cap \text{Co}(H)$, $\text{Av}(HHD)$, and $\text{Av}(HD)$. We first apply Theorem 2.1 to get

$$\begin{aligned} \text{Av}_H(HH) \cap \text{Co}(H) &= \{H\} \times \text{Av}(H) \\ \text{Av}_H(HHD) &= \{H\} \times \text{Av}(HD) \\ \text{Av}_H(HD) &= \{H\} \times \text{Av}(D). \end{aligned} \quad (12)$$

Note, $\text{Av}(D)$ is precisely $\{H\}^*$.

We then apply Theorem 2.2 to get

$$\begin{aligned}
\text{Av}_U(HH) \cap \text{Co}(H) &= \text{Av}_U(-HH, H-H, HH-) \cap \text{Co}_U(-H, H-) \\
&= (\text{Av}_U(H-, -HH) \cap \text{Co}_U(-H)) \sqcup (\text{Av}_U(-H, HH-) \cap \text{Co}_U(H-)) \\
\text{Av}_U(HD) &= \text{Av}_U(-HD, H-D, HD-) \\
&= \text{Av}_U(H-, -HD) \sqcup (\text{Av}_U(-D, HD-) \cap \text{Co}_U(H-)) \\
\text{Av}_U(HHD) &= \text{Av}_U(-HHD, H-HD, HH-D, HHD-) \\
&= \text{Av}_U(H-, -HHD) \sqcup (\text{Av}_U(-HD, HH-D, HHD-) \cap \text{Co}_U(H-)) \\
&= \text{Av}_U(H-, -HHD) \sqcup (\text{Av}_U(-HD, HH-) \cap \text{Co}_U(H-)) \\
&\quad \sqcup (\text{Av}_U(-D, HHD-) \cap \text{Co}_U(HH-)).
\end{aligned} \tag{13}$$

By a similar argument as before the sets $\text{Av}_U(-D, HD-) \cap \text{Co}_U(H-)$ and $\text{Av}_U(-D, HHD-) \cap \text{Co}_U(HH-)$ are the empty set. For the remaining all of the patterns are local crossing patterns so we can apply Theorem 2.3.

$$\begin{aligned}
\text{Av}_U(H-, -HH) \cap \text{Co}_U(-H) &= \{UD\} \times \text{Av}(H) \times (\text{Av}(HH) \cap \text{Co}(H)) \\
\text{Av}_U(-H, HH-) \cap \text{Co}_U(H-) &= \{UD\} \times (\text{Av}(HH) \cap \text{Co}(H)) \times \text{Av}(H) \\
\text{Av}_U(H-, -HD) &= \{UD\} \times \text{Av}(H) \times \text{Av}(HD) \\
\text{Av}_U(H-, -HHD) &= \{UD\} \times \text{Av}(H) \times \text{Av}(HHD) \\
\text{Av}_U(-HD, HH-) \cap \text{Co}_U(H-) &= \{UD\} \times (\text{Av}(HH) \cap \text{Co}(H)) \times \text{Av}(HD)
\end{aligned} \tag{14}$$

The Equations (8), (9), (10), (11), (12), (13), and (14) give a combinatorial specification for $\text{Av}(UHHD)$, and can be used directly to get the generating function

$$\frac{1 - 3x - 4x^2 + 12x^3 - (1 - 3x - 4x^2 + 8x^3)\sqrt{1 - 4x^2}}{2x^2(1 - 2x - 3x^2 + 8x^3 - 4x^4)}$$

for this set. The coefficients of this generating function are the sequence A347036 in the OEIS [6]. We ran our algorithm on many sets of pattern-avoiding Motzkin paths. We list a few which have connections to sequences in the OEIS [6].

The generating function for $\text{Av}(UDH)$ is

$$\frac{1 - 2x - \sqrt{1 - 4x^2}}{2x(2x - 1)}$$

which shows that there are $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ paths of length n in this set, i.e. sequence A001405 in the OEIS [6].

The set $\text{Av}(UDH, UHD)$ has the generating function

$$\frac{1 - \sqrt{1 - 4x^2}}{2x^2(x - 1)}$$

whose coefficients are the sequence A110199 in the OEIS [6]. This tells us that
 145 there are $\sum_{k=0}^n C_n$ many paths of lengths $2n$ and $2n + 1$ in $\text{Av}(UDH, UHD)$.
 This follows from the fact that the paths in this set can be described as some
 Dyck path prepended with an arbitrary number of H steps.

The generating functions for $\text{Av}(UUDD)$ and $\text{Av}(UDUDUD, UUDDUD, UUDUDD)$
 are

$$\frac{1 - 4x + 7x^2 - 6x^3 + 3x^4}{(1 - x)^5} \text{ and } \frac{1 - 4x + 6x^2 - 4x^3 + 2x^4}{(1 - x)^3(1 - 2x)}$$

whose coefficients are the sequences A000127 and A084634 in the OEIS [6].

In a few cases, we found the sequences for the coefficients at odd indices
 in the generating functions appeared in the OEIS [6]. For example, if we let
 a_n be the number of length n Motzkin paths in $\text{Av}(HHUD, HUHD, UHHD)$
 then the generating function for the sequence of odd length paths in this set,
 i.e., $\sum_{n \geq 0} a_{2n+1}x^n$, is

$$\frac{1 - 4x - \sqrt{1 - 4x}}{-2x(1 - 5x + 4x^2)}.$$

The coefficients are the sequence A079309 in the OEIS [6]. Similarly, the se-
 quence given by the number of odd length paths in $\text{Av}(UDHH)$ has the gener-
 ating function

$$\frac{1 - 6x + 8x^2 - (1 - 3x)\sqrt{1 - 4x}}{-x + 8x^2 - 16x^3}$$

which appears to be the sequence A194460 in the OEIS [6].

150 3. Proof of Theorem 1.1

In this section, we give a proof of Theorem 1.1. Analogously to what has been done in Bacher et al. [4] for Dyck paths, our strategy is the following. First, we describe functional equations satisfied by some bivariate generating functions of certain Motzkin prefixes, where the relevant statistics are the length and the
155 final height. These are then used to find functional equations for the generating functions of Motzkin paths avoiding a single pattern. The derived equations will clearly show that such generating functions are rational over x and $C(x)$, as desired.

A *Motzkin prefix* is defined exactly like a Motzkin path, except that the final
160 point of the path has nonnegative height (so it is not required that the path ends on the x -axis, but it can end at every point having nonnegative integer coordinates). Denote with \mathcal{MP} the set of all Motzkin prefixes. For a given non-empty Motzkin prefix p , let p^- be the Motzkin prefix obtained from p by removing its last step. Given a set P of Motzkin prefixes, let $\text{MinCo}(P) = \{p \in$
165 $\mathcal{MP} \mid p \text{ contains } P \text{ and } p^- \text{ avoids } P\}$. In other words, an element of $\text{MinCo}(P)$ is a smallest Motzkin prefix containing P . In the sequel, we will be interested in the case where $P = \{q\}$, for a certain Motzkin prefix q .

Denote with $\Gamma_q(x, y)$ the bivariate generating function of the smallest Motzkin prefixes containing q , where x keeps track of the length and y keeps track of
170 the final height. For instance, choosing $q = UH$, the generic smallest Motzkin prefix containing q is obtained by concatenating a sequence of letters H with a non-empty Dyck prefix followed by an H . Thus, recalling the expression of the bivariate generating function $\mathcal{DP}(x, y) = \frac{2}{1-2xy+\sqrt{1-4x^2}}$ of Dyck prefixes, we get

$$\Gamma_q(x, y) = \frac{1}{1-x} \left(\frac{2}{1-2xy+\sqrt{1-4x^2}} - 1 \right) x.$$

175 The following result gives a recursive procedure to compute $\Gamma_q(x, y)$. In the statement below, ϵ is the empty path; moreover, given $q \in \mathcal{MP}$ and $X \in \{U, H, D\}$, qX is the Motzkin prefix obtained by appending the step X to q .

Proposition 3.1. For any given Motzkin prefix q , we have:

$$\Gamma_\epsilon(x, y) = 1, \quad (15)$$

$$\Gamma_{qU}(x, y) = \frac{xy}{(1-x)(x-y(1-x))} \left(x\Gamma_q\left(x, \frac{x}{1-x}\right) - y(1-x)\Gamma_q(x, y) \right), \quad (16)$$

$$\Gamma_{qH}(x, y) = \frac{2x}{(1-2xy + \sqrt{1-4x^2})(y-xC(x))} (y\Gamma_q(x, y) - xC(x)\Gamma_q(x, xC(x))), \quad (17)$$

$$\Gamma_{qD}(x, y) = \frac{x}{y} \left(\frac{1}{1-xy-x}\Gamma_q(x, y) - \frac{1}{1-x}\Gamma_q(x, 0) \right). \quad (18)$$

Proof. Clearly, the only smallest Motzkin prefix containing the empty path is ϵ itself, which gives $\Gamma_\epsilon(x, y) = 1$.

180 Let $\pi \in \text{MinCo}(qU)$ and denote with π' the smallest prefix of π containing q . Moreover, we indicate with h the final height of π' . Then, π can be factorized as

$$\pi = \pi' \beta^{(h)} U, \quad (19)$$

where $\beta^{(h)}$ is a path starting at height h (which is the height of the final point of π') using only H steps and D steps and not crossing the x -axis (i.e., the final height i of $\beta^{(h)}$ is such that $0 \leq i \leq h$). Clearly, the path $\beta^{(h)}$ is the reverse of a path α starting at the origin, using only U steps and H steps, with final height less than or equal to h . It is not difficult to compute the bivariate generating function $A(x, y)$ of such paths α , where x and y track the length and the final height of α , respectively. Indeed, such a path α , if not empty, can be obtained either by taking an H step followed by a pattern of the same kind or by taking a U step followed by a pattern of the same kind. This leads to the functional equation:

$$A(x, y) = 1 + xA(x, y) + xyA(x, y),$$

hence

$$A(x, y) = \frac{1}{1-x-xy} = \frac{1}{1-x} \sum_{n \geq 0} \left(\frac{x}{1-x} \right)^n y^n, \quad (20)$$

If $B^{(h)}(x, y)$ denotes the generating function of the paths $\beta^{(h)}$ (where x and y have the same role as in $A(x, y)$), using essentially the same argument as above, we have

$$B^{(h)}(x, y) = \sum_{i=0}^h ([y^{h-i}] A(x, y)) y^i;$$

hence, in terms of generating functions, relation (19) becomes:

$$\begin{aligned} \Gamma_{qU}(x, y) &= \left(\sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) B^{(h)}(x, y) \right) xy \\ &= \left(\sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) \sum_{i=0}^h ([y^{h-i}] A(x, y)) y^i \right) xy. \end{aligned} \quad (21)$$

185 We note that, referring to (19), the term $\sum_{h \geq 0} [y^h] \Gamma_q(x, y)$ in (21) records the prefix π' , while the term xy tracks the step U . By using (20) for the coefficient $[y^{h-i}] A(x, y)$, expression (21) can be reduced to (16).

Similarly, let $\pi \in \text{MinCo}(qH)$ and let π' be the smallest prefix of π containing q . We denote with $\delta^{(h)}$ a Dyck factor starting at height h (i.e., a sequence of U and D steps which does not cross the x -axis) and with $D^{(h)}(x, y)$ the bivariate generating function for such paths. We have:

$$\pi = \pi' \delta^{(h)} H.$$

Therefore,

$$\Gamma_{qH}(x, y) = \left(\sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) D^{(h)}(x, y) \right) x. \quad (22)$$

As far as $D^{(h)}(x, y)$ is concerned, denoting with γ a generic Dyck prefix and with γ_i a Dyck path, we observe that a Dyck factor $\delta^{(h)}$ can be factorized as
190 $\delta^{(h)} = (\gamma_1 D)(\gamma_2 D) \dots (\gamma_r D) \gamma$, with $0 \leq r \leq h$, where the first D step reaching height $h - i$ is highlighted, for each $i = 1, 2, \dots, r$. From the above construction, in terms of generating functions we have:

$$D^{(h)}(x, y) = \mathcal{DP}(x, y)y^h + C(x)x\mathcal{DP}(x, y)y^{h-1} +$$

$$C(x)^2x^2\mathcal{DP}(x, y)y^{h-2} + \dots + C(x)^hx^h\mathcal{DP}(x, y)y^{h-h}$$

$$= \mathcal{DP}(x, y) \sum_{i=0}^h x^i y^{h-i} C(x)^i,$$

leading to

$$D^{(h)}(x, y) = \frac{2}{1 - 2xy + \sqrt{1 - 4x^2}} \cdot \frac{y^{h+1} - x^{h+1}C(x)^{h+1}}{y - xC(x)}. \quad (23)$$

Plugging (23) into (22) we obtain:

$$\Gamma_{qH}(x, y) = \frac{2x}{(1 - 2xy + \sqrt{1 - 4x^2})(y - xC(x))} \sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) (y^{h+1} - x^{h+1}C(x)^{h+1}(x))$$

195 which boils down to (17).

Finally, let $\pi \in \text{MinCo}(qD)$ and let π' be the smallest prefix of π containing q . Then

$$\pi = \pi' \alpha D$$

where α is, as before, a path starting at the origin, using only U and H steps, and with the additional restriction that, if π' ends at height $h = 0$, then $\alpha \neq H^r$, $r \geq 0$ (otherwise π would not be a Motzkin prefix, since it would terminate below the x -axis). In terms of generating functions, recalling the expression (20) for

200 the bivariate generating function of the paths α , we then have:

$$\begin{aligned} \Gamma_{qD}(x, y) &= \left(\sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) y^h A(x, y) \right) xy^{-1} - ([y^0] \Gamma_q(x, y)) \sum_{i \geq 0} x^i xy^{-1} \\ &= \frac{x}{y} A(x, y) \Gamma_q(x, y) - \frac{x}{y} \Gamma_q(x, 0) \frac{1}{1 - x} \\ &= \frac{x}{y} \left(\frac{1}{1 - x - xy} \Gamma_q(x, y) - \frac{1}{1 - x} \Gamma_q(x, 0) \right), \end{aligned}$$

which is equal to (18). \square

Let $\Delta_q(x)$ be the generating function of Motzkin paths avoiding a Motzkin prefix q with respect to the length. The following result gives a recursive procedure to compute $\Delta_q(x)$.

205 **Proposition 3.2.** For any Motzkin prefix q , the generating function $\Delta_q(x)$ is given by:

$$\Delta_\epsilon(x) = 0 \tag{24}$$

$$\Delta_{qD}(x) = \Delta_q(x) + \Gamma_q(x, 0) \frac{1}{1-x} \tag{25}$$

$$\Delta_{qH}(x) = \Delta_q(x) + C(x) \cdot \Gamma_q(x, xC(x)) \tag{26}$$

$$\Delta_{qU}(x) = \Delta_q(x) + \frac{1}{1-x} \Gamma_q\left(x, \frac{x}{1-x}\right). \tag{27}$$

Proof. Every Motzkin path contains the empty path ϵ , hence $\Delta_\epsilon(x) = 0$. Let $\pi \in \text{Av}(qD)$. There are two cases: either π avoids q , and such paths π are counted by $\Delta_q(x)$, or π contains q but avoids qD . In the latter case, let π' be the smallest prefix of π containing q . Obviously π' cannot be followed by any D step in any position, otherwise the path π would contain qD . Hence the only possibility is that π' has final height equal to 0 and is followed by a certain number of consecutive H steps. In other words, π can be factorized as

$$\pi = \pi' H^i,$$

with $i \geq 0$. In terms of generating functions, the above argument leads to:

$$\Delta_{qD}(x) = \Delta_q(x) + ([y^0] \Gamma_q(x, y)) \sum_{i \geq 0} x^i = \Delta_q(x) + \Gamma_q(x, 0) \frac{1}{1-x}$$

which is equation (25).

Suppose that $\pi \in \text{Av}(qH)$. If π also avoids q , then, as in the previous case, we obtain the generating function $\Delta_q(x)$. Otherwise, π can be decomposed as

its smallest prefix π' containing q , ending at height $h \geq 0$, followed by a path starting from height h , using only U and D steps and ending on the x -axis. This path is easily seen to be the reverse of a Dyck prefix having final height h , hence:

$$\Delta_{qH}(x) = \Delta_q(x) + \sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) ([y^h] \mathcal{DP}(x, y)) . \quad (28)$$

Since we have

$$[y^h] \mathcal{DP}(x, y) = \frac{2}{1 + \sqrt{1 - 4x^2}} \left(\frac{2x}{1 + \sqrt{1 - 4x^2}} \right)^h ,$$

plugging the above expression into (28), and observing that $\frac{2}{1 + \sqrt{1 - 4x^2}} = C(x)$, we get:

$$\Delta_{qH}(x) = \Delta_q(x) + C(x) \sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) (xC(x))^h$$

which is equivalent to (26).

Finally, let $\pi \in \text{Av}(qU)$. If π contains q , as usual let π' be the smallest prefix of π containing q . The path π can be written as π' , which ends at height $h \geq 0$, followed by a path starting from height h and using only D and H steps. This latter path is the reverse of a path α starting at the origin, using only U steps and H steps and ending at height h . Recalling once more the expression (20) of the bivariate generating function $A(x, y)$ of such paths, we obtain:

$$\Delta_{qU}(x) = \Delta_q(x) + \sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) ([y^h] A(x, y)) .$$

Since $[y^h] A(x, y) = \frac{1}{1-x} \left(\frac{x}{1-x} \right)^h$, we get:

$$\Delta_{qU}(x) = \Delta_q(x) + \frac{1}{1-x} \sum_{h \geq 0} ([y^h] \Gamma_q(x, y)) \left(\frac{x}{1-x} \right)^h ,$$

210 which is (27). □

As a consequence of Propositions 3.1 and 3.2, we get that, for a given pattern q , the generating function $\Delta_q(x)$ of Motzkin paths avoiding q is rational over x and $C(x)$, which is the statement of Theorem 1.1.

4. Conclusion

215 The main results of the present paper, namely an algorithm to determine the
generating function of Motzkin paths avoiding a set of patterns, and the proof
that such generating function is rational over x and $C(x)$ (at least in the case of
a single pattern), may be seen as a further step towards a deeper investigation of
pattern avoidance in lattice paths. For instance, the same approach developed
220 here can be pursued for Schröder paths, for which pattern avoidance has been
first studied in Cioni and Ferrari [10], thus getting completely analogous results
(in particular, the same technique described in Section 3 can be exploited to
show that the generating function of Schröder paths avoiding a single pattern is
also rational over x and $C(x)$). It would then be interesting to find analogous
225 results in the case of an arbitrary set of steps.

Another issue that seems worth investigation is the asymptotic behavior of
classes of pattern-avoiding Motzkin paths. In the case of Dyck paths, in Bacher
et al. [4] it is shown that, regardless of the specific pattern to be avoided, the
asymptotic behavior of all classes of Dyck paths avoiding a single pattern is
230 the same (and it is polynomial). Having a similar result for pattern-avoiding
Motzkin paths would be desirable.

There are some papers, such as Asinowski et al. [2] and Asinowski et al. [11],
which develop a methodology based on automata and a variant of the kernel
method to study lattice paths avoiding a *consecutive* pattern (that is a pattern
whose elements are adjacent in the path). It seems conceivable that a similar
235 approach could be fruitful also in the case of generic patterns.

References

- [1] R. Donaghey, L. W. Shapiro, Motzkin numbers, J. Combinatorial Theory
Ser. A 23 (3) (1977) 291–301.
- 240 [2] A. Asinowski, A. Bacher, C. Banderier, B. Gittenberger, Analytic combina-
torics of lattice paths with forbidden patterns, the vectorial kernel method,

- and generating functions for pushdown automata, *Algorithmica* 82 (2020) 386–428.
- [3] A. Sapounakis, I. Tasoulas, P. Tsikouras, Counting strings in Dyck paths,
245 *Discrete Math.* 307 (2007) 2909–2924.
- [4] A. Bacher, A. Bernini, L. Ferrari, B. Gunby, R. Pinzani, J. West, The Dyck pattern poset, *Discrete Math.* 321 (2014) 12–23.
- [5] P. Flajolet, R. Sedgewick, *Analytic combinatorics*, Cambridge University Press, Cambridge, 2009.
- 250 [6] N. J. A. Sloane, The on-line encyclopedia of integer sequences.
URL oeis.org
- [7] C. Bean, Finding structure in permutation sets, Ph.D. thesis, Reykjavik University (June 2018).
- [8] C. Bean, J. S. Eliasson, E. Nadeau, J. Pantone, H. Ulfarsson, Permutatri-
255 angle/comb_spec_searcher: Version 4.0.0 (June 2021).
URL <https://doi.org/10.5281/zenodo.4946832>
- [9] C. Bean, Permutatriangle/motzkin: Version 1.0.0 (August 2021).
URL <https://doi.org/10.5281/zenodo.5159935>
- [10] L. Cioni, L. Ferrari, Enumerative results on the Schröder pattern poset,
260 in: A. Dennunzio, E. Formenti, L. Manzoni, A. Porreca (Eds.), *Cellular Automata and Discrete Complex Systems. AUTOMATA 2017*, Vol. 10248 of *Lecture Notes in Computer Science*, Springer, Cham, 2017, pp. 56–67.
- [11] A. Asinowski, C. Banderier, V. Roitner, Generating functions for lattice paths with several forbidden patterns, *Sém. Lothar. Combin.* 84B (2020)
265 Article #95, 12 pp.