

SYSTEMATIC REVIEW

Community health services in European literature: A systematic review of their features, outcomes, and nursing contribution to care

Valeria Caponnetto RN, MSN, PhD, Assistant Professor¹  | Angelo Dante RN, MSN, PhD, Associate Professor¹  | Khadija El Aoufy RN, MSN, PhD, Research Fellow²  | Maria Ramona Melis RN, MSN, PhD³  | Giulia Ottonello MSN, RN, PhD^{4,5,6}  | Francesca Napolitano RN, MSN, PhD^{4,7}  | Fabio Ferraiuolo RN, MSN, PhD¹  | Francesco Camero RN, MSN, PhD¹  | Angela Cuoco RN, MSN, PhD^{8,9}  | Ilaria Erba RN, MSN, PhD, Research Fellow¹⁰  | Laura Rasero RN, CCN, MSN, Associate Professor^{2,11} | Loredana Sasso RN, MSN, MEdSc, FFNMRCISI, Full Professor, Director of Scientific Committee CERSI-FNOPI^{4,11}  | Annamaria Bagnasco RN, MSN, PhD, MEdSc, FFNMRCISI, Full Professor^{4,11}  | Rosaria Alvaro RN, MSN, FESC, FAAN, Full Professor, Pro-Rector^{8,11}  | Duilio Fiorenzo Manara RN, MSN, Associate Professor, Director of Center of Nursing Research and Innovation of Milan^{11,12}  | Gennaro Rocco RN, MSN, PhD, FAAN, FFNMRCISI, Director of Center of Excellence for Nursing Scholarship of Rome^{13,14}  | Maurizio Zega RN, MSN, PhD^{13,15} | Giancarlo Cicolini RN, MSN, PhD, Full Professor^{15,16}  | Beatrice Mazzoleni RN, MSN, PhD^{15,17}  | Loreto Lancia RN, MSN, Full Professor^{1,11} 

¹Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy²Department of Health Sciences, University of Florence, Florence, Italy³Staff Training Department, Careggi University Hospital, Florence, Italy⁴Department of Health Sciences, University of Genoa, Genoa, Italy⁵Ingram School of Nursing, McGill University, Montreal, Canada⁶Direction of Health Professionals, "IRCCS Istituto Giannina Gaslini," Genoa, Italy⁷Department of Emergency and Admission, Policlinic Hospital "IRCSS San Martino," Genoa, Italy⁸Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy⁹Orthopedic and Traumatology Clinic, Orthopedic Institute "IRCSS Rizzoli," Bologna, Italy¹⁰Bachelor of Science in Nursing, Saint Camillus International University of Health and Medical Sciences, Rome, Italy¹¹Scientific Committee CERSI-FNOPI, Federazione Nazionale Ordini Professioni Infermieristiche, Rome, Italy¹²Center for Nursing Research and Innovation, Vita-Salute San Raffaele University, Milan, Italy¹³Center of Excellence for Nursing Scholarship, Rome, Italy¹⁴Faculty of Medicine, University "Our Lady of the Good Counsel", Tirana, Albania¹⁵FNOPI, Federazione Nazionale Ordini Professioni Infermieristiche, Rome, Italy¹⁶Section of Nursing and Midwifery, Department of Innovative Technologies in Medicine & Dentistry, University "G. d'Annunzio" Chieti - Pescara, Chieti, Italy¹⁷Department of Biomedical Sciences, Humanitas University, Milan, Italy

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *International Nursing Review* published by John Wiley & Sons Ltd on behalf of International Council of Nurses.

Correspondence

Angelo Dante, Department of Health, Life, and Environmental Sciences, University of L'Aquila, Edificio Rita Levi Montalcini, Via Giuseppe Petroni 67100 L'Aquila, Italy.
Email: angelo.dante@univaq.it

Abstract

Background: To meet the population's needs, community care should be customized and continuous, adequately equipped, and monitored.

Introduction: Considering their fragmented and heterogeneous nature, a summary of community healthcare services described in European literature is needed. The aim of this study was to summarize their organizational models, outcomes, nursing contribution to care, and nursing-related determinants of outcomes.

Methods: A systematic review was performed by searching PubMed, CINAHL, Scopus, and Embase in October 2022 and October 2023 (for updated results). Quantitative studies investigating the effects of community care, including nursing contribution, on patient outcomes were included and summarized. Reporting followed the PRISMA checklist. The review protocol was registered on PROSPERO (CRD42022383856).

Results: Twenty-three studies describing six types of community care services were included, which are heterogeneous in terms of target population, country, interventions, organizational characteristics, and investigated outcomes. Heterogeneous services' effects were observed for access to emergency services, satisfaction, and compliance with treatment. Services revealed a potential to reduce rehospitalizations of people with long-term conditions, frail or older persons, children, and heart failure patients. Models are mainly multidisciplinary and, although staffing and workload may also have an impact on provided care, this was not enough investigated.

Discussion: Community health services described in European literature in the last decade are in line with population needs and suggest different suitable models and settings according to different care needs. Community care should be strengthened in health systems, although the influence of staffing, workload, and work environment on nursing care should be investigated by developing new management models.

Conclusions and implications for health policy: Community care models are heterogeneous across Europe, and the optimum organizational structure is not clear yet. Future policies should consider the impact of community care on both health and economic outcomes and enhance nursing contributions to care.

KEYWORDS

Community care services, home care services, nursing determinants, nursing homes, patient outcomes, primary care, remote health, systematic review, transitional care

INTRODUCTION

In the last two decades, there have been epidemiological changes, pharmaceutical and technological progress, financial restraints, and a lack of human resources all over the world. This has led to an increased incidence of noncommunicable diseases, longer life expectancy and years lived with disability, higher incidence of frailty (OECD, 2021; WHO, 2022a, 2022c; McKee & Healy, 2001), and a reduction in in-hospital length of stay (McKee & Healy, 2001; OECD, 2021). Noncommunicable diseases need special and multi-professional attention (OECD, 2021; WHO, 2022c, 2023), often after the patient's discharge from the hospital (Coffey et al., 2019; Veenstra & Gautun, 2021).

Most of a population's needs could be effectively managed in community care settings by ensuring the availability of good-quality healthcare services and adequately equipped facilities, such as general practice, family nursing, nursing homes, community hospitals, district care, continuity of care units, and palliative care networks (Davidson et al., 2019; Spasova et al., 2018). Current organizational models for com-

munity healthcare services across Europe are manifold, and there are still challenges that need to be overcome regarding care organization and evaluation (Spasova et al., 2018). Besides the lack of standardized care and quality indicators across countries (Joling et al., 2018; Veldhuizen et al., 2021), the main issues in this field include service fragmentation and scarcity (WHO, 2022b). Ideally, community care services should be person-centered, ensure continuity of care and availability of adequate equipment, and be evaluated over time (Spasova et al., 2018).

Considering some emerging population needs also in the aftermath of the COVID-19 pandemic, community healthcare services in Italy are undergoing a reorganization. In this process, nurses play a pivotal role in the organization and delivery of care (Ministero della Salute, 2022). Hence, the need to gain insights into this reorganization and evaluate nursing contribution in delivering community care is paramount. This would be ensured by a deep investigation of existing community care in Italy (Bagnasco et al., 2023), as well as features and outcomes of community health services documented in European literature.

Considering that knowledge of in-hospital optimum organizational models, staffing, and skill mix of healthcare professionals has been fundamental to a better understanding of care outcomes (Aiken et al., 2017), this review aimed to describe the organizational models of community healthcare documented in European literature, their outcomes on service users, the contribution of nursing to care, and nursing-related determinants of those outcomes.

Based on the available European literature, this systematic review addressed the following questions: (i) What are the main features of community healthcare services documented in European literature in the last decade? (ii) What are the outcomes of these services? (iii) What is the contribution of nursing to care in these services? (iv) What is the role of nursing-related determinants on community care service outcomes? By addressing these questions, this study would provide insights into services documented in European literature and their outcomes in the last decade, to inform future policies and research.

METHODS

Study design

The Italian Centre of Excellence for Research and Nursing Development (CERSI) conducted a systematic literature review based on recommendations of the Joanna Briggs Institute (JBI) Manual for Evidence Synthesis (Aromataris & Munn, 2020; Tufanaru et al., 2020), providing a report by following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist (Page et al., 2021).

Sources of information and search strategy

PICO (population, comparison, intervention, and outcome) was applied to conduct the electronic search strategy:

- Population: patients receiving community health services.
- Intervention: community health services where nurses operate either independently or as part of a multidisciplinary team. These services included primary care, district nursing, community nursing, family nursing, continuity of care, transitional care, home care, community hospital, palliative home care, and school health services. Intervention also accounted for nursing-related factors that could influence patient outcomes, i.e., staffing levels, nurse-to-patient ratios, skill and qualification mix, workload, caseload, and care models.
- Comparison: any type of health services provided as an alternative to the investigated community health service, including usual care. Additionally, intra-group comparisons were considered, i.e., outcome indicators changing after implementing the service detailed in the intervention. As an inclusive strategy, this component of the PICO was

not included in search strings to ensure including studies without a control group.

- Outcomes: quality indicators linked with nursing interventions or care models within community health services, i.e., unplanned rehospitalization, access to emergency medical services, missed care, self-care, empowerment, self-efficacy, therapeutic adherence, health literacy, and patient satisfaction.

To individuate community health services denominations and relevant quality indicators (outcomes), a discussion was undertaken among the authors, after independently consulting literature on this topic. Afterward, a preliminary search was performed on PubMed to retrieve search terms. On October 27, 2022, PubMed, CINAHL, Scopus, and Embase databases were searched for relevant studies conducted in the last 10 years (Supplementary Tables S1 and Table S2). Considering the broad coverage provided by these databases, no additional searches were done, which was contrary to the review protocol published in PROSPERO. Since study selection and data extraction lasted almost one year, on October 18, 2023, search strings were relaunched and studies published after October 27, 2022, were examined. The retrieved literature was managed using EndNote X7.8 (Thomson Reuters, New York). Where necessary, university libraries or paper authors were contacted to retrieve full texts.

Eligibility and inclusion criteria

The following inclusion criteria were considered both at the eligibility (title and abstract analysis) and inclusion (full-text examination) stages: quantitative original studies, published in peer-reviewed journals, conducted in European countries, published in English or Italian, examining the impact of community care models including the contribution of nursing to patient outcomes described in the PICO. Hence, all the considered community care included health services. The description of nursing-related factors influencing patient outcomes was not utilized as an inclusion criterion. Instead, these factors were summarized solely for studies reporting them. References with no abstract or full text available at the eligibility or inclusion stage were excluded.

Study selection, data extraction, and analysis

Two researchers independently examined titles, abstracts, and eligible full texts. They also extracted data using a standardized and piloted Microsoft Excel® template. This template included author, year of publication, country, research design, population, sample size, sample characteristics, study purpose, intervention(s), comparison (if any), outcome(s), measurement tools, nursing contribution to the model, nursing-related determinants of outcomes, statistical

methods, measures of association, and covariates. Nursing contribution was considered as part of a “community care” model, i.e., nurses working independently or in a multidisciplinary team (Ministero della Salute, 2022). Nursing-related determinants of patient outcomes (staffing level, nurse-to-patient ratio, skill mix, qualification mix, workload, caseload, and care model) were considered as an integral part of community nursing (Alghamdi, 2016; Butler et al., 2019; Cunningham et al., 2019; Ministero della Salute, 2021; Veldhuizen et al., 2021). Moreover, to provide a framework regarding summarized research, the economical features (i.e., whether public or private) of documented interventions were extracted, if reported by authors. For data extraction, disagreements were resolved through discussion with a third researcher.

When summarizing results, studies that reported similar services were clustered and described together under a distinctive label, closely resembling labels identified by the authors of the included manuscripts. The label was shaped by considering the settings for service provision and the main care needs of the patients, regardless of their underlying pathology. Services performed remotely using technology were included in a separate category. This approach was implemented to ensure summarizing together features and outcomes of similar interventions. Due to data heterogeneity, meta-analyses were not feasible. Consequently, general information of studies, types of community care services and features (setting, population, main purpose of the service, and whether the care model was multidisciplinary), outcomes on service users, the contribution of nursing to care, and nursing-related determinants of those outcomes, were described using tables and the narrative form. Results were described and critically discussed according to study design, type of comparison performed, and studies’ methodological quality. Each study was assigned an ID (Results section; Supplementary Table S3), which is cited when describing results.

Methodological quality

Two researchers independently evaluated the methodological quality of the included studies using the JBI critical appraisal tools (Aromataris & Munn, 2020). Disagreements were resolved through discussion with a third researcher. Considering the explorative nature of the review, methodological quality was not considered as an exclusion criterion; hence, low-quality studies were not excluded.

Ethical approval, informed consent, and registration

Ethical approval and informed consent were not applicable. The systematic review protocol was registered in PROSPERO (CRD42022383856) on December 25, 2022.

RESULTS

Overall, 8880 references were retrieved, and 7901 of them were examined after removing duplicates. A total of 266 full texts were examined and 23 studies were included (Augestad et al., 2020—A01; Bertelsen et al., 2017—A02; Campbell et al., 2015—A03; Carter-Stephens, 2020—A04; de Stampa et al., 2014—A05; Facultad & Lee, 2019—A06; Ferrara et al., 2015—A07; Fournaise et al., 2023—A08; González-Franco et al., 2022—A09; Karlsson et al., 2013—A10 and A11; McGloin et al., 2020—A12; Ng et al., 2014—A13; Phelan et al., 2018—A14; Profili et al., 2017—A15; Röhrich et al., 2017—A16; Senek et al., 2020—A17; Smits et al., 2020—A18; Vainieri et al., 2018—A19; Vianello et al., 2016—A20; Villani et al., 2014—A21; Zimmermann et al., 2016—A22; Zúñiga et al., 2015—A23) (Figure 1).

Characteristics of included studies

As given in Supplementary Table S3, included studies were conducted in the United Kingdom ($n = 6$, 26.1%) (A03, A04, A06, A13, A16, and A17), Italy ($n = 5$, 21.7%) (A07, A15, and A19–A21), Spain (A09, A11), Ireland (A12 and A14), Germany (A08 and A22) ($n = 2$, 8.7% each), and other European countries ($n = 6$, 26.1%) (A01, A02, A05, A10, A18, and A23). Years of publication ranged from 2013 to 2023.

Eleven studies (43.5%) were observational (A04, A07, A09, A10, A13–A17, A19, and A23), seven (30.4%) were experimental (A01–03, A08, and A20–22), three (13.0%) were quasi-experimental (A05, A11, and A18), two (9.7%) were descriptive (A06 and A19), and one (4.3%) was mixed-method (A12). Multicentric studies were less than half of the total included studies ($n = 11$, 47.8%) (A02, A03, A08–A11, A16, A17, A20, and A23). Sampling was consecutive ($n = 6$, 26.1%) (A05, A10–A12, A20, and A22), random ($n = 5$, 21.7%) (A02, A03, A19, A21, and A23), included the whole reference population ($n = 5$, 21.7%) (A06–A09, A18), convenient ($n = 1$, 4.4%) (A16), purposive ($n = 1$, 4.4%) (A14), or unclear ($n = 5$, 21.7%) (A01, A04, A13, A15, and A17).

Most of the studies included people with different diseases and characteristics ($n = 20$, 87.0%, total = 44 269 persons) (A01–A16, A18–A22), while three studies (13.0%) (A14, A17, and A23) included healthcare professionals (total = 6332). The mean number of participants was 2213.5 (SD = 4985.4, minimum = 33, maximum = 16 972) and 2110.7 (SD = 2037.2, minimum = 283, maximum = 4307) for users and professionals, respectively.

Types of community care services and related outcomes

Six types of community care services were identified (Table 1), namely, home care ($n = 8$, 34.8%) (A05–A11, and A13), remote health ($n = 5$, 21.7%) (A01, A03, A12, A20, and A21),




FIGURE 1 PRISMA flow diagram.

community health ($n = 4$, 17.4%) (A04, A14, A17, and A19), primary healthcare ($n = 4$, 17.4%) (A15, A16, A18, and A22), nursing homes (A23), and transitional care (A02) ($n = 1$, 4.5% both). All community services provided healthcare, although they were heterogeneous in terms of the target population, interventions, and organizational characteristics. Moreover, they were delivered at home (e.g., home care), in institutions (e.g., nursing homes), or through facilities and devices, with professionals (e.g., community health, primary healthcare, remote health, and transitional care) acting as a bridge between home setting and institutions. A detailed description of provided services and interventions is reported in Supplementary Table S4, while detailed and summarized descriptions of outcomes are reported in Supplementary Table S5 and Table 2, respectively.

Home care services

Home care services were provided at home by multidisciplinary teams for early identification of home care clients at risk of hospitalization (A08) and to manage frailty due to aging (A05, A07, and A10), long-term conditions (e.g., heart failure and diabetes) (A09 and A11), or special health needs of children and critically ill patients (A06 and A13). Multidisciplinary interventions consisted of comprehensive clinical assessment and care planning (A05 and A11), monitoring and

follow-up (A05, A06, A08–A11, and A13), patients' stabilization at home (A06, A09, A11, and A13), education and self-care promotion (A09, A11, and A13), and comprehensive clinical and social services (A07).

Nursing interventions included assessment and care planning (A05), home visits and consultations (A09, A10), supervision and educational activities (A13), care coordination (A05), and decision-making (A08), although they were not always fully described.

These services were largely provided to older people. In a quasi-experimental study, authors compared 105 very frail older people, who were cared for by a two-person case manager team (i.e., a nurse and a physician) supported by a geriatrician, and 323 similar persons receiving usual care. The experimental group showed lower unplanned hospital admissions after one year (adjusted OR = 0.39, 95%CI = 0.16–0.98) (A05). Also, in a large randomized controlled trial (RCT) involving 2464 older people and in which a prediction algorithm was applied by nurses to prevent hospital readmission, the intervention group showed a significantly lower incidence of readmissions compared with the usual care group (incident rate ratio IRR = 0.41, 95%CI = 0.24–0.68, $p = 0.0007$) (A08). Finally, a quasi-experimental study involved 101 older people affected by long-term conditions who were cared for through a multidisciplinary integrated care pathway comprising communication, care coordination, and home-based care, instead of usual care (A11). Results highlighted a significant reduction

TABLE 1 Type and general characteristics of community care services and number of studies including these services.

Service	No. of studies	Characteristics	Financial model (ID)
Home care	8	Encompass a comprehensive and personalized range of health care interventions provided by multidisciplinary teams to patients in their own homes. These services are generally provided to face chronic health issues, patients' frailty, or special health needs.	Public (A08, A05, A07, A10, A11, and A13) Not reported (A06 and A09)
Remote health	5	Health services delivered by healthcare professionals through electronic communication technologies. These services bridge the gap between geographically distant healthcare professionals and patients, enabling regular monitoring, assessment, and management of conditions, ensuring timely interventions and support, while promoting convenience, accessibility, patient engagement, self-care, and overall satisfaction with the healthcare experience.	Not reported (A01, A03, A12, A20, and A21)
Community health	4	Integrated and accessible healthcare services provided within local communities (e.g., clinics, districts, ambulatories), encompassing preventive care, patient education, and support to enhance health outcomes, satisfaction, and self-care.	Public (A14 and A17) Not reported (A04 and A19)
Primary health care	4	Comprehensive and person-centered healthcare services, generally coordinated or supervised by general practitioners, encompass a multidisciplinary approach, and are delivered in clinics or at patients' homes, ensuring holistic care aimed at promoting self-care, satisfaction, and reduce access to emergency services.	Public (A15) Not reported (A16, A18, and A22)
Nursing homes	1	Are residential facilities that offer a range of services for individuals requiring long-term care, short-term stays, adult day care, and post-acute care, providing a dedicated living environment for individuals with chronic or terminally ill conditions. With a focus on multidisciplinary interventions, including rehabilitation and nursing care, nursing homes strive to meet the needs of residents, enhance their quality of life, and ensure their well-being in a residential setting, catering to both their ongoing care and support requirements.	Mixed ^a (A23)
Transitional care	1	Encompass a coordinated and multidisciplinary approach to care, specifically targeting patients recently discharged from the hospital after experiencing acute health problems. These services aim to facilitate a seamless transition from the hospital to home or community settings by offering a range of interventions, including health education, physical exercise, smoking cessation support, dietary advice, clinical evaluations, medical visits, with the ultimate objective of enhancing patients' treatment adherence and optimizing their overall health outcomes throughout the transitional period.	Public (A02)

^a Mixed: Nursing homes financed with an approximately equal proportion of public, private, or privately subsidized (operating under private law but guaranteed by the public sector) systems.

in the average number of accesses to emergency services ($p \leq 0.001$) and repeated hospitalizations ($p = 0.008$).

Similar results were obtained in an observational study of 809 patients with heart failure, who were included in an intensive and protocolized follow-up program involving a comprehensive clinical approach, an in-person and telephone follow-up, educational interventions, and care in case of decompensation. A significantly lower risk of rehospitalizations due to heart failure was documented ($p < 0.001$) compared with usual care patients ($n = 2053$) (A09). Moreover, 33 acutely ill children were observed after being managed at home by a nursing matron who may consult with a pediatrician. A 17.3% reduction in repeated hospitalization and a 5% reduction in access to emergency services were observed, although no statistical tests were performed in this observational study (A13).

Another highly investigated outcome was user satisfaction. Among 417 older people included in a cross-sectional study to observe the outcome of the provision of home clinical and social services, 67.2% declared their trust in the model, with higher percentages among city residents (72.1%) compared with people living in peripheral areas (43.0%) (A07). Although overall user satisfaction is highly valued,

these services seem to be more appreciated by older people who receive them at home rather than in special accommodations ($p = 0.004$), as revealed in a cross-sectional study involving 166 older persons who received at least two home/accommodation visits by home care nurses (A10).

Finally, in other populations, satisfaction was high in 206 persons with various acute ill who received intensive care treatments at home (99.0% satisfied or very satisfied) (A06) and in 33 acutely ill children managed at home by a nursing matron who may consult with a pediatrician (94% described as excellent the care received and 6% as good) (A13).

Remote health services

These services were delivered by healthcare professionals using digital communication technologies to people affected by a variety of surgical and medical conditions (e.g., postsurgical stoma, heart failure, diabetes, and COPD) (A01, A03, A12, A20, and A21).

Vital signs and other clinical data were daily transmitted to multidisciplinary professionals (mainly physicians or nurses) who decided either to modify therapy, plan a nurse visit at

TABLE 2 Studies results according to the care model and study design.

Outcome		Home care	Remote health	Community health	Primary health	Nursing homes	Transitional care
Hard	Access to emergency	Quasi-experimental + Observational +	Randomized controlled trial –	Observational +	Observational –	–	–
	Repeated hospitalization	Randomized controlled trial + Quasi-experimental + Observational +	Randomized controlled trial +	–	–	–	–
	Hospital admission due to HF	Observational +	–	–	–	–	–
	Unplanned hospital admission	Quasi-experimental +	–	–	–	–	–
PROs	Adherence to treatment	–	Randomized Controlled trial +	–	–	–	Randomized controlled trial +
	Empowerment	–	Mixed-method +	–	–	–	–
	Patient satisfaction	Observational + Cross-sectional + Descriptive +	Randomized Controlled trial – Mixed-method +	Descriptive –	Quasi-experimental + Observational +	–	–
	Self-efficacy	–	–	–	Randomized controlled trial +	–	–
	Self-care	–	Randomized controlled trial –	Descriptive –	–	–	–
NROs	Missed nursing care associated with staffing	–	–	Cross-sectional +	–	Cross-sectional +	–

Abbreviations: NROs: nurses reported outcomes; PROs: patients reported outcomes.

Note: Each + indicates a study demonstrating improved outcome pre-post or compared with control.

Each – indicates a study demonstrating no outcome differences pre-post or compared with control.

home, or refer patients to specialists, including psychologists (A20 and A21). Main nursing interventions were follow-up through teleconsultation (A01, A03, and A12), remote monitoring of clinical conditions (A12, A20, and A21), and arranging appointments with medical specialists (A01 and A03). Other nursing activities included providing self-care recommendations (A03), collecting clients' requests (A21), performing clinical examinations (A01), and home visits (A20).

One RCT evaluated the effects of multidisciplinary telemonitoring on repeated hospitalizations and access to emergency in a sample of 230 persons with COPD. Results showed a positive impact of the service on repeated hospitalization (IRR usual care vs. intervention group = 0.46, 95% CI = 0.24–0.89) as opposed to access to emergency (IRR usual care vs. intervention group = 0.94, 95% CI = 0.76–1.18) (A20).

Compared with 3-monthly in-person visits, multidisciplinary telemonitoring showed better compliance to treat-

ment in a sample of 80 patients with chronic heart failure (intervention group = 3.8 ± 0.5 vs. control group = 3.0 ± 0.8 ; $p < 0.05$) (A21). Also, empowerment significantly improved in a sample of 39 patients affected by type II diabetes after including telemonitoring, led by a clinical nurse specialist, to usual care (intervention group pre vs. post at 3 months, mean difference = +0.72, $p < 0.001$) (A12). Compared with traditional hospital nurse-led follow-up, teleconsultation performed by stoma nurses did not impact differently on self-care in a sample of 110 persons with postsurgical stoma ($p = 0.825$) (A01).

Finally, client satisfaction was high (i.e., scored above 4 out of 5) for telemonitoring performed by a clinical nurse specialist on 39 patients with type II diabetes (A12), although another study involving 110 people with postsurgical stoma in an RCT found no differences between nursing teleconsultation and usual care in a hospital setting (good overall experience in the intervention group $n = 103$, 49.9% vs. control

group $n = 149$, 59.1%; $p = 1.000$) (A01). Also, in a trial including 16 211 persons in primary care, general practitioner-led telephone triage and usual care showed higher user satisfaction compared with nurse-led computer-supported telephone triage (mean difference: + 3.94, 95% CI = 1.88–5.99 and 2.60, 95% CI = 0.58–4.63, respectively) (A03).

Community health services

Community health services were delivered through community clinics, districts, outpatient services, or directly at home, if necessary (A04, A14, A17, and A19), to manage long-term conditions, such as diabetes and heart failure (A19), or to address the healthcare needs of people receiving enteral tube feeding at home (A04). These services included nursing care (A04, A14, A17, and A19), physiotherapy, dietetic recommendations, and medical specialist visits (A19). Where described, nursing interventions consisted of providing educational activities and support (A04 and A19), scheduling specialist visits and follow-ups (A19), and managing complications (A04).

In an observational study involving an unknown number of people receiving nurse-led enteral tube feeding, the authors reported a considerable reduction (–93%) in the number of accesses to emergency services, which translated to better user experience (A04). In a descriptive study involving 1300 patients with single and multiple long-term conditions, a chronic care model, compared with standard care, did not influence patient satisfaction (people with single long-term conditions OR = 0.91, $p = 0.63$; people with multiple long-term conditions OR = 1.09, $p = 0.61$) or self-care (people with single long-term conditions OR = 1.88, $p = 0.23$; people with multiple long-term conditions OR = 0.53, $p = 0.12$). The model was introduced in community services and applied by a multidisciplinary team (GP, nurses, physiotherapists, dieticians, and medical specialists). The authors documented that during the observational period of the project, changes occurred in the GPs' practice based on the model itself (A19).

Finally, two cross-sectional studies involving 283 and 1742 community/public health nurses documented a significant association between organizational factors or staffing and missed care (A14) and an increased prevalence of missed nursing care in understaffed services (A17). Younger community nurses were more likely to miss several direct and indirect nursing activities ($p < 0.05$) and those without a bachelor's degree were more likely to miss "report writing" ($p < 0.05$). Moreover, detected differences in missed nursing care between adequately staffed and understaffed services ranged from +5.1% to +15.5% (A17).

Primary healthcare services

Primary healthcare included multidisciplinary services coordinated or supervised by general practitioners (A15, A16, A18,

and A22). These services were generally delivered, in clinics or at home (if required), to persons affected by mental disorders (A16, A22), long-term conditions (A15), or low-complexity health problems (A18).

Due to the diverse target populations that require these services, multidisciplinary and nursing interventions were heterogeneous. Educational programs (A15 and A16), activities promoting self-management of disease (A15, A22), care planning (A15, A16), as well as monitoring and follow-up (A15, A16, A22), were the most provided multidisciplinary interventions. Nurses were responsible for measuring vital signs and planning routine services (A15), administering medications, scheduling lifestyle services, educating patients (A16), and visiting them at home (A18).

A chronic care model for 16 972 type II diabetic patients comprising multiprofessional intervention, active patient involvement in the educational program, personalized evidence-based care, and scheduled follow-up showed no differences in accessing emergency services (IRR = 0.94, 95% CI = 0.84–1.01) compared to traditional care based on the essential level of assistance (A15). In an RCT involving 325 patients with psychiatric disorders, a primary care-based complex intervention to promote self-management and enhance self-efficacy demonstrated a significant increase in patient self-efficacy when compared with usual care led by general practitioners ($p = 0.004$) (A22).

Finally, clients with low-complexity health problems showed higher satisfaction ($n = 217$) when receiving evening and weekend home visits by trained nurses compared with 412 clients receiving visits only from their general practitioners (8.6/10, 95%CI = 8.5–8.8; vs. 8.3/10 95%CI = 8.2–8.4) (A18). Also, 126 psychiatric patients involved in an enhanced primary care pathway based on a multidisciplinary collaboration between healthcare providers and educational activities reported high satisfaction (from 8.8/10 to 9.4/10) (A16).

Nursing homes

Nursing homes were investigated only in one cross-sectional study including 4307 direct care workers employed in 156 nursing homes in Switzerland. They included a variety of residential services offering long-term care, short stays, adult day care, and post-acute care (A23). Nursing homes were accessed mainly by people with dementia, older people with psychiatric disorders, or persons needing palliative care. Even if few details were reported in the study, these services were based on multidisciplinary interventions, such as rehabilitation and nursing care. Results showed a significant relationship between the adequacy of staffing levels, workload, and a positive healthcare environment with implicit rationing of nursing care ($p < 0.05$) (A23).

Transitional care services

Transitional care services were assessed in one RCT conducted in Denmark on a sample of 212 patients exposed to shared care cardiac rehabilitation after acute coronary syndrome. The services involved a set of multidisciplinary interventions to ensure continual care for patients discharged from the hospital and included health education, physical exercise, abstinence from smoking, dietary advice, clinical evaluation, and medical visits (A02). Specifically, they were provided after hospitalization due to acute coronary syndrome in community services or in hospital outpatient clinics to improve adherence to treatment. Compared to patients exposed to a hospital cardiac-based rehabilitation program, no significant difference in outcome was reported (RR = 0.98, 95% CI = 0.73–1.32) (A02).

Nursing determinants

The most frequently reported nursing-related determinant of outcomes was skill mix ($n = 7$, 30.4%) (A01, A02, A04, A05, A07, A12, and A22), followed by staffing models ($n = 6$, 26.1%) (A01, A04, A06, A07, A22, and A23), type of contract ($n = 3$, 13.0%) (A16, A14, and A22), caseload ($n = 3$, 13.0%) (A14, A22, and A23), and nurse-to-patient ratio ($n = 2$, 8.7%) (A05, A22). Workload was not always reported in the included studies, but when it was, the authors did not always describe its association with patient outcomes (A14 and A17) (Supplementary Table S6).

Methodological quality

Methodological quality was rated from low to high in the included studies, with high variability among studies according to the study design. The main concerns were related to treatment concealment and blinding of patients, providers, and outcome assessors in RCTs; assessment of confounding factors and handling of incomplete follow-up in quasi-experimental studies; and standardization of outcome assessment and handling of confounding factors in cross-sectional studies (Supplementary Table S7).

DISCUSSION

Considering the lessons learned from the COVID-19 pandemic and the need to improve the quality of community care services, we performed this review to summarize the organizational models of community healthcare documented in European literature, their outcomes on service users, the contribution of nursing to care, and nursing-related determinants of outcomes reported. The ultimate goal of this review was to contribute to the reorganization of community ser-

vices in Italy (Ministero della Salute, 2022) and inform future European policies.

The results highlight a stable scientific production on the topic over the last decade, except for the year 2020 (the COVID-19 year), when more than one-quarter (26.1%) of included studies were published, perhaps pushed by the need to find innovative and more resilient models to provide community care. Studies were conducted mainly in the United Kingdom and Italy, with the rest conducted in other European countries, highlighting the need for further studies in various contexts.

Almost half of the studies were multicentric, ensuring good representativeness of involved contexts. Finally, despite the limited number of experimental studies, only a quite small number of included studies utilized unclear or high-risk sampling methods (30.4%). Nevertheless, the results of this review should be interpreted and generalized with caution since some observational studies showed methodological weaknesses, especially regarding the management of confounding factors. As a further limitation, it should be considered that evaluated outcomes do not encompass all those that may be influenced by community care services. To gather a comprehensive overview of the impact of these services on outcomes, quantitative findings should be integrated with qualitative research involving stakeholders, as this can highlight relevant aspects or factors. Examples of such factors include team dynamics, features of team-patient relationship, potential health system barriers, patient caseload, and attitudes of team members (Lalani et al., 2019; Leask et al., 2020). Lastly, study heterogeneity did not allow meta-analysis of data. Instead, the strengths of this review include rigorous methodology based on international standards, which ensured internal consistency, reproducibility, and accuracy of results.

Community care documented in European literature tends to be very complex and heterogeneous. Different types of services were described, although these services could be clustered considering the provision setting and main needs of patients accessing them.

Almost all studies reporting financial coverage details (Table 1, $n = 10$) were free of charge for patients. Even if not explicitly stated, it is reasonable to assume that in the other included studies, services were also free of charge as they were provided within the context of experimental research and/or in countries known for their universal healthcare coverage. This made it difficult to assess differences according to contextual factors, such as economic coverage (public or private), regarding outcomes. Hence, it is advisable that policymakers develop services informed by the results of this review, local needs, and available resources. Moreover, policymakers should endorse additional research on health and economic outcomes of services addressing predominantly emerging needs within their local context, fostering a direct and reciprocal exchange between research and practice.

In line with emerging population needs (OECD, 2021; WHO, 2022a, 2022c) and technological advancements, most of the investigated services were home care and remote health. Moreover, almost all services were multidisciplinary,

though including different activities and tasks performed by nurses, such as assessment and care planning, home visits and consultations, support and education, care coordination, continuous follow-up, administration of medications, and monitoring of clinical conditions. Most of the studies refer to very specific types of patients and each type of service often refers to the same type of patients. For instance, home care services were mainly explored in regard to older people, remote services mainly targeted patients with specific diseases, such as heart failure or COPD, and primary care services mainly considered low-complexity patients. Outcomes included “hard” indicators (i.e., access to emergency services, repeated hospitalization, hospital readmissions due to heart failure, and unplanned hospital admission) and “soft” indicators reported by either users (i.e., satisfaction, therapeutic adherence, empowerment, self-efficacy, and self-care) or professionals (i.e., missed nursing care). The impact of community care on access to emergency services, satisfaction, and adherence to treatment remains unclear. Community care with significant nursing contribution reduced rehospitalization of people with long-term conditions, frail or older persons, children, and patients with heart failure. Through different delivery types and modalities, appropriate community care prevents unnecessary access to and treatment in hospitals (MacNeill et al., 2021). Considering the pivotal role of clinical surveillance and early interventions, patient and caregiver education is fundamental as patients and caregivers are required to have high autonomy and self-care abilities to manage a disease.

Factors confirmed to reduce repeated hospitalization in community care settings include continuous follow-up and monitoring (Masotta et al., 2024). Other factors that appear to reduce this outcome are home visits and clinical stabilization at home, education on self-care, and empowerment programs. An additional strategy not delineated in the included studies, but which has demonstrated the potential to enhance both public health management and clinical outcomes, is nursing prescription (Laurant et al., 2018; Maier, 2019; Weeks et al., 2016). In this context, the World Health Organization advocates for the adoption of strategies that promote collaboration among healthcare professionals to address prescription requirements (WHO, 2020). This approach may enable nurses to fully utilize their professional capabilities while optimizing both patient and system outcomes (WHO, 2020).

Similar to in-hospital settings (Chiappinotto et al., 2022; Imam et al., 2023), missed nursing care seems to be associated with nurse staffing, workload, and work environment, although this relationship has not been investigated adequately.

Implications for nursing and health policy

Community care with significant nursing contribution should be largely boosted across Europe, while investigating its outcomes, in order to individuate the optimum organizational structure in each context and for each target population. In

fact, available data suggest that each target population may be effectively cared for in specific settings and through organized ad hoc services.

Organizational and nursing determinants should be accurately considered and investigated when developing community services, considering the impact of multidisciplinary and nursing interventions on population outcomes.

CONCLUSIONS

In the last decade, research on community care in Europe has been undertaken through complex and heterogeneous services targeting different types of patients, and by evaluating heterogeneous outcomes. In this complex context, the following main insights have been revealed:

- Community services are organized according to a specific type of setting, population needs, or technology used.
- Target populations considered in the literature are in line with worldwide epidemiology and worldwide unmet population needs.
- Community services can be multidisciplinary or delivered by one professional.
- Community services have a positive effect on rehospitalization, while conflicting results were retrieved for other outcomes.
- Staffing, workload, and work environment of community services appear to influence nursing care, although this relationship is not sufficiently investigated in the literature.

Future research should include both hard and soft outcomes and investigate their relationship with the organizational determinants of care. In addition, an improvement in the methodological quality of studies is recommended.

AUTHOR CONTRIBUTIONS

Study design: Angelo Dante, Laura Rasero, Loredana Sasso, Annamaria Bagnasco, Rosaria Alvaro, Duilio Fiorenzo Manara, Gennaro Rocco, Maurizio Zega, Giancarlo Cicolini, Beatrice Mazzoleni, and Loreto Lancia. *Data collection:* Valeria Caponnetto, Angelo Dante, Khadija El Aoufy, Maria Ramona Melis, Giulia Ottonello, Francesca Napolitano, Fabio Ferraiuolo, Francesco Camero, Angela Cuoco, and Ilaria Erba. *Data analysis:* Valeria Caponnetto and Angelo Dante. *Study supervision:* Angelo Dante, Laura Rasero, Loredana Sasso, Annamaria Bagnasco, Rosaria Alvaro, Duilio Fiorenzo Manara, Gennaro Rocco, Maurizio Zega, Giancarlo Cicolini, Beatrice Mazzoleni, and Loreto Lancia. *Manuscript writing:* Valeria Caponnetto, Angelo Dante, Khadija El Aoufy, Maria Ramona Melis, Giulia Ottonello, Francesca Napolitano, Fabio Ferraiuolo, Francesco Camero, Angela Cuoco, and Ilaria Erba. *Critical revisions for important intellectual content:* Angelo Dante, Valeria Caponnetto, Laura Rasero, Loredana Sasso, Annamaria Bagnasco, Rosaria Alvaro, Duilio Fiorenzo Manara, Gennaro Rocco, Maurizio Zega, Giancarlo Cicolini, Beatrice Mazzoleni, and Loreto Lancia.

ACKNOWLEDGMENTS

The authors would like to express their gratitude for their invaluable external support in bringing this project to fruition to: Michela Calzolari and Maria Emma Musio from the University of Genova; Luca Bertocchi, Alessia Marcotullio, and Vittorio Masotta from the University of L'Aquila; Maddalena De Maria, Mariachiara Figura, Loredana Piervisani, Gianluca Pucciarelli, Francesco Zaghini, and Valentina Zeffiro from the University of Rome "Tor Vergata." The authors also thank Giuseppe Aleo from the University of Genova for language editing of the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

FUNDING INFORMATION

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

DATA AVAILABILITY STATEMENT

The authors make the original data available upon reasonable request.

ORCID

Valeria Caponnetto RN, MSN, PhD  <https://orcid.org/0000-0003-4573-6110>

Angelo Dante RN, MSN, PhD  <https://orcid.org/0000-0001-5875-3353>

Khadija El Aoufy RN, MSN, PhD  <https://orcid.org/0000-0003-3278-8296>

Maria Ramona Melis RN, MSN, PhD  <https://orcid.org/0000-0002-4720-345X>

Giulia Ottonello MSN, RN, PhD  <https://orcid.org/0000-0002-3785-7847>

Francesca Napolitano RN, MSN, PhD  <https://orcid.org/0000-0002-9204-0899>

Fabio Ferraiuolo RN, MSN, PhD  <https://orcid.org/0000-0003-2203-8715>

Francesco Camero RN, MSN, PhD  <https://orcid.org/0009-0003-0070-3494>

Angela Cuoco RN, MSN, PhD  <https://orcid.org/0000-0002-2631-6287>

Ilaria Erba RN, MSN, PhD  <https://orcid.org/0000-0002-0756-1949>

Loredana Sasso RN, MSN, MEdSc, FFMRCISI  <https://orcid.org/0000-0001-5886-5937>

Annamaria Bagnasco RN, MSN, PhD, MEdSc, FFMRCISI  <https://orcid.org/0000-0002-9079-8460>

Rosaria Alvaro RN, MSN, FESC, FAAN  <https://orcid.org/0000-0002-4659-1569>

Duilio Fiorenzo Manara RN, MSN  <https://orcid.org/0000-0001-5182-6994>

Gennaro Rocco RN, MSN, PhD, FAAN, FFMRCISI  <https://orcid.org/0000-0003-3508-2352>

Giancarlo Cicolini RN, MSN, PhD  <https://orcid.org/0000-0002-2736-1792>

Beatrice Mazzoleni RN, MSN, PhD  <https://orcid.org/0000-0002-3870-1238>

Loreto Lancia RN, MSN  <https://orcid.org/0000-0002-2345-1731>

REFERENCES

- Aiken, L.H. et al. (2017) Nursing skill mix in European hospitals: cross-sectional study of the association with mortality, patient ratings, and quality of care. *BMJ Quality & Safety*, 26, 559–568. <https://doi.org/10.1136/bmjqs-2016-005567>
- Alghamdi, M.G. (2016) Nursing workload: a concept analysis. *Journal of Nursing Management*, 24, 449–457. <https://doi.org/10.1111/jonm.12354>
- Aromataris, E. & Munn, Z. (2020) JBI manual for evidence synthesis. Available from: <https://synthesismanual.jbi.global> [Accessed 14th December 2023].
- Augustad, K.M., Sneve, A.M. & Lindsetmo, R.O. (2020) Telemedicine in post-operative follow-up of STOMA Patients: a randomized clinical trial (the STOMPA trial). *British Journal of Surgery*, 107, 509–518. <https://doi.org/10.1002/bjs.11491>
- Bagnasco, A., Alvaro, R., Lancia, L., Manara, D.F., Zega, M., Rocco, G., Rasero, L., Mazzoleni, B. & Sasso, L. (2023) AIDOMUS-IT Collaborator Group. Protocol for evaluating quality and safety for the public through home care nursing in Italy: a multicentre cross-sectional descriptive observational study (AIDOMUS-IT). *British Medical Journal Open*, 13, e071155. <https://doi.org/10.1136/bmjopen-2022-071155>
- Bertelsen, J.B. et al. (2017) Cardiac rehabilitation after acute coronary syndrome comparing adherence and risk factor modification in a community-based shared care model versus hospital-based care in a randomised controlled trial with 12 months of follow-up. *European Journal of Cardiovascular Nursing*, 16, 334–343. <https://doi.org/10.1177/1474515116666781>
- Butler, M. et al. (2019) Hospital nurse-staffing models and patient- and staff-related outcomes. *Cochrane Database of Systematic Reviews*, 4, Cd007019. <https://doi.org/10.1002/14651858.CD007019.pub3>
- Campbell, J.L. et al. (2015) The clinical effectiveness and cost-effectiveness of telephone triage for managing same-day consultation requests in general practice: a cluster randomised controlled trial comparing general practitioner-led and nurse-led management systems with usual care (the ESTEEM trial). *Health Technology Assessment*, 19, 1–212, vii–viii. <https://doi.org/10.3310/hta19130>
- Carter-Stephens, R. (2020) Improving the experience of patients who receive home enteral tube feeding. *British Journal of Community Nursing*, 25, 178–183. <https://doi.org/10.12968/bjcn.2020.25.4.178>
- Chiappinotto, S. et al. (2022) Antecedents of unfinished nursing care: a systematic review of the literature. *BMC Nursing*, 21, 137. <https://doi.org/10.1186/s12912-022-00890-6>
- Coffey, A. et al. (2019) Interventions to promote early discharge and avoid inappropriate hospital (Re)admission: a systematic review. *International Journal of Environmental Research and Public Health*, 16, 2457. <https://doi.org/10.3390/ijerph16142457>
- Cunningham, J., O'Toole, T., White, M. & Wells, J.S.G. (2019) Conceptualizing skill mix in nursing and health care: an analysis. *Journal of Nursing Management*, 27, 256–263. <https://doi.org/10.1111/jonm.12673>
- Davidson, D. et al. (2019) Health services and delivery research. *Analysis of the profile, characteristics, patient experience and community value of community hospitals: a multimethod study*. Southampton, UK: NIHR Journals Library.
- de Stampa, M. et al. (2014) Impact on hospital admissions of an integrated primary care model for very frail elderly patients. *Archives of Gerontology and Geriatrics*, 58, 350–355. <https://doi.org/10.1016/j.archger.2014.01.005>
- Facultad, J. & Lee, G.A. (2019) Patient satisfaction with a hospital-in-the-home service. *British Journal of Community Nursing*, 24, 179–185. <https://doi.org/10.12968/bjcn.2019.24.4.179>

- Ferrara, M. et al. (2015) Quality of life and level of satisfaction of elderly persons receiving home health care: a cross-sectional study. *Igiene e Sanità Pubblica*, 71, 39–50.
- Fournaise, A. et al. (2023) Structured decision support to prevent hospitalisations of community-dwelling older adults in Denmark (PATINA): an open-label, stepped-wedge, cluster-randomised controlled trial. *The Lancet Healthy Longevity*, 4, e132–e142. [https://doi.org/10.1016/s2666-7568\(23\)00023-5](https://doi.org/10.1016/s2666-7568(23)00023-5)
- González-Franco, Á. et al. (2022) Morbidity and mortality in elderly patients with heart failure managed with a comprehensive care model vs. usual care: the UMIPIC program. *Revista Clínica Española*, 222, 123–130. <https://doi.org/10.1016/j.rceng.2021.05.007>
- Imam, A. et al. (2023) Missed nursing care in acute care hospital settings in low-income and middle-income countries: a systematic review. *Human Resources for Health*, 21, 19. <https://doi.org/10.1186/s12960-023-00807-7>
- Joling, K.J. et al. (2018) Quality indicators for community care for older people: a systematic review. *PLoS ONE*, 13, e0190298. <https://doi.org/10.1371/journal.pone.0190298>
- Karlsson, S., Edberg, A.K., Jakobsso, U. & Hallberg, I.R. (2013) Care satisfaction among older people receiving public care and service at home or in special accommodation. *Journal of Clinical Nursing*, 22, 318–330. <https://doi.org/10.1111/jocn.12115>
- Lalani, M., Fernandes, J., Fradgley, R., Ogunsola, C. & Marshall, M. (2019) Transforming community nursing services in the UK; lessons from a participatory evaluation of the implementation of a new community nursing model in East London based on the principles of the Dutch Buurtzorg model. *BMC Health Services Research*, 19(1), 945. <https://doi.org/10.1186/s12913-019-4804-8>
- Laurant, M. et al. (2018) Nurses as substitutes for doctors in primary care. *The Cochrane Database for Systematic Reviews*, 7(7), CD001271. <https://doi.org/10.1002/14651858.CD001271.pub3>
- Leask, C.F., Bell, J. & Murray, F. (2020) Acceptability of delivering an adapted Buurtzorg model in the Scottish care context. *Public Health*, 179, 111–117. <https://doi.org/10.1016/j.puhe.2019.10.011>
- MacNeill, A.J., McGain, F. & Sherman, J.D. (2021) Planetary health care: a framework for sustainable health systems. *The Lancet Planetary Health*, 5, e66–e68. [https://doi.org/10.1016/S2542-5196\(21\)00005-X](https://doi.org/10.1016/S2542-5196(21)00005-X)
- Maier, C.B. (2019) Nurse prescribing of medicines in 13 European countries. *Human Resources for Health*, 17(1), 95. <https://doi.org/10.1186/s12960-019-0429-6>
- Masotta, V. et al. (2024) Telehealth care and remote monitoring strategies in heart failure patients: a systematic review and meta-analysis. *Heart & Lung*, 64, 149–167. <https://doi.org/10.1016/j.hrtlng.2024.01.003>
- Mateo-Abad, M. et al. (2020) Impact of the care well integrated care model for older patients with multimorbidity: a quasi-experimental controlled study in the Basque country. *BMC Health Services Research*, 20, 613. <https://doi.org/10.1186/s12913-020-05473-2>
- McGloin, H. et al. (2020) Patient empowerment using electronic telemonitoring with telephone support in the transition to insulin therapy in adults with type 2 diabetes: observational, pre-post, mixed methods study. *Journal of Medical Internet Research*, 22, e16161. <https://doi.org/10.2196/16161>
- McKee, M. & Healy, J. (2001) The changing role of the hospital in Europe: causes and consequences. *Clinical Medicine*, 1, 299–304. <https://doi.org/10.7861/clinmedicine.1-4-299>
- Ministero della Salute. (2021) Programma Nazionale Esiti. Available at: https://pne.agenas.it/assets/documentazione/report/agenas_pne_report_2021.pdf
- Ministero della Salute. (2022) Decreto 23 maggio 2022, n. 77. Regolamento recante la definizione di modelli e standard per lo sviluppo dell'assistenza territoriale nel Servizio sanitario nazionale. GU Serie Generale n.144 del 22-06-2022. Available at: <https://www.gazzettaufficiale.it/eli/id/2022/06/22/22G00085/sg>
- Ng, S.M., Mariguzzi, S., Coward, S. & Middleton, H. (2014) Paediatric community home nursing: a model of acute care. *British Journal of Nursing*, 23, 209–212. <https://doi.org/10.12968/bjon.2014.23.4.209>
- Organisation for Economic Cooperation and Development (OECD). (2021) Health at a glance 2021. Available at: https://www.oecd-ilibrary.org/social-issues-migration-health/health-at-a-glance-2021_ae3016b9-en
- Page, M.J. et al. (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Journal of Clinical Epidemiology*, 134, 178–189. <https://doi.org/10.1016/j.jclinepi.2021.03.001>
- Phelan, A., McCarthy, S. & Adams, E. (2018) Examining missed care in community nursing: a cross-section survey design. *Journal of Advanced Nursing*, 74, 626–636. <https://doi.org/10.1111/jan.13466>
- Profilì, F. et al. (2017) Changes in diabetes care introduced by a chronic care model-based programme in Tuscany: a 4-year cohort study. *European Journal of Public Health*, 27, 14–19. <https://doi.org/10.1093/eurpub/ckw181>
- Röhrich, F. et al. (2017) Implementation of a novel primary care pathway for patients with severe and enduring mental illness. *BJPsych Bull*, 41, 314–319. <https://doi.org/10.1192/pb.bp.116.055830>
- Senek, M., et al. (2020) Nursing care left undone in community settings: results from a UK cross-sectional survey. *Journal of Nursing Management*, 28, 1968–1974. <https://doi.org/10.1111/jonm.12995>
- Smits, M., et al. (2020) Substitution of general practitioners by nurse practitioners in out-of-hours primary care home visits: a quasi-experimental study. *International Journal of Nursing Studies*, 104, 103445. <https://doi.org/10.1016/j.ijnurstu.2019.103445>
- Spasova, S., Baeten, R. & Vanhercke, B. (2018) Challenges in long-term care in Europe. *Eurohealth*, 24, 7–12.
- Tufanaru, C. et al. (2020) Chapter 3: systematic reviews of effectiveness. JBI manual for evidence synthesis. JBI; 2020. Available from: <https://synthesismanual.jbi.global>
- Vainieri, M. et al. (2018) Reported experience of patients with single or multiple chronic diseases: empirical evidence from Italy. *BMC Health Services Research*, 18, 659. <https://doi.org/10.1186/s12913-018-3431-0>
- Veenstra, M. & Gautun, H. (2021) Nurses' assessments of staffing adequacy in care services for older patients following hospital discharge. *Journal of Advanced Nursing*, 77, 805–818. <https://doi.org/10.1111/jan.14636>
- Veldhuizen, J.D. et al. (2021) Evidence-based interventions and nurse-sensitive outcomes in district nursing care: a systematic review. *International Journal of Nursing Studies Advances*, 3, 100053. <https://doi.org/10.1016/j.ijnsa.2021.100053>
- Veldhuizen, J.D. et al. (2021) Nurse-sensitive outcomes in district nursing care: a Delphi study. *PLoS ONE*, 16, e0251546. <https://doi.org/10.1371/journal.pone.0251546>
- Vianello, A. et al. (2016) Home telemonitoring for patients with acute exacerbation of chronic obstructive pulmonary disease: a randomized controlled trial. *BMC Pulmonary Medicine*, 16, 157. <https://doi.org/10.1186/s12890-016-0321-2>
- Villani, A. et al. (2014) Clinical and psychological telemonitoring and telecare of high risk heart failure patients. *Journal of Telemedicine and Telecare*, 20, 468–475. <https://doi.org/10.1177/1357633x14555644>
- Weeks, G., George, J., Maclure, K. & Stewart, D. (2016) Non-medical prescribing versus medical prescribing for acute and chronic disease management in primary and secondary care. *The Cochrane Database of Systematic Reviews*, 11(11), CD011227. <https://doi.org/10.1002/14651858.CD011227>
- World Health Organization (WHO). (2023) Noncommunicable diseases. Geneva: World Health Organization. Available at: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases> [Accessed 30th November 2023].
- World Health Organization (WHO). (2022a) Ageing and health. Geneva: World Health Organization. Available at: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health> [Accessed 30th November 2023].
- World Health Organization (WHO). (2022b) Long-term care. Geneva: World Health Organization. Available at: <https://www.who.int/europe/news-room/questions-and-answers/item/long-term-care> [Accessed 14th December 2023].
- World Health Organization (WHO). (2022c) World health statistics 2022: monitoring health for the SDGs, sustainable development goals. Geneva: World Health Organization; 2022. Available at: <https://www.who.int/publications/i/item/9789240051157>



- World Health Organization (WHO). (2020) State of the world's nursing 2020: investing in education, jobs and leadership. Geneva: World Health Organization; 2020. Available at: <https://iris.who.int/bitstream/handle/10665/331677/9789240003279-eng.pdf?sequence=1> [Accessed 24th January 2024].
- Zimmermann, T. et al. (2016) Collaborative nurse-led self-management support for primary care patients with anxiety, depressive or somatic symptoms: cluster-randomised controlled trial (findings of the SMADS study). *International Journal of Nursing Studies*, 63, 101–111. <https://doi.org/10.1016/j.ijnurstu.2016.08.007>
- Zúñiga, F. et al. (2015) The relationship of staffing and work environment with implicit rationing of nursing care in Swiss nursing homes: a cross-sectional study. *International Journal of Nursing Studies*, 52, 1463–1474. <https://doi.org/10.1016/j.ijnurstu.2015.05.0>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Caponnetto, V., Dante, A., El Aoufy, K., Melis, M.R., Ottonello, G., Napolitano, F. et al. (2024) Community health services in European literature: A systematic review of their features, outcomes, and nursing contribution to care. *International Nursing Review*, 71, 716–728. <https://doi.org/10.1111/inr.13033>