



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Full rotational symmetry from reflections or rotational symmetries in finitely many subspaces

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Full rotational symmetry from reflections or rotational symmetries in finitely many subspaces / Gabriele Bianchi; Richard J. Gardner; Paolo Gronchi. - In: INDIANA UNIVERSITY MATHEMATICS JOURNAL. - ISSN 0022-2518. - STAMPA. - 71:(2022), pp. 767-784. [10.1512/iumj.2022.71.9818]

Availability:

This version is available at: 2158/1257516 since: 2024-12-14T15:49:27Z

Published version:

DOI: 10.1512/iumj.2022.71.9818

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

(Article begins on next page)

May 6, 2022

FULL ROTATIONAL SYMMETRY FROM REFLECTIONS OR ROTATIONAL SYMMETRIES IN FINITELY MANY SUBSPACES

GABRIELE BIANCHI, RICHARD J. GARDNER, AND PAOLO GRONCHI

ABSTRACT. Two related questions are discussed. The first is when reflection symmetry in a finite set of i -dimensional subspaces, $i \in \{1, \dots, n-1\}$, implies full rotational symmetry, i.e., the closure of the group generated by the reflections equals $O(n)$. For $i = n-1$, this has essentially been solved by Burchard, Chambers, and Dranovski, but new results are obtained for $i \in \{1, \dots, n-2\}$. The second question, to which an essentially complete answer is given, is when (full) rotational symmetry with respect to a finite set of i -dimensional subspaces, $i \in \{1, \dots, n-2\}$, implies full rotational symmetry, i.e., the closure of the group generated by all the rotations about each of the subspaces equals $SO(n)$. The latter result also shows that a closed set in \mathbb{R}^n that is invariant under rotations about more than one axis must be a union of spheres with their centers at the origin.

1. INTRODUCTION

The question of when symmetry in a restricted set of subspaces implies full rotational symmetry arises naturally in several contexts. Radial functions are particularly amenable to analysis, and a function on \mathbb{R}^n is radial if and only if it is symmetric with respect to reflection in any hyperplane through the origin. A basic geometrical or mechanical problem asks whether a body that is rotationally symmetric in more than one axis must be a ball. Our motivation comes from the widely applied method of Steiner symmetrization, in which iterated Steiner symmetrals of a compact set, if taken with respect to a suitable sequence of hyperplanes, converge to a ball. For more information about the latter, see [2], where we find use for all the main results obtained here.

Let $i \in \{1, \dots, n-1\}$. We say that a set G of reflections in i -dimensional subspaces of \mathbb{R}^n generates $O(n)$ if the closure of the group $\langle G \rangle$ generated by G equals $O(n)$. For the applications we have in mind, it is particularly desirable to find finite sets G with this property, and this is the focus of most of the paper, in Section 3. In Section 4, we address the corresponding question as to when G generates $SO(n)$ if G is the set of all the rotations about each of several subspaces, each with dimension between 1 and $n-2$.

In the case of reflections, prior work mainly addresses the case when $i = n-1$, i.e., of reflections in hyperplanes through the origin. When G is infinite, Eaton and Perlman [4]

2020 *Mathematics Subject Classification.* 20F55, 22E15, 51F15, 51F25, 52A20.

Key words and phrases. reflection, Coxeter group, rotational symmetry, body of revolution.

First and third author supported in part by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). Second author supported in part by U.S. National Science Foundation Grant DMS-1402929.

proved that G generates $O(n)$ if it is irreducible, i.e., there is no nontrivial proper subspace S of \mathbb{R}^n such that $gS \subset S$ for $g \in G$, and it is clear that this sufficient condition is also necessary. The problem for arbitrary sets G was solved by Burchard, Chambers, and Dranovski [3]. In [3, Proposition 4.2], they show that for a set of reflections to generate $O(n)$, it suffices that the unit normal vectors to the corresponding hyperplanes satisfy three conditions: (i) at least two are at an angle that is an irrational multiple of π ; (ii) they span \mathbb{R}^n ; and (iii) they cannot be partitioned into two mutually orthogonal subsets. As these authors point out on [3, p. 1189], the first condition can be replaced by (i'): the set of reflections do not belong to a finite Coxeter group. The latter is weaker than (i), since, when $n > 2$, it is enough that at least two unit normal vectors are at an angle which is not an integer multiple of $\pi/3$, $\pi/4$, or $\pi/5$. Conditions (i'), (ii), and (iii) are necessary as well as sufficient.

We obtain the following results for reflections. Theorem 3.2 deals with the relatively simple case when $i = 1$, and proves that the conditions (i), (ii), and (iii) stated above suffice, where the unit vectors are now taken parallel to the subspaces. When $i \in \{2, \dots, n-2\}$, the situation is inherently more complicated. One of the main ingredients in our approach is the quite difficult but general Lemma 3.7, which we call the symmetry extension lemma. This assumes that H and L are subspaces of \mathbb{R}^n such that $H \cap L^\perp = \{o\}$ and $\dim H < \dim L$, and that G is a closed subgroup of $O(n)$ that acts transitively on $S^{n-1} \cap (L + x)$ for each $x \in S^{n-1}$. (Here $+$ denotes the Minkowski or vector sum; see Section 2 for this and other notation and terminology.) The conclusion is that the closure of the subgroup generated by G and the reflection R_H in H acts transitively on $S^{n-1} \cap (H + L + x)$ for each $x \in S^{n-1}$. Using the symmetry extension lemma and other tools such as Kronecker's approximation theorem, we are able in Theorem 3.8 to provide sufficient conditions for a finite set of reflections in i -dimensional subspaces to generate $O(n)$. In Corollary 3.9, we conclude that if

$$k = \begin{cases} n, & \text{if } i = 1 \text{ or } i = n-1, \\ \lceil n/i \rceil + 1, & \text{if } 1 < i \leq n/2, \\ \lceil n/(n-i) \rceil + 1, & \text{if } n/2 \leq i < n-1, \end{cases}$$

then there are i -dimensional subspaces H_1, \dots, H_k , such that the reflections R_{H_1}, \dots, R_{H_k} generate $O(n)$, or, equivalently, are such that if $E \subset S^{n-1}$ is nonempty, closed, and satisfies $R_{H_j}E = E$ for $j = 1, \dots, k$, then $E = S^{n-1}$. This in turn implies that if K is a convex body in \mathbb{R}^n satisfying $R_{H_j}K = K$ for $j = 1, \dots, k$, then K is a ball with center at the origin.

The previous result prompts the question as to what happens if reflections in subspaces are replaced by (full) rotational symmetries in subspaces, and we answer this in Section 4. (See [6] for a result in \mathbb{R}^3 in the same spirit.) There are easy examples of nonempty, closed, proper subsets of S^{n-1} that are left invariant under all rotations about more than one subspace; for example, the Cartesian product in \mathbb{R}^4 of two orthogonal copies of the circle with center at the origin and radius $1/\sqrt{2}$ is invariant under any rotation about each of the two-dimensional subspaces orthogonal to the circles. Similarly, the Cartesian product in \mathbb{R}^4 of two orthogonal disks centered at the origin (a so-called duocylinder) is a nonspherical convex body of revolution about each of the two-dimensional subspaces orthogonal to the disks. Theorem 4.1 states

a necessary and sufficient pair of conditions for subspaces H_1, \dots, H_k with $1 \leq \dim H_j \leq n-2$ to be such that if a nonempty closed subset of S^{n-1} is invariant under any rotation about each H_j , $j = 1, \dots, k$, then $E = S^{n-1}$. The latter is equivalent to saying that a closed set in \mathbb{R}^n that is invariant under rotations that fix H_j , $j = 1, \dots, k$, must be a union of spheres with their centers at the origin, or to saying that if a convex body K in \mathbb{R}^n is rotationally symmetric in H_j , $j = 1, \dots, k$, then K must be a ball with center at the origin. The conditions are (a) $H_1^\perp + \dots + H_k^\perp = \mathbb{R}^n$ and (b) $\{H_1^\perp, \dots, H_k^\perp\}$ cannot be partitioned into two mutually orthogonal nonempty subsets. Corollary 4.2 draws the conclusion that if $1 \leq i \leq n-2$ and $k = \lceil n/(n-i) \rceil$, then there are i -dimensional subspaces H_1, \dots, H_k , that satisfy (a) and (b). Corollary 4.3 shows that among closed sets in \mathbb{R}^n , $n \geq 3$, only unions of spheres with their centers at the origin are invariant under rotations about two different axes.

We are very grateful to Fabio Podestà for valuable discussions, and to a referee who suggested extracting the present paper from the original version of [2] and whose thorough reading and extensive knowledge led to many improvements.

2. PRELIMINARIES

As usual, S^{n-1} denotes the unit sphere and o the origin in Euclidean n -space \mathbb{R}^n with Euclidean norm $\|\cdot\|$. We assume throughout that $n \geq 2$. The term *ball* in \mathbb{R}^n will always mean an n -dimensional ball unless otherwise stated. The unit ball in \mathbb{R}^n will be denoted by B^n . If $x, y \in \mathbb{R}^n$, we write $x \cdot y$ for the inner product.

Let X and Y be sets in \mathbb{R}^n . We denote by $\text{lin } X$ and $\text{cl } X$ the *linear hull* and *closure* of X , respectively. If $t \in \mathbb{R}$, then $tX = \{tx : x \in X\}$ is the *dilate* of X by the factor t and

$$X + Y = \{x + y : x \in X, y \in Y\}$$

denotes the *Minkowski sum* of X and Y .

If $x \in \mathbb{R}^n \setminus \{o\}$, then x^\perp is the $(n-1)$ -dimensional subspace orthogonal to x . Throughout the paper, the term *subspace* means a linear subspace. If H is a subspace of \mathbb{R}^n , then $\dim H$ is its *dimension*, $X|H$ is the (orthogonal) projection of a set X on H , and $x|H$ is the projection of a vector $x \in \mathbb{R}^n$ on H .

The Grassmannian of k -dimensional subspaces in \mathbb{R}^n is denoted by $\mathcal{G}(n, k)$.

As usual, $O(n)$ and $SO(n)$ denote the *orthogonal group* and *special orthogonal group*, respectively, of isometries of \mathbb{R}^n . If ϕ is an isometry and H is a subspace of \mathbb{R}^n , we say that ϕ *fixes* H (or H is *fixed* by ϕ) if ϕ acts as the identity on H . (Note the difference between this and invariance; as usual, we say that a set X is *invariant* under ϕ if $\phi X = X$.) By $O(n)_H$ (or $SO(n)_H$) we mean the *pointwise stabilizer* of H , i.e. the subgroup of isometries in $O(n)$ (or $SO(n)$, respectively), that fix H . If G is a subgroup of $O(n)$ and $x \in S^{n-1}$, we denote by $G(x) = \{\phi x : \phi \in G\}$ the *orbit* of x .

If H is a subspace of \mathbb{R}^n , we write $R_H X$ for the *reflection* of X in H , i.e., the image of X under the map that takes $x \in \mathbb{R}^n$ to $2(x|H) - x$. If $R_H X = X$, we say X is *H-symmetric*. If $H = \{o\}$, we instead write $-X = (-1)X$ for the reflection of X in the origin and *o-symmetric* for $\{o\}$ -symmetric. A set X is called *rotationally symmetric* with respect to the i -dimensional subspace H if for all $x \in H$, $X \cap (H^\perp + x)$ is a union of $(n-i-1)$ -dimensional spheres, each

with center at x . These are precisely the sets that are invariant under each element of $O(n)_H$ or of $SO(n)_H$. If $\dim H = n - 1$, then X is rotationally symmetric with respect to H if and only if it is H -symmetric.

As usual, $\alpha_1, \dots, \alpha_l \in \mathbb{R}$ are called *linearly independent over \mathbb{Q}* if $\sum_{j=1}^l q_j \alpha_j = 0$ with $q_1, \dots, q_l \in \mathbb{Q}$ implies that $q_1 = \dots = q_l = 0$.

Finally, \mathcal{K}_n^n signifies the class of *convex bodies*, i.e., compact convex sets with interior points. This notation follows that of Schneider's classic text [8].

3. FULL ROTATIONAL SYMMETRY VIA REFLECTIONS IN FINITELY MANY SUBSPACES

This section focuses on finding finite sets of i -dimensional subspaces such that reflections in these subspaces generate full rotational symmetry.

Our first main result, Theorem 3.2 below, relies essentially on the next lemma, a special case of the symmetry extension lemma, Lemma 3.7. It is closely related to [7, Lemma 4], a result stated using the notion of dimension of a subgroup of a Lie group and proved using group theory. We provide a simple topological proof. Note that since $E \subset S^{n-1}$, requiring E to be invariant with respect to every element of $O(n)_H$ is the same as requiring E to be rotationally symmetric with respect to H .

Lemma 3.1. *Let $H \in \mathcal{G}(n, 1)$, $n \geq 3$, and let $O(n)_H$ be its pointwise stabilizer in $O(n)$. Let $\phi \in O(n)$ be such that $\phi H \neq H$. If $E \subset S^{n-1}$ is nonempty, closed, and invariant with respect to ϕ and to all the elements of $O(n)_H$, then $E = S^{n-1}$.*

Proof. Suppose that $E \neq S^{n-1}$. The set $S^{n-1} \setminus H$ contains at least one point in E , for otherwise $E \subset H$ and E could not be invariant with respect to ϕ . It also contains at least one point not in E , or else E , being closed, would coincide with S^{n-1} . Since any continuous curve joining a point of E to a point not in E contains a point of $\text{relbd } E$, there exists a point $v \in S^{n-1} \setminus H$ with $v \in \text{relbd } E$, where here and for the remainder of the proof, relbd denotes the boundary relative to S^{n-1} .

Note that if $\psi \in O(n)$ and $\psi E = E$, then $\psi(\text{relbd } E) = \text{relbd } E$. It follows that the orbit $C = O(n)_H(v)$ of v under $O(n)_H$ is a hyperplane section of S^{n-1} contained in $\text{relbd } E$. It is connected because $n \geq 3$. The image ϕC of C under ϕ is also a hyperplane section of S^{n-1} contained in $\text{relbd } E$. The projection $(\phi C)|H$ of ϕC onto H is an interval I , since ϕC is connected. See Figure 1. The set I is not a single point, because the hyperplane containing ϕC is not orthogonal to H , due to $\phi H \neq H$. Let

$$(1) \quad D = \{O(n)_H(x) : x \in \phi C\} = \{u \in S^{n-1} : u|H \in I\}.$$

Let D' be the interior of D relative to S^{n-1} and let I' be the interior of I relative to H . Then (1) yields $D' = \{u \in S^{n-1} : u|H \in I'\} \neq \emptyset$. However, $D \subset \text{relbd } E$, and since E is closed, the interior of $\text{relbd } E$ relative to S^{n-1} is empty, so $D' = \emptyset$, a contradiction. \square

Theorem 3.2. *Let $H_j \in \mathcal{G}(n, 1)$, $j = 1, \dots, n$, be such that*

- (i) *at least two of them form an angle that is an irrational multiple of π ,*
- (ii) *$H_1 + \dots + H_n = \mathbb{R}^n$, and*




FIGURE 1.

(iii) $\{H_1, \dots, H_n\}$ cannot be partitioned into two mutually orthogonal nonempty subsets.

If $E \subset S^{n-1}$ is nonempty, closed, and such that $R_{H_j}E = E$, $j = 1, \dots, n$, then $E = S^{n-1}$.

Proof. We claim that there is an ordering of H_1, \dots, H_n so that for $m = 2, \dots, n$ and all $v \in E$, we have

$$(2) \quad S^{n-1} \cap (H_1 + \dots + H_m + v) \subset E.$$

When $m = n$, this yields the theorem, by hypothesis (ii).

We argue by induction on m . By reordering, if necessary, we may, by hypothesis (i), assume that H_1 and H_2 form an angle that is an irrational multiple of π . The composition $R_{H_2}R_{H_1}$ maps the 2-dimensional plane $H_1 + H_2 + v$ to itself. Moreover, $R_{H_2}R_{H_1}$ restricted to $H_1 + H_2 + v$ is a rotation by an irrational multiple of π . By Kronecker's approximation theorem [1, Theorem 7.7], the set $\{(R_{H_2}R_{H_1})^m v : m \in \mathbb{N}\}$ is dense in $S^{n-1} \cap (H_1 + H_2 + v)$, so the invariance of E with respect to $R_{H_2}R_{H_1}$ and its being closed imply that $S^{n-1} \cap (H_1 + H_2 + v) \subset E$. This proves (2) when $m = 2$.

Let $m \in \{2, \dots, n-1\}$ and assume that (2) holds for all $v \in E$. Define $P_k = H_1 + \dots + H_k$ for $k = 2, \dots, n$. By hypothesis (iii), among the lines H_{m+1}, \dots, H_n there is at least one that is neither orthogonal to nor contained in P_m . By reordering, if necessary, we may assume that this line is H_{m+1} .

Let $v \in E$. We want to prove that

$$(3) \quad S^{n-1} \cap (P_{m+1} + v) = S^{n-1} \cap (P_m + H_{m+1} + v) \subset E.$$

We apply Lemma 3.1, where $P_{m+1} + v$ plays the role of the ambient space \mathbb{R}^n (so that $S^{n-1} \cap (P_{m+1} + v)$ acts as the unit sphere), E is replaced by $E \cap (P_{m+1} + v)$, H is the line $P_m^\perp \cap (P_{m+1} + v)$, $O(n)_H$ is replaced by the isometries of $P_{m+1} + v$ to itself that fix H , and $\phi = R_{H_1}R_{H_{m+1}}$. Note

that $E \cap (P_{m+1} + v)$ is invariant with respect to ϕ since

$$R_{H_{m+1}}(E \cap (P_{m+1} + v)) = E \cap (P_{m+1} - v)$$

and

$$R_{H_1}(E \cap (P_{m+1} - v)) = E \cap (P_{m+1} + v).$$

The other hypotheses of Lemma 3.1 are also satisfied. Indeed, $\dim(P_{m+1} + v) \geq 3$, and $\phi H \neq H$, because H_{m+1} is neither orthogonal to nor contained in P_m ; and the assumption that $E \cap (P_{m+1} + v)$ is invariant under each isometry of $P_{m+1} + v$ to itself that fixes H comes from the inductive hypothesis (2), which says that $S^{n-1} \cap (P_m + w) \subset E$ for each $w \in E$. With all this in hand, Lemma 3.1 gives (3). \square

Burchard, Chambers, and Dranovski [3, Proposition 4.2] prove a result similar to the previous theorem for reflections in hyperplanes through the origin. More precisely, they prove that if $H_j \in \mathcal{G}(n, n-1)$, $j = 1, \dots, n$, then under the same assumptions on H_j^\perp , $j = 1, \dots, n$, as in Theorem 3.2, a nonempty closed subset of S^{n-1} that is symmetric with respect to H_j , $j = 1, \dots, n$, coincides with S^{n-1} . In fact, in [3, p. 1189] they state without proof a stronger result, which is also a consequence of the main theorem of Eaton and Perlman [4]. The latter states that if G is a closed subgroup of $O(n)$ generated by reflections in $(n-1)$ -dimensional subspaces that is infinite and irreducible (i.e., there is no nontrivial proper subspace S of \mathbb{R}^n such that $gS \subset S$ for $g \in G$), then $G = O(n)$. Consequently, instead of condition (i) of Theorem 3.2, one need only assume that the subgroup generated by the reflections in $(n-1)$ -dimensional subspaces is not a finite Coxeter subgroup of $O(n)$. It appears to be unknown whether the result in [4] extends to reflections in lower-dimensional subspaces; in this connection, see Problem 5.1.

Remark 3.3. The similarity between Theorem 3.2 and [3, Proposition 4.2] suggests that there may be some duality at play. To investigate this possibility, observe that if H is a subspace of \mathbb{R}^n , then clearly

$$(4) \quad R_{H^\perp} = -Id \circ R_H = R_H \circ -Id,$$

where Id is the identity map. Now let $H_j \in \mathcal{G}(n, 1)$, $j = 1, \dots, n$, satisfy the hypotheses of Theorem 3.2. Then $H_j^\perp \in \mathcal{G}(n, n-1)$, $j = 1, \dots, n$, satisfy the hypotheses of [3, Proposition 4.2]. Suppose that $E \subset S^{n-1}$ is nonempty, closed, and such that $R_{H_j}E = E$, $j = 1, \dots, n$. If E is o -symmetric, (4) implies that $R_{H_j^\perp}E = E$, $j = 1, \dots, n$, and then [3, Proposition 4.2] gives $E = S^{n-1}$. Thus [3, Proposition 4.2] implies Theorem 3.2 when E is o -symmetric, and a similar argument shows that Theorem 3.2 implies [3, Proposition 4.2] when E is o -symmetric. For general sets $E \subset S^{n-1}$, however, the relationship between these results is not obvious; see Problem 5.2.

To obtain results similar to Theorem 3.2 for the case when $i \in \{2, \dots, n-2\}$, some preliminary observations will be useful. Let H_1 and H_2 be subspaces of \mathbb{R}^n of dimension k and i , respectively, with $k \geq i$. There exists a unique increasing sequence $\alpha_1, \dots, \alpha_i$ in $[0, \pi/2]$ and

an orthonormal basis e_1, \dots, e_n for \mathbb{R}^n such that

$$\begin{aligned} H_1 &= \text{lin} \{e_1, \dots, e_k\} \quad \text{and} \\ H_2 &= \text{lin} \{\cos \alpha_j e_j + \sin \alpha_j e_{i+k+1-j} : j = 1, \dots, i\} \\ (5) \quad &= \text{lin} \{\cos \alpha_1 e_1 + \sin \alpha_1 e_{k+i}, \dots, \cos \alpha_i e_i + \sin \alpha_i e_{k+1}\}, \end{aligned}$$

where the first $i+k-n$ angles are zero if $i+k > n$. Then it is easy to see that any $x \in \mathbb{R}^n$ can be written as

$$(6) \quad x = \sum_{j=1}^i \rho_j (\cos \theta_j e_j + \sin \theta_j e_{i+k+1-j}) + y_1 + y_2$$

for suitable $\rho_1, \dots, \rho_i \in \mathbb{R}$, $\theta_1, \dots, \theta_i \in [0, 2\pi)$, $y_1 \in \text{lin} \{e_{i+1}, \dots, e_k\}$ (where $y_1 = 0$ when $i = k$) and $y_2 \in (H_1 + H_2)^\perp$.

Lemma 3.4. *Let the subspaces H_1 and H_2 , of dimension k and $i \leq k$, respectively, be as in (5) and let the point $x \in \mathbb{R}^n$ be specified by (6). Then*

$$(7) \quad R_{H_1} x = \sum_{j=1}^i \rho_j (\cos(-\theta_j) e_j + \sin(-\theta_j) e_{i+k+1-j}) + y_1 - y_2,$$

$$(8) \quad R_{H_2} x = \sum_{j=1}^i \rho_j (\cos(2\alpha_j - \theta_j) e_j + \sin(2\alpha_j - \theta_j) e_{i+k+1-j}) - y_1 - y_2,$$

and for $m \in \mathbb{N}$,

$$(9) \quad (R_{H_2} R_{H_1})^m x = \sum_{j=1}^i \rho_j (\cos(2m\alpha_j + \theta_j) e_j + \sin(2m\alpha_j + \theta_j) e_{i+k+1-j}) + (-1)^m y_1 + y_2.$$

Proof. To obtain (7), note that R_{H_1} keeps the component of x in $\text{lin} \{e_1, \dots, e_k\}$ unchanged and changes the sign of the component in $\text{lin} \{e_{k+1}, \dots, e_n\}$.

To prove (8), we first compute $x|_{H_2}$. Since the vectors in the representation of H_2 are orthonormal, we have

$$x|_{H_2} = \sum_{j=1}^i (x \cdot (\cos \alpha_j e_j + \sin \alpha_j e_{i+k+1-j})) (\cos \alpha_j e_j + \sin \alpha_j e_{i+k+1-j}).$$

If x is as in (6), then

$$x \cdot (\cos \alpha_j e_j + \sin \alpha_j e_{i+k+1-j}) = \rho_j \cos(\alpha_j - \theta_j)$$

and therefore

$$x|_{H_2} = \sum_{j=1}^i \rho_j \cos(\alpha_j - \theta_j) (\cos \alpha_j e_j + \sin \alpha_j e_{i+k+1-j}).$$

Since $R_{H_2}x = 2(x|H_2) - x$, we obtain

$$R_{H_2}x = \sum_{j=1}^i \rho_j (2 \cos(\alpha_j - \theta_j) \cos \alpha_j - \cos \theta_j) e_j + \\ + \sum_{j=1}^i \rho_j (2 \cos(\alpha_j - \theta_j) \sin \alpha_j - \sin \theta_j) e_{i+k+1-j} - y_1 - y_2.$$

Then the relations

$$2 \cos(\alpha_j - \theta_j) \cos \alpha_j - \cos \theta_j = \cos(2\alpha_j - \theta_j)$$

and

$$2 \cos(\alpha_j - \theta_j) \sin \alpha_j - \sin \theta_j = \sin(2\alpha_j - \theta_j)$$

yield (8). Finally, (9) is an immediate consequence of (7) and (8). \square

It follows from (4) that if H_1 and H_2 are subspaces of \mathbb{R}^n , then

$$(10) \quad R_{H_2^\perp} R_{H_1^\perp} = R_{H_2} R_{H_1}.$$

Recall that if $1 \leq p \neq q \leq n$, then $SO(n)_{\text{lin}\{e_p, e_q\}^\perp}$ denotes the isometries in $SO(n)$ that act as the identity on $\text{lin}\{e_p, e_q\}^\perp$, the $(n-2)$ -dimensional subspace orthogonal to e_p and e_q .

Lemma 3.5. *Let $i \leq n/2$ and let $\alpha_1, \dots, \alpha_i$ be an increasing sequence in $(0, \pi/2)$ such that $\pi, \alpha_1, \dots, \alpha_i$ are linearly independent over \mathbb{Q} . Let $H_1, H_2 \in \mathcal{G}(n, i)$ have representations as in (5). Then both*

$$(11) \quad \{(R_{H_2} R_{H_1})^m : m \in \mathbb{N}\} \quad \text{and} \quad \{(R_{H_2^\perp} R_{H_1^\perp})^m : m \in \mathbb{N}\}$$

are dense in $SO(n)_{\text{lin}\{e_j, e_{2i-j+1}\}^\perp}$, $j = 1, \dots, i$.

Proof. Let $j \in \{1, \dots, i\}$. In (6), let $x = e_j$, so that $y_1 = y_2 = 0$, $\rho_j = 1$, $\rho_l = 0$ for $l \neq j$, and $\theta_j = 0$. With $k = i$, (9) yields

$$(12) \quad (R_{H_2} R_{H_1})^m e_j = \cos 2m\alpha_j e_j + \sin 2m\alpha_j e_{2i-j+1}.$$

The claim regarding the set on the left in (11) is then a direct consequence of the linear independence of $\pi, \alpha_1, \dots, \alpha_i$ over \mathbb{Q} and Kronecker's approximation theorem [1, Theorem 7.7]. Indeed, the latter implies that because α_j is not a rational multiple of π , the set $\{(\cos 2m\alpha_j, \sin 2m\alpha_j) : m \in \mathbb{N}\}$ is dense in S^1 . By (12), it follows that $\{(R_{H_2} R_{H_1})^m e_j : m \in \mathbb{N}\}$ is dense in $S^{n-1} \cap \text{lin}\{e_j, e_{2i-j+1}\}$ and hence, since $\dim(\text{lin}\{e_j, e_{2i-j+1}\}) = 2$, that $\{(R_{H_2} R_{H_1})^m : m \in \mathbb{N}\}$ is dense in $SO(n)_{\text{lin}\{e_j, e_{2i-j+1}\}^\perp}$. This also proves the claim regarding the set on the right in (11), since by (10), the two sets in (11) are equal. \square

Lemma 3.6. *Let $2 \leq i \leq n/2$ and let $\alpha_1, \dots, \alpha_i$ be an increasing sequence in $(0, \pi/2)$ such that $\pi, \alpha_1, \dots, \alpha_i$ are linearly independent over \mathbb{Q} . Let $H_1, H_2, H_3 \in \mathcal{G}(n, i)$ be defined by*

$$(13) \quad \begin{aligned} H_1 &= \text{lin} \{e_1, e_3, \dots, e_{2i-1}\}, \\ H_2 &= \text{lin} \{\cos \alpha_j e_{2j-1} + \sin \alpha_j e_{2j} : j = 1, \dots, i\}, \quad \text{and} \\ H_3 &= \text{lin} \{\{\cos \alpha_1 e_1 + \sin \alpha_1 e_{2i}\} \cup \{\cos \alpha_j e_{2j-1} + \sin \alpha_j e_{2j-2}\} : j = 2, \dots, i\}. \end{aligned}$$

Let $E \subset S^{n-1}$ be nonempty, closed, and such that either $R_{H_j} E = E$, $j = 1, 2, 3$, or $R_{H_j^\perp} E = E$, $j = 1, 2, 3$. Then if $x \in E$, we have $S^{n-1} \cap (\text{lin} \{e_1, \dots, e_{2i}\} + x) \subset E$.

Proof. Assume that $R_{H_j} E = E$ for $j = 1, 2, 3$. Since E is closed, it is invariant with respect to each element of $\text{cl} \{(R_{H_2} R_{H_1})^m : m \in \mathbb{N}\} \cup \text{cl} \{(R_{H_3} R_{H_1})^m : m \in \mathbb{N}\}$. The same conclusion holds if we assume instead that $R_{H_j^\perp} E = E$ for $j = 1, 2, 3$, since by (10), $R_{H_2} R_{H_1} = R_{H_2^\perp} R_{H_1^\perp}$ and $R_{H_3} R_{H_1} = R_{H_3^\perp} R_{H_1^\perp}$. By Lemma 3.5, modified suitably in view of the different expressions for H_1 and H_2 in (5) with $k = i$ and in (13), E is invariant with respect to each element of

$$V_1 = \cup \{SO(n)_{\text{lin} \{e_{2j-1}, e_{2j}\}^\perp} : j = 1, \dots, i\}.$$

By using Lemma 3.5 similarly but with H_2 replaced by H_3 , we see that E is also invariant with respect to each element of

$$V_2 = \cup \{SO(n)_{\text{lin} \{e_{2j-1}, e_{2j-2}\}^\perp} : j = 2, \dots, i\}.$$

(The factor $SO(n)_{\text{lin} \{e_1, e_{2i}\}^\perp}$ could also be included in the latter union, but we do not require this for the remaining argument.) Consequently, E is invariant with respect to each element of the subgroup G of $SO(n)$ generated by $V_1 \cup V_2$.

Suppose that $x \in E$ and choose $y \in \text{lin} \{e_1, \dots, e_{2i}\}^\perp$ such that

$$S^{n-1} \cap (\text{lin} \{e_1, \dots, e_{2i}\} + y) = S^{n-1} \cap (\text{lin} \{e_1, \dots, e_{2i}\} + x).$$

It clearly suffices to prove that G acts transitively on $S^{n-1} \cap (\text{lin} \{e_1, \dots, e_{2i}\} + y)$. To see this, let $u \in S^{n-1} \cap (\text{lin} \{e_1, \dots, e_{2i}\} + y)$ and choose $z = (z_1, \dots, z_{2i}) \in \text{lin} \{e_1, \dots, e_{2i}\}$ such that $u = y + z$. Then there is a $\phi_1 \in SO(n)_{\text{lin} \{e_{2i-1}, e_{2i}\}^\perp} \subset V_1$ such that

$$\phi_1 u = \phi_1 y + \phi_1 z = y + \left(z_1, z_2, \dots, z_{2i-2}, \sqrt{z_{2i-1}^2 + z_{2i}^2}, 0 \right)$$

and hence a $\phi_2 \in SO(n)_{\text{lin} \{e_{2i-1}, e_{2i-2}\}^\perp} \subset V_2$ such that

$$\phi_2 \phi_1 u = \phi_2 y + \phi_2 \phi_1 z = y + \left(z_1, z_2, \dots, z_{2i-3}, \sqrt{z_{2i-2}^2 + z_{2i-1}^2 + z_{2i}^2}, 0, 0 \right).$$

Observe that the graph with vertices $\{1, 2, \dots, 2i\}$ and edges joining $2j-1$ to $2j$, $j = 1, \dots, i$ (corresponding to V_1), and joining $2j-1$ to $2j-2$, $j = 2, \dots, i$ (corresponding to V_2), is connected, since each vertex is joined to the preceding one and the following one. Using this property, we may choose $\phi_3, \dots, \phi_{2i-1} \in V_1 \cup V_2 \subset G$, in the same way that we chose ϕ_1 and ϕ_2 , so that

$$\phi_{2i-1} \cdots \phi_1 u = y + (\|z\|, 0, \dots, 0) = y + \left(\sqrt{1 - \|y\|^2}, 0, \dots, 0 \right).$$

Now if $u' \in S^{n-1} \cap (\text{lin}\{e_1, \dots, e_{2i}\} + y)$, we can similarly find $\psi_3, \dots, \psi_{2i-1} \in G$ such that

$$\psi_{2i-1} \cdots \psi_1 u' = y + \left(\sqrt{1 - \|y\|^2}, 0, \dots, 0 \right) = \phi_{2i-1} \cdots \phi_1 u.$$

Thus there is a $g \in G$ with $gu' = u$, so G acts transitively on $S^{n-1} \cap (\text{lin}\{e_1, \dots, e_{2i}\} + y)$. \square

The next lemma says that if H and L are subspaces satisfying the stated hypotheses and a nonempty closed set $E \subset S^{n-1}$ is H -symmetric and rotationally symmetric with respect to L^\perp , then E is also rotationally symmetric with respect to $(H + L)^\perp$. A couple of small modifications to the proof show that the lemma also holds when instead of $\dim H < \dim L$ it is assumed that $\dim H = \dim L$ and $H \cap L \neq \{o\}$, but we have no use for this case in the sequel.

Lemma 3.7. (Symmetry extension lemma.) *Let H and L be subspaces of \mathbb{R}^n such that $H \cap L^\perp = \{o\}$ and $\dim H < \dim L$. If $E \subset S^{n-1}$ is nonempty, closed, and invariant under R_H and under any element of $O(n)_{L^\perp}$, then E is invariant under any element of $O(n)_{(H+L)^\perp}$. The same conclusion holds if invariance under R_H is replaced by invariance under R_{H^\perp} .*

Proof. Let G be the closure of the subgroup of $O(n)$ generated by $O(n)_{L^\perp}$ and R_H . The first conclusion of the lemma is equivalent to the statement that for all $x \in E$, the orbit $G(x)$ contains the orbit $O(n)_{(H+L)^\perp}(x)$. The latter orbit is $S^{n-1} \cap (H + L + x)$, and we claim that it coincides with $S_x = S^{n-1} \cap (M + x)$, where $M = L + R_H L$. (See Figure 2, where for the sake of illustration, both H and L are 1-dimensional.) It suffices to prove that $M = H + L$. To this end, we use the easily-proved identity $A^\perp \cap B^\perp = (A + B)^\perp$, where A and B are subspaces of \mathbb{R}^n , to conclude that the assumption $H \cap L^\perp = \{o\}$ is equivalent to $H^\perp + L = \mathbb{R}^n$. Then $L|H = (H^\perp + L)|H = \mathbb{R}^n|H = H$. Hence, if $z \in H + L$, then $z = x|H + y$ for some $x, y \in L$. Thus

$$z = (x/2 + y) + (x|H - x/2) = (x/2 + y) + R_H(x/2) \in L + R_H L.$$

This shows that $H + L \subset M$. On the other hand, the definition of R_H implies that $R_H L \subset (L|H) + L \subset H + L$ and hence that $M \subset H + L$. This proves the claim.

Let $x \in E$. We have to show that $S_x \subset G(x)$. If $x \in M^\perp$, then $S_x = \{x\}$ and there is nothing to prove. Therefore we may assume that $x \notin M^\perp$, in which case S_x is of dimension $\dim M - 1$. We may also assume that $\dim H > 0$, since otherwise $H + L = L$ and the first conclusion of the lemma is trivially true. Then $\dim L > 1$, which in turn yields $\dim M = \dim(H + L) > 1$, so S_x is connected. In order to prove that $G(x)$ contains S_x , we shall prove that $G(x) \cap S_x$ is both closed and open relative to S_x .

To prove that $G(x) \cap S_x$ is closed relative to S_x , let $\{y_k\}$ be a sequence of points in $G(x) \cap S_x$ converging to y . By the definition of $G(x)$, for each k , there is a $\phi_k \in O(n)$ such that $\phi_k(x) = y_k$. Since $O(n)$ is compact, there exists a subsequence $\{\phi_{k_m}\}$ of $\{\phi_k\}$ converging to $\phi \in O(n)$. Clearly $\phi(x) = y$, so $y \in G(x)$, and $y \in S_x$ because S_x is closed.

It remains to show that $G(x) \cap S_x$ is open relative to S_x . By the closed-subgroup theorem (see, for example, [9, Theorem 3.42]), G is a Lie group, because it is a closed subgroup of a Lie group. Since G is a C^∞ manifold and the map $a : G \rightarrow G(x)$ given by $a(g) = g(x)$ is C^∞ , the orbit $G(x)$ is also a C^∞ manifold. Moreover, $G(x) \cap S_x$ is also a manifold, because

Next, we observe that the following equations hold:

$$\begin{aligned} \dim L - \dim(L \cap p^\perp) &= \begin{cases} 0, & \text{if } p \in L^\perp \\ 1, & \text{if } p \notin L^\perp, \end{cases} \\ \dim R_H L - \dim(R_H L \cap p^\perp) &= \begin{cases} 0, & \text{if } p \in R_H L^\perp \\ 1, & \text{if } p \notin R_H L^\perp, \end{cases} \\ \dim(L \cap R_H L) - \dim(L \cap R_H L \cap p^\perp) &= \begin{cases} 0, & \text{if } L \cap R_H L \subset p^\perp \\ 1, & \text{if } L \cap R_H L \not\subset p^\perp. \end{cases} \end{aligned}$$

It follows that (15) holds for any p such that $L \cap R_H L \not\subset p^\perp$, or equivalently, for any $p \notin L^\perp + R_H L^\perp$.

We have $x \notin M^\perp = L^\perp \cap R_H L^\perp$, and thus either $x \notin L^\perp$ or $x \notin R_H L^\perp$. We claim that if $x \notin L^\perp$, then $S^{n-1} \cap (L + x)$ contains such a p , while if $x \notin R_H L^\perp$, then $S^{n-1} \cap (R_H L + x)$ contains such a p . It is enough to deal with the first case, since the proof for the second one is similar. Thus we need to prove that if $x \notin L^\perp$ then

$$(16) \quad S^{n-1} \cap (L + x) \not\subset L^\perp + R_H L^\perp.$$

Suppose that (16) is false. Then $L^\perp + R_H L^\perp$ contains all vectors which are differences of points in $S^{n-1} \cap (L + x)$. The assumption $x \notin L^\perp$ implies that $S^{n-1} \cap (L + x)$ has dimension $\dim L - 1 > 0$. It follows that $L^\perp + R_H L^\perp$ contains L . Therefore $L^\perp + R_H L^\perp = \mathbb{R}^n$, since $L^\perp + R_H L^\perp$ contains both L and L^\perp . From $H \cap L^\perp = \{o\}$ and $\dim H < \dim L$, we obtain

$$\dim(H^\perp \cap L) = \dim H^\perp + \dim L - \dim(H^\perp + L) = n - \dim H + \dim L - n > 0,$$

so there exists $w \neq o$ in $H^\perp \cap L$. Clearly, $R_H w = -w$ and $w \in L \cap R_H L$. Thus $L \cap R_H L \neq \{o\}$ and $L^\perp + R_H L^\perp = (L \cap R_H L)^\perp \neq \mathbb{R}^n$. This contradiction proves (16).

When $R_{H^\perp} E = E$, the proof is substantially identical. Due to (4), we have $R_H L = R_{H^\perp} L$, allowing us to repeat the arguments above with R_H replaced by R_{H^\perp} . \square

The next result provides for $i \in \{2, \dots, n-2\}$ a possible choice of i -dimensional subspaces H_1, \dots, H_k such that the reflections R_{H_1}, \dots, R_{H_k} generate $O(n)$.

Theorem 3.8. *Let $k \geq 3$ and let $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$. If $2 \leq i \leq n/2$, assume that*

- (i) H_1, H_2 , and H_3 are as in (13),
- (ii) $H_1 + \dots + H_k = \mathbb{R}^n$, and
- (iii) for each $j = 3, \dots, k-1$,

$$H_{j+1} \cap (H_1 + \dots + H_j)^\perp = \{o\}.$$

If $n/2 < i \leq n-2$, assume that (i)–(iii) are satisfied with each H_j replaced by H_j^\perp . If $E \subset S^{n-1}$ is nonempty, closed, and such that $R_{H_j} E = E$ for $j = 1, \dots, k$, then $E = S^{n-1}$.

Proof. Suppose that $2 \leq i \leq n/2$. Note that $H_1 + H_2 + H_3 = \text{lin}\{e_1, \dots, e_{2i}\}$. Therefore the assumptions on E and hypothesis (i), together with Lemma 3.6, imply that E is invariant

under any element of $O(n)_{(H_1+H_2+H_3)^\perp}$. Let $3 \leq j \leq k-1$ and suppose that E is invariant under any element of $O(n)_{(H_1+\dots+H_j)^\perp}$. Since

$$\dim H_{j+1} = i < 2i = \dim(H_1 + H_2 + H_3) \leq \dim(H_1 + \dots + H_j),$$

we may by hypothesis (iii) apply Lemma 3.7 with H and L replaced by H_{j+1} and $H_1 + \dots + H_j$, respectively, to conclude that E is invariant under any element of $O(n)_{(H_1+\dots+H_{j+1})^\perp}$. In view of hypothesis (ii), the proof is completed by induction on j . When $n/2 < i \leq n-2$, the previous argument can be repeated with each H_j replaced by H_j^\perp . \square

Corollary 3.9. *Let $1 \leq i \leq n-1$ and let*

$$k = \begin{cases} n, & \text{if } i = 1 \text{ or } i = n-1, \\ \lceil n/i \rceil + 1, & \text{if } 1 < i \leq n/2, \\ \lceil n/(n-i) \rceil + 1, & \text{if } n/2 \leq i < n-1. \end{cases}$$

There exist $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$, such that if $E \subset S^{n-1}$ is nonempty, closed, and such that $R_{H_j}E = E$ for $j = 1, \dots, k$, then $E = S^{n-1}$. Hence, if $F \subset \mathbb{R}^n$ is closed and $R_{H_j}F = F$ for $j = 1, \dots, k$ (or $K \in \mathcal{K}_n^n$ and $R_{H_j}K = K$ for $j = 1, \dots, k$), then F is a union of o -symmetric spheres (or K is an o -symmetric ball, respectively).

Proof. Let $E \subset S^{n-1}$ satisfy the stated hypotheses. If $i = 1$ or $i = n-1$, $E = S^{n-1}$ is a consequence of Theorem 3.2 or of [3, Proposition 4.2], respectively. Suppose that $1 < i \leq n/2$ and $k = \lceil n/i \rceil + 1$. Then if $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$, are as in Theorem 3.8, hypothesis (iii) there and $\dim(H_1 + H_2 + H_3) = 2i$ imply that

$$\dim(H_1 + \dots + H_k) = 2i + (k-3)i \geq 2i + (n/i - 2)i = n$$

and hence hypothesis (ii), so it follows from that theorem that $E = S^{n-1}$. Suppose that $n/2 \leq i < n-1$ and $k \geq \lceil n/(n-i) \rceil + 1$. Choose $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$, so that $H_j^\perp \in \mathcal{G}(n, n-i)$, $j = 1, \dots, k$, are as in Theorem 3.8. Then $E = S^{n-1}$ follows from the already established result for $1 < i \leq n/2$ on replacing i by $n-i$.

Let $F \subset \mathbb{R}^n$ be closed and satisfy $R_{H_j}F = F$ for $j = 1, \dots, k$. If F is not a union of o -symmetric spheres, there is an $r > 0$ such that if $A = F \cap rS^{n-1}$, then A is a nonempty, closed, proper subset of rS^{n-1} . It follows that the set $E = (1/r)A \subset S^{n-1}$ satisfies $R_{H_j}E = E$ for $j = 1, \dots, k$, but $E \neq S^{n-1}$, contradicting what has been proved. The proof is completed by noting that if $K \in \mathcal{K}_n^n$ is a union of o -symmetric spheres, it must be an o -symmetric ball. \square

4. FULL ROTATIONAL SYMMETRY VIA ROTATIONAL SYMMETRIES IN FINITELY MANY SUBSPACES

This section focuses on finding finite sets of i -dimensional subspaces such that (full) rotational symmetries in these subspaces generate full rotational symmetry.

Theorem 4.1. *Let H_1, \dots, H_k be subspaces in \mathbb{R}^n such that $1 \leq \dim H_j \leq n-2$ for $j = 1, \dots, k$. The following statements are equivalent.*

- (i) $H_1^\perp + \cdots + H_k^\perp = \mathbb{R}^n$ and $\{H_1^\perp, \dots, H_k^\perp\}$ cannot be partitioned into two mutually orthogonal nonempty subsets.
- (ii) If $E \subset S^{n-1}$ is nonempty and closed, and for each $j = 1, \dots, k$ and $x \in E$, we have $S^{n-1} \cap (H_j^\perp + x) \subset E$, then $E = S^{n-1}$.
- (iii) If $F \subset \mathbb{R}^n$ is closed and invariant under any rotation that fixes H_j for each $j = 1, \dots, k$, then F is a union of o -symmetric spheres.
- (iv) If $K \in \mathcal{K}_n^n$ is rotationally symmetric with respect to H_j for each $j = 1, \dots, k$, then K is an o -symmetric ball.

Proof. (i) \Rightarrow (ii) Suppose that (i) holds. If E satisfies the hypotheses in (ii), then every section of E with a translate of H_1^\perp is either empty or a sphere. By (i), there must be a $j_0 \in \{2, \dots, k\}$ such that $H_{j_0}^\perp$ is not contained in H_1 , for otherwise $\{H_1\}$ and $\{H_2, \dots, H_k\}$ would be a partition of $\{H_1, \dots, H_k\}$ rendering (i) false. Without loss of generality, assume that $j_0 = 2$ and choose an orthogonal basis v_1, \dots, v_l in H_2^\perp such that $v_1 \notin H_1$, i.e., v_1 is not orthogonal to H_1^\perp . If $x \in S^{n-1}$, then

$$R_{v_1^\perp} x \in \text{lin}\{v_1\} + x \subset H_2^\perp + x.$$

The hypothesis on E in (ii) with $j = 2$ thus implies that if $x \in E$, then $R_{v_1^\perp} x \in E$, that is, that E is symmetric with respect to the hyperplane v_1^\perp . By Lemma 3.7 with $H = \text{lin}\{v_1\}$ and $L = H_1^\perp$, every section of E with a translate of $H_1^\perp + \text{lin}\{v_1\}$ is either empty or a sphere. For $p = 1, \dots, l$, define $w_p = v_1 + \cdots + v_p$. It is easy to see that w_p is not orthogonal to w_1, \dots, w_{p-1} . Let $F_p = \text{lin}\{w_1, \dots, w_p\}$, $p = 1, \dots, l$. Using Lemma 3.7 iteratively with $H = \text{lin}\{w_p\}$ and $L = H_1^\perp + F_{p-1}$, $p = 1, \dots, l$, we conclude that every section of E with a translate of $H_1^\perp + F_l = H_1^\perp + H_2^\perp$ is either empty or a sphere.

Suppose we have shown that every section of E with a translate of $J_r = H_1^\perp + \cdots + H_r^\perp$ is either empty or a sphere, where $2 \leq r \leq k-1$. By (i), there must be a $j_1 \in \{r+1, \dots, k\}$ such that $H_{j_1}^\perp$ is not contained in J_r , for otherwise (i) would be false via the partition $\{H_1, \dots, H_r\}$ and $\{H_{r+1}, \dots, H_k\}$. Without loss of generality, assume that $j_1 = r+1$ and apply the argument in the previous paragraph, with H_1^\perp and H_2^\perp replaced by J_r and H_{r+1}^\perp , respectively. This yields that every section of E with a translate of J_{r+1} is either empty or a sphere. By induction on r , we arrive at the conclusion that every section of E with a translate of J_k is either empty or a sphere. But $J_k = H_1^\perp + \cdots + H_k^\perp = \mathbb{R}^n$, so $E = S^{n-1}$.

(ii) \Rightarrow (iii) \Rightarrow (iv) These implications are proved by the arguments in the last paragraph of the proof of Corollary 3.9.

(iv) \Rightarrow (i) Suppose that (i) is false and $H_1^\perp + \cdots + H_k^\perp = \mathbb{R}^n$. Then there is a partition of $\{1, \dots, k\}$ into two nonempty disjoint sets I_1 and I_2 such that the subspaces $S_m = \sum\{H_j^\perp : j \in I_m\}$, $m = 1, 2$, are orthogonal. Let $K = (B^n \cap S_1) \times (B^n \cap S_2)$ and note that $K \in \mathcal{K}_n^n$ as

$$S_1 + S_2 = H_1^\perp + \cdots + H_k^\perp = \mathbb{R}^n.$$

Let $j \in \{1, \dots, k\}$ and suppose without loss of generality that $j \in I_1$. Since H_j^\perp is orthogonal to S_2 , we have $K \cap (H_j^\perp + x) = B^n \cap (H_j^\perp + x)$, if $x|_{S_2} \in B^n \cap S_2$, and $K \cap (H_j^\perp + x) = \emptyset$, otherwise. Therefore K is rotationally symmetric with respect to H_j and since K is not a

ball, (iv) is false. If $H_1^\perp + \dots + H_k^\perp = S \neq \mathbb{R}^n$, then $K = (B^n \cap S) \times (B^n \cap S^\perp) \in \mathcal{K}_n^n$ is not a ball and again (iv) is false. \square

Corollary 4.2. *Let $1 \leq i \leq n - 2$ and let $k = \lceil n/(n - i) \rceil$. There exist $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$, such that the statements in Theorem 4.1 hold.*

Proof. It suffices to ensure that Theorem 4.1(i) holds. This is easily accomplished by choosing $H_j \in \mathcal{G}(n, i)$ inductively such that for $k = 1, 2, \dots$, we have $H_{k+1}^\perp \not\subset (H_1^\perp + \dots + H_k^\perp)^\perp$ and

$$\dim(H_1^\perp + \dots + H_k^\perp) \geq \min\{k(n - i), n\},$$

until $k(n - i) \geq n$, i.e., $k = \lceil n/(n - i) \rceil$. \square

In the narrower context of surfaces in \mathbb{R}^3 , the first statement in the following simple corollary was observed without proof in [5, p. 10] and proved in [6, Theorem A].

Corollary 4.3. *Let $n \geq 3$. If $H_1, H_2 \in \mathcal{G}(n, 1)$ with $H_1 \neq H_2$, the statements in Theorem 4.1 hold.*

No finite k exists such that if H_j , $j = 1, \dots, k$, are different subspaces with $2 \leq \dim H_j \leq n - 2$ for $j = 1, \dots, k$, the statements in Theorem 4.1 hold.

Proof. Let $H_1, H_2 \in \mathcal{G}(n, 1)$ with $H_1 \neq H_2$. Since $\dim H_1^\perp = \dim H_2^\perp = n - 1$ and $H_1^\perp \neq H_2^\perp$, Theorem 4.1(i) holds.

Suppose that $k \in \mathbb{N}$. The assumption that $2 \leq \dim H_j \leq n - 2$ for $j = 2, \dots, k$ allows us to choose different H_j 's such that $H_1^\perp, \dots, H_k^\perp$ are all contained in the same $(n - 1)$ -dimensional subspace. The first condition in Theorem 4.1(i) then fails.

With these observations in hand, the result follows from Theorem 4.1. \square

5. OPEN PROBLEMS

Problem 5.1. *For $i \in \{1, \dots, n - 2\}$, find necessary and sufficient conditions for a set \mathcal{F} of i -dimensional subspaces such that reflection symmetry with respect to each subspace in \mathcal{F} implies full rotational symmetry.*

As was mentioned after Theorem 3.2, for $i = n - 1$ such conditions are a consequence of the results in [3, Proposition 4.2] and [4].

Problem 5.2. *Let $i \in \{1, \dots, n - 1\}$, and let $H_j \in \mathcal{G}(n, i)$, $j = 1, \dots, k$, be such that reflection symmetry with respect to each H_j implies full rotational symmetry. Is it true that reflection symmetry with respect to each H_j^\perp implies full rotational symmetry?*

As in Remark 3.3, a positive answer to Problem 5.2 for α -symmetric sets follows from (4), but in general the latter equality does not help in any simple way. For example, if E is the set of vertices of a regular simplex T inscribed in S^2 , and $H_j \in \mathcal{G}(n, 1)$, $j = 1, \dots, 4$, are the lines through the midpoints of opposite pairs of edges of T , then $R_{H_j}E = E$, $j = 1, \dots, 4$, but $R_{H_j^\perp}E \neq E$, $j = 1, \dots, 4$.

REFERENCES

- [1] T. M. Apostol, *Modular Functions and Dirichlet Series in Number Theory*, second edition, Springer, New York, 1990.
- [2] G. Bianchi, R. J. Gardner, and P. Gronchi, Convergence of symmetrization processes, *Indiana Univ. Math. J.* **71** (2022), 785–817.
- [3] A. Burchard, G. R. Chambers, and A. Dranovski, Ergodic properties of folding maps on spheres, *Discrete Contin. Dyn. Syst.* **37** (2017), 1183–1200.
- [4] M. L. Eaton and M. Perlman, Generating $O(n)$ with reflections, *Pacific J. Math.* **73** (1977), 73–80.
- [5] D. Hilbert and S. Cohn-Vossen, *Geometry and the Imagination*, Chelsea Publishing Company, New York, NY, 1952.
- [6] D.-S. Kim and Y. H. Kim, Surfaces of revolution with more than one axis, *J. Korean Soc. Math. Educ. Ser. B Pure Appl. Math.* **19** (2012), 1–5.
- [7] D. Montgomery and H. Samelson, Transformation groups of spheres, *Ann. of Math.* **44** (1943), 454–470.
- [8] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, second edition, Cambridge University Press, Cambridge, 2014.
- [9] F. W. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Springer, New York–Berlin, 1983.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, FIRENZE, ITALY I-50134

Email address: gabriele.bianchi@unifi.it

DEPARTMENT OF MATHEMATICS, WESTERN WASHINGTON UNIVERSITY, BELLINGHAM, WA 98225-9063

Email address: richard.gardner@wwu.edu

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, PIAZZA Ghiberti 27, FIRENZE, ITALY I-50122

Email address: paolo.gronchi@unifi.it