

Former charcoal platforms in Mediterranean forest areas: a hostile microhabitat for the recolonization by woody species

Elisa Carrari⁽¹⁾, Evy Ampoorter⁽²⁾,
Kris Verheyen⁽²⁾, Andrea Coppi⁽³⁾,
Federico Selvi⁽¹⁾

Production of wood charcoal is a traditional form of forest use that lasted for millennia in the Mediterranean countries. Following their almost complete abandonment in the last century, thousands of old charcoal platforms remain in present-day forest landscapes. These sites are characterized by peculiar ecological conditions, whose effects on the recolonization by woody plants are still unknown. We examined 61 platforms in evergreen sclerophyllous woodlands and deciduous broadleaf forests with oaks and beech, spread over a wide geographic range in Tuscany (Italy). At each site, one kiln plot (on charcoal platform) and one control plot (in the adjacent stand) were established, and soil, light conditions and herb cover were measured. We examined species richness and composition of trees and shrubs in the understorey layer (<1.3 m) and in the “established regeneration” layer (> 1.3-4 m). In the latter, structural parameters such as number of stools, dbh, mean height and number of stems were compared. The density of seedlings of dominant tree species in the understorey was also measured in a subsample of sites per forest type. In the understorey, a general positive effect of kiln platforms was found on species richness at both the habitat and plot-scale level, as well as on species composition, especially in oak forests. Increased light availability, total C content and soil pH were positively related with species richness, while N content was a negative predictor. Density of seedlings was not substantially affected. Contrastingly, woody species richness in the established regeneration layer was considerably lower in the kiln plots of all three forest types. In sclerophyllous forests, all species in this layer were taller, denser and with a higher basal area compared to control plots, while regeneration was completely lacking on platforms of the two other forest types. Soil N content had a positive influence on structural parameters, while total C content resulting from charcoal accumulation had a negative influence. We conclude that charcoal platforms are a favorable microhabitat only in the first regeneration stages of woody species, as their further growth is hindered by long-term effects that should be investigated with an experimental approach.

Keywords: Charcoal Platforms, Diversity, Forest Recolonization, Mediterranean Area, Tree Regeneration, Species Composition, Woody Species

Introduction

Wood charcoal production is one of the oldest forms of anthropic forest use in most countries of temperate regions (Antal & Gronli 2003) and the main source of energy since the Iron Age until the 19th cen-

tury (Blondel 2006). Charcoal production was carried out in the so-called “charcoal kilns” or “charcoal hearths”, which are small terrace-like platforms (35-50 m²) with circular or elliptical shape, commonly established in wind-sheltered forest sites not

far from water courses, and connected by ad-hoc footpaths across hill and mountain slopes (Cantiani 1955, Attwood et al. 1981, Landi & Piussi 1988). A detailed description of the different production phases and the procedures used for charcoal kiln construction are reported in Landi & Piussi (1988), Powell (2008) and Giorgerini (2009), and are briefly illustrated in Fig. S1 (Supplementary Material).

In northern and central Europe this practice was generally abandoned in the 19th century due to the rapidly increasing and widespread use of coal (Deforce et al. 2013), while in most Mediterranean countries the production of wood charcoal increased during the industrial revolution because of the lack of other fuel sources. It generally vanished only in the 1950s, though in some mountain areas it is still in use. As a result, remnants of charcoal platforms are nowadays commonly found in many forest landscapes (Blondel 2006, Pèlachs et al. 2009). They are characterized

□ (1) University of Firenze, Dept. of AgriFood Production and Environmental Sciences, Laboratories of Applied and Environmental Botany, P.le Cascine 28, I-50144 Firenze (Italy); (2) Ghent University, Dept. of Forest and Water Management, Forest & Nature Lab (ForNaLab), Geraardsbergenweg 267, B-9090 Gontrode (Belgium); (3) University of Firenze, Dept. of Biology, Botanical Laboratories, v. G. La Pira 4, I-50121 Firenze (Italy)


@ Elisa Carrari (elisa.carrari@unifi.it)

Received: May 07, 2015 - Accepted: Jun 23, 2016

Citation: Carrari E, Ampoorter E, Verheyen K, Coppi A, Selvi F (2016). Former charcoal platforms in Mediterranean forest areas: a hostile microhabitat for the recolonization by woody species. iForest 10: 136-144. - doi: [10.3832/ifor1701-009](https://doi.org/10.3832/ifor1701-009) [online 2016-10-06]

Communicated by: Davide Ascoli

Fig. 1 - Geographic location of the forest stands and the investigated charcoal platforms in Tuscany (central Italy).



by clear alterations of color and texture of the soil material due to the charred woody remains (Montanari et al. 2000). In fact, the condensed aromatic structure of wood charcoal allows fragments and particles to persist in soils and other sedimentary records over millennial time-scales (Cheng et al. 2008, Zimmerman 2010). Anthropogenic deposits of charcoal dating back to the Neolithic period have been documented in Germany and Italy (Cremaschi et al. 2006, Ludemann 2010) and very old samples (> 8000 years BP) originating from wildfires have been found almost unaltered in forest soils of the Italian Alps (Valese et al. 2014).

Several anthracological studies have been carried out to reconstruct the former tree species composition of forests through the identification of the woody taxa used for charcoal production, thus highlighting the usefulness of charcoal platforms in detecting vegetation changes through time (Montanari et al. 2000, Ludemann 2003, Nelle 2003, Ludemann et al. 2004). However, the present-day vegetation of these sites has been rarely investigated in Europe (Wittig et al. 1999, Carrari et al. 2016a), and no studies focused on the tree regeneration and forest recolonization processes in charcoal platforms.

Successful recolonization of charcoal platforms by woody species is expected because of the apparently favorable conditions such as the low degree of rockiness, flat terrain and often a higher availability of light (Carrari et al. 2016a). On the other hand, adult trees are usually absent in most of the platforms, even in those abandoned several decades ago. This suggests that tree regeneration and establishment is hampered by unknown persistent biotic or abiotic factors, which represent a legacy of

the past forest use still affecting ecosystem dynamics at the “microhabitat” scale.

In order to test this hypothesis, we analyzed the diversity, density and structural parameters of trees and shrub species in the regeneration layer (< 4 m) of these platforms, and compared them with control sites located in the adjacent forest environment. Differences were then related to variation in light, soil conditions and herb cover measured at the same sites. Using such a paired design and an extensive sampling in three major forest types of central Italy, we could provide the first insights into the long-term effects of the old charcoal platforms on woody species regeneration in the Mediterranean area, and on the possible driving factors.

Material and methods

Study area

This study was performed in the forests of Tuscany (central Italy - Fig. 1). The study area includes three main climate and forest types, according to the EEA classification system (European Environment Agency 2007): (1) meso-Mediterranean evergreen forest dominated by *Quercus ilex* L. and sclerophyllous shrubs, along the Tyrrhenian coast (hereafter: “sclerophyllous” forests); (2) supra-Mediterranean thermophilous mixed communities dominated by deciduous oaks (*Quercus cerris* L., *Q. petraea* (Matt.) Liebl., *Q. pubescens* Willd.) in the hilly areas of central Tuscany (hereafter: “oak” forests); (3) montane forests with beech (*Fagus sylvatica* L.), on the Apennine chain (Casentino and Mugello areas) and the volcanic massif of Mt. Amiata (hereafter: “beech” forests). Mean annual rainfall and temperature in this area vary from 780.7

mm and 15.4 °C along the coast to 1.641 mm and 8.7 °C on the Apennines (period 1961–1990 – Regione Toscana 2016). The investigated area is characterized by a variety of geological formations and soil conditions, but cambisols are the prevalent type according to the Soil Atlas of Europe (European Soil Bureau Network 2005).

Field sampling design and data collection

An extensive search for abandoned charcoal platforms was carried out with *ad-hoc* field trips in representative areas of the three main forest types described above. Charcoal platforms were easily recognized due to the characteristics of the ground surface (Attwood et al. 1981, Landi & Piussi 1988, Ludemann 2012). One-hundred fifty-four different platforms were identified, and their position recorded with a GPS device. At each site, altitude, slope, aspect, soil type and tree species composition of the adjacent forest stands were also recorded. Sixty-one out of the above 154 sites were randomly selected among those that were not affected by recent anthropogenic or animal disturbance: (i) 18 “sclerophyllous” sites in 4 geographically distant forest stands; (ii) 22 “oak” sites in 3 stands; (iii) 21 “beech” sites in 3 stands. In the two areas (Volterra hills and Farma-Merse) both sclerophyllous and oak stands were present (Fig. 1, Tab. 1). The main geographical and environmental variables of these sites are given in Tab. 1. All but two sites are included in protected areas of the Natura 2000 network, Nature Reserves and National Parks. Charcoal production in all sites lasted for centuries and was abandoned at least 60 years ago, while the coppice-with-standards management for firewood production was abandoned later (ca. 45–50 years ago) in both oak and beech forests. In the sclerophyllous stands, this management type was abandoned more recently or it is still in use (Carrari & Selvi, personal observation).

At each selected site, a 3×3 m quadrat (hereafter: “kiln plot”, KP) was established in the center of the charcoal kiln platform. The relatively small size and the general shape of the platforms did not allow the adoption of larger quadrats, which would have included the kiln-forest edge. Light conditions (Photosynthetic Active Radiation, PAR), herbaceous cover and soil factors were measured following the protocol described in Carrari et al. (2016a); observations on the soil profile were also conducted in each quadrat.

Tree regeneration was analyzed in two distinct layers: (1) the understorey vegetation layer (< 1.3 m – hereafter: “UVL”) and the established tree regeneration layer (1.3–4 m – “ETL”). In the UVL, seedlings of all tree and shrub species were identified and their maximum height was recorded. The ground cover percentage of each of these species was quantified as the projection of their crown foliage on the soil sur-

Tab. 1 - Main geographical and environmental variables of the studied sites, with number of charcoal kiln (KP) and control plots (CP) for each of the examined forest stands. (*): Variables considered in the starting mixed model as predictors for the response variables.

District	Lat Long	No KP	No CP	Forest Type*	Elev. Range (m a.s.l.)	Aspect*	Parent rock material	Slope inclination	
								kiln	control
Colla di Casaglia	44°01' 57" N 11°29' 01" E	5	5	Beech	972-1029	SE/E/NE	marl-sandstone	0	10-50%
Mt. Amiata	42°52' 10" N 11°35' 03" E	11	11	Beech	846-1268	N/NE/E/SE/ S/SW/W/NW	trachyte	0	20-50%
Casentino	43°48' 19" N 11°52' 09" E	5	5	Beech	1040-1223	S/SE/E	marl-sandstone	0	5-45%
Volterra hills	43°25' 55" N 11°00' 02" E	7	7	Oaks/ Sclerophyllous	382-967	E/SE/N/ NW/NE	diabase/limestone/ sandstone	0	3-40%
Fiesole hills	43°48' 15" N 11°20' 27" E	2	2	Oaks	242-347	W/E	marl-calcareous	0	3-5%
Farma-Merse	43°05' 22" N 11°10' 46" E	15	15	Oaks/ Sclerophyllous	265-511	N/NE/E/SE/ S/SW/W/NW	quartzitic sandstone	0	3-40%
Mt. Leoni	42°56' 27" N 11°10' 58" E	5	5	Sclerophyllous	155-437	S/SW/-/W/E	quartzitic sandstone	0	3-35%
Magona Forest/ Mt. Massoncello	43°15' 50" N 10°37' 54" E	7	7	Sclerophyllous	157-201	W/NW/N/SE	marl-clay/sandstone	0	3-20%

face. According to Pignatti (1982), we considered as trees all woody angiosperms or gymnosperms belonging to the Raunkiaer's life forms "Pscap" (scapose phanerophyte) and "Pcaesp" (caespitose phanerophyte). In addition, the number of seedlings (height < 1.3 m) of the dominant species (all belonging to the Fagaceae family) was counted within the quadrat in a sub-sample of five randomly selected platforms for each forest type (15 plots in total - Tab. S4 in Supplementary Material).

In the "ETL", we recorded the species richness of trees and shrubs and the following structural parameters: number of trees/stools, number of stems per stool, diameter at breast height (dbh, cm) of each stem (> 0.5 cm), and mean height of the stems for each stool.

For each of the 61 kiln plots, the same measurements mentioned above were taken in a control plot (CP) of the same size randomly established in the stand adjacent to the kiln platform at a distance of 10-20 m from the edge of the KP (depending on local topographic and ground conditions); downhill locations were excluded to avoid potential charcoal "contamination" by runoff. This allowed to minimize the variation of the major environmental factors (altitude, parent rock material, slope inclination and slope aspect) between each kiln plot and its respective control plot.

Data analysis

All statistical analyses were performed in R 3.1.2 (R Core Team 2014). Soil factors, PAR values and herbaceous cover were compared between KP and CP using the non-parametric Mann-Whitney U test, due to the violation of assumptions of normality and homogeneity of the variance ("wilcox.test" function). Differences in woody species composition of the UVL between KP and CP in each forest type were assessed using the non-metric multi-

dimensional scaling (NMDS) based on the cover-weighted Bray-Curtis dissimilarity index ("metaMDS" function of the "vegan" package - Oksanen et al. 2015). Significance of the differences were tested by permutational multivariate analysis of variance (PERMANOVA) using the "adonis" function with 999 permutations. To verify that such differences were related to the effect of the factor kiln/control (i.e., to dissimilarities in composition between kiln and control plots) and not to a dispersion effect (i.e., dissimilarities within each of the two plot types), we tested for multivariate homogeneity of dispersion using the "betadisper" function, a multivariate analogue of the Levene's test for homogeneity of variances (Anderson 2001, Oksanen et al. 2015).

Density of seedlings of the locally dominant tree species was compared between KP and CP using the non-parametric Mann-Whitney U test. The effects of the charcoal kiln platforms on species richness (SR) at the habitat level (γ -diversity of KP and CP) and plot level (mean SR for KP and CP) in the UVL and in the ETL were also evaluated.

A mixed model approach was adopted in order to examine all tree regeneration parameters (seedling density of dominant species, SR in UVL and ETL and structural parameters) as response variables to the following predictor variables: forest type (levels: sclerophyllous, oak, beech), environmental variables (altitude, parent rock, slope inclination and slope aspect), soil and light factors (N, C, pH and PAR) and herbaceous cover. First, predictor variables were tested for collinearity using Pearson's and Spearman's correlation analysis to obtain quantitative and ordinal correlation coefficients, respectively. The association between nominal variables or between a nominal and a continuous variable was tested by chi-square or Student's t-test, respec-

tively. As a result, parent rock material, altitude and pH (associated with forest type) were excluded from further analyses.

The variable "forest stand" was included in the model as random factor to account for possible effects of environmental and/or management-related factors on the tree regeneration variables specific to each forest area.

Tree seedling density and SR in the under-story were initially analyzed using the following model (eqn. 1):

$$y = \alpha + \beta_1 FT \cdot HC + \beta_2 FT \cdot PAR + \beta_3 FT \cdot C + \beta_4 FT \cdot N + \beta_5 SA + \beta_6 SI + 1|FS + \varepsilon$$

where y is the response variable, α is the intercept, β is the parameter to be estimated for each factor, FT is the forest type, HC is the herb cover, PAR is the photosynthetic active radiation measured, C is the soil carbon content, N is the soil nitrogen content, SA is the slope aspect, SI is the slope inclination, $1|FS$ indicates the forest stand as random factor and ε is the error. The model was tested using the "glmer" function in R with a Poisson error distribution and the "loglink" function of the "lme4" package (Bates et al. 2014). For SR in the ETL, HC was excluded from the model as not relevant for woody species growth in this layer; light was also excluded since PAR values were measured at 1 m above the forest floor.

Starting from these full models, we looked for model parsimony following the approach of Zuur et al. (2009). First, we deleted the random variation across sites while keeping the fixed effect term, using the "glm" function from the "stats" package in R, with a Poisson error distribution, log link and parameter estimation via maximum likelihood. The model yielding the lowest value of the Akaike's Information Criterion (AIC - Akaike 1973) was considered as the best fitting to the data. Once an optimal random structure was found,

Tab. 2 - Mean values (\pm standard deviation) of soil carbon content (C), soil nitrogen content (N), pH and photosynthetic active radiation (PAR) and herbaceous cover (HC) of kiln and control plots (KP, CP) for each forest type (Sclerophyllous, Oaks, Beech).

Parameter	Sclerophyllous		Oaks		Beech				
	KP	CP	Prob	KP	CP	Prob	KP	CP	Prob
C (%)	11.22 \pm 2.92	6.03 \pm 3.12	<0.001	12.69 \pm 3.700	6.15 \pm 3.32	<0.001	7.47 \pm 1.68	5.14 \pm 2.56	0.003
N (%)	0.51 \pm 0.14	0.40 \pm 0.21	n.s.	0.57 \pm 0.18	0.45 \pm 0.24	0.02	0.43 \pm 0.08	0.43 \pm 0.16	n.s.
pH	6.22 \pm 0.68	5.97 \pm 0.76	n.s.	6.39 \pm 0.94	6.15 \pm 1.02	n.s.	5.54 \pm 0.46	5.35 \pm 0.56	0.023
PAR ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	43.55 \pm 47.98	21.83 \pm 14.57	n.s.	55.59 \pm 90.26	13.40 \pm 9.27	0.001	32.85 \pm 38.13	26.71 \pm 38.61	n.s.
HC (%)	1.74 \pm 1.97	1.66 \pm 3.36	n.s.	5.13 \pm 5.10	1.91 \pm 3.17	<0.001	1.39 \pm 1.84	1.26 \pm 3.11	n.s.

we searched for the optimal fixed effect structure by comparing the AIC of models with the same random effect structure, but a different fixed effect structure (here parameter values were maximum likelihood estimates). In case of over-dispersion, the standard errors were corrected using a quasi-GLM model (Zuur et al. 2009). Accordingly to Zuur et al. (2009) models were validated looking at response residuals (observed minus fitted values – Fig. S2 and Fig. S3 in Supplementary Material).

Using a similar mixed model approach we tested the effects of the relevant predictors and their interactions on the structural parameters (number of trees/stools, number of stems per plot and their basal area and mean height in the ETL) using the following model (eqn. 2):

$$y = \alpha + \beta_1 FT \cdot C + \beta_2 FT \cdot N \\ + \beta_3 SA + \beta_4 SI + 1 | FS + \varepsilon$$

The number of trees/stools and the number of stems were fitted using the “glmer” function with a Poisson error distribution, as for the species richness models. Basal area and mean height were fitted allowing for random variation across “forest stands” using the “lmer” with a Gaussian error distribution. In this case, fixed effect models were tested using the “gls” from the “nlme” package (Pinheiro et al. 2013) with a Gaussian error distribution, and parameter estimation was calculated with a restricted maximum likelihood. Basal area was preliminary log-transformed to match the requirements of normality and homoscedasticity.

For the optimal models selected for each variable, we calculated the R-squared (R^2),

i.e., the fraction of the total variation in the response variable accounted for by the model. For models with fixed effects only, the adjusted R^2 of the linear model was reported; for models that also contained random effects, a conditional R^2 was calculated using the “MuMIn” package (Barton 2013), according to Nakagawa & Schielzeth (2013).

Results

Soil, light and herbaceous cover

All kiln plots (KP) were characterized by a well distinct layer, 10-46 cm thick, of black soil formed by the accumulation of charcoal, with fragments of medium-large size. This layer was sharply delimited from the underlying mineral soil layer, whose color depended mainly on the parent rock material. The total carbon content in the charcoal layer of KP was almost two-fold higher than in CP (control plots) in sclerophyllous and oak forests, and significantly higher in KP of beech forests, though to a lower extent (Tab. 2). The same trend was generally recorded for N, but such differences were significant for oak forests only. Regarding pH, higher values were found in KP of the beech forest, while no significant differences were detected for the other two forest types. Both light availability (PAR) and herbaceous cover (HC) were higher in KP compared with CP in oak forests, while no significant differences could be found in the other two forest types (Tab. 2).

Seedlings density

In general, seedling density of the dominant tree species (Fagaceae) increased

with light availability, decreased with herb cover, and was affected by forest type and by its interaction with the above variables. Moreover, density decreased on steeper slopes and increased on south-facing slopes. No significant differences between KP and CP were found, with the exception of *Quercus pubescens* in sclerophyllous forests ($p = 0.04$ – Tab. 3). However, total C content had a weak negative effect on seedling density, while the latter was positively influenced by the N content (+1.68). The random effect of forest stands was not included in the selected model (Tab. 4), which explained 57% of the total variation.

Composition and species richness

Concerning the composition of woody species in the UVL, NMDS analysis yielded different results for the three forest types. Significant differences between KP and CP were found in oak forests ($p_{perm} = 0.008$), with a similar level of compositional variation within these two plot types ($p_{disp} = 0.541$ – Fig. 2). Beech and sclerophyllous understorey showed no differences in species composition between the two plot types.

In total, 45 woody species (26 trees and 19 shrubs) were recorded in the UVL of the 122 (= 61 \times 2) plots. In this layer, γ -diversity was always higher in KP compared to CP for all three forest types (22 vs. 14 in sclerophyllous forests; 28 vs. 22 in oak forests; 13 vs. 9 in beech forests – Fig. 3a). In sclerophyllous forests, nine species were unique to KP, while only one to CP; in oak forests, eight species were unique to KP and two to CP; in beech forests, four species were unique to KP (*Acer platanoides*, *Castanea sativa*) and one to CP (*Prunus avium* – Tab. S1 in Supplementary Material). Species richness in the UVL was significantly higher in KP compared with CP in oak forests; the same trend was also recorded for the other two forest types, though without significant differences (Fig. 3a).

The ETL included a total of eight tree and nine shrub species. In contrast to the UVL, γ -diversity was larger in CP than in KP (Fig. 3b). In the KP of sclerophyllous forests, this layer included four species (present in 16.7% of the plots – Tab. S1 in Supplementary Material), compared to seven species in the total sample of CP (Fig. 3b). In oak forests, no species were recorded in KP, while eight were present in this layer in 72.7% of the CP plots (Fig. 3b, Tab. S2). No

Tab. 3 - Density of seedlings of the dominant tree species in the charcoal kiln (KP) and control plots (CP) of the three forests types, with significance of the differences (Prob) after Mann-Whitney U-test.

Species	Forest type	Dominant seedling density/plot		Prob
		KP (n=5)	CP (n = 5)	
<i>Quercus petraea</i>	Oak	2.50	2.83	n.s.
<i>Quercus ilex</i>	Oak	0.83	2.83	n.s.
<i>Quercus cerris</i>	Oak	4.67	7.33	n.s.
<i>Quercus pubescens</i>	Oak	0.50	0.50	n.s.
<i>Quercus ilex</i>	Sclerophyllous	5.00	18.40	n.s.
<i>Quercus cerris</i>	Sclerophyllous	0.80	0.80	n.s.
<i>Quercus pubescens</i>	Sclerophyllous	3.00	7.00	0.041*
<i>Quercus suber</i>	Sclerophyllous	3.00	1.40	n.s.
<i>Fagus sylvatica</i>	Beech	29.00	28.20	n.s.

Tab. 4 - Optimal fixed-effects models for tree seedling density (SD), woody species richness (SR) at the plot level in the understorey (UVL) and established regeneration layer (ETL) based on AIC selection. The random factor “forest stand” was not significant. For each model factor, the relative change in the response variable (\pm standard error) is indicated, as compared to the reference level for the predictor forest type (beech forest) or for a unit increment of the continuous predictors. R^2 refers to the fraction of the variation explained by the optimal model structure. (HC): herbaceous cover; (C): soil carbon content; (N): soil nitrogen content; (SA): slope aspect; (SI): slope inclination; (df): degrees of freedom; (***) $: p<0.001$; (**) $: p<0.01$; (*) $: p<0.05$. (1): corrected for overdispersion with quasi-Poisson distributions.

Model factors	Level	SD ¹	SR-UVL	SR-ETL
Intercept	-	3.59 ± 0.47	0.66 ± 0.20	-4.20 ± 1.50
FT	Oaks	$-1.36 \pm 0.64^{***}$	$1.01 \pm 0.18^{***}$	$4.20 \pm 1.49^{**}$
	Sclerophyllous	0.33 ± 0.51	$0.76 \pm 0.19^{***}$	$4.78 \pm 1.49^{**}$
HC	-	$-0.07 \pm 0.27^{**}$	$0.14 \pm 0.04^{***}$	-
C	-	$-0.13 \pm 0.09^{***}$	$0.06 \pm 0.02^*$	$-0.27 \pm 0.07^{***}$
N	-	1.68 ± 2.44	$-1.35 \pm 0.49^*$	$3.32 \pm 1.29^*$
PAR	-	0.002 ± 0.004	0.00 ± 0.00	-
SA	-	0.007 ± 0.27	-	-
SI	-	-0.02 ± 0.10	-	0.73 ± 0.41
FTxPAR	Oaks	0.001 ± 0.01	0.01 ± 0.00	-
	Sclerophyllous	-0.02 ± 0.03	0.01 ± 0.00	-
FTxHC	Oaks	$0.14 \pm 0.08^{***}$	$-0.09 \pm 0.04^*$	-
	Sclerophyllous	-0.04 ± 0.40	$-0.08 \pm 0.04^*$	-
FTxSI	Oaks	$0.18 \pm 0.17^*$	-	-0.55 ± 0.43
	Sclerophyllous	-0.35 ± 0.76	-	$-1.00 \pm 0.47^*$
df	-	121	121	31
R ²	-	0.658	0.658	0.574
Overdispersion	-	0.747	0.747	2.47

tree species were found in the ETL of KP in beech forests (Fig. 3b), while beech was present in 14.3% of the CP (Tab. S3). Consequently, species richness at the plot-level in ETL was always significantly higher in CP compared with KP in all forest types (Fig. 3b).

Mixed model analysis showed that the random variation across the examined forest stands was not relevant for species richness in the two layers, while the fixed effect of forest type was always significant, explaining 66% of the total variation in the UVL and a lower proportion (26%) in ETL (Tab. 4). In the UVL, C content affected positively SR, while N had a negative effect. Moreover, light and herb cover also increased SR, though to a varying extent





Fig. 2 - Scatter plot from NMDS based on Bray Curtis dissimilarity index of woody species in the understorey (UVL) of oak forests. Differences in woody species composition of the understorey of kiln (KP) and control plots (CP) were significant. p_{perm} indicates the combined significance of the location and dispersion effect, based on PERMANOVA with 999 permutations; p_{disp} indicates the significance of the dispersion effect.






Fig. 4 - Differences in structural variables of the ETL layer (1.30-4 m) between kiln (KP) and control plots (CP) for each forest type. (A): mean number of stools per plot; (B): mean number of stems per plot; (C) mean basal area (G, cm^2); (D) mean height of tree stems (m). Error bars represent the standard error. (***): $p < 0.001$; (ns): $p > 0.05$.

depending on the forest type (Tab. 4). In the ETL, SR increase was mainly due to N (+3.1) and slope inclination (+0.73), with differences among forest types, while its decrease was mainly associated with C content (Tab. 4).

Structural parameters

Regarding the structural variables in the ETL, large differences were found between

the three forest types. As expected, the density of trees and shrubs in the sclerophyllous and oak stands was considerably higher than in the beech stands (Fig. 4). Remarkable differences were found comparing KP with CP. No stools or stems were present in KP of oak and beech forests (Fig. 4a, Fig. 4b). In sclerophyll forests, though a few species were present, the mean basal area of KP was always much

lower than in CP (Fig. 4c). The height of these species was also significantly lower in KP (Fig. 4d).

Looking at the effect of the measured variables on such parameters, FT (forest type), C and N were the main predictors. In particular, C content was the main factor negatively affecting the number of stools and stems, especially in oak forests. Contrastingly, N content was strongly and pos-

Tab. 5 - Optimal mixed-effects models for the structural variables in established tree regeneration layer, based on AIC selection. The random factor “forest stand” was significant for number of stools and stems. For each model factor, the relative change in the response variable (\pm standard error) is indicated, as compared to the reference level for the predictor forest type (beech forest) or for a unit increment of the continuous predictors. R^2 refers to the fraction of the variation explained by the optimal model structure. Over-dispersion is shown for Poisson distribution. (C): soil carbon content; (N): soil nitrogen content; (SA): slope aspect; (SI): slope inclination; (df): degrees of freedom; (1): Parameters corrected for over-dispersion with quasi-Poisson distributions; (2): Significant random effect; (3): Response variable log-transformed; (***): $p < 0.001$; (**): $p < 0.01$; (*): $p < 0.05$.

Model factors	Levels	No. Stools ¹⁻²	No. Stems ¹⁻²	Basal area ³	Stem height
Intercept	-	-0.08 \pm 1.07	0.30 \pm 2.10	-0.69 \pm 1.32	-0.88 \pm 0.65
FT	oaks	2.08 \pm 1.32***	2.21 \pm 2.58**	3.62 \pm 2.43	2.77 \pm 0.82***
	sclerophyllous	3.77 \pm 1.39***	8.78 \pm 2.73**	8.27 \pm 2.57***	2.68 \pm 0.88***
C	-	-0.04 \pm 0.20***	-0.11 \pm 0.41*	0.11 \pm 0.25	0.05 \pm 0.09
N	-	1.19 \pm 3.83***	1.50 \pm 7.85*	-	-
FTxC	Oaks	-0.13 \pm 0.22	-0.17 \pm 0.46	-0.28 \pm 0.28	-0.20 \pm 0.11*
	Sclerophyllous	1.19 \pm 0.24	-0.82 \pm 0.50	-0.53 \pm 0.31*	-0.17 \pm 0.11
FTxN	Oaks	0.17 \pm 4.49	1.19 \pm 9.18	-	-
	Sclerophyllous	1.61 \pm 4.95	6.02 \pm 10.01	-	-
SA	-	-0.002 \pm 0.2	-	-	-
SI	-	-	-	0.28 \pm 0.41	-
FTxSI	Oaks	-	-	1.09 \pm 0.73	-
	Sclerophyllous	-	-	1.13 \pm 0.92	-
Variance	Forest stand	0.37 \pm 0.60	0.33 \pm 0.57	-	-
df	-	121	121	121	121
R ²	-	0.424	0.438	0.291	0.211
Overdispersion	-	1.62	3.43	-	-

itively related with number of stools and stems, both alone and in the interaction with forest type, while it was not significant for the other two variables. Moreover, slope inclination and aspect had other minor effects on structural variables (Tab. 5). The models for the number of stools and stems explained up to the 42% and 44% of the variation, respectively, while the models for the mean basal area and height explained not more than 30% of the total variation. The best model for number of stools and stems included the random factor of the forest stand location (Tab. 5).

Discussion

The results of this study, obtained by analyzing three distinct forest types, demonstrated that former charcoal platforms represent a peculiar microhabitat for the regeneration of woody species.

The effect of kiln platforms on the γ -diversity in the understorey layer was clearly positive, especially in oak forests, where richness was also enhanced at the plot-level. In the latter forest type, kiln platforms also affected the composition in woody species, in line with previous studies on the understorey in mixed oak stands of the same area (Carrari et al. 2016a), as well as in beech forests of Germany (Wittig et al. 1999). Interestingly, no selection effect seems to occur at this early stage of regeneration, since the species recorded have different ecological requirements and functional traits, especially in thermophilous oak forests (Carrari et al. 2016b). Regeneration of pioneer shrub species such as *Cytisus scoparius*, *Crataegus monogyna* and *Prunus spinosa* occurred in the understorey mixed with those of early- and late-successional trees, such as *Fraxinus ornus* and *Quercus ilex*, respectively, depending on the forest type.

Recent findings have proven that kiln platforms are microhabitats which favor species diversity in the understorey (Carrari et al. 2016a). The results of our models revealed that various factors significantly affect woody species composition in the understorey, among which herb cover. Facilitation mechanisms among plants (Callaway 1995) may explain the positive influence of herbaceous cover on tree species richness in platforms of oak forests. On the other hand, herb cover negatively influenced the seedling density of dominant trees, likely due to the higher competitive ability of herbaceous plants for soil nutrients (Lyon & Sharpe 2003). Moreover, other important factors affecting woody species richness on the platforms are the higher C content in the soil and the increased light availability, the latter being a major driver of understorey diversity and productivity in European forests (Axmanová et al. 2011, Ewald 2008). Indeed, charcoal platforms may initially act as small canopy gaps, increasing the light available on the forest ground and positively affecting the early regeneration stages (Poulson

& Platt 1989, Beckage et al. 2008). It is well documented that canopy gaps can provide recruitment opportunities for tree seedlings and thus increase the number of species, promoting forest diversity (Busing & White 1997). In beech forests, the soil pH was higher in our platforms than in control plots, confirming previous findings in other kiln sites in North America and Europe (Mikan & Abrams 1995, 1996, Young et al. 1996, Wittig et al. 1999, Criscuoli et al. 2014). This is likely another positive factor for species richness, also based on evidence from deciduous woodlands in central Europe (Chytry et al. 2003, Axmanová et al. 2011). On the contrary, the higher N availability resulted to negatively affect woody species richness, since most plants are outcompeted by few species able to better exploit this resource (Clark et al. 2007).

Regarding the effects of kiln platforms on the established regeneration layer (1.3–4 m), the response of woody species was generally largely negative. Remarkably, a similar detrimental influence on the woody species was found in the old charcoal platforms of south-eastern Pennsylvania (Mikan & Abrams 1995) and the Appalachian mountains in north America (Young et al. 1996). In our study, both total (γ -diversity) and plot-level species richness were generally lower on the platforms than in the control plots for both oak and sclerophyllous forests, clearly indicating less suitable conditions for trees and shrubs at later stages of their development.

A similar negative effect was detected for most structural variables. Although seedling density in the understorey was similar between platforms and control plots, the number of stools and stems per plot above 1.3 m (ETL) was much lower in platforms than in the adjacent stands of sclerophyllous forests, or even stools and stems are completely lacking in oak and beech forests.

Overall, these findings suggest that such contrasting “kiln platform effect” is not simply a “forest gap effect”, but is driven by peculiar factors that are likely associated with the long-lasting production of wood charcoal at the same sites. According to previous works (Mikan & Abrams 1995, 1996, Carrari et al. 2016a), these include abiotic conditions, most likely in the soil, and possibly other components of biotic nature. The results of our models allow to exclude the increased soil nitrogen content as a negative factor for species richness, as well as the number of stems and stools. Instead, there is evidence for a general negative effect of soil carbon content on species richness and most structural parameters, with differences among forest types. Hence, there is a contrasting response of woody species to increased soil C (positive in the UVL and negative in ETL) and N content (negative in the UVL and positive in ETL).

The high C content in the platform soil is

mainly due to the large proportion of charred fragments, as recently demonstrated for a hearth in the Eastern Alps (Criscuoli et al. 2014). Thus, the negative effect on the established regeneration layer are likely associated with the charcoal accumulation in the soil, rather than to its carbon content per se. Consequences of charcoal accumulation on tree growth are still poorly known. A positive influence has been reported by authors who support the “biochar” practice to promote forest restoration (Thomas & Gale 2015). Indeed, previous studies suggested that the higher availability of some macro- and micronutrients in these soils could exert a positive effects on microbial communities, as in the case of the south American Terra Preta soils (Glaeser et al. 2001, 2002, Lehmann et al. 2003, Kim et al. 2007) and charcoal hearths in the Alpine region (Criscuoli et al. 2014). In this context, our results are in line with the positive effect of charcoal on nutrient availability (especially due to the reduced leaching of nitrogen) reported in several agricultural biochar studies (Bell & Worrall 2011, Knowles et al. 2011, Ventura et al. 2012).

On the other hand, less favourable effects of charcoal on plant growth have been reported in the literature, such as an altered nutrition as a consequence of the poor development of mycorrhizal fungi. For example, experimental “biochar” studies on agricultural crops showed its negative effect on arbuscular mycorrhizal fungi (Warnock et al. 2007), and a decrease in both bio-available organic C and N in their ectomycorrhizal system (Wallstedt et al. 2002). Further, Gaur & Adholeya (2000) found that the biochar media hampered the P uptake by host plants, indicating that in some cases charcoal may reduce the formation of mycorrhiza by decreasing the nutrient availability or creating unfavorable nutrient ratios in the soil (Wallstedt et al. 2002).

Another consequence of charcoal addition is the alteration of the soil water retention. Although charcoal is known to positively affect soil water retention due to its porosity, the real availability of charcoal-adsorbed water to plants still needs to be assessed (Karhu et al. 2011).

Finally, we cannot exclude that other factors not considered in this study could partly explain the “kiln platform effect”, such as the repeated soil heating on the platforms. It is well documented that slash pile burning in forests can cause cascading effects on soil structure (e.g., porosity) and chemical properties (e.g., pH, nutrient availability) and has a negative impact on the populations of arbuscular mycorrhizal fungi (Korb et al. 2004) and microbial communities (Jiménez Esquilín et al. 2007). In addition, Mikan & Abrams (1995) suggested the presence of potentially harmful soluble salts in the soil of charcoal platforms that may cause physiological drought. In our study, the presence of drought-tolerant species in the overstorey of the char-

coal sites (e.g., *Fraxinus ornus*, *Arbutus unedo*) and the lack of some more mesophytic species found in the overstorey of the controls (e.g., *Acer campestre*, *Carpinus betulus*) may lend circumstantial support to this hypothesis.

Conclusions

The extensive networks of old charcoal platforms in Mediterranean forests provide a natural experimental setting to investigate the legacy effects of an ancient form of forest use on tree regeneration dynamics. In this work, we demonstrated a general positive influence of kiln platforms on woody species at their first stage of regeneration in three different forest types. On the other hand, we found that the further growth and development of trees is negatively affected by modified soil and environmental conditions that prevent their access and establishment into the higher layers, thus causing a substantial lack of forest recolonization. This effect hinders or at least slows down the recolonization of even the oldest kiln platforms, unlike in forest gaps of different origin. Whether charcoal accumulation *per se* has a negative impact on tree growth remains an open issue which should be investigated on different species with an experimental approach.

List of abbreviations

KP: charcoal kiln plot; CP: control plot; UVL: understorey vegetation layer; ETL: established tree regeneration layer; SR: species richness.

Acknowledgements

Authors wish to thank Giacomo Certini (University of Firenze, Italy), Giovanni Mastrolonardo (University of Bologna, Italy) and Davide Ascoli (University of Torino, Italy) for discussion and comments that improved the first version of the manuscript; Giovanni Quilghini (Corpo Forestale dello Stato, UTB Pratovecchio, AR, Italy) for allowing field sampling in the Badia Prataglia forest, and Andrea Barzaghi for his help during the field work. This study was supported by funds from the Italian Ministry of University and Scientific Research and the University of Firenze to FS; the research of KV is supported by the ERC Consolidator Grant 614839 - PASTFORWARD.

References

- Akaike H (1973). Information theory and an extension of the maximum likelihood principle. In: Proceedings of "2nd International Symposium on Information Theory" (Petrov BN, Paki FC eds). Tsahkadsor (Armenia, USSR) 2-8 Sep 1971. Akademia Kiado, Budapest, Hungary, pp. 267-281.
- Anderson M (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26: 32-46. - doi: [10.1111/j.1442-9993.2001.01070.pp.x](https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x)
- Antal MJ, Gronli M (2003). The art, science, and technology of charcoal production. *Industrial and Engineering Chemistry Research* 42 (8): 1619-1640. - doi: [10.1021/ie0207919](https://doi.org/10.1021/ie0207919)
- Attwood M, Iavicoli C, Seppilli T (1981). Il Carbonao, una tecnica tradizionale di preparazione del carbone da legna nella montagna Cortonese [The charcoal burner, a traditional technique of charcoal production on the mountains around Cortona]. La Zincografica Fiorentina, Firenze, Italy, pp. 47. [in Italian]
- Axmanová I, Zelený D, Li CF, Chytrý M (2011). Environmental factors influencing herb layer productivity in Central European oak forests: insights from soil and biomass analyses and a phytometer experiment. *Plant and Soil* 342 (1-2): 183-194. - doi: [10.1007/s11104-010-0683-9](https://doi.org/10.1007/s11104-010-0683-9)
- Bates D, Maechler M, Bolker B, Walker S, Christensen RHB, Singmann H, Dai B, Eigen C, Rcpp L (2014). Package "lme4". R Foundation for Statistical Computing, Vienna, Website. [online] URL: <http://cran.r-project.org/web/packages/lme4/lme4.pdf>
- Barton K (2013). MuMIn: multi-model inference. R package version 1(5), R Project for Statistical Computing, Vienna, Austria, Website. [online] URL: <http://cran.r-project.org/web/packages/MuMIn/MuMIn.pdf>
- Beckage B, Kloepel BD, Yeakley JA, Taylor SF, Coleman DC (2008). Differential effects of understory and overstory gaps on tree regeneration. *The Journal of the Torrey Botanical Society* 135: 1-11. - doi: [10.3159/07-RA-034.1](https://doi.org/10.3159/07-RA-034.1)
- Bell MJ, Worrall F (2011). Charcoal addition to soils in NE England: a carbon sink with environmental co-benefits? *Science of Total Environment* 409: 1704-1714. - doi: [10.1016/j.scitotenv.2011.01.031](https://doi.org/10.1016/j.scitotenv.2011.01.031)
- Blondel J (2006). The "design" of Mediterranean landscapes: a millennial story of humans and ecological systems during the historic period. *Human Ecology* 34: 713-729. - doi: [10.1007/s10745-006-9030-4](https://doi.org/10.1007/s10745-006-9030-4)
- Busing RT, White PS (1997). Species diversity and small-scale disturbance in an old growth temperate forest: a consideration of gap partitioning concepts. *Oikos* 78: 562-568. - doi: [10.2307/3545618](https://doi.org/10.2307/3545618)
- Callaway RM (1995). Positive interactions among plants. *The Botanical Review* 61 (4): 306-349. - doi: [10.1007/BF02912621](https://doi.org/10.1007/BF02912621)
- Cantiani M (1955). Ricerche sperimentali sulla carbonizzazione della legna di cerro e di faggio [Experimental researches on the charcoalizing process of the Turkey oak and beech timber]. *Italia Forestale e Montana* 10: 125-129. [in Italian]
- Carrari E, Ampoorter E, Coppi A, Verheyen K, Selvi F (2016a). Former charcoal kiln platforms serve as microhabitats that increase the understorey diversity in Mediterranean forests. *Applied Vegetation Science* 19: 486-497. - doi: [10.1111/avsc.12238](https://doi.org/10.1111/avsc.12238)
- Carrari E, Ampoorter E, Coppi A, Selvi F (2016b). Diversity of secondary woody species in relation to species richness and cover of dominant trees in thermophilous deciduous forests. *Scandinavian Journal of Forest Research* 31: 484-494. - doi: [10.1080/02827581.2015.1081981](https://doi.org/10.1080/02827581.2015.1081981)
- Cheng CH, Lehmann J, Thies JE, Burton SD (2008). Stability of black carbon in soils across a climatic gradient. *Journal of Geophysical Research* 113: 2005-2012. - doi: [10.1029/2007JC004106](https://doi.org/10.1029/2007JC004106)
- and Engineering Chemistry Research 42 (8): 1619-1640. - doi: [10.1021/ie0207919](https://doi.org/10.1021/ie0207919)
- Chytrý M, Tichý L, Roleček J (2003). Local and regional patterns of species richness in Central European vegetation types along the pH/calcium gradient. *Folia Geobotanica* 38 (4): 429-442. - doi: [10.1007/BF02803250](https://doi.org/10.1007/BF02803250)
- Clark CM, Cleland EE, Collins SL, Fargione JE, Gough L, Gross KL, Pennings SC, Suding KN, Grace JB (2007). Environmental and plant community determinants of species loss following nitrogen enrichment. *Ecology Letters* 10 (7): 596-607. - doi: [10.1111/j.1461-0248.2007.01053.x](https://doi.org/10.1111/j.1461-0248.2007.01053.x)
- Cremaschi M, Pizzi C, Valsecchi V (2006). Water management and land use in the terramare and a possible climatic co-factor in their abandonment: the case study of the terramara of Poviglio Santa Rosa (Northern Italy). *Quaternary International* 151: 87-98. - doi: [10.1016/j.quaint.2006.01.020](https://doi.org/10.1016/j.quaint.2006.01.020)
- Crisciuoli I, Alberti G, Baronti S, Favilli F, Martinez C, Calzolari C, Pesceddu E, Rumpel C, Viola R, Miglietta F (2014). Carbon sequestration and fertility after centennial time scale incorporation of charcoal into soil. *PLoS ONE* 9: e91114. - doi: [10.1371/journal.pone.0091114](https://doi.org/10.1371/journal.pone.0091114)
- Deforce K, Boeren I, Adriaenssens S, Bastiaens J, De Keersmaecker L, Haneca K, Tys D, Vandekerhove K (2013). Selective woodland exploitation for charcoal production. A detailed analysis of charcoal kiln remains (ca. 1300-1900 AD) from Zoersel (northern Belgium). *Journal of Archaeological Science* 40 (1): 681-689. - doi: [10.1016/j.jas.2012.07.009](https://doi.org/10.1016/j.jas.2012.07.009)
- European Environment Agency (2007). European forest types. Categories and types for sustainable forest management reporting and policy (2nd edn). EEA Technical Report 09/2006, European Environment Agency, Copenhagen, Denmark, pp. 111.
- European Soil Bureau Network (2005). Soil atlas of Europe. Office for Official Publications of the European Communities, European Commission, L-2995 Luxembourg, pp. 128.
- Ewald J (2008). Plant species richness in mountain forests of the Bavarian Alps. *Plant Biosystems* 142 (3): 594-603. - doi: [10.1080/11263500802410942](https://doi.org/10.1080/11263500802410942)
- Gaur A, Adholeya A (2000). Effects of the particle size of soilless substrates upon AM fungus inoculum production. *Mycorrhiza* 10: 43-48. - doi: [10.1007/s005720050286](https://doi.org/10.1007/s005720050286)
- Giorgerini C (2009). Carbone e carboniae [Charcoal and charcoal kilns]. I Quaderni del Circolo, Sassetta ed., Livorno, Italy, pp. 20. [in Italian]
- Glaser B, Lehmann J, Zech W (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal - a review. *Biology and Fertility of Soils* 35: 219-230. - doi: [10.1007/s00374-002-0466-4](https://doi.org/10.1007/s00374-002-0466-4)
- Glaser B, Haumaier L, Guggenberger G, Zech W (2001). The "Terra preta" phenomenon: a model for sustainable agriculture in the humid tropics. *Naturwissenschaften* 88 (1): 37-41. - doi: [10.1007/s001140000193](https://doi.org/10.1007/s001140000193)
- Jiménez Esquilín AE, Stromberger ME, Massman WJ, Frank JM, Shepperd WD (2007). Microbial community structure and activity in a Colorado Rocky Mountain forest soil scarred by slash pile burning. *Soil Biology and Biochemistry* 39: 1111-1120. - doi: [10.1016/j.soilbio.2006.12.020](https://doi.org/10.1016/j.soilbio.2006.12.020)
- Karhu K, Mattila T, Bergström I, Regina K (2011).

106

- Biochar addition to agricultural soil increased CH₄ uptake and water holding capacity - results from a short-term pilot field study. *Agriculture, Ecosystems and Environment* 140: 309-313. - doi: [10.1016/j.agee.2010.12.005](https://doi.org/10.1016/j.agee.2010.12.005)
- Kim J-S, Sparovek G, Longo RM, De Melo WJ, Crowley D (2007). Bacterial diversity of Terra Preta and pristine forest soil from the Western Amazon. *Soil Biology and Biochemistry* 39: 684-690. - doi: [10.1016/j.soilbio.2006.08.010](https://doi.org/10.1016/j.soilbio.2006.08.010)
- Knowles OA, Robinson BH, Contangelo A, Clucas L (2011). Biochar for the mitigation of nitrate leaching from soil amended with biosolids. *Science of Total Environment* 409: 3206-3210. - doi: [10.1016/j.scitotenv.2011.05.011](https://doi.org/10.1016/j.scitotenv.2011.05.011)
- Korb EJ, Johnson NC, Covington WW (2004). Slash pile burning effects on soil biotic and chemical properties and plant establishment: recommendations for amelioration. *Restoration Ecology* 12: 52-62. - doi: [10.1111/j.1061-2971.2004.00304.x](https://doi.org/10.1111/j.1061-2971.2004.00304.x)
- Landi M, Püssi P (1988). Il lavoro nei boschi. Boscaioli e carbonai a Luco e Grezzano tra il 1939 e il 1950 [The work in the forest. Lumberjacks and charcoal burners at Luco and Grezzano between 1939 and 1950]. Gruppo d'Erci, Grezzano, Borgo San Lorenzo, Italy, pp. 72. [in Italian]
- Lehmann J, Pereira da Silva J, Steiner C, Nehls T, Zech W, Glaser B (2003). Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal ramendments. *Plant Soil* 249: 343-357. - doi: [10.1023/A:1022833116184](https://doi.org/10.1023/A:1022833116184)
- Ludemann T (2003). Large-scale reconstruction of ancient forest vegetation by anthracology - a contribution from the Black Forest. *Phytocoenologia* 33 (4): 645-666. - doi: [10.1127/0340-269X/2003/0033-0645](https://doi.org/10.1127/0340-269X/2003/0033-0645)
- Ludemann T, Michiels HG, Nölken W (2004). Spatial patterns of past wood exploitation, natural wood supply and growth conditions: indications of natural tree species distribution by anthracological studies of charcoal-burning remains. *European Journal of Forest Research* 123 (4): 283-292. - doi: [10.1007/s10342-004-0049-z](https://doi.org/10.1007/s10342-004-0049-z)
- Ludemann T (2010). Past fuel wood exploitation and natural forest vegetation in the Black Forest, the Vosges and neighbouring regions in the western Central Europe. *Palaeogeography Palaeoclimatology and Palaeoecology* 291: 154-165. - doi: [10.1016/j.palaeo.2009.09.013](https://doi.org/10.1016/j.palaeo.2009.09.013)
- Ludemann T (2012). Airborn laser scanning of historical wood charcoal production sites - a new tool of kiln site anthracology at the landscape level. *SAGVNVM Extra* 13: 247-252.
- Lyon J, Sharpe WE (2003). Impacts of hay-scented fern on nutrition of northern Red Oak seedlings. *Journal of Plant Nutrition* 26: 487-502. - doi: [10.1081/PLN-120017661](https://doi.org/10.1081/PLN-120017661)
- Mikan CJ, Abrams MD (1996). Mechanisms inhibiting the forest development of historic charcoal hearths in southeastern Pennsylvania. Canadian Journal of Forest Research 26: 1893-1898. - doi: [10.1139/x26-213](https://doi.org/10.1139/x26-213)
- Mikan CJ, Abrams MD (1995). Altered forest composition and soil properties of historic charcoal hearths in southeastern Pennsylvania. Canadian Journal of Forest Research 25 (5): 687-696. - doi: [10.1139/x95-076](https://doi.org/10.1139/x95-076)
- Montanari C, Prono P, Scipioni S (2000). The study of charcoal-burning sites in the Apennine Mountains of Liguria (NW Italy) as a tool for forest history. In: "Methods and Approaches in Forest History" (Agnoletti M, Anderson S eds). CABI Publishing, CAB International, Wallingford, UK, pp. 79-91.
- Nakagawa S, Schielzeth H (2013). A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4: 133-142. - doi: [10.1111/j.2041-210X.2012.00261.x](https://doi.org/10.1111/j.2041-210X.2012.00261.x)
- Nelle O (2003). Woodland history of the last 500 years revealed by anthracological studies of charcoal kiln sites in the Bavarian Forest, Germany. *Phytocoenologia* 33 (4): 667-682. - doi: [10.1127/0340-269X/2003/0033-0667](https://doi.org/10.1127/0340-269X/2003/0033-0667)
- Oksanen J, Blanchet F G, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson G L, Solymos M, Stevens M H H, Wagner H (2015). Package "vegan". Community ecology package, version 2.3-5. R Project for Statistical Computing, Vienna, Austria, Website. [online] URL: <http://cran.r-project.org/web/packages/vegan/vegan.pdf>
- Pélachs A, Nadal J, Soriano JM, Molina D, Cunill R (2009). Changes in Pyrenean woodlands as a result of the intensity of human exploitation: 2,000 years of metallurgy in Vallferrera, northeast Iberian Peninsula. *Vegetation History and Archaeobotany* 18: 403-416. - doi: [10.1007/s00340-009-0218-6](https://doi.org/10.1007/s00340-009-0218-6)
- Pignatti S (1982). Flora d'Italia. Edagricole, Bologna, Italy, pp. 2324. [in Italian]
- Pinheiro J, Bates D, DebRoy S, Sarkar D (2013). "nlme": linear and non linear mixed effect models. R Project for Statistical Computing, Vienna, Austria, Website. [online] URL: <http://cran.r-project.org/web/packages/nlme/nlme.pdf>
- Poulson TL, Platt TL (1989). Gap light regime influences canopy tree diversity. *Ecology* 70: 553-555. - doi: [10.2307/1940202](https://doi.org/10.2307/1940202)
- Powell A (2008). The magnetic signatures of medieval and post-medieval pyrotechnical industries, PhD thesis, University of Bradford, UK, pp. 172.
- R Core Team (2014). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [online] URL: <http://www.R-project.org>
- Regione Toscana (2016). Settore idrogeologico regionale. Regione Toscana, Firenze, Italy, Website. [in Italian] [online] URL: <http://www.sir.toscana.it/>
- Thomas SC, Gale N (2015). Biochar and forest restoration: a review and meta-analysis of tree growth responses. *New Forests* 46 (5-6): 931-946. - doi: [10.1007/s11056-015-9491-7](https://doi.org/10.1007/s11056-015-9491-7)
- Valese E, Conedera M, Held AC, Ascoli D (2014). Fire, humans and landscape in the European Alpine region during the Holocene. *Anthropocene* 6: 63-74. - doi: [10.1016/j.ancene.2014.06.006](https://doi.org/10.1016/j.ancene.2014.06.006)
- Ventura M, Sorrenti G, Panzacchi P, Tonon G (2012). Biochar reduces nitrate leaching in an apple orchard. In: Proceedings of the "17th Nitrogen Workshop - Innovations for Sustainable Use of Nitrogen Resources" (Richards KG, Fenton O, Watson CJ eds). Springer Verlag, Wexford, Ireland, pp. 42-43.
- Wallstedt A, Coughlan A, Munson AD, Nilsson MC, Margolis HA (2002). Mechanisms of interaction between *Kalmia angustifolia* cover and *Picea mariana* seedlings. *Canadian Journal of Forest Research* 32: 2022-2031. - doi: [10.1139/x02-124](https://doi.org/10.1139/x02-124)
- Warnock DD, Lehmann J, Kuyper TW, Rillig MC (2007). Mycorrhizal responses to biochar in soil - concepts and mechanisms. *Plant and Soil* 300: 9-20. - doi: [10.1007/s11104-007-9391-5](https://doi.org/10.1007/s11104-007-9391-5)
- Wittig R, Walter S, Xie Y (1999). Ehemalige Meilerplätze als Mikrostandorte des *Galio odorati-Fagetum* im Luzulo-Fagetum [Former kiln sites as microsites of *Galio odorati-Fagetum* in Luzulo-Fagetum]. *Acta Biologica Benrodis* 10: 57-68. [in German]
- Young MJ, Johnson JE, Abrams MD (1996). Vegetative and edaphic characteristics on relic charcoal hearths in the Appalachian Mountains. *Vegetatio* 125 (1): 43-50. - doi: [10.1007/BF00045203](https://doi.org/10.1007/BF00045203)
- Zimmerman AR (2010). Abiotic and microbial oxidation of laboratory produced black carbon (biochar). *Environmental Science and Technology* 44: 1295-1301. - doi: [10.1021/es903140c](https://doi.org/10.1021/es903140c)
- Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM (2009). Mixed effects models and extensions in ecology with R. Springer, New York, USA, pp. 574.

Supplementary Material

- Fig. S1** - Building process of a charcoal kiln.
- Tab. S1** - Woody species in sclerophyllous forests.
- Tab. S2** - Woody species in oak forests.
- Tab. S3** - Woody species in beech forests.
- Fig. S2** - Model residuals for species richness and tree seedling density.
- Fig. S3** - Model residuals for structural parameters.
- Tab. S4** - Environmental variables for the seedling density plot subsample.
- Link:** Carrari_1701@suppl001.pdf