



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

First evidence of microplastic ingestion by fishes from the Amazon River estuary

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

First evidence of microplastic ingestion by fishes from the Amazon River estuary / Pegado T.D.S.E.S, Schmid K., Winemiller K.O., David Chelazzi, Alessandra Cincinelli, Luigi Dei, Giarrizzo, T.. - In: MARINE POLLUTION BULLETIN. - ISSN 0025-326X. - STAMPA. - 133:(2018), pp. 814-821.
[10.1016/j.marpolbul.2018.06.035]

Availability:

The webpage <https://hdl.handle.net/2158/1137695> of the repository was last updated on 2022-03-20T16:45:07Z

Published version:

DOI: 10.1016/j.marpolbul.2018.06.035

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

1 **First evidence of microplastic ingestion by fishes from the Amazon River estuary**

2 Tamyrís de Souza e Silva Pegado¹, Kurt Schmid¹, Kirk O. Winemiller², David Chelazzi³,
3 Alessandra Cincinelli^{3,4}, Luigi Dei^{3,4}, Tommaso Giarrizzo¹.

4 ¹Laboratório de Biologia Pesqueira e Manejo dos Recursos Aquáticos, Universidade Federal do
5 Pará, Av. Perimetral 2651, Terra Firme, 66077830 Belém, Pará, Brazil.

6 ²Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX,
7 77843-2258, USA.

8 ³Consorzio Interuniversitario per lo Sviluppo dei Sistemi a Grande Interfase (CSGI), Via della
9 Lastruccia, 3, 50019 Sesto Fiorentino, Florence, Italy.

10 ⁴Department of Chemistry “Ugo Schiff”, University of Florence, Via della Lastruccia, 3, 50019
11 Sesto Fiorentino, Florence, Italy.

12

13 **Abstract**

14 This study investigated occurrence of microplastic particles in digestive tracts of fishes from the
15 Amazon River estuary. A total of 189 fish specimens representing 46 species from 22 families
16 was sampled from bycatch of the shrimp fishery. Microplastic particles removed from fish
17 gastrointestinal tracts were identified using Attenuated Total Reflectance – Fourier Transform
18 Infrared (ATR-FTIR). In total, 228 microplastic particles were removed from gastrointestinal
19 tracts of 26 specimens representing 14 species (30% of those examined). Microplastic particles
20 were categorized as pellets (97.4%), sheets (1.3%), fragments (0.4%) and threads (0.9%), with
21 size ranging from 0.38 to 4.16 mm. There was a positive correlation between fish standard length
22 and number of particles found in gastrointestinal tracts. The main polymers identified by ATR-
23 FTIR were polyamide, rayon and polyethylene. These findings provide the first evidence of
24 microplastic contamination of biota from the Amazon estuary and northern coast of Brazil.

25 **Keywords:** Brazil, Bycatch, Pollution, Trophic level

26 **Introduction**

27 During recent decades, changes in manufacturing and consumer behavior together with
28 insufficient waste management have resulted in accumulation of plastic debris in oceans
29 throughout the world (e.g., Costa and Barletta, 2015; Jambeck et al., 2015), with plastic now
30 composing between 60% and 80% of all marine debris (Barnes et al., 2009). It has been estimated
31 that nearly half of all plastic products are discarded in less than 12 months after production
32 (Hopewell et al., 2009). Once introduced into marine ecosystems, plastic waste becomes
33 fragmented as it disperses via wind and oceanic currents (Barnes et al., 2009; Lebreton et al.,
34 2012) and is distributed throughout the water column (Bellas et al., 2016). Plastic debris

1 accumulates not only in the open ocean, but also on beaches, mangrove forests and other coastal
2 habitats (Ivar do Sul et al., 2007). Although some plastic debris is dumped directly into marine
3 waters, rivers accumulate discarded material throughout their watersheds and transport it to the
4 oceans (Lechner et al., 2014; Vendel et al., 2017). Unfortunately, rivers and estuaries have
5 received relatively little attention with regard to the plastic pollution problem (Costa and Barletta,
6 2015), especially within the southern hemisphere (Cannon et al., 2016).

7 Reports of interactions between marine fauna and plastic debris have increased by 75%
8 over the last two decades, including 267 species reported in 1997 (Laist, 1997) and 693 species
9 reported in 2015 (Gall and Thompson, 2015). Plastic waste in the environment negatively
10 impacts biota, including entanglement of animals within large items (macroplastics) and
11 ingestion of microplastics (particles < 5 mm) by organisms, with subsequent transfer within the
12 food web (Fossi et al., 2012; Cole et al., 2013; Ivar do Sul and Costa, 2014). Ingestion of plastic
13 can affect organisms both physically and physiologically, including direct mortality from
14 entanglement and choking as well as sub-lethal effects, such as compromised feeding, digestion,
15 and reproduction activities (Gregory, 2009; Vendel et al., 2017). Exposure to chemical pollutants
16 that bind to plastic particles has become a major concern, especially when chemicals
17 bioaccumulate in fish destined for human consumption (Teuten et al., 2009). The effects of
18 human consumption of organisms that contain microplastics are still poorly understood. Some
19 evidence has been reported that plastic particles may cause immunotoxic responses, resulting
20 either from chemical exposure or particle-induced mechanical stress (Seltenrich, 2015).

21 In aquatic and marine environments, plastics undergo a continuous process of
22 disintegration from the action of water and wind causing abrasion from contact with solid
23 particles, and through chemical decomposition by exposure to solar radiation (Moore, 2008;
24 Barnes et al., 2009). Plastic debris is classified as macroplastics (particle diameter >25 mm),
25 microplastics (diameter <5 mm) (GESAMP, 2015) or mesoplastics (5-25 mm) (Jabeen et al.,
26 2017). Microplastics are further classified according to their origin. Primary microplastics are
27 resin pellets and microbeads used in cleaning products, cosmetics, medicines and other products;
28 secondary microplastics are formed from the fragmentation of larger meso- and macroplastics
29 (Cole et al., 2011). Plastic pellets are used worldwide as a raw material in the production of
30 plastic products (Ogata et al., 2009). With exposure to solar radiation, plastic pellets often lose or
31 change their initial white or translucent coloration and many anthropogenic and biogenic
32 chemicals can be adsorbed by their surface (Endo et al., 2005; Miranda et al., 2016).

1 Hydrophobic characteristics of plastics allow them to function as vectors for organic
2 contaminants and heavy metals (Colabuono et al., 2010; Holmes et al., 2012).

3 Many fishes ingest tiny plastic particles either intentionally or accidentally while feeding
4 in the water column or the benthos (Browne et al., 2010). Most investigations of microplastic
5 ingestion by wild fish have been conducted in the northern hemisphere (e.g. Boerger et al., 2010;
6 Phillips and Bonner, 2015), especially in Europe (e.g. Neves et al., 2015; Bellas et al., 2016;
7 McGoran et al., 2017) and North America (e. g. Carson, 2013; Petters e Bratton, 2016).
8 Microplastic ingestion by fishes in the Southern Hemisphere has been documented by studies
9 performed in Africa (e.g. Biginagwa et al., 2016; Naidoo et al., 2016), Australia (e.g. Cannon et
10 al., 2016), Easter Island (e.g. Ory et al., 2017), Indonesia (e.g. Rochman et al., 2015), and South
11 America (Mizraji et al., 2017; Ory et al., 2017). Studies in Brazil have been conducted in the
12 northeastern and southeastern regions (e.g. Possatto et al., 2011; Ferreira et al., 2016; Silva-
13 Cavalcanti et al., 2017), with no investigations as yet for the northern region that includes the
14 Amazon River estuary.

15 Brazil's northern coastline has low human population density and contains the world's
16 second-longest, continuous area of largely undisturbed mangrove forest (ca. 7,000 km²)
17 (Giarrizzo and Krumme, 2008). In 2016, an extensive and biodiverse reef system (~9,500 km²)
18 was discovered offshore from the mouth of the Amazon River (Moura et al., 2016). This
19 discovery, paired with the fact that 20% of Brazil's fisheries landings come from the northern
20 coast (Krumme et al., 2015), lends urgency to the need to improve knowledge about plastic
21 pollution in the region. Based on experiences in estuaries from northeastern Brazil, Costa and
22 Barletta (2015) identified the Amazon River estuary as a priority area for future studies on marine
23 plastic pollution. The goal of our study was to investigate the presence of microplastics ingestion
24 by fishes from the Amazon River estuary on the coast of Brazil. We hypothesized that quantity
25 and size of the ingested particles increases with fish body size, weight and vertical trophic
26 position within the estuarine food web.

27


28 **Material and Methods**

29 Study area

30 Brazil's North Coast extends over 1,400 km along the states of Amapá and Pará, covering
31 an area of approximately 488,000 km² and a variety of ecosystems including mesophotic reefs,
32 islands, tidal flats, and estuaries with extensive mangrove forests (Marceniuk et al., 2013) (Fig.

1 1). The region's equatorial humid climate (Kottek et al., 2006) has annual rainfall up to 3,300
2 mm and average annual temperatures of 27.5 to 29.5 °C (Pereira et al., 2009).

3



4

5 **Fig.1.** Location of the Amazon River estuary in northeastern Brazil (inset) showing the
6 survey area (yellow shaded area).

7

8 The region includes the estuary of the world's largest river, the Amazon, with its mean
9 annual discharge of 6.3 trillion m³ of freshwater, 1.2 billion tons of sediments and 290 million
10 tons of solutes that flow onto the continental shelf (Oltman, 1968; Meade et al., 1985; Nittrouer et
11 al., 1995). The Amazon's freshwater plume can seasonally expand up to 120 km from the river
12 mouth to the open ocean where salinities can close to zero.

13 The Amazon's massive freshwater discharge affects oceanographic processes, creating
14 dynamic system of currents and tidal fluxes. The large sediment discharge contributes to high
15 primary and secondary productivity, sustaining important artisanal and commercial fisheries
16 (Neiva and Moura 1977, Wolff et al. 2000). Estuarine fishes and crustaceans have great economic
17 importance and many of them interact with substrates, influencing physical and chemical
18 processes, including nutrient dynamics (Lana et al., 1996).

19

1 Fish sampling

2 Fish specimens analyzed in this study were obtained from bycatch of the southern brown-
3 shrimp fishery, which is monitored by the Center for Research and Management of Fishing
4 Resources of Brazil's North Coast (CEPNOR) of the Brazilian Ministry of the Environment
5 (MMA). Samples were obtained from 104 bottom trawls carried out between July 2015 and
6 August 2016 at depths varying from 35 to 85 m. Due to ship's limited freezer space, a sample of
7 the most abundant fish species were collected from each haul, frozen immediately, and stored for
8 up to 2 weeks at -20 °C in the laboratory prior to processing.

9

10 Sample processing

11 In the laboratory, the fishes were identified to species level and each specimen was
12 measured for standard length (SL) with a caliper (0.01 cm precision) and weighed with a digital
13 scale (Marte BL3200H; 0.01 g precision). For each species trophic level was assigned according
14 to values provided in FishBase (Froese and Pauly, 2017). To remove the digestive tract, a
15 longitudinal incision was made in the abdominal area using surgical forceps and a scalpel. Each
16 digestive tract (stomach and intestine) was cut longitudinally, and contents were washed into a
17 Petri dish using 70% ethanol.

18 Stomach and intestine contents were examined under a stereo-microscope (Opton Tim-
19 2b) at 6.5× to 50× magnification. Plastic particles were separated from other ingested particles,
20 counted, classified according to shape and colour, measured (diameter in longest dimension to
21 0.001 mm precision), and photographed using a ZEISS SteREO Discovery V12 stereo
22 microscope with the Zen software (blue edition, v2.0, Zeiss, Oberkochen, Germany).

23 Before and after each procedure for each specimen, all work surfaces and instruments
24 were thoroughly cleaned with 70% ethanol and a new, clean pair of latex gloves was worn. To
25 test for potential presence of airborne plastic fibers at the work station, a clean (using 70%
26 ethanol) glass Petri dish that was placed at the work station at the start of each day was examined
27 with the stereo-microscope at the end of the day.

28 A sample of each microplastic type recorded in the study was randomly selected for
29 polymer identification. Recent studies showed that Fourier Transform Infrared (FTIR)
30 spectroscopy is the most reliable method to identify the composition of maritime plastic debris,
31 using either single-element or Focal Plane Array (FPA) detectors (Srinivasa Reddy et al., 2006;
32 Mecozzi et al., 2016; Cincinelli et al., 2017). In this study, FTIR spectra were collected in

1 Attenuated Total Reflectance (ATR) mode, using a single-element MCT detector, which was
2 deemed as the optimal set up, given the type and morphology of the plastic samples. The ATR-
3 FTIR analysis of the samples was carried out using a Cary 620-670 FTIR microscope, equipped
4 with a GeATR crystal (Agilent Technologies). The spectra were recorded directly on the samples
5 with a spectral resolution of 8 cm^{-1} , acquiring 128 scans for each spectrum in the $4000\text{-}650\text{ cm}^{-1}$
6 spectral range.

7

8 Statistical analysis

9 The percentage of frequency of occurrence of microplastics within digestive tracts was
10 calculated using the following formula: $\text{FO}\% = (\text{Ni}/\text{N}) \times 100$, where FO% = frequency of
11 occurrence of microplastic particles; Ni = number of gastrointestinal tracts that contained
12 microplastic particles; N = total number of gastrointestinal tracts examined.

13 Kendall's rank correlation was performed to assess the association between i) number of
14 ingested microplastic particles and fish body size, ii) number of ingested microplastic particles
15 and fish weight, iii) number of ingested microplastic particles and fish trophic level, and iv) the
16 size of microplastic particles and fish body size. Whenever the correlation was significant, a
17 linear regression was performed to derive an equation predicting the relationship between the
18 response variable and independent variable. Statistical tests only included specimens that had
19 microplastic particles in the gastrointestinal tract, and were performed using the R statistical
20 package (R Core Team, 2017).

21

22 Results

23 Overall, 189 fish specimens representing 46 species from 22 families were analyzed
24 (Table 1). Standard length (\pm SD) averaged $24.7 (\pm 13.3)$ cm, varying from 10.2 to 92 cm, and
25 average body weight was $558.3\text{ g} (\pm 1429.2)$ ranging from 28.2 g to 10000g. All fishes are
26 carnivorous and the species trophic level (TL), based on values reported in FishBase, ranged
27 from ranged from 3.1 to 4.5.

Table 1. Summary information for fishes examined for microplastic particles.

Family	Species	Trophic level	N° of fish	N° of fish with microplastic	Frequency of occurrence (%)	N° of microplastic particles in gastrointestinal tract	Average N° of microplastic particles per specimen**	N° of fish with microplastic in stomach	N° of microplastic particles in stomach	N° of fish with microplastic in intestine	N° of microplastic particles in intestine
Ariidae	<i>Bagre bagre</i> *	4	7	5	71.4	64	12.8	5	62	2	2
	<i>Bagre marinus</i> *	3.5	4	4	100	31	7.8	4	31	0	0
	<i>Notarius grandicassis</i>	4	4	0	0	0	0	0	0	0	0
Batrachoididae	<i>Batrachoides surinamensis</i>	3.7	1	0	0	0	0	0	0	0	0
Carangidae	<i>Caranx crysos</i>	4.1	3	0	0	0	0	0	0	0	0
	<i>Caranx hippos</i> *	3.6	3	3	100	92	30.7	3	92	0	0
	<i>Selene setapinnis</i>	3.7	1	0	0	0	0	0	0	0	0
	<i>Selene vomer</i> *	4.3	2	1	50	2	2.0	1	2	0	0
Ehippididae	<i>Chaetodipterus faber</i>	4.5	5	0	0	0	0	0	0	0	0
Haemulidae	<i>Anisotremus surinamensis</i>	3.6	1	0	0	0	0	0	0	0	0
	<i>Anisotremus virginicus</i>	3.6	1	0	0	0	0	0	0	0	0
	<i>Conodon nobilis</i>	3.6	8	0	0	0	0	0	0	0	0
	<i>Genyatremus luteus</i>	3.5	8	0	0	0	0	0	0	0	0
	<i>Haemulon plumieri</i>	3.8	13	0	0	0	0	0	0	0	0
	<i>Haemulon steindachneri</i>	3.7	5	0	0	0	0	0	0	0	0
	<i>Orthopristis ruber</i>	3.6	2	0	0	0	0	0	0	0	0
Lutjanidae	<i>Lutjanus analis</i> *	3.9	3	1	33.3	1	1.0	0	0	1	1
	<i>Lutjanus synagris</i> *	3.8	2	1	50	1	1.0	0	0	2	1
Muraenesocidae	<i>Cynoponticus savanna</i>	3.5	1	0	0	0	0	0	0	0	0
Muraenidae	<i>Gymnothorax ocellatus</i>	4.1	1	0	0	0	0	0	0	0	0
Myliobatidae	<i>Rhinoptera bonasus</i>	3.2	1	0	0	0	0	0	0	0	0
Narcinidae	<i>Narcine brasiliensis</i> *	3.2	6	1	16.7	3	3.0	0	0	1	3
Ophichthidae	<i>Ophichthus cylindroideus</i>	4	1	0	0	0	0	0	0	0	0
	<i>Ophichthus ophis</i>	4.5	1	0	0	0	0	0	0	0	0
Polynemidae	<i>Polydactylus oligodon</i> *	3.7	1	1	100	3	3.0	1	3	0	0

	<i>Polydactylus virginicus</i>	3.7	13	0	0	0	0	0	0	0	0
Pomatomidae	<i>Pomatomus saltatrix</i>	4.5	2	0	0	0	0	0	0	0	0
Rachycentridae	<i>Rachycentron canadum</i>	4	1	0	0	0	0	0	0	0	0
Sciaenidae	<i>Bairdiella ronchus</i>	3.5	4	0	0	0	0	0	0	0	0
	<i>Ctenosciaena gracilicirrhus</i>	3.9	11	0	0	0	0	0	0	0	0
	<i>Cynoscion jamaicensis</i>	3.8	3	0	0	0	0	0	0	0	0
	<i>Cynoscion leiarchus</i> *	3.1	2	1	50	2	2.0	1	2	0	0
	<i>Cynoscion microlepidotus</i> *	4	16	3	18.7	4	1.3	3	3	1	1
	<i>Cynoscion virescens</i> *	4	7	1	14.3	3	3.0	1	3	0	0
	<i>Macrodon ancylodon</i> *	3.9	13	1	7.7	2	2.0	1	2	0	0
	<i>Menticirrhus americanus</i>	3.5	1	0	0	0	0	0	0	0	0
	<i>Micropogonias furnieri</i>	3.1	6	0	0	0	0	0	0	0	0
	<i>Paralonchurus brasiliensis</i>	3.4	6	0	0	0	0	0	0	0	0
Scombridae	<i>Scomberomorus brasiliensis</i>	3.3	1	0	0	0	0	0	0	0	0
Serranidae	<i>Epinephelus itajara</i>	4.1	2	0	0	0	0	0	0	0	0
Sphyrnidae	<i>Sphyrna tiburo</i> *	3.9	2	2	100	18	9.0	1	8	2	10
Stromateidae	<i>Peprilus paru</i>	4.5	2	0	0	0	0	0	0	0	0
Tetraodontidae	<i>Colomesus psittacus</i>	3.6	2	0	0	0	0	0	0	0	0
Triakidae	<i>Mustelus canis</i>	3.6	2	0	0	0	0	0	0	0	0
	<i>Mustelus higmani</i>	3.6	3	0	0	0	0	0	0	0	0
Trichiuridae	<i>Trichiurus lepturus</i> *	4.4	5	1	20	2	2.0	1	2	0	0
TOTAL			189	26	-	228	1.75	22	210	9	18


* Fishes that ingested microplastic, ** Sample based only on specimens that ingested microplastic. Species in bold font support important commercial fisheries.

1

2 In total, 228 microplastic particles were recovered from gastrointestinal tracts of 26
3 specimens belonging to 14 species (Table 1), which represented 13.7% of the total abundance
4 and 30.4% of the total species richness in samples. From the total number of microplastic
5 particles, 210 (92.1%) were found in the stomachs of 21 specimens representing 11 species, and
6 18 particles (7.9%) were recovered from the intestines of 13 specimens representing six species
7 (Table 1). On average, 1.2 (\pm 5.0) particles per fish were found. The standard length of fishes
8 with ingested microplastics varied from 16.0 to 57.5 cm (mean SL = 32.7 cm \pm 11.7), and weight
9 varied between 72.7 and 10,000 g (mean = 1,531.9 g (\pm 3,132.7)). Size, weight and number of
10 microplastic particles found in gastrointestinal tract for each fish are provided as supplementary
11 material (Appendix 1).

12 The greatest number of ingested microplastics per specimen (50 particles) was recorded
13 for *C. hippos*. This species was responsible for 40.3% of all recovered microplastic particles. The
14 frequency of occurrence (FO%) of microplastics per species, among those with three or more
15 specimens examined, varied between 18.7% (*Cynoscion microlepidotus*) and 100% (*Bagre*
16 *marinus*, *Caranx hippos*).

17 A positive correlation was found between fish SL and number of microplastic particles in
18 the gastrointestinal tract (Kendall's Tau= 0.41, $p=$ 0.003). Linear regression analysis indicated
19 that each additional centimeter of SL was associated with an additional 0.58 ingested particles
20 ($r^2=$ 0.37; $p=$ 0.0008; $y=$ -10.374+0.586x) (Fig. 2). In contrast, no significant correlation ($p=$
21 0.43) was found between the quantity of ingested particles and trophic level, between the size
22 microplastic particles and fish SL ($p=$ 0.48), or between the quantity of ingested particles and
23 fishes weight ($p=$ 0.15).




1
2

3 **Fig. 2.** Scatter plot showing the relationship between the number of ingested microplastic
4 particles and SL among fishes that had microplastic particles in their gastrointestinal tracts
5 (n=26). Shaded area indicates 95% confidence interval for the linear regression.

6
7 The size of plastic particles recovered from fish gastrointestinal tracts was always < 5 mm
8 (range= 0.38–4.16 mm, average (\pm SD) = 1.82 (\pm 0.68)). Consequently, all particles are classified
9 as microplastics (Arthur et al., 2009). Particles were recorded as four shape categories: pellets
10 (97.4%), sheets (1.3%), fragments (0.4%) and threads (0.9%). Colours were either clear, yellow,
11 orange or blue (Fig. 3).

12



1
2
3 **Fig. 3.** ATR-FTIR spectra of the microplastic particles: (a) blue thread (polyethylene); (b)
4 yellow fragment (polyethylene); (c) transparent sheet (rayon); (d) pellet I (polyamide); (e) pellet
5 II (polyamide). Highlighted in red the FTIR inset showing the oxidation bands of polyethylene in
6 spectra a and b.

7
8 Microplastic polymers were identified using ATR-FTIR and comparison with reference
9 spectra (Fig. 3). Samples classified as blue thread and yellow fragments had absorption spectra
10 consistent with polyethylene (Fig. 03a,b): 2920 (ν_{as} CH₂), 2850 (ν_s CH₂), 1471 (bending
11 deformation), 1373 (δ_s CH₃) and 720 cm⁻¹ (rocking deformation) (Gulmine et al., 2002). Among
12 blue and yellow fragments, some spectral bands also could be consistent with presence of
13 adsorbed algae: ca. 3350 (N-H, O-H stretching), 1650 (amide I) and 1540 cm⁻¹ (amide II) (Kong
14 and Yu, 2007). Among blue fragments, some weak bands and shoulders suggest the presence of
15 oxygenated groups that form during the abiotic oxidation of polyethylene (Gewert et al., 2015):
16 1715 (C=O stretching ketones, carboxylic acids) and 1738 cm⁻¹ (C=O stretching esters) (Gardette
17 et al., 2013). Transparent sheets (Fig. 03c) showed spectral bands characteristic of Rayon at 3330
18 (O-H stretching), 2901 (CH, CH₂ stretching), 1648 (HOH bending of water), 1445 (H-C-H and H-

1 O-C bending), 1373 and 1320 (H-C-C, H-C-O, and H-O-C bend), 1110 and 1045 (C-C and C-O-
2 C stretching), and 901cm^{-1} (C-O-C in plane, symmetric) (Kaur et al., 2013; Li-Ling, 2007).
3 Finally, samples classified as pellet I and pellet II (Fig. 4e,d) had spectra indicative of nylon at
4 3285 (N-H stretching), 3060 (C-H stretching, asymm., N-H band), 2953 (CH_2 asymm.
5 stretching), 1640 (amide I), 1529 (amide II), 1447 and 1400 (CH_2 bending), and 1242 cm^{-1}
6 (amide III) (Mahdi, 2011; Charles et al., 2009; Fayemi et al., 2016). For the pellet I sample, the
7 broad band around 1045 cm^{-1} (C-O-C stretching) suggests the presence of adsorbed cellulosic
8 fibers.

9

10 Discussion

11 This study provides the first evidence of ingestion of microplastic particles by fishes from
12 the Amazon River estuary. In recent studies conducted in the northern Atlantic Ocean, 19.80%
13 (Neves et al., 2015), 17.50% (Bellas et al., 2016) and 2.98% (Brate et al., 2016) of fishes had
14 microplastics in their gastrointestinal tracts. Microplastics were reported from guts of 50% of
15 marine fishes near the city of Salvador (Miranda et al., 2016) and in 23% of fishes from the
16 Goiana estuary (Possatto et al., 2011), both in Northeast Brazil. Another study of fishes from the
17 Goiana estuary (Ramos et al., 2012) found levels of microplastic ingestion (13.4%) similar to the
18 level we found for fishes the Amazon estuary (13.7%).

19 The greatest number of ingested microplastics (50 particles) was encountered in a
20 Crevalle jack (*Caranx hippos*) a predatory fish that can attain 124 cm SL and 32 kg (Meyer et al.,
21 2001; Froese and Pauly, 2017). Considered generalists, Crevalle jack feed on abundant prey, such
22 as schooling fish and crustaceans (Kwei, 1978; Sancho, 2000). A previous study had confirmed
23 the relationship between the ingestion of microplastics and the feeding strategies of demersal
24 fishes in the Eastern Mediterranean (Anastasopoulou et al., 2013). Romeo et al. (2015) proposed
25 that microplastics are ingested most frequently by generalist foragers that target abundant small
26 prey - a characteristic consistent with Crevalle jack feeding habits. The authors further proposed
27 that microplastics also are ingested secondarily when their prey already contain microplastics.
28 Given that microplastics tend to accumulate in the benthos (Bellas et al., 2016), biomagnification
29 may be accelerated in predators that target bottom-feeding fishes and crustaceans. Ingestion of
30 microplastics by the Crevalle jack is cause for concern, given the species' ecological importance
31 in many tropical and subtropical marine ecosystems. Additionally, fishes of the Carangidae
32 family are among the most important in tropical fisheries (Reuben et al., 1992, Crabtree et al.,

1 2002). Our finding of a significant correlation between fish length and the quantity of ingested
2 microplastic particles corroborates those of Alomar et al. (2017) for goatfish, *Mullus surmuletus*,
3 in the Western Mediterranean. Our failure to find a relationship between the number of ingested
4 microplastics and trophic level may have been influenced by the limited range of trophic levels
5 (3.1 to 4.5) among the species surveyed.

6 Several investigations of marine vertebrates and invertebrates have found that microfibers
7 are the most commonly ingested microplastic particles (Cole et al., 2013; Bellas et al., 2016;
8 Mizraji et al., 2017). In our study, plastic pellets comprised more than 97% of the microplastics
9 recovered from fish gastrointestinal tracts. This corroborates the findings of Miranda et al. (2016)
10 that fishes from the Brazil's northeastern coast near the city of Salvador had exclusively ingested
11 pellets. This region is home to the largest petrochemical complex in Latin America, which could
12 contribute to plastic pollution of rivers and coastal habitats (Ogata et al., 2009).

13 The color of the recovered pellets (predominately shades of yellow) was similar to that
14 observed by Miranda et al. (2016) and indicative of persistent organic pollutants (POPs) attached
15 to the surface of plastics (Endo et al., 2005). Furthermore, the yellowish color suggests that the
16 pellets experienced oxidation and photo-oxidation while adrift in the sea (Ogata et al., 2009;
17 Gewert et al., 2015), implying the original introduction into the environment might have occurred
18 far from our study area. Yellow colour also can be caused by interaction with digestive enzymes
19 (Miranda et al., 2016).

20 As estimated by ATR-FTIR analysis, polyamide (Nylon) represented 97.4% of all
21 microplastics ingested by fishes in our sample from the Amazon estuary. We conclude that the
22 high density (1.13–1.15 g/m³) of polyamide (GESAMP, 2015), paired with the round shape of
23 pellets, contributed to their accessibility to fishes in the study area. Most fishing gear is
24 manufactured from nylon (Timmers et al., 2005), and the fishing industry has been estimated to
25 contribute approximately 18% of all plastic debris found in the oceans (Andrady, 2011).

26 Apart from the nylon pellets, blue thread, yellow fragments and transparent sheets each
27 contributed < 2% to the total of plastic particles recovered from fish gastrointestinal tracts. Blue
28 thread and yellow fragments were identified as polyethylene (PE), a commonly produced
29 polymer (Jambeck et al. 2015) used to make plastic bags and storage containers (GESAMP,
30 2015). Globally, PE is the most abundant polymer found in the environment (Andrady and Neal,
31 2009; Hidalgo-Ruz et al., 2012). Relative to nylon pellets, PE microplastics were uncommon in
32 fish gastrointestinal tracts, and this could be related to their relatively low density (0.91–0.96

1 g/cm³) (Coutinho et al., 2003; GESAMP, 2015). Floating PE particles, therefore, would be less
2 available to demersal fishes. Transparent sheets were made of rayon, a man-made, semi-synthetic
3 polymer that is used as an artificial textile material (Kauffman, 1993; Woodall et al., 2014).
4 Possible sources are clothing, furniture, and personal hygiene products in sewage effluents
5 (Lusher et al., 2012). In the Amazon estuary, this rayon contributed insignificantly, probably due
6 to this material's rapid rate of degradation (Park, et al., 2004).

7 Some marine systems, such as enclosed bodies (e.g., Mediterranean Sea) and zones of
8 gyre convergence, appear to accumulate large amounts of microplastics (Barnes et al. 2009; Ryan
9 et al. 2009), and rivers are considered major contributors to pollution in these regions (Jambeck et
10 al., 2015). Schmidt et al. (2017) demonstrated that rivers transport microplastics more efficiently
11 than macroplastics, and that the concentration of microplastic particles is positively correlated to
12 river size. Along with mismanagement of solid waste disposal, wastewater discharge, inland
13 navigation and industrial pollution all contribute to entry of microplastics into fluvial ecosystems
14 (Lechner et al., 2014). The Amazon River discharges the world's greatest volume of freshwater.
15 Lebreton et al. (2017) estimated that 38,900 tons of plastic wastes are transported from the
16 Amazon River into the Atlantic Ocean annually, suggesting that most of the microplastics found
17 in our study could have originated in the river.

18 Microplastics, more than macroplastics, are influenced by advection and circulation
19 patterns, which contributed to the accumulation of particles in deep-sea environments (Woodall
20 et al., 2014; Van Cauwenberghe et al., 2015). Offshore convection, saline subduction, and other
21 oceanographic processes along the coast near the Amazon River Mouth could contribute to
22 accumulate microplastics in marine sediments (Talley, 2002; Stabholz et al., 2013).

23 The United Nations Environment Program (UNEP) listed plastic pollution as a critical
24 problem, comparable to climate change (UNEP, 2014). Research that identifies and quantifies
25 microplastics within the environment and bodies of organisms will help us to understand the
26 magnitude and scope of microplastic pollution, and pave the way to manage this problem
27 (Boerguer et al., 2010; Alomar and Deudero, 2017). Our findings indicate significant ingestion of
28 microplastics by diverse fishes inhabiting the Amazon estuary and half of the species, including
29 many that support important fisheries (Table 1). This raises human health concerns, because
30 ingestion of fish that consume plastics has the potential to increase the body burden of hazardous
31 chemicals that adsorb to plastics in the environment (Rochman et al., 2015) and subsequently

1 bioaccumulate (Oehlmann et al., 2009; Rochman et al., 2013). The degree to which microplastics
2 and associated compounds biomagnify in food chains is poorly understood at present.

3

4 **Acknowledgments**

5 The authors are grateful to A. Klautau and J. Romão Jr (CEPNOR) Center for Research and
6 Management of Fishing Resources of Brazil's North Coast of the Brazilian Ministry of the
7 Environment (MMA) in partnership with the Aquatic Ecology Group of the Federal University of
8 Pará (UFPA), for their logistical support during the offshore sampling campaigns in the "Shrimp
9 NEN" research project (#445766/2015-8). We also like to thank the students of the Aquatic
10 Ecology Group (GEA) at the Federal University of Pará (UFPA) for their assistance during the
11 laboratory processing and analysis. We thank the editor and the two referees for their valuable
12 comments on the manuscript. The first (TP) and second (KS) authors were funded by the
13 Coordenação de Aperfeiçoamento do Pessoal de Nível Superior (CAPES). The third author (KW)
14 received support from the International Sportfish Fund and National Science Foundation grant
15 DEB 1257813. The fourth author (DC) and fifth author (AC) gratefully acknowledge CSGI
16 (Consorzio Interuniversitario per lo Sviluppo dei Sistemi a Grande Interfase), Florence (Italy) for
17 financial support, and the European Union's Horizon 2020 research and innovation program
18 under grant agreement No 646063 for partial funding. The seventh author (TG) received a
19 productivity grant (CNPq # 310299/2016-0) and funding from Fundacao de Apoio á Pesquisa,
20 Extensão e Ensino em Ciências Agrárias (FUNPEA) with the project Estudos Ecológicos e
21 Ecotoxicológicos em Peixes da Região Norte e Nordeste do Brasil".

22

References

- Alomar, C.; Sureda, A.; Capó, X.; Guijarro, B.; Tejada, S.; Deudero, S. Microplastic ingestion by *Mullus surmuletus* Linnaeus, 1758 fish and its potential for causing oxidative stress. 2017. *Environmental Research*, 159, 135–142. <http://dx.doi.org/10.1016/j.envres.2017.07.043>
- Anastasopoulou, A.; Mytilineou, C.; Smith, C. J.; Papadopoulou, K. P. Plastic debris ingested by deep-water fish of the Ionian Sea (Eastern Mediterranean). 2013. *Deep-Sea Research I*, 74, 11–13. <http://dx.doi.org/10.1016/j.dsr.2012.12.008>
- Andrady, A. L.; Neal, M. A. Applications and societal benefits of plastics. 2009. *Philosophical Transactions of The Royal Society B*, 364, 1977–1984. doi:10.1098/rstb.2008.0304
- Andrady, A. L. Microplastics in the marine environment. 2011. *Marine Pollution Bulletin*, 62, 1596–1605. doi:10.1016/j.marpolbul.2011.05.030
- Arthur, C., J. Baker and H. Bamford (eds). 2009. Proceedings of the International Research Workshop on the Occurrence, Effects and Fate of Microplastic Marine Debris. Sept 9-11, 2008. NOAA Technical Memorandum NOS-OR&R-30. <https://repository.library.noaa.gov/view/noaa/2509> Accessed on January 15th 2018
- Barnes, D.K.A.; Galgani, F.; Thompson, R.C.; Barlaz, M. Accumulation and fragmentation of plastic debris in global environments. 2009. *Philosophical Transactions of The Royal Society of London Series B Biological Sciences*, 364, 1985–1998. doi:10.1098/rstb.2008.0205
- Bellas, J.; Martínez-Armental, J.; Martínez-Cámara, A.; Besada, V.; Martínez-Gomes, C. Ingestion of microplastics by demersal fish from the Spanish Atlantic and Mediterranean coasts. 2016. *Marine Pollution Bulletin*, 109, 55–60. <http://dx.doi.org/10.1016/j.marpolbul.2016.06.026>
- Biginagwa, F. J.; Mayoma, B. S.; Shashoua, Y.; Syberg, K.; Khan, F. R. First evidence of microplastics in the African Great Lakes: recovery from Lake Victoria Nile perch and Nile tilapia. 2016. *Journal of Great Lakes Research*, 42, 146-149. <http://dx.doi.org/10.1016/j.jglr.2015.10.012>.
- Boerger, C. M.; Lattin, G. L.; Moore, S. L.; Moore, C. J. Plastic ingestion by planktivorous fish in the North Pacific Central Gyre. 2010. *Marine Pollution Bulletin*, 60, 2275–2278. doi:10.1016/j.marpolbul.2010.08.007
- Brate, I. L. N.; Eidsvoll, D. P.; Steindal, C. C.; Thomas, K. V. Plastic ingestion by Atlantic cod (*Gadus morhua*) from the Norwegian coast. 2016. *Marine Pollution Bulletin*, 112, 105-110. <http://dx.doi.org/10.1016/j.marpolbul.2016.08.034>.
- Browne, M. A.; Galloway, T. S.; Thompson, R. C. Spatial patterns of plastic debris along estuarine shorelines. 2010. *Environmental Science & Technology*, 44, 3404–3409. DOI:10.1021/es903784e

Cannon, S. M. E.; Lavers, J. L.; Figueiredo, B. Plastic ingestion by fish in the Southern Hemisphere: A baseline study and review of methods. 2016. *Marine Pollution Bulletin*, 107, 286-291. <http://dx.doi.org/10.1016/j.marpolbul.2016.03.057>.

Charles J.; Ramkumaar G. R.; Azhagiri S.; Gunsakeran, S. FTIR and Thermal Studies on Nylon-66 and 30% Glass Fibre Reinforced Nylon-66. 2009. *E-Journal of Chemistry*, 6, 23-33. <http://dx.doi.org/10.1155/2009/909017>

Cincinelli, A.; Scopetani, C.; Chelazzi, D.; Lombardini, E., Martellini, T., Katsoyannis, A.; Fossi, M.C.; Corsolini, S. 2017. Microplastic in the surface waters of the Ross Sea (Antarctica): occurrence, distribution and characterization by FTIR. *Chemosphere*, 175, 391-400. DOI: 10.1016/j.chemosphere.2017.02.024

Colabuono, F. E.; Taniguchi, S.; Montone, R. C. Polychlorinated biphenyls and organochlorine pesticides in plastics ingested by seabirds. 2010. *Marine Pollution Bulletin*, 60, 630–634. DOI: 10.1016/j.marpolbul.2010.01.018

Cole, M.; Lindeque, P.; Halsband, C.; Galloway, T. S. Microplastics as contaminants in the marine environment: a review. 2011. *Marine Pollution Bulletin* 62, 2588–2597. doi:10.1016/j.marpolbul.2011.09.025

Cole, M.; Lindeque, P.; Fileman, E.; Halsband, C.; Goodhead, R.; Moger, J.; Galloway, T. S. Microplastic ingestion by zooplankton. 2013. *Environmental Science & Technology*, 47, 6646–6655. DOI: 10.1021/es400663f

Costa, M. F.; Barletta, M. Microplastics in Coastal and Marine Environments of the Western Tropical and Sub-Tropical Atlantic Ocean. 2015. *Environmental Science: Processes & Impacts*, 17, 1868-1879. DOI: 10.1039/C5EM00158G.

Coutinho, F. M. B.; Mello, I. L.; de Santa Maria, L. C. Polietileno: Principais tipos, propriedades e aplicações. 2003. *Polímeros: Ciência e Tecnologia*, 13, 1-13.

Crabtree, R.; Hood, P. B.; Snodgrass D. Age, growth and reproduction of permit (*Trachinotus falcatus*) in Florida waters. 2002. *Fishery Bulletin – National Oceanic and Atmospheric Administration*, 100, 26-34.

Endo, S.; Takizawa, R.; Okuda, K.; Takada, H.; Chiba, K.; Kanehiro, H.; Ogi, H.; Yamashita, R.; Date, T. Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets: variability among individual particles and regional differences. 2005. *Marine Pollution Bulletin*, 50, 1103–1114. doi:10.1016/j.marpolbul.2005.04.030

Fayemi, O. E.; Adekunle, A. S.; Ebenso E. E. A sensor for the Determination of Lindane Using PANI/Zn, Fe(III) Oxides and Nylon 6,6/MWCNT/Zn, Fe(III) Oxides Nanofibers Modified Glassy Carbon Electrode. 2016. *Journal of Nanomaterials*. 2016(1):1-10. <http://dx.doi.org/10.1155/2016/4049730>

Ferreira, G. V. B.; Barletta, M.; Lima, A. R. A.; Dantas, D. V.; Justino, A. K. S.; Costa, M. F. Plastic debris contamination in the life cycle of Acoupa weakfish (*Cynoscion acoupa*) in a tropical estuary. 2016. ICES Journal of Marine Science, 73, 2695-2707. doi:10.1093/icesjms/fsw108.

Fossi, M. C.; Panti, C.; Guerranti, C.; Coppola, D.; Giannetti, M.; Marsili, L.; Minutoli, R. Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). 2012. Marine Pollution Bulletin, 64, 2374–2379. doi: 10.1016/j.marpolbul.2012.08.013.

Froese, R. and Pauly, D. Editors. 2017. FishBase. World Wide Web electronic publication. www.fishbase.org, version (02/2017).

Gall, S.; Thompson, R. The impact of debris on marine life. 2015. Marine Pollution Bulletin, 92, 170–179. <http://dx.doi.org/10.1016/j.marpolbul.2014.12.041>

Gardette, M.; Perthue, A.; Gardette, J.L.; Janecska, T.; Földes, E.; Pukànszky, B.; Therias, S.. Photo- and thermal-oxidation of polyethylene: comparison of mechanisms and influence of unsaturation content. 2013. Polymer Degradation and Stability, 98, 2383-2390. DOI: [10.1016/j.polymdegradstab.2013.07.017](http://dx.doi.org/10.1016/j.polymdegradstab.2013.07.017)

GESAMP. Sources, fate and effects of microplastics in the marine environment: A global assessment. Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection. Retrieved from: http://ec.europa.eu/environment/marine/good-environmental-status/descriptor-10/pdf/GESAMP_microplastics%20full%20study.pdf. Accessed on December 12, 2017.

Gewert, B.; Plassmann, M. M.; MacLeod, M.. Pathways for degradation of plastic polymers floating in the marine environment. 2015. Environmental. Science.: Processes & Impacts, 17,1513–1521. doi: 10.1039/c5em00207a

Gregory, M.R. Environmental implications of plastic debris in marine settings—entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions. 2009. Philosophical Transactions of the Royal Society B: Biological Sciences, 364 , 2013–2025. DOI: 10.1098/rstb.2008.0265

Giarrizzo, T.; U. Krumme. Heterogeneity in intertidal fish fauna assemblages along the world's longest mangrove area in northern Brazil. 2008. Journal of Fish Biology, 72: 773-779. DOI: 10.1111/j.1095-8649.2007.01728.x

Gulmine, J. V.; Janisse k, P. R.; Heise, H. M.; Akcelrud, L. Polyethylene characterization by FTIR. 2002. Polymer Testing, 21, 5, 557-563. doi.org/10.1016/S0142-9418(01)00124-6.

Hidalgo-Ruz, V.; Gutow, L.; Thompson, R. C.; Thiel, M. Microplastics in the marine environment: A review of the methods used for identification and quantification. 2012. Environmental Science & Technology, 46, 3060–3075. doi:10.1021/es2031505.

Holmes, L. A.; Turner, A.; Thompson, R. C. Adsorption of trace metals to plastic resin pellets in the marine environment. 2012. *Environmental Pollution*, 160, 42-48. <https://doi.org/10.1016/j.envpol.2011.08.052>.

Hopewell, J.; Dvorak, R.; Kosior, E. Plastics recycling: challenges and opportunities. 2009. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364, 2115–2126. doi: 10.1098/rstb.2008.0311

Ivar do Sul, J. A.; Costa, M. F. Marine debris review for Latin America and the Wider Caribbean Region: From the 1970s until now, and where do we go from here? 2007. *Marine Pollution Bulletin*, 54, 1087-1104. Doi:10.1016/j.marpolbul.2007.05.004

Ivar do Sul, J. A.; Costa, D. M. F. The present and future of microplastic pollution in the marine environment. 2014. *Environmental Pollution*, 185, 352–364. <https://doi.org/10.1016/j.envpol.2013.10.036>

Jabeen, K.; Lei, S.; Li, J.; Dongqi, Y.; Tong, C.; Jingli, M.; Shi, H. Microplastics and mesoplastics in fish from coastal and fresh waters of China. 2017. *Environmental Pollution*, 221, 141-149. <http://dx.doi.org/10.1016/j.envpol.2016.11.055>.

Jambeck, J. R.; Geyer, R.; Wilcox, C.; Siegler, T. R.; Perryman, M.; Andrady, A.; Narayan, R.; Law, K. L. Plastic waste inputs from land into the ocean. 2015. *Science*, 347, 768-771. DOI: 10.1126/science.1260352

Kauffman, G. B. Rayon. The first Semi-Synthetic Fiber Product. *Products of chemistry* (1993), 70, 11.

Kaur, I., Sharma, N., Kumari, V. Modification of fiber properties through grafting of acrylonitrile to rayon by chemical and radiation methods. 2013. *Journal of Advanced Research*, 4, 547-557. <https://doi.org/10.1016/j.jare.2012.11.003>

Kong, J.; Yu, S. Fourier transform infrared spectroscopic analysis of protein secondary structures. 2007. *Acta biochimica et biophysica Sinica*, 39, 549-559. DOI: 10.1111/j.1745-7270.2007.00320.x

Kottek, M.; Grieser, J.; Beck, C.; Rudolf, B.; Rubel, F. World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, 15, 259-263. DOI: 10.1127/0941-2948/2006/0130

Krumme, U.; Giarrizzo, T.; Pereira, R.; Silva de Jesus, A. J.; Schaub, C.; Saint-Paul, U. Airborne synthetic-aperture radar (SAR) imaging to help assess impacts of stationary fishing gear on the north Brazilian mangrove coast. 2015. *ICES Journal of Marine Science*, 72, 939–951. doi:10.1093/icesjms/fsu188.

Kwei, E. Food and spawning activity of *Caranx hippos* off the coast of Ghana. 1978. *Journal of Natural History*, 12, 195-215. <https://doi.org/10.1080/00222937800770081>

Laist, D.W. Impacts of marine debris entanglement of marine life in marine debris including a comprehensive list of species with entanglement and ingestion records. 1997. In: Coe, J.M., Rogers, D.B. (Eds.), *Marine Debris—Sources, Impact and Solutions*. Springer-Verlag, New York, pp. 99–139. DOI: https://doi.org/10.1007/978-1-4613-8486-1_10

Lana, P. C.; Camargo, M. G.; Brogim, R. A.; Isaac, V. J. Os Bentos da Costa Brasileira: avaliação crítica e levantamento bibliográfico, 1858-1996. 1996. Rio de Janeiro: FEMAR, 431p.

Lebreton, L. C. M.; Greer, S. D.; Borrero, J. C. Numerical modelling of floating debris in the world's oceans. 2012. *Marine Pollution Bulletin*, 64, 653–661. <https://doi.org/10.1016/j.marpolbul.2011.10.027>

Lebreton, L. C. M.; Van der Zwet, J.; Damsteeg, J. W.; Slat, B.; Andrady, A.; Reisser, J. River plastic emissions to the world's oceans. 2017. *Nature Communications*, 8, 15611, doi: 10.1038/ncomms15611

Lechner, A.; Keckeis, H.; Lumesberger-Loisl, F.; Zens, B.; Krusch, R.; Tritthart, M.; Glas, M.; Schludermann, E. The Danube so colourful: a potpourri of plastic litter outnumbers fish larvae in Europe's second largest river. 2014. *Environmental Pollution*, 188, 177–181. <http://dx.doi.org/10.1016/j.envpol.2014.02.006>.

Li-Ling, C., Identification of textile fiber by Raman microspectroscopy. 2007. *Forensic Science Journal*, 6, 55-62. Available online at: fsjournal.cpu.edu.tw

Lusher, A. L.; McHugh, M.; Thompson, R. C. Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel. 2012. *Marine Pollution Bulletin*, 67, 94-99. <http://dx.doi.org/10.1016/j.marpolbul.2012.11.028>.

Mahdi, H.A. An FTIR Study of Characterization of Neat and UV Stabilized Nylon 6,6 Polymer Films. 2011. *Pure & Applied Science*, 24.

Marceniuk, A. P.; Caires, R. A.; Wosiacki, W. B.; Di Dario, F. Knowledge and conservation of the marine and estuarine fishes (Chondrichthyes and Teleostei) of the north coast of Brazil. 2013. *Biota Neotropica*, 13, 251-259. <http://www.biotaneotropica.org.br/v13n4/en/abstract?inventory+bn02613042013>

McGoran, A. R.; Clark, P. F.; Morritt, D. Presence of microplastic in the digestive tracts of European flounder, *Platichthys flesus*, and European smelt, *Osmerus eperlanus*, from the River Thames. 2017. *Environmental Pollution* 220, 744 e 751. <https://doi.org/10.1016/j.envpol.2016.09.078>

Mecozzi, M.; Pietroletti, M.; Monakhova, Y.B.. FTIR spectroscopy supported by statistical techniques for the structural characterization of plastic debris in the marine environment: Application to monitoring studies. 2016. *Marine Pollution Bulletin*, 106, 155-161. DOI: [10.1016/j.marpolbul.2016.03.012](https://doi.org/10.1016/j.marpolbul.2016.03.012).

Meyer, C. G.; Holland, K. N.; Wetherbee, B. M.; Lowe, C. G. Diet, resource partitioning and gear vulnerability of Hawaiian jacks captured in fishing tournaments. 2001. Fisheries Research, 53, 105-113. [https://doi.org/10.1016/S0165-7836\(00\)00285-X](https://doi.org/10.1016/S0165-7836(00)00285-X)

Miranda, D. A.; Carvalho-Souza, G. F. Are we eating plastic-ingesting fish?. 2015. Marine Pollution Bulletin, 103, 109-114. <http://dx.doi.org/10.1016/j.marpolbul.2015.12.035>.

Mizraji, R.; Ahrendt, C.; Pérez-Venegas, D.; Vargas, J.; Pulgar, J.; Aldana, M.; Ojeda, F. P.; Duarte, C.; Galbán-Malagón, C. Is the feeding type related with content of microplastics in intertidal fish gut?. 2017. Marine Pollution Bulletin, 116, 498-500. <http://dx.doi.org/10.1016/j.marpolbul.2017.01.008>

Moore, C. J. Synthetic polymers in the marine environment: a rapidly increasing long-term threat. 2008. Environmental Research. 108, 131-139. <https://doi.org/10.1016/j.envres.2008.07.025>

Moura, R. L.; Amado-Filho, G. M.; Moraes, F. C.; Brasileiro, P. S.; Salomon, P. S.; Mahiques, M. M.; Bastos, A. C.; Almeida, M. G.; Silva-Jr, J. M.; Araújo, B. F.; Brito, F. P.; Oliveira, B. C. V.; Bahia, R. G.; Paranhos, R. P.; Dias, R. J. S.; Siegle, R.; Figueiredo-Jr, A. B. G.; Pereira, R. C.; Leal, C. V.; Hajdu, E.; Asp, N. E.; Gregoracci, G. B.; Neumann-Leitão, S.; Yager, P. L.; Francini-Filho, R. B.; Fróes, A.; Campeão, M.; Silva, B. S.; Moreira, A. P. B.; Oliveira, L.; Soares, A. C.; Araújo, L.; Oliveira, N.L.; Teixeira, J. B.; Valle, R. A. B.; Thompson, C. C.; Rezende, C. E.; Thompson, F. L. An extensive reef system at the Amazon River mouth. 2016. Science Advances, 2, e1501252. DOI: 10.1126/sciadv.1501252

Naidoo, T.; Glassom, D.; Smit, A. J. Plastic ingestion by estuarine mullet, *Mugil cephalus* (L. 1758), in an urban harbour, KwaZulu-Natal, South Africa. 2016. African Journal of Marine Science, 38, 145–149. <https://doi.org/10.2989/1814232X.2016.1159616>

Neiva G.S.; Moura, S.J.C., 1977. Sumário sobre exploração de recursos marinhos do litoral brasileiro: situação atual e perspectivas. Programa de Pesquisa e Desenvolvimento Pesqueiro do Brasil. Série Documentos Opcionais, Brasília, 27, 1-48.

Neves, D.; Sobral, P.; Ferreira, J. L.; Pereira, T. Ingestion of microplastics by commercial fish off the Portuguese coast. 2015. Marine Pollution Bulletin, 101, 119-126. <http://dx.doi.org/10.1016/j.marpolbul.2015.11.008>.

Nittrouer, C. A.; Brunskill, G. J.; Figueiredo, A.G. Importance of tropical coastal environments. 1995. Geo-Marine Letters, 15, 121-126. DOI: 10.1007/BF01204452

Oehlmann, J.; Schulte-Oehlmann, U.; Kloas, W.; Jagnytsch, O.; Lutz, I.; Kusk, K.O.; Wollenberger, L.; Santos, E.M.; Paull, G.C.; Van Look, K.J.W.; Tyler, C.R. A critical analysis of the biological impacts of plasticizers on wildlife. 2009. Philosophical Transactions of the Royal Society B: Biological Sciences, 364, 2047-2062. <https://doi.org/10.1098/rstb.2008.0242>

Ogata, Y.; Takada, H.; Mizukawa, K.; Hiraia, H.; Iwasaa, S.; Endo, S.; Mato, Y.; Saha, M.; Okuda, K.; Nakashima, A.; Murakami, M.; Zurcher, N.; Booyatumanondo, R.; Zakaria, M. P.; Dung, L. Q.; Gordon, M.; Miguez, C.; Suzuki, S.; Moore, C.; Karapanagiotik, H. K.; Weerts, S.; McClurg, T.; Burrell, E.; Smith, W.; Van Velkenburg, M.; Lang, J. S.; Lang, R. C.; Laursen, D.; Danner, B.; Stewardson, N.; Thompson, R. C. International PelletWatch: global monitoring of

persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs. 2009. *Marine Pollution Bulletin*, 58, 1437–1446. doi: 10.1016/j.marpolbul.2009.06.014

Oltman, R.E. Reconnaissance investigation of discharge and water quality of the Amazon River. 1968. US. Geological Survey Circular 552, Washington DC, p.16.

Ory, N. C.; Sobral, P.; Ferreira, J. L.; Thiel, M. Amberstripe scad *Decapterus muroadsi* (Carangidae) fish ingest blue microplastics resembling their copepod prey along the coast of Rapa Nui (Easter Island) in the South Pacific subtropical gyre. 2017. *Science of the Total Environment* 586, 430–437. doi: 10.1016/j.scitotenv.2017.01.175

Park, C. H.; Kang, Y. K.; Im, S. S. Biodegradability of cellulose fabrics. 2004. *Journal of Applied Polymer Science*, 94, 248–253. <http://dx.doi.org/10.1002/app.20879>.

Pereira, L. C. C; Dias, J. A.; Do Carmo, J. A.; Polette, M. A. Zona Costeira Amazônica Brasileira/ The Brazilian Amazon Coastal Zone. 2009. *Revista de Gestão Costeira Integrada / Journal of Integrated Coastal Zone Management*, 9,3-7. DOI: 10.5894/rgci172

Petters, C. A.; Bratton, S. P. Urbanization is a major influence on microplastic ingestion by sunfish in the Brazos River Basin, Central Texas, USA. 2016. *Environmental Pollution*, 210, 380–387. doi: 10.1016/j.envpol.2016.01.018.

Phillips, M. B.; Bonner, T. Occurrence and amount of microplastic ingested by fishes in watersheds of the Gulf of Mexico. 2015. *Marine Pollution Bulletin*, 100, 264–269. <https://doi.org/10.1016/j.marpolbul.2015.08.041>

Possatto, F. E.; Barletta, M. Costa, M. F.; Ivar do Sul, J. A.; Dantas, D. V. Plastic debris ingestion by marine catfish: An unexpected fisheries impact. 2011. *Marine Pollution Bulletin*, 62, 1098–1102. <https://doi.org/10.1016/j.marpolbul.2011.01.036>

R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Ramos, J. A. A.; Barletta, M.; Costa, M. F. Ingestion of nylon threads by Gerreidae while using a tropical estuary as foraging grounds. 2012. *Aquatic Biology*, 17, 29–34. doi: 10.3354/ab00461.

Reuben, S.; Kasim, H. M.; Sivakami, S.; Radhakrishnan, P. N.; Kurup, K. N.; Sivadas, M.; Noble, A.; Nair, K. V. S.; Raje, S. G. Fishery, biology and stock assessment of Carangid resources from the Indian seas. 1992. *Indian Journal of Fisheries*, 39,195-234. <http://eprints.cmfri.org.in/id/eprint/526>

Rochman, C.M.; Hoh, E.; Kurobe, T.; Teh, S.J. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. 2013. *Scientific Reports*, 3, 3263. <https://doi.org/10.1038/srep03263>

Rochman, C. M.; Tahir, A.; Williams, S. L.; Baxa, D. V.; Lam, R.; Miller, J. T.; Teh, F. C.; Werorilangi, S.; Teh, S. L. Anthropogenic debris in seafood: Plastic debris and fibers from textiles

in fish and bivalves sold for human consumption. 2015. *Scientific Reports*, 5, 14340. DOI: 10.1038/srep14340.

Romeo, T.; Battaglia, P.; Peda, C.; Consoli, P.; Andaloro, F.; Fossi, M. C. First evidence of presence of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. 2015. *Marine Pollution Bulletin*, 95, 358-361. <http://dx.doi.org/10.1016/j.marpolbul.2015.04.048>.

Ryan, P. G.; Moore, C. J.; Van Franeker, J. A.; Moloney, C. L. Monitoring the abundance of plastic debris in the marine environment. 2009. *Philosophical Transactions of the Royal Society B* 364, 1999–2012. doi:10.1098/rstb.2008.0207

Sancho, G. Predatory behaviors of *Caranx melampygus* (Carangidae) feeding on spawning reef fishes: a novel ambushing strategy. 2000. *Bulletin of Marine Science* 66, 487–496. <https://www.researchgate.net/publication/233698439>

Seltenrich, N. New link in the food chain? Marine plastic pollution and seafood safety. 2015. *Environmental Health Perspectives*, 123, A34. . doi: 10.1289/ehp.123-A34.

Stabholz, M.; Durrieu de Madron, X.; Canals, M.; Khripounoff, A.; Taupier-Letage, I.; Testor, P.; Heussner, S.; Kerhervé, P.; Delsaut, N.; Houpert, L.; Lastras, G.; Dennielou, B. Impact of open-ocean convection on particle fluxes and sediment dynamics in the deep margin of the Gulf of Lions. 2013. *Biogeosciences*, 10, 1097–1116. doi:10.5194/bg-10-1097-2013

Silva-Cavalcanti, J. S.; Silva, J. D. B.; França, E. J.; Araújo, M. C. B.; Gusmão, F. Microplastics ingestion by a common tropical freshwater fishing resource. 2017. *Environmental Pollution* 221, 218-226. <https://doi.org/10.1016/j.envpol.2016.11.068>

Schmidt, C.; Krauth, T.; Wagner, S. Export of plastic debris by rivers into the sea. 2017. *Environmental Science & Technology*, 51, 12246-12253. DOI: 10.1021/acs.est.7b02368

Srinivasa Reddy, M.; Basha, S.; Adimurthy, S.; Ramachandraiah, G. 2006. Description of the small plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India. *Estuarine, Coastal and Shelf Science*, 68, 656-660. DOI: [10.1016/j.ecss.2006.03.018](https://doi.org/10.1016/j.ecss.2006.03.018)

Talley, J. W.; Ghosh, U.; Tucker, S. G.; Furey, J. S.; Luthy, R. G. Particle-scale understanding of the bioavailability of PAHs in sediments. 2002. *Environmental Science & Technology*, 36, 477-483. DOI: 10.1021/es010897f

Teuten, E. L.; Saquing, J. M.; Knappe, D. R. U.; Barlaz, M. A.; Jonsson, S.; Björn, A.; Rowland, S. J.; Thompson, R. C.; Galloway, T. S.; Yamashita, R.; Ochi, D.; Watanuki, Y.; Moore, C.; Viet, P. H.; Tana, T. S.; Prudente, M.; Boonyatyanond, R.; Zakaria, M. P.; Akkhavong, K.; Ogata, Y.; Hirai, H.; Iwasa, S.; Mizukawa, K.; Hagino, Y.; Imamura, A.; Saha, M.; Takada, H. Transport and release of chemicals from plastics to the environment and to wildlife. 2009. *Philosophical Transactions of the Royal Society B*, 364, 2027–2045 doi:10.1098/rstb.2008.0284

Timmers, M. A.; Kistner, C. A.; Donohue, M. J. Marine Debris of the Northwestern Hawaiian Islands: Ghost Net Identification. 2005. U.S. Dept of Commerce, National Oceanic and

Atmospheric Administration, National Sea Grant College Program, 31 pages. ark:/86086/n20864gx_cuwr_conservancy_001_3714.

UNEP. 2014. UNEP Year Book 2014: Emerging issues update. United Nations Environment Programme, Nairobi, Kenya.

Van Cauwenberghe, L.; Claessens, M.; Vandegheuchte, M. B.; Janssen, C. R. Microplastics are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats. 2015. Environmental Pollution, 199, 10-17. <http://dx.doi.org/10.1016/j.envpol.2015.01.008>

Vendel, A. L.; Bessa, F.; Alves, V. E. N.; Amorim, A. L. A.; Patrício, J.; Palma, A. R. T. Widespread microplastic ingestion by fish assemblages in tropical estuaries subjected to anthropogenic pressures. 2017. Marine Pollution Bulletin, 117, 448-455. <http://dx.doi.org/10.1016/j.marpolbul.2017.01.081>.

Woodall, L. C.; Sanchez-Vidal, A.; Canals, M.; Paterson, G. L. J.; Coppock, R.; Sleight, V.; Calafat, A.; Rogers, A. D.; Narayanaswamy, B. E.; Thompson, R. C. The deep sea is a major sink for microplastic debris. 2014. The Royal Society Open Science, 1, 140317. DOI: 10.1098/rsos.140317

Wolff, M; Koch, V.; Isaac, V. A trophic flow model of the Caeté mangrove estuary (North Brazil) with considerations for the sustainable management of its resources. 2000. Estuarine, Coastal and Shelf Science, 50, 789-803. <https://doi.org/10.1006/ecss.2000.0611>