



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Antinociceptive profile of SM21: a novel analgesic with a presynaptic cholinergic mechanism of action

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Antinociceptive profile of SM21: a novel analgesic with a presynaptic cholinergic mechanism of action / C. GHELARDINI; N. GALEOTTI; F. GUALTIERI; C. BELLUCCI; D. MANETTI; A. GIOTTI; P. MALMBERG; A. GALLI; A. BARTOLINI. - In: THE JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS. - ISSN 0022-3565. - STAMPA. - 282:(1997), pp. 430-439.

Availability:

This version is available at: 2158/310583 since:

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Antinociceptive Profile of 3- α -tropanyl-(2-Cl)-acid phenoxybutyrate (SM-21): A Novel Analgesic with a Presynaptic Cholinergic Mechanism of Action¹

CARLA GHELARDINI, NICOLETTA GALEOTTI, FULVIO GUALTIERI, CRISTINA BELLUCCI, DINA MANETTI, ALBERTO GIOTTI, PETRA MALMBERG-AIELLO, ALESSANDRO GALLI and ALESSANDRO BARTOLINI

Department of Pharmacology (C.G., N.G., A.G., P.M.A., A.G., A.B.), University of Florence and Department of Pharmaceutical Sciences (F.G., C.B., D.M.), University of Florence, Florence, Italy

Accepted for publication March 18, 1997

ABSTRACT

The antinociceptive effect of (\pm)-3- α -tropanyl-(2-Cl)-acid phenoxybutyrate (SM-21) (10–40 mg kg⁻¹ s.c., 10–30 mg kg⁻¹ i.p., 20–60 mg kg⁻¹ p.o., 3–20 mg kg⁻¹ i.v. and 5–20 μ g per mouse i.c.v.) was examined in rodents and guinea pigs by using the hot-plate, abdominal constriction, tail-flick and paw-pressure tests. The antinociception produced by (\pm)-SM-21 was prevented by atropine, pirenzepine and hemicholinium-3 but not by quinpirole, R-(α)-methylhistamine, [1-[2(methylsulfonyl)amino]ethyl]-4-piperidiny]methyl-5-floro-2-methoxy-1H-indole-3-carboxylate hydrochloride, N⁶-cyclopentyladenosine, 1-(2-methoxyphenyl)-4-[4-(2-phthalimido)butyl]piperazine hydrobromide, naloxone, 3-aminopropyl-diethoxy-methyl-phos-

phinic acid or reserpine. On the basis of the above data, it can be postulated that (\pm)-SM-21 exerted an antinociceptive effect mediated by a central potentiation of cholinergic transmission. Affinity profiles of (\pm)-SM-21 for muscarinic receptor subtypes, determined by functional studies (rabbit vas deferens for M₁, guinea pig atrium for M₂, guinea pig ileum for M₃ and immature guinea pig uterus for putative M₄) have shown a selectivity ratio M₂/M₁ of 4.6 that, although very low, might be responsible for the antinociception induced by (\pm)-SM-21 through an increase in ACh extracellular levels. In the antinociceptive dose range, (\pm)-SM-21 did not impair mouse performance evaluated by the rota-rod and hole-board tests.

Ghelardini *et al.* (1990) have reported that the antimuscarinic compound atropine, at very low doses, was able to induce a central cholinergic antinociception in rodents regardless of the route of administration and the noxious stimulus applied. Furthermore, this antinociceptive activity was not accompanied by the typical cholinergic symptomatology. The atropine-induced increase in the pain threshold was attributable to the R-(+)-enantiomer of atropine, R-(+)-hyoscyamine, because S-(–)-hyoscyamine was ineffective in all antinociceptive tests used (Ghelardini *et al.*, 1992). More recently, Bartolini *et al.* (1994), investigating the antinociceptive effect of atropine, demonstrated, using microdialysis

techniques, that R-(+)-hyoscyamine, at analgesic doses, produced an increase in the release of ACh from the rat cerebral cortex *in vivo*. On the bases of the above-mentioned results, the racemate (Gualtieri *et al.*, 1994) and the enantiomers (Romanelli *et al.*, 1995) of the compound labeled SM-21 (fig. 1), which is structurally related to atropine, have been synthesized in order to obtain a new cholinergic amplifier endowed with more intensive antinociceptive activity than atropine but as lacking as atropine in cholinergic side effects. To this end, we investigated (\pm)-SM-21 antinociceptive properties by using the hot-plate, abdominal constriction, paw-pressure and tail-flick tests, and the incidence of behavioral side effects was detected by the rota-rod and hole-board tests.

Received for publication July 16, 1996.

¹This research was supported by grants from Fidia S.p.A. (Abano Terme, Italy) and from Ministero dell'Università e della Ricerca Scientifica e Tecnologia (MURST). Preliminary data were presented at the XIII International Symposium on Medicinal Chemistry, Paris, September 19–23, 1994; at the XXVII Meeting of Italian Pharmacological Society, Turin, September 25–29, 1994 and at the VI International Symposium on Subtypes of Muscarinic Receptors, Fort Lauderdale, Florida, November 9–12, 1994.

Materials and Methods

Animals. Male Swiss albino mice (23–30 g), Wistar rats (200–300 g) from Morini (San Polo d'Enza, Italy), Fisher 344 rats (200–300 g) from Charles River (Calco, Italy) and guinea pigs (150–200 g) from Rodentia (Bergamo, Italy) breeding farms were used. Fifteen mice

ABBREVIATIONS: SM-21, 3- α -tropanyl-(2-Cl)-acid phenoxybutyrate; HC-3, hemicholinium-3; NAN 190, 1-(2-methoxyphenyl)-4-[4-(2-phthalimido)butyl]piperazine hydrobromide; McN-A-343, 4-(N-[3-chlorophenyl]-carbamoyloxy)-2-butynyl-trimethylammonium chloride; GR 125487, [1-[2(methylsulfonyl)amino]ethyl]-4-piperidiny] methyl-5-fluoro-2-methoxy-1H-indole-3-carboxylate hydrochloride; AFDX-116, 11,2-(diethylamino)methyl-1-piperidiniol acetyl-5,11-dihydro-6H-pyrido 2,3-b 1,4 benzodiazepine-6-one; CGP 35348, 3-aminopropyl-diethoxy-methyl-phosphinic acid; RAMH, (R)- α -methylhistamine.




Fig. 1. Chemical structure of SM-21 racemate and its enantiomers.

and four rats or guinea pigs were housed per cage. The cages were placed in the experimental room 24 h before the test for acclimatization. The animals were kept at $23 \pm 1^\circ\text{C}$ with a 12-h light/dark cycle, light on at 7 A.M., with food and water *ad libitum*. All experiments were carried out according to the guidelines of the European Community Council.

Analgesic Tests

Hot-plate test. The method adopted was described by O'Callaghan and Holtzman (1975). Mice were placed inside a stainless steel container thermostatically set at $52.5 \pm 0.1^\circ\text{C}$ in a precision water bath from KW Mechanical Workshop, Siena, Italy. Reaction times (s), were measured with a stopwatch before treatment and at regular intervals up to a maximum of 60 min after treatment. The endpoint used was the licking of the fore or hind paws. Those mice scoring below 12 and over 18 s in the pretest were rejected (30%). An arbitrary cut-off time of 45 s was adopted.

Abdominal constriction test. Mice were injected i.p. with a 0.6% solution of acetic acid (10 ml kg^{-1}), according to Koster *et al.* (1959). The number of stretching movements was counted for 10 min, starting 5 min after acetic acid injection.

Paw-pressure test. The nociceptive threshold in the rat and guinea-pig was determined with an analgesimeter (Ugo Basile, Varese, Italy), according to the method described by Leighton *et al.* (1988). Threshold pressure was measured before treatment and 15, 30 and 45 min after treatment. Rats and guinea-pigs scoring below 30 g or over 85 g during the test before drug administration were rejected (25%). An arbitrary cutoff value of 250 g was adopted.

Tail-flick test. An analgesimeter from Ugo Basile (Varese, Italy) was used to perform the tail-flick test described by D'Amour and Smith (1941). The light from a project bulb situated beneath the platform where the animal was placed was focused through a small hole on the ventral part of the tail at a point about 4 cm from the tip. Withdrawal of the tail exposed a photocell to the light, which turned off the thermal stimulus and automatically stopped the clock. The intensity was regulated so that the reaction time varied between 2 and 4 s. The analgesia was tested before treatment and 15, 30 and 45 min after treatment. Each value was derived from the mean of three consecutive readings in which the light was focused on three adjacent points of the tail.

Anti-inflammatory Test

Carrageenan-induced paw edema. Rats paw volumes were measured using a plethysmometer (Ugo Basile, Varese, Italy). Rats received (\pm)-SM 21, indomethacin or saline 90 min after a 0.1-ml injection of 0.5% carrageenan in the right hind paw. Two hours after the injection of carrageenan, the paw volume of the right hind paw was measured and compared with saline-treated controls.

Additional Behavioural Tests

Hole-board test. The hole-board test consists of a 40-cm-square plane with 16 flush-mounted cylindrical holes (diameter 3 cm) distributed 4 by 4 in an equidistant, grid-like manner. Mice were placed on the center of the board one by one and left to move about freely for a period of 10 min each. Two electric eyes, crossing the plane from midpoint to midpoint of opposite sides, thus dividing the plane into 4 equal quadrants, automatically signaled the movement of the animals on the surface of the plane. Miniature photoelectric cells in each of the 16 holes recorded the exploration of the holes (head plunging activity) by the mice.

Rota-rod test. The apparatus consisted of a base platform and a rotating rod 3 cm in diameter with a nonslippery surface. The rod was placed at a height of 15 cm from the base. The rod, 30 cm in length, was divided into 5 equal sections by 6 disks. Thus up to 5 mice were tested simultaneously on the apparatus, with a rod-rotating speed of 16 rpm. The integrity of motor coordination was assessed on the basis of the number of falls from the rod in 30 s according to Vaught *et al.* (1985). The performance time was measured before treatment and 15, 30 and 45 min after treatment.

In Vitro Functional Studies

Isolated rabbit vas deferens. Experiments on isolated rabbit vas deferens were performed according to the method described by Eltze (1988) and modified by Dei *et al.* (1995). The preparations were maintained at 32°C , and tissues were stimulated through platinum electrodes by square-wave pulses (2 ms, 0.1 Hz, 10–30 V). Contractions were measured isometrically after tissues had been equilibrated for 1 h, and then a cumulative dose-response curve for the inhibitory effect of McN-A-343 was plotted.

Isolated guinea pig left atria. Isolated left atria were prepared according to the method described by Eltze *et al.* (1985) and modified by Dei *et al.* (1995). Bath fluid temperature was maintained at 30°C . Atria were electrically stimulated (1 Hz, 1 ms, 4–10 V) by means of two platinum electrodes. Carbachol negative inotropic effects on isometric atria contractions were recorded before and 1 h after perfusion with antagonists.

Guinea pig isolated ileum. Isolated ileum fragments were prepared according to Eltze and Figala (1988). Bath fluid temperature was maintained at 37°C . Isotonic ileum contractions induced by ACh were recorded before and 1 h after perfusion with antagonists.

Guinea pig isolated uterus. Experiments on isolated immature guinea pig uterus were performed according to Dörje *et al.* (1990). The preparations were maintained at 30°C , and after a 1-h equilibration period, isotonic contractions to carbachol were recorded. Initially the tissues were exposed to a single-concentration of carbachol (3 nmol l^{-1}) to check the responsiveness to the agonist, and a dose-response curve for carbachol was obtained.

Determination of antagonist affinities. After a stabilization time of 30 to 60 min, agonist concentration-response curves were plotted before and after equilibration with antagonists. In separate control experiments, no significant changes in tissue sensitivity to the agonist were observed over the period required for the determination of two concentration-response curves. The antagonists were allowed to equilibrate for 60 min. No more than two concentrations of antagonist were tested in the same preparation. Agonist EC_{50} values in the absence and presence of antagonists were determined graphically for the calculation of dose ratios.




Fig. 2. Dose-response curves of (\pm)-SM-21 administered s.c. (panel A), i.c.v. (panel B), p.o. (panel C) and i.v. (panel D) in the mouse hot-plate test. The doses are expressed as mg kg^{-1} s.c., p.o. and i.v. and as μg per mouse i.c.v. Vertical lines show S.E.M. $\wedge P < .05$; $* P < .01$ in comparison with saline controls. Each point represents the mean of at least 10 mice.

AChE activity. AChE activity was assayed according to Ellman *et al.* (1961), using 0.5 mM acetylthiocholine iodide as substrate. The (\pm)-SM-21 inhibitory effect was tested at various concentrations on a purified preparation of AChE from the electric eel.

Drugs. SM-21 racemate was prepared according to Gualtieri *et al.* (1994); R-(+)-SM-21 and S-(-)-SM-21 were prepared according to Romanelli *et al.* (1995); R-(+)-hyoscyamine was prepared according to Gualtieri *et al.* (1991). Also used were atropine sulfate, carbamylcholine chloride, carrageenan, physostigmine hemisulfate and yohimbine hydrochloride (Sigma, Milan, Italy), HC-3, pirenzepine dihydrochloride, naloxone hydrochloride, quinpirole hydrochloride, (R)- α -methylhistamine dihydrochloride, indomethacin, N^6 -cyclopentyladenosine, NAN 190, McN-A-343 (R.B.I., Milan, Italy); ACh chloride (Merck, Florence, Italy); GR 125487 (Boehringer Ingelheim, Milan, Italy); morphine hydrochloride (U.S.L. 10/D, Florence, Italy), diphenhydramine hydrochloride and AFDX-116 (De Angeli, Milan, Italy); clomipramine hydrochloride (anafranil), CGP 35348 and reserpine (Ciba Geigy, Basel, Switzerland); (+)-amphetamine sulfate (Recordati, Rome, Italy). Other chemicals were of the highest quality commercially available. All drugs were dissolved in isotonic (NaCl 0.9%) saline solution or dispersed in sodium carboxymethylcellulose 1% immediately before use, except reserpine, which was dissolved in a 20% solution of ascorbic acid, and R-(+)-hyoscyamine, which was dissolved in 0.1 M HCl and then diluted with saline (1:10). Drug

concentrations were prepared in such a way that the necessary dose could be administered in a volume of 10 ml kg^{-1} by s.c., i.p., and p.o. route or 5 ml kg^{-1} by i.v. route. Intracerebroventricular administration was performed under ether anaesthesia using isotonic saline as solvent, according to the method described by Haley and McCormick (1957) for mice and that we adapted for rats. Briefly, during anaesthesia, mice and rats were grasped firmly by the loose skin behind the head. A hypodermic needle 0.4 mm in external diameter, attached to a $10\text{-}\mu\text{l}$ syringe, was inserted perpendicularly through the skull at a depth of no more than 2 mm into the brain of the mouse and 4 mm into the brain of the rat, where $5 \mu\text{l}$ (mice) or $10 \mu\text{l}$ (rats) were then administered. The injection site was 1.5 mm (mice) or 2.5 mm (rats) from either side of the midline on a line drawn through to the anterior base of the ears. To ensure that the drugs were administered exactly into the cerebral ventricle, some mice and rats were injected i.c.v. with 5 to $10 \mu\text{l}$ of diluted 1:10 Indian ink and their brains examined macroscopically after sectioning. Intraplantar injections of carrageenan were performed by injecting $100 \mu\text{l}$ of a suspension in sterile saline solution of 0.5% carrageenan in the rat hind paw.

Statistical analysis. Results are given as the mean \pm S.E.M.; analysis of variance (ANOVA), followed by Fisher's PLSD procedure for *post-hoc* comparison, was used to verify the significance of the difference between two means. P values of less than .05 were con-

sidered significant. Data were analyzed with the StatView for the Macintosh computer program (1992).

Results

Antinociceptive activity of SM-21. (\pm)-SM-21, as shown in figure 2, produced a dose-dependent increase in the pain threshold in the mouse hot-plate test after s.c. (10–40 mg kg⁻¹; panel A), i.c.v. (5–20 μ g per mouse; panel B), p.o. (20–60 mg kg⁻¹; panel C) and i.v. (3–20 mg kg⁻¹; panel D) administration. The antinociceptive effect of (\pm)-SM-21 peaked 15 min after s.c. and i.c.v. administration and then slowly diminished. (\pm)-SM-21, after p.o. and i.v. administration, reached its maximum analgesic effect respectively 30 and 10 min after injection. Figure 3 (panels A and B) illustrates the analgesic effect of (\pm)-SM-21 in the mouse acetic acid abdominal constriction test. (\pm)-SM-21 induced an increase in the pain threshold in a dose-dependent manner starting from the dose of 10 mg kg⁻¹ s.c. (fig. 3, panel A). (\pm)-SM-21 showed antinociceptive properties also after the injection of 1 and 5 μ g per mouse i.c.v., reaching its maxi-

mum effect between 15 and 30 min after administration (fig. 3, panel B).

(\pm)-SM-21 was able to produce an increase in the pain threshold not only in mice but also in rats and guinea pigs. In the paw pressure test (\pm)-SM-21 administered i.p. at the dose of 20–30 mg kg⁻¹ in the rat, and 20 mg kg⁻¹ in guinea pigs, induced antinociception starting 15 min after injection, reaching a maximum after 30 min and persisting up to 45 min (table 1). The analgesic profile of (\pm)-SM-21 was also investigated in Wistar and Fisher 344 rat strains by using the tail flick test (fig. 4). In both rat strains used (\pm)-SM-21 exhibited a similar antinociceptive activity 30 min after i.p. injection of 20–30 mg kg⁻¹ (fig. 4).

The analgesic effect of the two enantiomers of (\pm)-SM-21, R-(+)-SM-21 and S(-)-SM-21, was evaluated in the mouse hot-plate test (fig. 5, panels A and C) and in the acetic acid abdominal constriction test (fig. 5, panels B and D). Both enantiomers dose-dependently were able to increase the pain threshold, even if R-(+)-SM-21 was slightly more effective than S(-)-SM-21.

The areas under the curve of the antinociception induced by (\pm)-SM-21 (30 mg kg⁻¹ i.p.), R-(+)-hyoscyamine (5 μ g kg⁻¹ i.p.), morphine (8 mg kg⁻¹ i.p.), diphenhydramine (20 mg kg⁻¹ i.p.) and clomipramine (25 mg kg⁻¹ i.p.) are reported in figure 6. The doses of the analgesic drugs chosen were the highest that did not impair rota-rod performance.

The potentiating effect on antinociception exerted by non-analgesic doses of physostigmine (0.05 mg kg⁻¹ i.p.) on AFDX-116 (6.3 ng per mouse i.c.v.) and R-(+)-hyoscyamine (5 μ g kg⁻¹ s.c.) in the mouse hot-plate test is illustrated in table 2. In the same experimental conditions physostigmine was not able to potentiate (\pm)-SM-21 (30 mg kg⁻¹ i.p.) antinociception (Table 2).

Antagonism of the (\pm)-SM-21 induced antinociception. In the mouse hot-plate test, the antinociceptive effect of (\pm)-SM-21 (30 mg kg⁻¹ s.c.) was not antagonized by naloxone (1 mg kg⁻¹ i.p.), CGP-35348 (2.5 μ g per mouse i.c.v.), (R)- α -methylhistamine (10 mg kg⁻¹ i.p.), quinpirole (0.1 mg kg⁻¹ i.p.), GR-48125 (20 mg kg⁻¹ i.p.), N⁶-cyclopentyladenosine (5 μ g per mouse i.c.v.), NAN 190 (0.5 μ g per mouse i.c.v.) (table 3) and, in the abdominal constriction test, by reserpine (2 mg kg⁻¹ i.p.) (fig. 3). Conversely, atropine (5 mg kg⁻¹ i.p.), pirenzepine (0.1 μ g per mouse i.c.v.) and hemicolinium-3 (1 μ g per mouse or rat i.c.v.) were able to completely prevent (\pm)-SM-21 antinociception in the mouse hot-plate (table 3), abdominal constriction (fig. 3) and the rat paw-pressure tests (table 1). All antagonists were injected 15 min before (\pm)-SM-21, with the exception of reserpine, injected twice 48 and 24 h before the test, N⁶-cyclopentyladenosine, administered simultaneously with (\pm)-SM-21 and CGP 35348, injected 5 min before (\pm)-SM-21.

Evaluation of the (\pm)-SM-21 effect in the carrageenan-induced paw edema test. (\pm)-SM-21 failed to suppress paw edema in response to carrageenan administration at the dose of 20 and 30 mg kg⁻¹ i.p. The positive control, indomethacin at the dose of 1 mg kg⁻¹ i.p., produced a significant inhibition over saline + carrageenan treated control animals (data not shown).

Evaluation of the SM-21 effect on spontaneous activity and motor coordination. The motor coordination of mice treated with (\pm)-SM-21, R-(+)-SM-21 and S(-)-SM-21 was evaluated by using the rota-rod test (table 4) while their




Fig. 3. A) Dose-response curves of (\pm)-SM-21 administered s.c. and effect of both HC-3 (1 μ g per mouse i.c.v.) and reserpine (2 mg kg⁻¹ i.p.) pretreatments on antinociception induced by (\pm)-SM-21 (10 mg kg⁻¹ s.c.) in the mouse abdominal constriction test induced by acetic acid 0.6%. The nociceptive responses were recorded 15 min after (\pm)-SM-21 administration. HC-3 was injected 5 h and reserpine 48 and 24 h before testing. B) Time course of (\pm)-SM-21 administered i.c.v. The nociceptive responses were recorded 15, 30, 45 and 60 min after (\pm)-SM-21 administration. Vertical lines show S.E.M. [^] P < .05; * P < .01 in comparison with saline controls. ^o P < .01 in comparison with (\pm)-SM-21 (10 mg kg⁻¹ s.c.). Numbers inside the columns indicate the number of mice.

TABLE 1

Antinociception exerted by (\pm)-SM-21 in the paw-pressure test: in the rat (A) and in the guinea pig (B) and antagonism by atropine and HC-3 on (\pm)-SM-21 antinociception in the rat

	Pretreatment	Treatment, i.p.	Dose (mg kg ⁻¹)	Paw Pressure (g)			
				Before Pretreatment	After Treatment		
					15 min	30 min	45 min
A)	Saline	Saline		58.3 ± 2.2 (18)	61.8 ± 2.8 (18)	60.2 ± 3.4 (18)	61.2 ± 2.6 (18)
	10 ml kg ⁻¹ i.p.	(\pm)-SM-21	10	60.3 ± 4.0 (5)	62.8 ± 4.4 (5)	59.4 ± 4.2 (5)	57.8 ± 4.0 (5)
	Saline	(\pm)-SM-21	20	60.4 ± 3.2 (16)	97.2 ± 4.2* (16)	100.4 ± 4.6* (16)	86.6 ± 3.8* (16)
		(\pm)-SM-21	30	62.0 ± 3.6 (8)	93.2 ± 5.2* (8)	95.4 ± 3.8* (8)	84.5 ± 3.6* (8)
	Atropine	Saline		58.8 ± 3.2 (10)	60.4 ± 4.2 (10)	59.7 ± 3.2 (10)	58.7 ± 3.0 (10)
	5 mg kg ⁻¹	(\pm)-SM-21	20	62.7 ± 2.8 (5)	63.2 ± 4.6° (5)	64.5 ± 4.0° (5)	61.1 ± 4.8° (5)
	i.p.	Saline		56.8 ± 3.4 (10)	62.4 ± 4.0 (10)	65.3 ± 4.2 (10)	62.3 ± 3.6 (10)
	HC-3	(\pm)-SM-21	20	60.6 ± 2.6 (5)	66.8 ± 4.4° (5)	68.5 ± 4.4° (5)	60.4 ± 4.5° (5)
	1 µg per rat	Saline		44.6 ± 3.6 (10)	41.4 ± 2.8 (10)	43.6 ± 3.8 (10)	42.0 ± 2.4 (10)
	i.c.v.	(\pm)-SM-21	20	40.0 ± 2.2 (5)	92.0 ± 4.0* (5)	110.0 ± 6.6* (5)	64.0 ± 6.2 (5)
B)	Saline		44.6 ± 3.6 (10)	41.4 ± 2.8 (10)	43.6 ± 3.8 (10)	42.0 ± 2.4 (10)	
	(\pm)-SM-21	20	40.0 ± 2.2 (5)	92.0 ± 4.0* (5)	110.0 ± 6.6* (5)	64.0 ± 6.2 (5)	

The number of rats and guinea pigs is shown in parentheses.

Atropine and HC-3 were injected 15 min and 5 h before (\pm)-SM-21, respectively.

* P < .01 in comparison with saline-saline; ° P < .01 vs. saline-(\pm)-SM-21.




Fig. 4. Antinociceptive effect of (\pm)-SM-21 in the rat tail-flick test. Nociceptive responses were recorded 30 min after drug administration. Vertical lines show S.E.M. * P < .01 in comparison with saline controls. Numbers inside the columns indicate the number of mice.

spontaneous activity was investigated by using the hole-board test. The rota-rod performance of mice treated with (\pm)-SM-21 at the dose of 30 and 40 mg kg⁻¹ s.c. and both enantiomers at the dose of 20 and 30 mg kg⁻¹ s.c. was not impaired in comparison with controls (table 4). On the contrary, (\pm)-SM-21 administered at higher doses (50 and 60 mg kg⁻¹ s.c.) produced a significant impairment of the rota-rod performance (table 4). The number of falls by control animals progressively decreased at every measurement since the mice learnt how to balance on the rotating rod. The spontaneous motility and exploratory behavior of mice was not

modified by treatment with (\pm)-SM-21 (20 and 40 mg kg⁻¹ s.c.) as revealed by the hole-board test (data not shown).

In vitro functional studies. (\pm)-SM-21 blocked the McN-A-343-induced inhibition of twitch contractions of the rabbit vas deferens ($pK_B = 5.97 \pm 0.11$), antagonized the negative inotropic carbachol-induced effect in guinea-pig left atrium ($pK_B = 6.63 \pm 0.10$), the contractile responses to acetylcholine in guinea-pig ileum ($pK_B = 6.35 \pm 0.04$) and to carbachol in immature guinea-pig uterus ($pK_B = 6.26 \pm 0.05$) as shown in table 5. Increasing concentrations of (\pm)-SM-21 produced parallel shifts of the agonist concentration-response curves progressively to the right and no appreciable change in basal tension or maximum agonist response was observed (data not shown). pA_2 values of R-(+)-hyoscyamine and AFDX-116, used as reference drugs, are shown in table 5. The selectivity ratios for (\pm)-SM-21, R-(+)-hyoscyamine and AFDX-116, obtained as differences between respectively pK_B or pA_2 values, are reported in table 5.

Finally, (\pm)-SM-21 was shown to be endowed with a weak antiacetylcholinesterase activity, its IC_{50} value on electrical eel acetylcholinesterase being $1.1 \cdot 10^{-4}$ M (data not shown).

Discussion

(\pm)-SM-21 was able to induce antinociception in mice, rats and guinea-pigs. Antinociception was elicited regardless of which noxious stimulus was used: thermal (hot-plate and tail flick tests), chemical (abdominal constriction test) and mechanical (paw pressure test). (\pm)-SM-21 antinociception was obtained without producing any visible modification of animal gross behavior. Moreover, (\pm)-SM-21 treated mice showed a complete integrity of motor coordination on the rota-rod test, normal spontaneous motility, as well as exploratory behavior as revealed by the hole-board test.




Fig. 5. Dose-response curves of R-(+)-SM-21 (panels A and C) and S(-)-SM-21 (panels B and D) in the mouse hot-plate test and abdominal constriction test. The doses are expressed as mg kg⁻¹ s.c. Vertical lines show S.E.M. ^ P < .05; * P < .01 in comparison with saline controls. Numbers inside the columns are the number of mice.

(±)-SM-21 exerted its antinociceptive effect by acting centrally. It was, in fact, possible to reach the same intensity of analgesia by injecting directly into the cerebral ventricles doses (5–20 μg per mouse) of (±)-SM-21 which were one thousand times lower than those needed parenterally. That the antinociception depends on a retrodiffusion of the drug from the cerebral ventricles to the periphery can thus be ruled out.

(±)-SM-21 antinociception was found to be dependent on central cholinergic activation since it was prevented by the non-selective muscarinic antagonist atropine, the M₁-antagonist pirenzepine and the ACh depletor HC-3. Taking into account that HC-3 and pirenzepine were able to antagonize (±)-SM-21 antinociception after i.c.v. injection, this supports the hypothesis that the analgesic site of action of (±)-SM-21 is localized in the CNS. A presynaptic mechanism facilitating cholinergic transmission is involved in (±)-SM-21 antinociception as revealed by the antagonism by HC-3. A postsynaptic mechanism of action can be ruled out since, as reported by Bartolini et al. (1987; 1992), HC-3 was not able to antagonize antinociception induced by agonists of postsynaptic muscarinic receptors such as oxotremorine, McN-A-343 and AF-102B.

The hypothesis of a presynaptic cholinergic mechanism for (±)-SM-21 is in agreement with previous results demonstrating, by microdialysis studies, an increase in ACh release from rat cerebral cortex induced by (±)-SM-21 administration (Bartolini et al., 1994). This effect occurred in the same range of doses (10 and 20 mg kg⁻¹ i.p.) in which the above-mentioned compound exerted its antinociceptive activity.

(±)-SM-21, like R-(+)-hyoscyamine (see introduction), demonstrated antinociceptive properties underlying a presynaptic cholinergic mechanism, but with greater efficacy than that exerted by R-(+)-hyoscyamine. The analgesic effect of (±)-SM-21 was also compared with the analgesia induced by some analgesic drugs such as morphine, diphenhydramine and clomipramine at the highest doses that did not impair the rota-rod performances. By comparing the areas under the curve, the antinociceptive efficacy of (±)-SM-21 (30 mg kg⁻¹ s.c.) resulted almost equal to that exerted by morphine (8 mg kg⁻¹ s.c.), but greater than those induced by diphenhydramine (20 mg kg⁻¹ s.c.) and clomipramine (25 mg kg⁻¹ s.c.).

(±)-SM-21 and R-(+)-hyoscyamine have been reported to increase the extracellular levels of ACh in cortical microdialysis studies (Bartolini et al., 1994; Romanelli et al., 1995). Since ACh release can be increased by blocking M₂/M₄ mus-




Fig. 6. Area under the curve of the analgesic effect of (\pm)-SM-21 in comparison with R-(+)-hyoscyamine, morphine, diphenhydramine and clomipramine in hot-plate test. The nociceptive responses were recorded from 15 min after drug injection to 90 min after drug administration. Numbers inside the columns indicate the number of mice.

carinic autoreceptors (Lapchak *et al.*, 1989; Töröcsik and Vizi, 1991; McKinney *et al.*, 1993; Stillman *et al.*, 1993) and R-(+)-hyoscyamine showed a very high affinity for the prepuberal guinea-pig uterus putative M_4 receptors (Ghelardini *et al.*, 1993), the (\pm)-SM-21 affinity profile towards muscarinic receptor subtypes was investigated in vitro. The affinity profile of (\pm)-SM-21 versus M_1 (rabbit vas deferens), M_2 (guinea-pig atrium), M_3 (guinea-pig ileum) and putative M_4 receptors (prepuberal guinea-pig uterus) was evaluated by in vitro functional studies. The M_4 muscarinic receptor subtype has been defined as putative since it has not been confirmed that the mRNA codifying M_4 is expressed in the prepuberal

uterus tissue. However, pharmacological and biochemical studies show that the M_4 putative receptor of prepuberal guinea-pig uterus has a pharmacological and biochemical profile identical to that of the muscarinic m_4 receptor subtype expressed in the rat striatum (McKinney *et al.*, 1991; Waelbroeck *et al.*, 1992) and in NG 108-15 cells (Leiber *et al.*, 1984; Marc *et al.*, 1986). (\pm)-SM-21 showed, unlike R-(+)-hyoscyamine, a very low M_4/M_1 muscarinic receptor subtype selectivity ratio (twice), but higher M_2/M_1 selectivity ratio (4.6 times). By comparing the affinity profile of the M_2 muscarinic antagonists: AFDX-116 (Giachetti *et al.*, 1986), methoctramine (Melchiorre *et al.*, 1987) and AQRA-741 (Doods *et al.*, 1991), a selectivity ratio M_2/M_1 lower than that showed by (\pm)-SM-21 can be observed. Moreover, all the above-mentioned M_2 antagonists, like (\pm)-SM-21, are also endowed with cholinergic presynaptic antinociceptive properties (Bartolini *et al.*, 1989; Gualtieri *et al.*, 1989; Ghelardini *et al.*, 1991) and AFDX-116 and methoctramine are able to increase the ACh release (Lapchak *et al.*, 1989; Töröcsik and Vizi, 1991). It seems, therefore, reasonable to suppose that a selectivity ratio of 4.6, even if small, may be high enough to enhance the pain threshold as a consequence of ACh release. The antinociception induced by (\pm)-SM-21 may be due to the antagonism of the M_2 muscarinic autoreceptor. The selectivity on blocking the M_2/M_4 towards M_1 was evaluated since Bartolini *et al.*, (1992) have demonstrated that the muscarinic postsynaptic receptor responsible for central cholinergic antinociception belongs to the M_1 subtype. The antinociceptive efficacy of (\pm)-SM-21 was greater than that of R-(+)-hyoscyamine, AFDX-116, methoctramine or AQRA-741. Therefore, we cannot exclude that other mechanisms able to potentiate the endogenous cholinergic system may be involved in the antinociception induced by (\pm)-SM-21.

Both enantiomers of SM-21, R-(+) and S(-), contrary to atropine in which the analgesic activity resides only in the R-(+) isomer (Ghelardini *et al.*, 1992), showed very similar antinociceptive properties in the presence of either a thermal or chemical stimulus. However, in both analgesic tests used,

TABLE 2

Effect of physostigmine on (\pm)-SM-21, AFDX-116 and R-(+)-hyoscyamine antinociception in the mouse hot-plate test

Treatment	Before Pretreatment	Licking Latency(s)		
		After Treatment		
		15 min	30 min	45 min
Saline	13.9 \pm 0.9 (8)	14.3 \pm 1.6 (8)	14.6 \pm 1.2 (8)	13.8 \pm 1.5 (8)
Physostigmine 0.05 mg kg ⁻¹ i.p.	15.7 \pm 1.1 (15)	16.2 \pm 1.5 (15)	14.1 \pm 1.2 (15)	13.6 \pm 1.4 (15)
(\pm)-SM-21 30 mg kg ⁻¹ s.c.	14.1 \pm 1.0 (18)	33.8 \pm 1.6* (18)	27.5 \pm 1.3* (18)	25.7 \pm 1.5* (18)
AFDX-116 6.3 ng/mouse i.c.v.	13.9 \pm 1.3 (12)	22.1 \pm 1.3* (12)	18.3 \pm 1.2^ (12)	15.2 \pm 1.2 (12)
R-(+)-hyoscyamine 5 µg kg ⁻¹ s.c.	14.2 \pm 0.9 (25)	25.2 \pm 0.7* (25)	20.4 \pm 0.8* (25)	15.1 \pm 1.1 (25)
Physostigmine + (\pm)-SM-21	14.5 \pm 0.8 (10)	32.7 \pm 1.3* (10)	28.4 \pm 1.6* (10)	23.8 \pm 1.3* (10)
Physostigmine + AFDX-116	14.4 \pm 0.5 (10)	31.6 \pm 2.1* ^o (10)	29.8 \pm 2.7* ^o (10)	22.4 \pm 1.9* ^o (10)
Physostigmine + R-(+)-hyoscyamine	13.8 \pm 0.9 (16)	35.2 \pm 1.4* ^o (16)	32.1 \pm 2.2* ^o (16)	23.3 \pm 1.3* ^o (16)

The number of mice is shown in parentheses. Physostigmine was injected 5 min before the other drugs. ^ P < .05; * P < .01 in comparison with saline controls. ° P < .05 vs. the corresponding group treated with the same analgesic drug.

TABLE 3

Effects of atropine, pirenzepine, HC-3, naloxone, CGP-35348, RAMH, quinpirole, GR-48125, N⁶-cyclopentyladenosine and NAN-190 on antinociception induced by (±)-SM-21 in the mouse hot-plate test

Pretreatment	Treatment	mg · kg ⁻¹ s.c.	n	Licking Latency(s)			
				Before Pretreatment	After Treatment		
					15 min	30 min	45 min
Saline 10 ml kg ⁻¹ i.p.	Saline		44	14.4 ± 0.6	14.1 ± 1.1	13.6 ± 1.2	14.0 ± 0.9
Saline 5 µl i.c.v.	Saline		28	13.7 ± 0.7	14.3 ± 1.0	14.5 ± 0.8	14.3 ± 0.7
Saline i.p. or i.c.v.	(±)-SM-21	30	41	15.8 ± 0.7	33.7 ± 0.2*	28.1 ± 0.7*	24.3 ± 0.6*
Atropine 5 mg kg ⁻¹ i.p.	Saline		15	14.3 ± 0.6	13.8 ± 0.8	14.1 ± 1.2	14.4 ± 1.6
	(±)-SM-21	30	18	14.9 ± 0.9	18.8 ± 1.7°	18.0 ± 1.5°	14.5 ± 1.6°
Pirenzepine 0.1 µg mouse i.c.v.	Saline		10	13.9 ± 1.3	14.5 ± 1.7	14.2 ± 2.0	15.0 ± 1.6
	(±)-SM-21	30	10	14.0 ± 1.0	18.3 ± 2.1°	18.2 ± 2.3°	17.6 ± 1.8°
HC-3 1 µg per mouse i.c.v.	Saline		14	14.8 ± 1.0	14.1 ± 1.7	13.8 ± 1.2	14.1 ± 2.1
	(±)-SM-21	30	11	13.6 ± 0.8	18.7 ± 2.2°	18.1 ± 1.9°	16.7 ± 1.6°
Naloxone 1 mg kg ⁻¹ i.p.	Saline		13	15.0 ± 0.8	13.9 ± 1.3	14.8 ± 1.3	15.0 ± 1.4
	(±)-SM-21	30	8	15.2 ± 1.2	30.8 ± 2.2*	27.6 ± 2.3*	26.1 ± 2.3*
CGP 35348 2.5 µg mouse i.c.v.	Saline		10	14.1 ± 0.7	12.6 ± 1.8 [^]	13.1 ± 2.1	12.8 ± 1.6
	(±)-SM-21	30	8	14.5 ± 1.0	35.1 ± 2.0*	31.2 ± 2.4*	26.0 ± 1.9*
RAMH 10 mg kg ⁻¹ i.p.	Saline		10	14.0 ± 0.7	12.8 ± 0.6	13.7 ± 1.1	15.8 ± 1.7
	(±)-SM-21	30	10	15.4 ± 0.6	33.5 ± 2.8*	28.6 ± 3.6*	25.7 ± 2.8*
Quinpirole 0.1 mg kg ⁻¹ i.p.	Saline		10	15.0 ± 1.0	14.2 ± 2.0	14.3 ± 1.8	14.1 ± 1.6
	(±)-SM-21	30	10	13.7 ± 1.2	34.1 ± 2.2*	27.1 ± 1.7*	25.6 ± 2.0*
GR 48125 20 mg kg ⁻¹ i.p.	Saline		10	13.7 ± 1.1	13.2 ± 1.9	14.1 ± 1.7	14.4 ± 1.5
	(±)-SM-21	30	10	14.6 ± 0.6	31.5 ± 2.3*	26.5 ± 1.9*	22.7 ± 2.2*
N ⁶ -CPA 5 µg mouse i.c.v.	Saline		12	13.9 ± 0.7	15.0 ± 0.9	15.7 ± 1.1	16.4 ± 1.7
	(±)-SM-21	30	8	15.2 ± 0.9	30.7 ± 3.0*	33.1 ± 3.0*	26.7 ± 2.1*
NAN-190 0.5 µg mouse i.c.v.	Saline		12	15.2 ± 1.3	14.7 ± 1.5	13.2 ± 1.5	14.9 ± 1.7
	(±)-SM-21	30	9	14.4 ± 0.8	34.7 ± 2.1*	29.5 ± 1.7*	22.1 ± 1.5*

* P < .01; [^] P < 0.05 in comparison with saline-saline; ° P < .01 vs. saline-(±)-SM-21-treated mice. N⁶-cyclopentyladenosine = N⁶ CPA.

TABLE 4

Effect of (±)-SM-21, R-(+)-SM-21 and S-(-)-SM-21 in the mouse rota-rod test

Treatment	Dose	Number of Falls in 30 s			
		Before Pretreatment	After Treatment		
			15 min	30 min	45 min
Saline	10 mg kg ⁻¹ s.c.	3.2 ± 0.3	1.8 ± 0.3	1.4 ± 0.2	0.8 ± 0.2
(±)-SM-21	30 mg kg ⁻¹ s.c.	2.8 ± 0.2	1.9 ± 0.3	1.5 ± 0.5	1.2 ± 0.5
(±)-SM-21	40 mg kg ⁻¹ s.c.	3.3 ± 0.4	1.6 ± 0.6	1.2 ± 0.4	1.2 ± 0.4
(±)-SM-21	50 mg kg ⁻¹ s.c.	2.8 ± 0.6	2.2 ± 0.4	2.6 ± 0.5*	2.4 ± 0.5*
(±)-SM-21	60 mg kg ⁻¹ s.c.	2.7 ± 0.4	3.5 ± 0.5*	4.3 ± 0.4*	4.4 ± 0.6*
R-(+)-SM-21	30 mg kg ⁻¹ s.c.	2.9 ± 0.3	2.0 ± 0.4	1.5 ± 0.3	1.4 ± 0.4
R-(+)-SM-21	40 mg kg ⁻¹ s.c.	3.1 ± 0.4	3.6 ± 0.5*	4.1 ± 0.4*	3.5 ± 0.5*
S-(-)-SM-21	30 mg kg ⁻¹ s.c.	3.0 ± 0.3	2.1 ± 0.4	1.6 ± 0.2	0.8 ± 0.4
S-(-)-SM-21	40 mg kg ⁻¹ s.c.	2.8 ± 0.4	3.4 ± 0.5*	3.4 ± 0.4*	2.5 ± 0.6*
Saline	5 µl i.c.v.	2.5 ± 0.4	1.8 ± 0.3	1.1 ± 0.3	0.7 ± 0.2
(±)-SM-21	20 µg i.c.v.	2.9 ± 0.5	2.2 ± 0.4	1.2 ± 0.3	1.0 ± 0.2
(±)-SM-21	40 µg i.c.v.	2.4 ± 0.4	3.5 ± 0.3*	3.3 ± 0.4*	2.2 ± 0.4*
Saline	10 ml kg ⁻¹ p.o.	2.8 ± 0.3	2.0 ± 0.3	1.1 ± 0.2	1.2 ± 0.4
(±)-SM-21	60 mg kg ⁻¹ p.o.	2.6 ± 0.4	1.5 ± 0.4	1.3 ± 0.4	0.6 ± 0.2
(±)-SM-21	90 mg kg ⁻¹ p.o.	3.0 ± 0.4	3.2 ± 0.5*	2.9 ± 0.3*	2.4 ± 0.4*
Saline	10 ml kg ⁻¹ i.v.	3.1 ± 0.3	2.8 ± 0.3	2.3 ± 0.4	1.2 ± 0.2
(±)-SM-21	20 ml kg ⁻¹ i.v.	2.8 ± 0.4	2.4 ± 0.3	1.3 ± 0.3	1.1 ± 0.3
(±)-SM-21	30 ml kg ⁻¹ i.v.	2.6 ± 0.3	3.9 ± 0.4*	3.3 ± 0.3*	2.4 ± 0.3*

Each value represents the mean of 5 to 10 mice.

* P < .01 in comparison with saline controls.

R-(+)-SM-21 resulted weakly more effective than S-(-)-SM-21. Since R-(+) and S-(-)-SM-21 were not endowed with a different analgesic profile, their in vitro selectivity towards the muscarinic receptor subtypes was not considered worth investigating.

It has been demonstrated that D₂ dopaminergic (Gorell and Czarnecki, 1986; Wedzony *et al.*, 1988; Scatton, 1992; Imperato *et al.*, 1993), A₁ adenosinergic (Jackisch *et al.*, 1984;

Carter *et al.*, 1995), H₃ histaminergic (Clapham and Kilpatrick, 1992), 5-HT₄ serotonergic heteroreceptors (Consolo *et al.*, 1994), all located on central cholinergic neurones, increase ACh release. The involvement of the above-mentioned heteroreceptors was, therefore, investigated. Quinpirole (D₂ agonist), N⁶-cyclopentyladenosine (A₁ agonist), R-(α)-methylhistamine (H₃ agonist), and GR-48125 (5-HT₄ antagonist), at doses able to prevent the antinociception in-

TABLE 5

Affinity profiles of (±)-SM-21, R-(+)-hyoscyamine and AFDX-116 at muscarinic M₁ receptors in rabbit vas deferens, M₂ receptors in guinea pig left atrium, M₃ receptors in guinea pig ileum and M₄ putative receptors in guinea pig uterus. The ratios of affinity constants are given as a measure of receptor selectivity

Drug	pA ₂ Values				Selectivity Ratios	
	M ₁ Rabbit Vas Deferens	M ₂ Rat Left Atrium	M ₃ Rat Ileum	M ₄ -putative Guinea Pig Uterus	M ₂ /M ₁	M ₄ /M ₁
(±)-SM-21	5.97 ± 0.11*	6.63 ± 0.10*	6.35 ± 0.04*	6.26 ± 0.05*	4.6	1.9
R-(+)-hyoscyamine	7.05 ± 0.05 ^a	7.25 ± 0.04 ^a	6.88 ± 0.05 ^a	9.56 ± 0.01 ^a	1.6	323
AFDX-116	6.84 ± 0.14 ^b	7.12 ± 0.11 ^b	6.34 ± 0.13 ^c	6.70 ± 0.06	1.9	0.7

* pK_B values are obtained with (±)-SM-21 1 μM. Each value is the mean of at least five experiments. pA₂ values are the mean ± S.E.M. of 12 to 16 preparations. Selectivity ratios were calculated as antilogs of the difference between pK_B or pA₂ values.

^a Ghelardini *et al.*, (1993); ^b Eltze (1988); ^c Eltze and Figala (1988).

duced respectively by haloperidol, caffeine (Ghelardini *et al.*, 1992), thioperamide (Malmberg-Aiello *et al.*, 1994), BIMU 1 and BIMU 8 (Ghelardini *et al.*, 1996), failed to prevent (±)-SM-21 antinociception. Previous data have shown that antinociception induced by (±)-SM-21 and its enantiomers was partially prevented by the 5-HT₄ antagonist SDZ 205-557 (Romanelli *et al.*, 1995). Since GR-48125 was more selective than SDZ 205-557 towards 5-HT₄ receptors and was not able to antagonize SM-21 antinociception, the prevention of the SM-21 effect produced by SDZ 205-557 was probably not related to an antagonism of 5-HT₄ receptors. It has also been observed that the activation of the serotonergic autoreceptor 5-HT_{1A} enhances ACh release from the guinea-pig cortex (Bianchi *et al.*, 1990). Pretreatment with the 5-HT_{1A} selective antagonist NAN 190 at doses which block the antinociception induced by 5-HT_{1A} agonists (Ghelardini *et al.*, 1994), did not prevent the enhancement of the pain threshold produced by (±)-SM-21 administration. The present data suggest that the above-mentioned receptors, even though they are able to increase ACh release, are not involved in (±)-SM-21 mechanism of analgesic action.

The antinociception induced by antagonists of the muscarinic autoreceptors, such as R-(+)-hyoscyamine and AFDX-116, was significantly potentiated by pretreatment with a subliminary non-analgesic dose of physostigmine. By contrast, in the same experimental conditions, (±)-SM-21 antinociception was not modified by physostigmine pretreatment. These observations may suggest that (±)-SM-21 is endowed with very low anticholinesterase activity, a hypothesis that seems to be in agreement with the *in vitro* evaluation of the IC₅₀ value of (±)-SM-21 (IC₅₀ = 1.1 · 10⁻⁴ M). It is possible that (±)-SM-21 is able to amplify cholinergic neurotransmission through the antagonism of the muscarinic autoreceptor and that this effect is in its turn potentiated by its low cholinesterase inhibitory activity.

Other neurotransmitter systems are not involved in (±)-SM-21 antinociception since the opioid antagonist naloxone, the GABA_B antagonist CGP-35348 and the polyamine depletor reserpine, were all unable to prevent the effect of (±)-SM-21. The doses and administration schedules of the above-mentioned drugs were ideal for preventing antinociceptions induced respectively by morphine (Ghelardini *et al.*, 1992), the GABA_B agonist baclofen (Malcangio *et al.*, 1991) and the antidepressant drugs clomipramine and amitriptyline (Galeotti *et al.*, 1995).

(±)-SM-21 at analgesic doses failed to suppress paw edema in response to carrageenan administration, suggesting that its antinociception is not due to an antiinflammatory action.

In summary, our results have shown that (±)-SM-21 is able to produce dose-dependent antinociception in rodents and guinea pigs, without impairing motor coordination, by potentiating endogenous cholinergic activity.

Acknowledgments

The authors wish to thank Giba Geigy for the gift of CGP-35348.

References

- BARTOLINI, A., GALLI, A., GHELARDINI, C., GIOTTI, A., MALCANGIO, M., MALMBERG-AIELLO, P. AND ZUCCHI, P. L.: Antinociception induced by systemic administration of local anaesthetics depends on a cholinergic mechanism. *Br. J. Pharmacol.* **92**: 711-721, 1987.
- BARTOLINI, A., GHELARDINI, C., FANTETTI, L., MALCANGIO, M., MALMBERG-AIELLO, P. AND GIOTTI, A.: Role of muscarinic receptor subtypes in central antinociception. *Br. J. Pharmacol.* **105**: 77-82, 1992.
- BARTOLINI, A., GHELARDINI, C., GIOVANNINI, M. G., CASAMENTI, F., MALMBERG-AIELLO, P., PEPEU, G. AND GIOTTI, A.: Modulators of ACh release as potent cognition enhancers and analgesics: Pharmacodynamic studies. Proceedings of the "XXVII Symposium of the Italian Pharmacological Society," p. 29, 1994.
- BARTOLINI, A., GHELARDINI, C., GUALTIERI, F., MALCANGIO, M., MALMBERG-AIELLO, P., ROMANELLI, M. AND GIOTTI, A.: I.c.v. AFDX 116 induces analgesia only when administered at very low doses. *Trends Pharmacol. Sci.* **99**: suppl. IV, 1989.
- BIANCHI, C., SINISCALCHI, A. AND BEANI, L.: 5-HT_{1A} agonists increase and 5-HT₃ agonist decrease acetylcholine efflux from the cerebral cortex of freely-moving guinea-pigs. *Br. J. Pharmacol.* **101**: 448-452, 1990.
- CARTER, A. D., O'CONNOR, W. T., CARTER, M. J. AND UNGERSTEDT, U.: Caffeine enhances acetylcholine release in the hippocampus *in vivo* by a selective interaction with adenosine A₁ receptors. *J. Pharmacol. Exp. Ther.* **273**: 637-642, 1995.
- CLAPHAM, J. AND KILPATRICK, G. J.: Histamine H₃ receptors modulate the release of [³H]-acetylcholine from slices of rat entorhinal cortex: Evidence for the possible existence of H₃ receptor subtypes. *Br. J. Pharmacol.* **107**: 919-923, 1992.
- CONSOLO, S., ARNABOLDI, S., GIORGI, S., RUSSI, G. AND LADINSKY, H.: 5-HT₄ receptor stimulation facilitates acetylcholine release in frontal cortex. *NeuroReport* **5**: 1230-1232, 1994.
- D'AMOUR, F. E. AND SMITH, D. L.: A method for determining loss of pain sensation. *J. Pharmacol. Exp. Ther.* **72**: 74-79, 1941.
- DEI, S., BELLUCCI, C., GUALTIERI, F., ROMANELLI, M. N., SCAPECCHI, S., TEODORI, E., BARTOLINI, A. AND GHELARDINI, C.: Analgesic, antimuscarinic activity and enantioselectivity of the four isomers of 3-quinclidinyl tropate as compared with the enantiomers of hyoscyamine. *Il Farmaco* **50**: 303-309, 1995.
- DOODS, H., ENTZEROTH, M. AND MAYER, N.: Cardiselectivity of AQ-RA 741, a novel tricyclic antimuscarinic drug. *Eur. J. Pharmacol.* **192**: 147-152, 1991.
- DÖRJE, F., FRIEBE, T., TACKE, R., MUTSCHLER, E. AND LAMBRECHT, G.: Novel pharmacological profile of muscarinic receptors mediating contraction of the guinea-pig uterus. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **342**: 284-289, 1990.
- ELLMAN, G. L., COURTNEY, K. D., ANDRES, V. JR AND FEATHERSTONE, R. M.: A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **7**: 88-95, 1961.
- ELTZE, M.: Muscarinic M₁ and M₂ receptors mediating opposite effects on neuromuscular transmission in rabbit vas deferens. *Eur. J. Pharmacol.* **151**: 205-221, 1988.
- ELTZE, M. AND FIGALA, V.: Affinity and selectivity of biperiden enantiomers for muscarinic receptor subtypes. *Eur. J. Pharmacol.* **158**: 11-19, 1988.
- ELTZE, M., GONNE, S., RIEDEL, R., SCHLOTKE, B., SCHUDT, C. AND SIMON, W. A.: Pharmacological evidence for selective inhibition of gastric acid secretion by telenezepine, a new antimuscarinic drug. *Eur. J. Pharmacol.* **112**: 211-224, 1985.

- GALEOTTI, N., GHELARDINI, C. AND BARTOLINI, A.: Involvement of the serotonergic system in the analgesic effect of tricyclic antidepressants. *Behav. Pharmacol.* **6**: (suppl. 1), 20, 1995.
- GHELARDINI, C., GALEOTTI, N., BALDINI, M., MEONI, P., GIOTTI, A. AND BARTOLINI, A.: The 5-HT_{1A} agonist 8-OH-DPAT induces antinociception through a cholinergic mechanism. *Can. J. Physiol. Pharmacol.* **72**: 1394-1400 (suppl. IUPHAR) p. 380, 1994.
- GHELARDINI, C., GALEOTTI, N., CASAMENTI, F., MALMBERG-AIELLO, P., PEPEU, G., GUALTIERI, F. AND BARTOLINI, A.: Central cholinergic antinociception induced by 5-HT₄ agonists: BIMU 1 and BIMU 8. *Life Sci.* **58**: 2297-2309, 1996.
- GHELARDINI, C., GIOTTI, A., GUALTIERI, F., MATUCCI, R., ROMANELLI, M. N., SCAPECCHI, S., TEODORI, E. AND BARTOLINI, A.: Presynaptic auto- and heteroreceptors in the cholinergic regulation of pain. *In Trends in Receptor Research*, ed. by P. Angeli, U. Gulini and W. Quaglia, pp. 95-114. Elsevier Science Publisher B. V., Amsterdam, NL, 1992.
- GHELARDINI, C., GIOTTI, A., MALMBERG-AIELLO, P. AND BARTOLINI, A.: Central cholinergic antinociception induced by the M₂ antagonist AQRA-741 (Abstract). *Proceedings of the 8th Camerino-Noordwijkerhout Symposium on "Trends in Receptors Research"*, Camerino, September 8-12, P. 40, p. 160, 1991.
- GHELARDINI, C., GUALTIERI, F., BALDINI, M., DEI, S., ROMANELLI, M. N., GIOTTI, A. AND BARTOLINI, A.: R-(+)-hyoscyamine: The first and selective antagonist for guinea-pig uterus muscarinic receptor subtypes. *Life Sci.* **52**: 569, 1993.
- GHELARDINI, C., MALMBERG-AIELLO, P., GIOTTI, A., MALCANGIO, M. AND BARTOLINI, A.: Investigation into atropine-induced antinociception. *Br. J. Pharmacol.* **101**: 49-54, 1990.
- GHICCHETTI, A., MICHELETTI, R. AND MONTAGNA, E.: Cardioselective profile of AF-DX 116: A muscarinic M₂ receptor antagonist. *Life Sci.* **38**: 1663-1672, 1986.
- GORELL, J. M. AND CZARNECKI, B.: Pharmacological evidence for direct dopaminergic regulation of striatal acetylcholine release. *Life Sci.* **38**: 2239-2246, 1986.
- GUALTIERI, F., CONTI, G., DEI, S., GIOVANNONI, P., NANNUCCI, F., ROMANELLI, M. N., SCAPECCHI, S., TEODORI, E., FANFANI, L., GHELARDINI, C., BARTOLINI, A. AND GIOTTI, A.: Presynaptic cholinergic modulators as potent nootropic and analgesic drugs. I. Tropic and 2-phenylpropionic acid esters. *J. Med. Chem.* **37**: 1704-1711, 1994.
- GUALTIERI, F., GHELARDINI, C., GIOTTI, A., MALCANGIO, M., MALMBERG-AIELLO, P. AND BARTOLINI, A.: Analgesia induced by the M₂ antagonist methoctramine administered i.c.v.. *Trends Pharmacol. Sci.* **suppl. IV**: 99, 1989.
- GUALTIERI, F., ROMANELLI, M. N., SCAPECCHI, S., TEODORI, E., BARTOLINI, A., FANTETTI, L., GHELARDINI, C. AND GIOTTI, A.: Muscarinic presynaptic autoreceptors and muscarinic postsynaptic receptors have opposite stereochemical requirements. *Med. Chem. Res.* **1**: 52-58, 1991.
- HALEY, T. J. AND MCCORMICK, W. G.: Pharmacological effects produced by intracerebral injection of drugs in the conscious mouse. *Br. J. Pharmacol. Chemother.* **12**: 12-15, 1957.
- IMPERATO, A., OBINU, M. C., CASU, A., MASCIA, S., DAZZI, L. AND GESSA, G. L.: Evidence that neuroleptics increase striatal acetylcholine release through stimulation of dopamine D₁ receptors. *J. Pharmacol. Exp. Ther.* **266**: 557-562, 1993.
- JACKISCH, R., STRITTMATTER, H., KASAKOV, I. AND HERTTING, G.: Endogenous adenosine as a modulator of hippocampal acetylcholine release. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **327**: 319-325, 1984.
- KOSTER, R., ANDERSON, M. AND DE BEER, E. J.: Acetic acid for analgesic screening. *Fed. Proc.* **18**: 412, 1959.
- LAPCHAK, P. A., ARAUJO, D. M., QUIRION, R. AND COLLIER, B.: Binding sites for [³H]AF-DX 116 and effect of AF-DX 116 on endogenous acetylcholine release from brain slices. *Brain Res.* **496**: 285-294, 1989.
- LEIBER, D., HARBON, S., GUILLET, J.-G., ANDRÉ, C. AND STROSBERG, D.: Monoclonal antibodies to purified muscarinic receptor display agonist-like activity. *Proc. Natl. Acad. Sci. U.S.A.* **81**: 4331-4334, 1984.
- LEIGHTON, G. E., RODRIGUEZ, R. E., HILL, R. G. AND HUGHES, J.: κ -Opioid agonists produce antinociception after i.v. and i.c.v. but not intrathecal administration in the rat. *Br. J. Pharmacol.* **93**: 553-560, 1988.
- MALCANGIO, M., GHELARDINI, C., GIOTTI, A., MALMBERG-AIELLO, P. AND BARTOLINI, A.: CGP 35348, a new GABA_B antagonist, prevents antinociception and muscle-relaxant effect induced by baclofen. *Br. J. Pharmacol.* **103**: 1303-1308, 1991.
- MALMBERG-AIELLO, P., LAMBERTI, C., GHELARDINI, C., GIOTTI, A. AND BARTOLINI, A.: Role of histamine in rodent antinociception. *Br. J. Pharmacol.* **111**: 1269-1279, 1994.
- MARC, S., LEIBER, D. AND HARBON, S.: Carbachol and oxytocin stimulate the generation of inositol phosphates in the guinea-pig myometrium. *FEBS Lett.* **201**: 9-14, 1986.
- McKINNEY, M., MILLER, J. H. AND AAGAARD, P. J.: Pharmacological characterization of the rat hippocampal muscarinic autoreceptor. *J. Pharmacol. Exp. Ther.* **264**: 74-78, 1993.
- McKINNEY, M., MILLER, J. H., GIBSON, V. A., NICKELSON, L. AND AKSOY, S.: Interaction of agonists with M₂ and M₄ muscarinic receptor subtypes mediating cyclic AMP inhibition. *Mol. Pharmacol.* **40**: 1014-1022, 1991.
- MELCHIORRE, C., CASSINELLI, A. AND QUAGLIA, W.: Differential blockade of muscarinic receptor subtypes by polymethylene tetraamines. Novel class of selective antagonists of cardiac M₂ muscarinic receptors. *J. Med. Chem.* **30**: 201-204, 1987.
- O'CALLAGHAN, J. P. AND HOLTZMAN, S. G.: Quantification of the analgesic activity of narcotic antagonists by a modified hot-plate procedure. *J. Pharmacol. Exp. Ther.* **192**: 497-505, 1975.
- ROMANELLI, M. N., BARTOLINI, A., BERTUCCI, C., DEI, S., GHELARDINI, C., GIOVANNINI, M. G., GUALTIERI, F., PEPEU, G., SCAPECCHI, S. AND TEODORI, E.: Synthesis, absolute configuration and cognition enhancing activity of SM-21, SM-32 and PG-9, new central ACh releasers. *Proceedings of II Int. Symposium on "Medicinal Chemistry Approaches to Alzheimer Disease"*, Strasbourg, August 28-30, 1995.
- SCATTON, B.: Further evidence for the involvement of D₂ but not D₁ dopamine receptors in dopaminergic control of striatal cholinergic transmission. *Life Sci.* **31**: 2883-2890, 1992.
- STILLMAN, M. J., SHUKITT-HALE, B., KONG, R. M., LEVY, A. AND LIEBERMAN, H. R.: Elevation of hippocampal extracellular acetylcholine levels by methoctramine. *Brain Res. Bull.* **32**: 385-389, 1993.
- TÖRÖCSIK, A. AND VIZI, E. S.: Presynaptic effects of methoctramine on release of acetylcholine. *Neuropharmacology* **30**: 293-298, 1991.
- VAUGHT, J., PELLEY, K., COSTA, L. G., SETHI, P. AND ENNA, S. J.: A comparison of the antinociceptive responses to GABA-receptor agonists THIP and baclofen. *Neuropharmacology* **24**: 211-216, 1985.
- WALBROECK, M., CAMUS, J., TASTENOY, M. AND CRISTOPHE, J.: Binding properties of nine 4-diphenyl-acetoxy-N-methyl-piperidine analogues to M₁, M₂, M₃ and putative M₄ muscarinic receptor subtypes. *Br. J. Pharmacol.* **105**: 97-102, 1992.
- WEDZONY, K., LIMBERGER, N., SPATH, L., WICHMAN, T. AND STARE, K.: Acetylcholine release in rat nucleus accumbens is regulated through dopamine D₂-receptors. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **338**: 250-255, 1988.

Send reprint requests to: Prof. Alessandro Bartolini, Department of Pharmacology, University of Florence, Viale G.B. Morgagni, 65, I-50134 Florence, Italy.