



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Antimicrobial resistance profile of Staphylococcus aureus isolated from raw meat: a research for methicillin resistant Staphylococcus

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Antimicrobial resistance profile of Staphylococcus aureus isolated from raw meat: a research for methicillin resistant Staphylococcus aureus (MRSA) / G.Pesavento; B.Ducci; N.Comodo; A.Lo Nostro. - In: FOOD CONTROL. - ISSN 0956-7135. - STAMPA. - 18(3):(2007), pp. 196-200.
[doi:10.1016/j.foodcont.2005.09.013]

Availability:

The webpage <https://hdl.handle.net/2158/355276> of the repository was last updated on

Published version:

DOI: doi:10.1016/j.foodcont.2005.09.013

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Antimicrobial resistance profile of *Staphylococcus aureus* isolated from raw meat: A research for methicillin resistant *Staphylococcus aureus* (MRSA)

G. Pesavento, B. Ducci, N. Comodo, A. Lo Nostro *

Department of Public Health, Food Safety Laboratory, University of Florence, Viale Morgagni 48, 50132 Florence, Italy

Received 3 May 2005; received in revised form 17 September 2005; accepted 19 September 2005

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) has emerged as a risk factor for patients in general population and particularly in immunocompromised patients. As a matter of fact, it can produce serious infections that may then evolve in septicemia. However, transmission of MRSA from food to people can represent a serious problem only for immunocompromised people. Vancomycin is the elective antimicrobial commonly used in case of MRSA infection, but *S. aureus* strains with reduced sensibility to vancomycin are emerging worldwide. We isolated 42 strains of *S. aureus* from 176 samples of raw meat (poultry, pork and beef) during a one-year survey. Each strain was tested against twelve antimicrobial to verify antibiotic resistance. We found no evidence of methicillin-, teicoplanin- or vancomycin-resistance, but a lot of multiresistant microorganisms, i.e. resistant to three or more antibiotics. These results confirm the hypothesis that antibiotics resistance is present not only in nosocomial bacteria, but also in community environments microorganisms.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: *Staphylococcus aureus*; Antimicrobial resistance; Food; Meat

1. Introduction

In the recent decades, antimicrobial resistance or/and sensitivity in bacteria have been considered as a major problem in Public Health. Colonization of patients with methicillin-resistance or/and reduced vancomycin-sensitivity in *Staphylococcus aureus* is a crucial factor in health care (Bradley, 1999; Brun-Buisson, 1998). Genes expressing antibiotic resistance in *S. aureus* can be either in chromosomal or in plasmidic DNA. In the case of plasmidic resistance, *S. aureus* cannot conjugate with another cell because of the lack of the “sexual pilus”, that is usually present in Gram negative bacteria. In such a condition, fagoc transduction or membrane binding are necessary (Courvalin, 1994).

The first evidence of *S. aureus* resistance to penicillin appeared in 1941, only after 2 years since its introduction in clinical therapy. Penicillin resistance is plasmidic, and therefore it spread out very quickly to several other strains, with the result that in the 1980s approximately 90% of *S. aureus*, isolated from patients, were resistant. Unlikely, methicillin-resistance is chromosomal, and therefore its diffusion is slower than the former, but it keeps going.

Because of the extended use and misuse of antibiotics (e.g. in agriculture, stock-farming and in the treatment of human diseases), the number of bacteria that are resistant to antimicrobial agents is rapidly increasing. Isolates from food have shown, in the last decade, a considerable increase in resistance against most antibiotics (Valsangiacomo, Dolina, Peduzzi, & Jäggl, 2000; Yücel, Citak, & Onder, 2005) and also against methicillin (Kitai et al., 2005).

In humans, the prevalence of methicillin-resistant *S. aureus* (MRSA) largely depends on the region, site of

* Corresponding author. Tel.: +39 55 4598 530; fax: +39 55 4598 935.
E-mail address: antonella.lonostro@unifi.it (A. Lo Nostro).

infection, and whether the infection was nosocomial or community onset (Acco, Ferreira, Henriques, & Tondo, 2003). The prevalence of MRSA isolated from nosocomial infections was greater than that from community infections, and spanned from 1.8% in Switzerland and 2% in The Netherlands, to 73.8% in Hong Kong (Diekema et al., 2001; Joo Lee et al., 2001; Naimi et al., 2001; Salmenlinna & Vuopio-Varkila, 2001). In Europe, prevalence of MRSA is increasing: from 0.1–1.5% for Denmark, Sweden and The Netherlands, to 30.3–34.4% for Spain, France and Italy (Voss, Milatovic, Wallrauch-Schwarz, Rosdahl, & Braveny, 1994). In the United States, the prevalence of methicillin resistance among nosocomial *S. aureus* extracts increased from 2.1% in 1975 to 35% in 1991 (Grosso et al., 2000) and it is greater than in community *S. aureus* isolates (Panlilio, Culver, & Gaynes, 1992).

Vancomycin is the antimicrobial commonly used in case of MRSA infection. In 1997, Hiramatsu et al. (1997) described the first clinical *S. aureus* isolate with intermediate resistance to vancomycin. Since then, other strains of *S. aureus* with reduced sensitivity to vancomycin were identified in Europe and in USA (CDCP, 1997). In June 2002, both *S. aureus* completely resistant to vancomycin and a vancomycin resistant *Enterococcus faecalis* were isolated from a patient in the United States (CDCP, 2002). The presence of *vanA* both in VRSA and in VRE suggested exchange of genetic material from enterococcus to staphylococcus. Importance of VRE is mainly due to the capacity of transferring vancomycin resistance genes to MRSA.

The prevalence of nosocomial VRE isolates has increased from 15% in the early 1990s to 24% in 1998 (Massachusetts Dep. Public Health, Div. of Epid. Immuniz., 2001): the use of antimicrobial avoparcin in pork and poultry meat manufacturing factories as a feed additive is correlated to the increase of VRE prevalence in food of animal origin, and consequently in community and in nosocomial isolates (Børgen, Sorum, Wasteson, & Kruse, 2001; Grosso et al., 2000; Müller, Ulrich, Ott, & Müller, 2001). The possibility of transmission of MRSA through food was unknown until 1994 (Kluytmans et al., 1995); however, today we know that when a few cells of *S. aureus* enter an immunocompetent organism, they are destroyed by the gastric juices, but when an immunocompromised patient's food contains cells of *S. aureus*, these can reach the circulatory system and cause infections that may evolve to septicaemia. Recent cases of MRSA infections have been reported in patients hospitalised in the Haematology Unit of the University Hospital in Rotterdam (The Netherlands), where the index case was recognized in a woman who had eaten a banana fruit infected from a nasal carriage of MRSA, and who had prepared the food for haematology patients (Kluytmans et al., 1995).

Another community acquired MRSA infection has been reported in 2001, when a family was involved in an outbreak from ingestion of MRSA with baked pork meat, contaminated from the food handler (Jones, Kellum, Porter, Bell, & Schaffner, 2002).

Food can be then considered an excellent way for introducing pathogenic microorganisms in general population and in immunocompromised people, and therefore it may transfer antibiotic-resistant bacteria to the intestinal tract of consumers very efficiently. It is exactly in the intestine that can occur the transfer of resistance genes between non-pathogenic bacteria and pathogenic or opportunist bacteria (Sorum & L'Abèe-Lund, 2002).

The present study was performed in order to evaluate the rates of food-isolated *S. aureus*, that was resistant to some of the antibiotics normally used in clinical therapy and particularly to methicillin and vancomycin (Monnet, 2000).

2. Methods

Strains of *S. aureus* were obtained from a one-year microbiological survey of raw meat of poultry, pork and beef, purchased from some butchers' shops and supermarkets in the Florence area (Italy). Meat items bought by butchers' shops usually come from local (Tuscany) or national breedings, while meat items sold by supermarkets usually derive from European countries, particularly from Italy, Germany, The Netherlands, and France.

Of the 176 meat samples that were analysed microbiologically: 42 came from poultry, 68 from beef and 66 from pork meat.

S. aureus isolated from Baird Parker agar (Oxoid, Milan, Italy), was confirmed by agglutination test (Staphylase, test, Oxoid, Milan, Italy), catalase test and biochemical tests (Api Staph, bioMérieux, Rome, Italy).

All isolates were tested against 12 antimicrobial agents: cefalotine, oxacillin, ampicillin, tetracycline, trimethoprim-sulfamethoxazole, erythromycin, clindamycin, gentamicin (Polibiodisc, Biolife, Italy), methicillin, teicoplanin, penicillin G, vancomycin (Oxoid, Milan, Italy).

Antimicrobial sensitivity was monitored with the disk diffusion assay (Kirby–Bauer) recommended by the National Committee for Clinical Laboratory Standards (NCCLS, 2000) on Muller Hinton agar (Oxoid, Milan, Italy); the zone of inhibition was interpreted according to NCCLS guidelines (Bauer, Kirby, Sherris, & Turck, 1966; NCCLS, 2000).

Reference strain used for quality control was *S. aureus* ATCC 25923 (Culti-Loops, Oxoid, Milan, Italy).

3. Statistics

Statistical analysis were done by StataCorp. 2003. Stata Statistical Software: Release 8.0. College Station, TX: Stata Corporation.

4. Results

During the one-year survey, we analysed 176 samples of raw meat purchased by some retailers in the Florence area.

Table 1
Antimicrobial sensitivity of *S. aureus* isolates ($n = 42$)

Agent	Resistance								Fisher's exact test
	Poultry (12)		Pork (10)		Beef (20)		Total (42)		
	<i>n</i>	(%)	<i>n</i>	(%)	<i>n</i>	(%)	<i>n</i>	(%)	
Cefalotine	0	0	1	10.00	0	0	1	2.38	0.238
Oxacillin	8	66.66	1	10.00	6	30.00	15	35.71	0.018
Ampicillin	7	58.33	2	20.00	9	45.00	18	42.86	0.204
Tetracycline	1	8.33	2	20.00	5	25.00	8	19.04	0.529
Trimethoprim-sulphamethoxazole	1	8.33	1	10.00	0	0	2	4.76	0.268
Eritromycin	1	8.33	2	20.00	5	25.00	8	19.04	0.529
Clindamycin	1	8.33	3	30.00	5	25.00	9	21.43	0.391
Gentamicin	2	16.66	1	10.00	1	5.00	4	9.52	0.674
Methicillin	0	0	0	0	0	0	0	0	1
Teicoplanin	0	0	0	0	0	0	0	0	1
Penicillin G	3	25.00	2	20.00	2	10.00	7	16.66	0.566
Vancomycin	0	0	0	0	0	0	0	0	1

S. aureus was present in 42 of the 176 samples (23.86%). Among them 12 (28.57%) were from poultry, 20 (29.41%) from beef, and 10 (15.15%) from pork meat.

All strains, tested for antibiotic sensitivity, were sensitive to methicillin, teicoplanin and vancomycin, and only one was resistant to cephalotone.

After the comparison of inhibition diameters and the relative frequencies, we could assess that the investigated strains had different patterns of antibiotic-resistance from those *S. aureus* isolated from nasal carriage or nosocomial infections. Table 1 shows the results of the antibiograms of all 42 *S. aureus* extracts; 13 strains (30.95%) were sensitive to all the tested antibiotics, 1 strain (2.38%) was sensible to all antibiotics except of oxacillin with an intermediate value, and 28 (66.67%) strains showed resistance to at least one antimicrobial substance.

We found similar resistance rates to those obtained in another study made in Tuscany (Italy) (Zanelli et al., 2002) from human nasal carriage only for a few antibiotics: the percentage of *S. aureus* resistant to gentamicin was 9.52% against a value of 13.2% coming from clinical samples, and no vancomycin-, teicoplanin- or methicillin-resistance was found. For the other tested antibiotics no similarity was found. Instead, we found a much stronger resistance against the other drugs, except for penicillin, where our results show lower levels. We found differences in antibiotic sensitivity also in another study made in Switzerland, where antibiotic sensitivity was studied in *S. aureus* isolated from food (Diekema et al., 2001; Valsangiacomo et al., 2000). Our strains present higher resistance levels for all the tested antimicrobial substances, except for ampicillin and penicillin G. Resistances were similar (Fisher's exact test >0.05) in the three kind of meat except for oxacillin, where resistance was seen more frequently in isolates from poultry as in isolates from pork and beef (Fisher's exact test = 0.018).

Although MRSA strain has always shown multiresistance (Salmenlinna & Vuopio-Varkila, 2001; Joo Lee et al., 2001), we found multiresistance also in our strains

of *S. aureus* methicillin sensible (MSSA). Only 11 (26.19%) of all the tested strains showed single resistance: 6 of them (14.28%) resistant to oxacillin, 3 (7.14%) to ampicillin, 1 (2.38%) to clindamycin, and 1 to penicillin G. Double resistances were present in 4 (9.52%) of the total strains isolated; the other 13 strains (30.95%) show multiple (≥ 3 antibiotics) resistances, 6 of them (14.28%) presented triple resistances, 3 (7.14%) quadruple resistances, 1 quintuple resistances and 3 sextuple resistances.

5. Discussion

Meat is an important vector for the transfer of antibiotic resistances from animals to humans, and antimicrobial resistance has always been a major concern for nosocomial infections in hospital environments. Such transfer can occur in three ways: by means of antibiotic residues in food, through the transfer of resistant food borne pathogens, or through the ingestion of resistant parts of the original food micro flora and resistance transfer to pathogenic microorganisms (Klein, 1999; Kruse & Sorum, 1994; Mayrhofer, Paulsen, Smulders, & Hilbert, 2004; Teuber, 1999).

The great use of avoparcin as a feed additive, at least in the last decade in Europe, in poultry and pork meat factories, has induced the occurrence of vancomycin-resistant enterococci and the potential transmission of this resistance to *S. aureus*. Today all European meat producers and breeders must comply with European directives that regulate the limits and kinds of additives and antimicrobial growth promoters admitted for breeding.

In our study, we analysed the major meat species (beef, poultry and pork) for the occurrence of *S. aureus* and antimicrobial resistance profile of isolates. The results showed a large presence of multiresistant *S. aureus* in food, with risks of infections and possibility of transmission of resistances to other bacteria. Indeed, these bacteria can reach immunocompromised people through food or a human carrier, cause infections and lead to failure of antibiotic therapies. Our study also showed that all *S. aureus* isolated

from food were sensitive to methicillin, vancomycin and teicoplanin. This fact demonstrates that these kinds of resistance are, actually, limited to bacteria in hospitals, that represent important reservoirs for multiresistant bacteria.

Resistance to ampicillin was high in all isolates (42.86%, Fisher's exact test = 0.204) in accordance with natural resistance for β lactams of staphylococcus induced by exposure to penicillins (Fish, Piscitelli, & Danziger, 1995). Resistance to oxacillin was extremely high in poultry meat probably also because of the extended utilize of avoparcin used as an aid in the prevention of necrotic enteritis and as growth promoter in broiler chickens (Cormican, Erwin, & Jones, 1997).

It is to notice that 30.95% of all isolates were multiresistant (resistant at least to three antibiotics) and although we did not find any methicillin, or teicoplanin or vancomycin resistant we hypothesize that it is necessary an accurate and active control against antibiotic resistances in food particularly on meat, community and in nosocomial infections, in order to detect any expansion and/or increment of new antibiotic resistance that can be transferred to other bacteria. Moreover, the microbiological safety of food has to be guaranteed in order to prevent the transmission of pathogen or opportunist microorganisms to the consumer, according to the hazard analysis and critical control point (HACCP) principles in food chain.

Acknowledgement

This study was supported by Department of Public Health, Food Safety Laboratory, University of Florence, Italy.

References

- Acco, M., Ferreira, F. S., Henriques, J. A. P., & Tondo, E. C. (2003). Identification of multiple strain of *Staphylococcus aureus* colonizing nasal mucosa of food handlers. *Food Microbiology*, 20, 489–493.
- Bauer, A. W., Kirby, W. M., Sherris, J. C., & Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology*, 45, 493–496.
- Børgen, K., Sorum, M., Wasteson, Y., & Kruse, H. (2001). *VanA*-type vancomycin-resistant enterococci (VRE) remain prevalent in poultry carcasses 3 year after avoparcin was banned. *International Journal of Food Microbiology*, 64, 89–94.
- Bradley, S. F. (1999). Methicillin-resistant *Staphylococcus aureus*: long-term care concerns. *American Journal of Medicine*, 106, 2–10.
- Brun-Buisson, C. (1998). Methicillin-resistant *Staphylococcus aureus*: evolution and epidemiology, clinical impact, and prevention. *Pathologie biologique*, 46, 227–234.
- Centre for Disease Control and Prevention (1997). Update: *Staphylococcus aureus* with reduced susceptibility to vancomycin – United States, 1997. *Morbidity and Mortality Weekly Report*, 46, 813–815.
- Centre for Disease Control and Prevention (2002). *Staphylococcus aureus* resistant to Vancomycin – United States, 2002. *Morbidity and Mortality Weekly Report*, 51, No.26.
- Cormican, M. G., Erwin, M. E., & Jones, R. N. (1997). Avoparcin, a glycopeptide used in animal foods: antimicrobial spectrum and potency tested against human isolates from the United States. *Diagnostic Microbiology of Infectious Disease*, 29, 241–248.
- Courvalin, P. (1994). Transfer of antibiotic resistance genes between gram-positive and gram-negative bacteria. *Antimicrobial Agents and Chemotherapies*, 38(7), 1447–1451.
- Diekema, D. J., Pfaller, M. A., Schmitz, F. J., Smayevsky, J., Bell, J., Lones, R. N., & Beachand, M. the SENTRY Participants Group. (2001). Survey of infections due to *Staphylococcus* species: frequency of occurrence and antimicrobial susceptibility of isolates collected in the United States, Canada, Latin America, Europe, and Western Pacific Region for the SENTRY Antimicrobial Surveillance Program, 1997–1999. *Clinical and Infectious Diseases*, 32(Suppl2), 114–132.
- Fish, D. N., Piscitelli, S. C., & Danziger, L. H. (1995). Development of resistance during antimicrobial therapy: a review of antibiotic classes and patient characteristics in 173 studies. *Pharmacotherapy*, 15, 279–291.
- Grosso, M. D., Caprioli, A., Chinzari, P., Fontana, M. C., Pezzotti, G., Manfrin, A., Giannatale, E. D., Goffredo, E., & Pantosti, A. (2000). Detection and characterization of vancomycin-resistant enterococci in farm animals and raw meat products in Italy. *Microbiological Drug Resistance*, 6, 313–318.
- Hiramatsu, K., Hanaki, H., Ino, T., Yabuta, K., Oguri, T., & Tenover, F. C. (1997). Methicillin-resistant *Staphylococcus aureus* clinical strain with reduced vancomycin susceptibility. *Antimicrobial Agents and Chemotherapies*, 40, 135–136.
- Jones, T. F., Kellum, M. E., Porter, S. S., Bell, M., & Schaffner, W. (2002). An outbreak of community-acquired foodborne illness caused by methicillin-resistant *Staphylococcus aureus*. *Emerging Infectious Disease*, 8, 82–84.
- Joo Lee, H., Tae Suh, J., Sic Kim, Y., Lenz, W., Bierbaum, G., & Schaal, K. P. (2001). Typing and antimicrobial susceptibilities of methicillin resistant *Staphylococcus aureus* (MRSA) strains isolated in a hospital in Korea. *Emerging Infectious Disease*, 16, 381–385.
- Kitai, S., Shimizu, A., Kawano, J., Sato, E., Nakano, C., Uji, T., & Kitagawa, H. (2005). Characterization of methicillin-resistant *Staphylococcus aureus* isolated from retail raw chicken meat in Japan. *Journal of Veterinary and Medical Science*, 67, 107–110.
- Klein, G. (1999). Food as a potential vector for antibiotic resistance. *Berline Munch Tierarztl Wochenschr*, 112, 365–369.
- Kluytmans, J., Van Leeuwen, W., Goessen, W., Hollis, R., Messer, S., Herwaldt, L., Bruining, H., Heck, M., Rost, J., Van Leeuwen, N., Van Belkum, A., & Verbrugh, H. (1995). Food-initiated outbreak of methicillin-resistant *Staphylococcus aureus* analysed by pheno- and genotyping. *Journal of Clinical Microbiology*, 33(5), 1121–1128.
- Kruse, H., & Sorum, H. (1994). Transfer of multiple drug resistance plasmids between bacteria of diverse origins in natural microenvironments. *Applied and Environmental Microbiology*, 60(11), 4015–4021.
- Massachusetts Department of Public Health, Division of Epidemiology and Immunization. (2001). Vancomycin-Resistant Enterococci, Infection Control Guidelines for Long-Term Care Facilities 617, 983–6800.
- Mayrhofer, S., Paulsen, P., Smulders, F. J., & Hilbert, F. (2004). Antimicrobial resistance profile of five major food-borne pathogens isolated from beef, pork and poultry. *International Journal of Food Microbiology*, 97, 23–29.
- Monnet, D. L. (2000). Toward multinational antimicrobial resistance surveillance systems in Europe. *International Journal of Antimicrobial Agents*, 15, 91–101.
- Müller, T., Ulrich, A., Ott, E. M., & Müller, M. (2001). Identification of plant-associated enterococci. *Journal of Applied Microbiology*, 91, 268–278.
- Naimi, T. S., LeDell, K. H., Boxrud, D. J., Groom, A. V., Steward, C. D., Johnson, S. K., Besser, J. M., O'Boyle, C., Danila, R. N., Cheek, J. E., Osterholm, M. T., Moore, K. A., & Smith, K. E. (2001). Epidemiology and clonality of community-acquired methicillin-resistant *Staphylococcus aureus* in Minnesota, 1996–1998. *Clinical Infectious Disease*, 33, 990–996.
- NCCLS. (2000). Performance standards for antimicrobial disk susceptibility tests. *Approved standard M2 A7* (7th ed. Vol. 20. No. 1).

- Panlilio, A. L., Culver, D. H., & Gaynes, R. P. (1992). Methicillin-resistant *Staphylococcus aureus* in US hospitals, 1975–1991. *Infectious Control of Hospital Epidemiology*, 13, 582–586.
- Salmenlinna, S., & Vuopio-Varkila, J. (2001). Recognition of two groups of methicillin-resistant *Staphylococcus aureus* strains based on epidemiology, antimicrobial susceptibility, hypervariable-region type, and ribotype in Finland. *Journal of Clinical Microbiology*, 39, 2243–2247.
- Sorum, H., & L'Abèe-Lund, T. M. (2002). Antibiotic resistance in food-related bacteria – a result of interfering with the global web of bacterial genetics. *International Journal of Food Microbiology*, 78, 43–56.
- Teuber, M. (1999). Spread of antibiotic resistance with food-borne pathogens. *Cellular and Molecular Life Science*, 56, 755–763.
- Valsangiacomo, C., Dolina, M., Peduzzi, R., & Jäggl, M. (2000). Antimicrobial susceptibility of *Staphylococcus aureus* isolates from hospitalised patients and dairy food (fresh cheese): a survey over a decade in southern Switzerland. *Clinical Microbiology and Infectious Diseases*, 6, 393–394.
- Voss, A., Milatovic, D., Wallrauch-Schwarz, C., Rosdahl, V. T., & Braveny, I. (1994). Methicillin resistant *Staphylococcus aureus* in Europe. *European Journal of Clinical Microbiology Infectious Disease*, 13, 50–55.
- Yücel, N., Citak, S., & Onder, M. (2005). Prevalence and antibiotic resistance of *Listeria* species in meat products in Ankara, Turkey. *Food Microbiology*, 22, 241–245.
- Zanelli, G., Sansoni, A., Zanchi, A., Cresti, S., Pollini, S., Rossolini, G. M., & Cellesi, C. (2002). *Staphylococcus aureus* nasal carriage in the community: a survey from central Italy. *Epidemiology and Infections*, 129, 417–420.