



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

The will-o'-the-wisp of genetics--hunting for the azoospermia factor gene.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

The will-o'-the-wisp of genetics--hunting for the azoospermia factor gene / Tyler-Smith C; Krausz C. - In: THE NEW ENGLAND JOURNAL OF MEDICINE. - ISSN 0028-4793. - STAMPA. - 360:(2009), pp. 925-927.

Availability:

This version is available at: 2158/388409 since:

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

of the dietary fat to carbohydrate ratio of isoenergetically dense diets: effect on food intake in feeding men ad libitum. *Int J Obes Relat Metab Disord* 1996;20:651-60.

9. Davis EM, Clark JM, Carrese JA, Gary TL, Cooper LA. Racial and socioeconomic differences in the weight-loss experiences of obese women. *Am J Public Health* 2005;95:1539-43.

10. Franz MJ, VanWormer JJ, Crain AL, et al. Weight-loss outcomes: a systematic review and meta-analysis of weight-loss

clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc* 2007;107:1755-67.

11. Romon M, Lommez A, Tafflet M, et al. Downward trends in the prevalence of childhood overweight in the setting of 12-year school- and community-based programmes. *Public Health Nutr* 2008 December 23 (Epub ahead of print).

12. Westley H. Thin living. *BMJ* 2007;335:1236-7.

Copyright © 2009 Massachusetts Medical Society.

The Will-o'-the-Wisp of Genetics — Hunting for the Azoospermia Factor Gene

Chris Tyler-Smith, Ph.D., and Csilla Krausz, M.D., Ph.D.

It has been known for decades that the Y chromosome carries genes required for spermatogenesis and male fertility. But it has been frustratingly difficult to pin down the relevant genes, to such an extent that a finding reported by Luddi et al.¹ in this issue of the *Journal* — that the ubiquitin-specific peptidase 9, Y-linked gene (*USP9Y*) is not essential for normal spermatogenesis — represents a noteworthy advance in the field.


Approximately 15% of couples have difficulty conceiving; the causes are many and include genetic factors. The most common known genetic causes of spermatogenic failure are karyotype anomalies and Y-chromosome microdeletions. Karyotype analysis and screening for Yq deletions are now routine diagnostic tests that are able to help elucidate both the cause of spermatogenic perturbations and their clinical consequences. It is therefore of some interest to know which genes are essential to spermatogenesis; with this information, a more specific diagnosis might be possible, and the clinical management — including genetic counseling — of infertile couples would be greatly improved.

The search for these genes began in 1976, when Tiepolo and Zuffardi² reported cytogenetically visible microdeletions in Yq in six azoospermic men. Such deletions must be very large to be visible and are therefore likely to involve the absence of many genes (and unlikely to reveal the causal gene), but this observation provided a starting point for a classic approach to identifying the gene or genes responsible for this phenotype. Smaller deletions within the Yq interval would be sought using molecular methods, leading to the identification of one or a few candidate genes, and inactivating mutations would then be used to confirm which of the candidates was responsible. This strategy

has proved successful in the identification of the causal genes underlying countless single-gene disorders since its initial application to chronic granulomatous disease in 1986.³

At first, the application of this approach for Y-chromosomal spermatogenic abnormalities seemed promising. Molecular analysis did indeed identify smaller deletions within Yq that were specific to men with spermatogenic failure. These were classified, on the basis of chromosomal location, as three azoospermia factor (*AZF*) regions: *AZFa*, *AZFb*, and *AZFc*.⁴ *AZFb* and *AZFc* were later shown to contain large segments of duplicated sequence and, in fact, to overlap each other. Given the challenge of tracking down specific loci in chromosomal regions enriched in sequence duplications, it is perhaps understandable that the gene or genes responsible for the *AZFb*–*AZFc* spermatogenesis phenotype have still not been identified.

In contrast, the *AZFa* interval consists of regular single-copy DNA (i.e., DNA that does not contain a supranormal level of repeated sequence) that should not present unusual complications for homing in on the key gene. And in 1999, Sun et al.⁵ described what seemed to be a definitive result. They sequenced the two genes in the *AZFa* region (*USP9Y* and the DEAD [Asp–Glu–Ala–Asp] box polypeptide 3, Y-linked gene, *DDX3Y*) in 576 infertile men and discovered exactly the kind of mutation they were looking for, in a man with spermatogenic failure: a de novo deletion of 4 bp within the *USP9Y* gene that destroyed a splice-donor site, resulting in the skipping of the next exon and truncation of approximately 90% of the peptide sequence (Fig. 1). Thus, it appeared that the loss of *USP9Y* was responsible for the observed azoospermia and that one gene underlying spermatogenic failure had been successfully found.



But subsequent findings complicated this simple conclusion. The phenotype associated with the loss of *USP9Y* turned out to be variable, ranging from normospermia in chimpanzees and bonobos, in which the gene became inactivated millions of years ago,⁸ to moderate oligoasthenoteratozoospermia in humans,⁷ to azoospermia.⁵ The report by Luddi et al. now adds normospermia in humans to this list of phenotypes. These observations have bearing on two related questions: Is the inactivation of *USP9Y* or *DDX3Y* responsible for the *AZF_a* azoospermic phenotype? and What conclusions should we draw from the variation in the *USP9Y*-associated phenotype?

There are a number of reasons why the first, apparently simple, question has been so difficult to resolve. First, *AZF_a* deletions are themselves rare. In literature reports, they represent less than 5% of all *AZF* deletions.⁹ Second, *AZF_a* deletions leading to azoospermia obviously cannot be inherited, and so all are de novo mutations. They usually arise by the same mechanism: homologous recombination between two misaligned human endogenous retrovirus sequences (HERVs, elements of viral origin inserted into the genome and inherited as part of the chromosome) spaced about 790 kb apart (Fig. 1), encompassing both

genes.¹⁰ Mutations due to HERV misalignment are therefore not informative for distinguishing between the contributions of the two genes. Years of study have identified just a handful of patients who carry nonstandard deletions that affect only one of the genes and are thus informative (Fig. 1). Third, all of the confirmed single-gene mutations affect *USP9Y*; we have no information about the phenotype associated with the loss of *DDX3Y* only.

The study by Luddi et al. clearly shows that complete *USP9Y* deletion is compatible with normal spermatogenesis and fertility. But previous studies⁵⁻⁷ show, equally clearly, that its loss can disrupt spermatogenesis to varying degrees and that natural transmission (fertility) is possible when the phenotype is mild.⁷ So we have to conclude that the phenotype associated with the loss of *USP9Y* varies according to the genetic or other background of the carrier, although the relevant modifying factors remain unknown. This variation in phenotype contrasts sharply with the effect of losing both genes, which invariably results in the Sertoli-cell-only syndrome (characterized by the lack of germ cells) and, consequently, azoospermia. This consistent effect strongly suggests a key role of *DDX3Y* in spermatogenesis, a conclusion supported by expression of the gene in

premeiotic spermatogonia (USP9Y is expressed in postmeiotic spermatids).¹¹ We predict that the loss of *DDX3Y* alone is associated with a more severe testicular phenotype, most likely azoospermia.

What are the clinical implications of these conclusions? Screening for *USP9Y* deletion should not be performed routinely until there is a better understanding of the factors that modify the phenotype. Luddi et al. suggest that revision of the European Academy of Andrology–European Molecular Genetics Quality Network screening guidelines may be warranted to ensure that *DDX3Y*-only deletions are detected. The lack of confirmed deletions of this type after many years of worldwide screening in research laboratories⁹ means that the cost of such a step may currently outweigh the clinical benefit, but we can look forward to a time when the thousand-dollar genome will make such information, and much more, readily available. In the meantime, the article by Luddi et al. focuses our attention even more strongly on understanding the role of *DDX3Y*.

No potential conflict of interest relevant to this article was reported.

From the Wellcome Trust Sanger Institute, Hinxton, United Kingdom (C.T.-S.); and the Andrology Unit, Department of Clinical Physiopathology, University of Florence, Florence, Italy (C.K.).

1. Luddi A, Margollicci M, Gambera L, et al. Spermatogenesis in a man with complete deletion of *USP9Y*. *N Engl J Med* 2009; 360:881-5.
2. Tiepolo L, Zuffardi O. Localization of factors controlling spermatogenesis in the nonfluorescent portion of the human Y chromosome long arm. *Hum Genet* 1976;34:119-24.
3. Royer-Pokora B, Kunkel LM, Monaco AP, et al. Cloning the gene for an inherited human disorder — chronic granulomatous disease — on the basis of its chromosomal location. *Nature* 1986;322:32-8.
4. Vogt PH, Edelmann A, Kirsch S, et al. Human Y chromosome azoospermia factors (AZF) mapped to different subregions in Yq11. *Hum Mol Genet* 1996;5:933-43.
5. Sun C, Skaletsky H, Birren B, et al. An azoospermic man with a *de novo* point mutation in the Y-chromosomal gene *USP9Y*. *Nat Genet* 1999;23:429-32.
6. Brown GM, Furlong RA, Sargent CA, et al. Characterisation of the coding sequence and fine mapping of the human *DFFRY* gene and comparative expression analysis and mapping to the Sxb interval of the mouse Y chromosome of the *Dffry* gene. *Hum Mol Genet* 1998;7:97-107.
7. Krausz C, Degl'Innocenti S, Nuti F, et al. Natural transmission of *USP9Y* gene mutations: a new perspective on the role of AZFa genes in male fertility. *Hum Mol Genet* 2006;15:2673-81.
8. Perry GH, Tito RY, Verrelli BC. The evolutionary history of human and chimpanzee Y-chromosome gene loss. *Mol Biol Evol* 2007;24:853-9.
9. Krausz C, Degl'Innocenti S. Y chromosome and male infertility: update, 2006. *Front Biosci* 2006;11:3049-61.
10. Kamp C, Hirschmann P, Voss H, Huellen K, Vogt PH. Two long homologous retroviral sequence blocks in proximal Yq11 cause AZFa microdeletions as a result of intrachromosomal recombination events. *Hum Mol Genet* 2000;9:2563-72.
11. Vogt PH, Falcao CL, Hanstein R, Zimmer J. The AZF proteins. *Int J Androl* 2008;31:383-94.

Copyright © 2009 Massachusetts Medical Society.