



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

S38G single-nucleotide polymorphism at the KCNE1 locus is associated with heart failure

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

S38G single-nucleotide polymorphism at the KCNE1 locus is associated with heart failure / C.Fatini; E.Sticchi; R.Marcucci; V.Verdiani; C.Nozzoli; C.Vassallo; M.Emdin; R.Abbate; G.F.Gensini. - In: HEART RHYTHM. - ISSN 1547-5271. - STAMPA. - 7:(2010), pp. 363-367. [10.1016/j.hrthm.2009.11.032]

Availability:

This version is available at: 2158/394747 since: 2020-05-13T12:40:30Z

Published version:

DOI: 10.1016/j.hrthm.2009.11.032

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze
(<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

S38G single-nucleotide polymorphism at the KCNE1 locus is associated with heart failure

Cinzia Fatini, MD, PhD,* Elena Sticchi, BS,* Rossella Marcucci, MD, PhD,* Valerio Verdiani, MD,† Carlo Nozzoli, MD,† Cristina Vassallo, BS,‡ Michele Emdin, MD, PhD,‡ Rosanna Abbate, MD,* Gian Franco Gensini, MD*†§

From the *Department of Medical and Surgical Critical Care, University of Florence, and Thrombosis Centre, Azienda Ospedaliero-Universitaria Careggi, Center for the Study at the Molecular and Clinical Level of Chronic, Degenerative, and Neoplastic Diseases to Develop Novel Therapies, University of Florence, and †Department of Heart and Vessels, Azienda Ospedaliero-Universitaria Careggi, Florence, ‡Department of Cardiovascular Medicine, G. Monasterio Foundation, National Research Council (CNR-Regione Toscana) Pisa, and §Fondazione Don Carlo Gnocchi Organizzazione Non Lucrativa di Utilità Sociale (ONLUS), Centro S. Maria degli Ulivi-Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS), Florence.

BACKGROUND Prolongation of the action potential duration, whose major determinants are the delayed-rectifier potassium currents, is a hallmark of failing ventricular myocardium. Genetic variants in the KCNE1 gene, encoding for the β -subunit (minK) of a slowly activated cardiac potassium channel (I_{Ks}), may impair myocardial repolarization. Experimental data demonstrated a higher KCNE1 expression in heart failure (HF).

OBJECTIVE The purpose of this study was to investigate the association between a KCNE1 S38G single-nucleotide polymorphism (SNP) and HF.

METHODS We genotyped 197 out of 323 previously investigated patients and 352 healthy controls comparable for age and sex. This study was replicated in 186 HF patients and in 200 healthy subjects comparable for age and sex and recruited from the Department of Cardiovascular Medicine of the National Research Council, Pisa, Italy.

RESULTS A significant difference in genotype distribution and allele frequency between patients and controls was observed for the KCNE1 S38G SNP ($P = .002$ and $P = .0008$, respectively). The KCNE1 38G variant was associated with a significant predisposition to HF under a dominant (odds ratio [OR] = 2.22 [1.23–3.28]; $P = .008$) and additive (OR = 2.13 [1.09–4.15]; $P = .03$) model, after adjustment for age, sex, and traditional cardiovascular risk fac-

tors. No difference in genotype distribution and allele frequency for the KCNE1 S38G SNP according to functional New York Heart Association class was found ($P = .4$ and $P = .3$, respectively). In the HF replication study, the KCNE1 38G allele frequency was significantly higher in comparison with that observed in the control population (38G = 0.59 vs. 0.49; $P = .004$). The 38G allele was associated with HF predisposition under the recessive (OR [95% confidence interval (CI)] = 2.49 [1.45–4.29]; $P = .001$) and additive models (OR [95% CI] = 2.63 [1.29–5.35]; $P = .008$), after adjustment for traditional risk factors.

CONCLUSION KCNE1 S38G SNP is associated with HF predisposition in two study populations. Nevertheless, further studies performed in larger populations and aimed to better define the role of this locus are required.

KEYWORDS Heart failure; KCNE1 gene; S38G SNP

ABBREVIATIONS **APD** = action potential duration; **CAD** = coronary artery disease; **EF** = ejection fraction; **HC** = hypertensive cardiomyopathy; **HF** = heart failure; **HC** = hypertensive cardiomyopathy; **LVEDD** = left ventricular end-diastolic diameter; **LVESD** = left ventricular end-systolic diameter; **NYHA** = New York Heart Association; **SNP** = single-nucleotide polymorphism

(Heart Rhythm 2010;7:363–367) © 2010 Heart Rhythm Society. All rights reserved.

Introduction

Heart failure (HF) results from any structural or functional cardiac disorder that impairs heart function. Ventricular arrhythmias are a common cause of sudden death in HF patients,¹ nevertheless, the underlying mechanism of arrhythmia is poorly understood. Prolongation of the action

potential duration (APD) is a hallmark of a failing ventricular myocardium,² in particular in the setting of an increased dispersion of repolarization.³ The rapid (I_{Kr}) and slow (I_{Ks}) components of the delayed-rectifier potassium current are the major determinants of APD.⁴ The slow component of the potassium current is related to minK and KCNQ1 channel subunits function. The minK protein is encoded by the KCNE1 gene located on chromosome 21q22.12, and mutations in this gene may impair the slowly activating cardiac delayed-rectifier (I_{Ks}) current,⁵ thus contributing to modulating myocardial repolarization. An experimental study performed on human endomyocardial

Address reprint requests and correspondence: Cinzia Fatini, M.D., Department of Medical and Surgical Critical Care, Thrombosis Centre, University of Florence, Viale Morgagni 85, 50134 Florence, Italy. E-mail address: cinzafatini@hotmail.com. (Received May 19, 2009; accepted November 29, 2009.)

biopsy samples demonstrated a higher *KCNE1* gene expression in HF patients.⁶

The *KCNE1* locus has been investigated in nonvalvular atrial fibrillation patients,^{7,8} but to the best of our knowledge no data concerning its influence on HF susceptibility are available. Accordingly, in the present study we investigated the association between the *KCNE1* S38G single-nucleotide polymorphism (SNP) and HF patients, and we validated data in a different HF population.

Methods

Study populations

Florence HF patients

One hundred ninety-seven (156 males/39 females, age 73 [20–95] years) HF Caucasian patients from Tuscany who were previously investigated⁹ were genotyped for the *KCNE1* S38G polymorphism. Inclusion criteria were diagnosis of HF according to the Framingham criteria¹⁰ and the presence of a left ventricular dysfunction defined as an ejection fraction (EF) <40% by echocardiography performed on hospital admission. The cause of HF was determined in each patient by clinical assessment and echocardiography. The etiology of HF was classified as coronary artery disease (CAD; n = 103), hypertensive cardiomyopathy (HC; n = 2), or idiopathic dilated cardiomyopathy (IDCM; n = 92) on the basis of disease history and coronary angiographic evaluation. Owing to the role of the *KCNE1* gene in influencing the susceptibility to atrial fibrillation,^{7,8} in this study atrial fibrillation patients (n = 39) were excluded. Moreover, the exclusion criterion for patients was the presence of familial cardiomyopathy (n = 1), as gene mutations involved in modulating arrhythmogenic disorders may be related to the pathogenesis of ventricular cardiomyopathies, in particular polymorphisms in the *KCNE1* gene.¹¹

Pisa HF patients

One hundred eighty-six HF Caucasian patients (134 males and 52 females, age 72 [24–97] years) from Tuscany, enrolled from the Department of Cardiovascular Medicine of the National Research Council, Pisa, were investigated to perform the replication study. Inclusion criteria were diagnosis of HF according to the Framingham criteria¹⁰ and the presence of a left ventricular dysfunction defined as an EF <40% by echocardiography performed on hospital admission. The cause of HF was determined in each patient by clinical assessment and echocardiography. The etiology of HF was classified as CAD (n = 99), HC (n = 14), or IDCM (n = 73) on the basis of disease history and coronary angiographic evaluation. Exclusion criteria were the presence of familial cardiomyopathy (n = 0) and atrial fibrillation (n = 46).

The subjects were considered to have hypertension according to the guidelines of the European Society of Hypertension/European Society of Cardiology¹² or if they were taking antihypertensive drugs. Dyslipidemia was defined according to the Third Report of the National Choles-

terol Education Program,¹³ and diabetes was defined in agreement with the American Diabetes Association.¹⁴ All subjects gave informed consent, and the study complies with the Declaration of Helsinki and was approved by the local ethics committee.

Control subjects (Florence and Pisa)

Three hundred fifty-two Caucasian control subjects, comparable for age and gender, were invited to participate in the study and consisted of partners or friends of patients and subjects from a population study, "Progetto Nutrizione per la Salute e la Prevenzione di Malattia," conducted between 2002 and 2004 and aimed to evaluate the lifestyle and dietary habits of clinically healthy persons living in Florence, Italy. Exclusion criteria for the controls were personal and family history of cardiovascular disease. A detailed interview addressed to personal and familial history was performed in the framework of a physical examination by expert physicians to identify symptom-free subjects and to exclude those who were suspected of having any form of vascular disease.

Two hundred Caucasian control subjects enrolled from healthy persons living in Pisa were used for the replication study. Exclusion criteria for the controls were personal and family history of cardiovascular disease. A detailed interview addressed to personal and familial history was performed in the framework of a physical examination by expert physicians to identify symptom-free subjects and to exclude those who were suspected of having any form of vascular disease.

Detection of *KCNE1* S38G polymorphism

The S38G (112A>G) *KCNE1* gene SNP was analyzed through polymerase chain reaction-Restriction Fragment Length Polymorphism (RFLP) analysis as described elsewhere.⁸

Statistical analysis

Statistical analysis was performed by using the SPSS (Statistical Package for Social Sciences, Chicago) software for Windows (version 11.5). The χ^2 -test was used to test for deviation of genotype distribution from Hardy-Weinberg equilibrium. The association between *KCNE1* SNP and HF was assessed using logistic regression analysis under a dominant, recessive, and additive genetic model.

The dominant genetic model compares individuals with one or more polymorphic alleles with a baseline group with no polymorphic alleles (e.g., *KCNE1* 38GG + GS vs. SS). The recessive genetic model compares the 38GG genotype with the combined GS + SS genotypes, which form the baseline group. The additive genetic model assumes that there is a linear gradient in risk between the 38GG, 38GS, and 38SS genotypes (38SS genotype baseline). This is equivalent to a comparison of the 38G allele versus the 38S allele (baseline). $P < .05$ was considered statistically significant (two-sided P -value).

We performed a replication study in different Caucasian HF and control populations to further strengthen the claim of predisposition for the *KCNE1* S38G polymorphism in HF

Table 1 Demographic and clinical characteristics of study populations (Florence HF and controls)

	Florence HF (n = 195)	Florence controls (n = 352)
Age, years, median (range)	73 (20–95)	72 (36–93)
Males/females	156/39	259/94
Hypertension	103 (52.8)	116 (33.0)
Dyslipidemia	58 (29.4)	72 (20.5)
Diabetes	58 (29.4)	25 (7.1)
Smoking habit	92 (46.7)	64 (18.2)
CAD	103 (52.2)	—
IDCM	92 (46.7)	—
HC	2 (1.01)	—
LVEF <40%	149 (75.6)	—
Chronic renal failure	33 (16.7)	—
NT-proBNP, pg/mL, median (range)	3564 (52–37,940)	—
NYHA class:		
II	51 (25.8)	—
III	73 (37.1)	—
IV	73 (37.1)	—
KCNE1 S38G genotype distribution ^a and allele frequency ^b :		
KCNE1 38GG	71 (36.0)	98 (27.8)
KCNE138SG	99 (50.3)	161 (45.7)
KCNE138SS	27 (13.7)	93 (26.4)
KCNE1 38G	0.61	0.51

Note: Values are n (%) unless otherwise indicated. LVEF: left ventricular ejection fraction.

^aP = .002.

^bP = .0008.

patients. The same healthy control population was used in this replication study.

Results

The demographic and clinical characteristics of the study populations are reported in **Tables 1** and **2**. The prevalence of traditional cardiovascular risk factors was significantly higher in Florence HF patients in comparison with that observed in the control group ($P <.05$). In the Pisa HF patients, we observed a significant difference in the prevalence of all traditional risk factors between patients and controls ($P <.05$).

No deviation from the expected population genotype proportions predicted by the Hardy-Weinberg equilibrium was detected at the *KCNE1* locus. A significant difference in genotype distribution and allele frequency between patients and controls was observed for the *KCNE1* S38G SNP in both the populations investigated (**Tables 1** and **2**).

Florence HF patients

The *KCNE1* S38G genotype distribution and allele frequency were investigated in relation to a HF clinical subset (**Table 3**). A significantly higher prevalence of the *KCNE1* 38G variant was observed in HF hypertensives, diabetics, smokers, and HF patients with a history of CAD than in controls (**Table 3**). The *KCNE1* 38G variant was associated with a significant predisposition to HF under all three genetic models of inheritance (dominant, recessive, and additive; **Table 4**). After adjustment for age, sex, and traditional

cardiovascular risk factors, the *KCNE1* 38G allele significantly and independently influenced the susceptibility to HF under a dominant and additive but not recessive model (**Table 4**).

No difference in genotype distribution and allele frequency for the *KCNE1* S38G SNP according to functional New York Heart Association (NYHA) class was found ($\chi^2 = 5.3$, $P = .4$ and $\chi^2 = 3.4$, $P = .3$, respectively).

The role of *KCNE1* S38G SNP has been analyzed according to parameters related to the severity of HF, such as EF, left ventricular end-diastolic diameter (LVEDD), and left ventricular end-systolic diameter (LVESD); a relationship, even not significant, between the *KCNE1* 38G allele and altered hemodynamic parameters (EF 38G = 0.62, LVEDD 38G = 0.64, and LVESD 38G = 0.63, respectively) was observed.

Pisa HF patients

To validate our results, we have replicated this study in different Caucasians HF and control populations enrolled from the Department of Cardiovascular Medicine of the National Research Council, Pisa. Data from the replication study demonstrated similar findings concerning the role of the *KCNE1* S38G polymorphism in modulating susceptibility to HF. The *KCNE1* 38G variant significantly influenced the predisposition to HF under the recessive and additive but not dominant model of inheritance after adjustment for age, sex, and cardiovascular risk factors (**Table 5**).

Table 2 Demographic and clinical characteristics of study populations (Pisa HF and controls)

	Pisa HF (n = 186)	Pisa controls (n = 200)
Age, years, median (range)	72 (24–97)	71 (36–96)
Males/females	134/52	129/71
Hypertension	79 (42.5)	65 (32.5)
Dyslipidemia	69 (37.0)	45 (22.5)
Diabetes	47 (25.3)	16 (8.0)
Smoking habit	72 (38.7)	35 (17.5)
CAD	99 (53)	—
IDCM	73 (39)	—
HC	14 (8)	—
LVEF <40%	143 (77)	—
Chronic renal failure	73 (39)	—
NT-proBNP, pg/mL, median (range)	1211 (10–38,200)	—
NYHA class:		
II	115 (62.0)	—
III	52 (28.0)	—
IV	19 (10)	—
KCNE1 S38G genotype distribution ^a and allele frequency ^b :		
KCNE1 38GG	65 (34.9)	47 (23.5)
KCNE138SG	90 (48.4)	101 (50.5)
KCNE138SS	31 (16.7)	52 (26.0)
KCNE1 38G	0.59	0.49

Note: Values are n (%) unless otherwise indicated. LVEF = left ventricular ejection fraction.

^aP = .01.

^bP = .004.

Table 3 KCNE1 S38G polymorphism genotype distribution and allele frequency according to subsets of Florence HF patients

	Hypertension, n = 103	Dyslipidemia, n = 58	Diabetes, n = 58	Smoking habit, n = 92	CAD, n = 103	IDCM, n = 92	LVEF<40%, n = 149	Chronic renal failure, n = 33
KCNE1 38GG	34 (33.0)	21 (36.2)	21 (36.2)	38 (41.3)	35 (34.0)	34 (37.0)	56 (37.6)	10 (30.3)
KCNE1 38SG	57 (55.3)	28 (48.3)	30 (51.7)	41 (44.6)	53 (51.5)	39 (42.4)	73 (49.0)	14 (42.4)
KCNE1 38SS	12 (11.7)	9 (15.5)	7 (12.1)	13 (14.1)	15 (14.6)	19 (20.6)	20 (13.4)	9 (27.3)
KCNE1 38G	0.61	0.60	0.62	0.64	0.60	0.58	0.62	0.51
Allele frequency (P-value) ^a	.01	.05	.02	.002	.02	.07	.001	.9
Genotype distribution (P-value) ^a	.007	.2	.05	.01	.04	.2	.003	.9

Note: Values are n (%) unless otherwise indicated. LVEF = left ventricular ejection fraction.

^aχ²-test: allele frequency and genotype distribution vs. controls.

Discussion

This is the first study in which the effect of the *KCNE1* gene has been evaluated as a potential predisposing factor to HF. The results of the present study provide evidence of its role in predisposing to HF apart from cardiovascular risk factors but not in modulating the severity of the disease. These findings have been replicated in a separate HF population with the same genetic background and recruited from the Department of Cardiovascular Medicine of Pisa.

Data from the replication study confirmed the role of the *KCNE1* 38G allele in influencing the predisposition to the disease, independently of the presence of traditional cardiovascular risk factors. The two study populations investigated were comparable for sex and age but not for the percentage of cardiovascular risk factors. Accordingly, these findings stressed the role of the *KCNE1* gene in modulating HF susceptibility *per se*, despite the presence or absence of traditional cardiovascular risk factors.

Little information is available on the genetic modulation of delayed-rectifier potassium currents in humans.^{15,6} Data from experimental studies performed in rabbit and human hearts demonstrated a prolongation of the APD in left ventricular myocytes isolated from failing rabbit ventricles¹⁶ and that *KCNE1* gene expression was significantly higher in myocytes from HF patients in comparison with controls, thus contributing to the prolongation of the QT interval

through reducing the net outward current during the plateau of the action potential.⁶

The *KCNE1* gene encodes for a single transmembrane domain protein (minK), which represents the β-subunit of a potassium channel able to conduct the slow component (I_{Ks}) of the delayed-rectifier current, which is relevant in the repolarization of cardiac myocytes. The minK protein cytoplasmatic domain is also able to interact with regulatory proteins, such as the sarcomeric Z-line component T-cap (telethonin), which is involved in cardiac muscle contraction. Indeed, it has been reported that the biomechanical stress during myocardium contraction-relaxation cycles may contribute to determining an altered ion channel localization, thus possibly influencing the ionic properties of cardiac tissue.^{17,18}

An SNP in the *KCNE1* gene (A to G at position 112), leading to a glycine substitution for serine at amino acid position 38, has been identified.¹⁹ The functional significance of the S38G polymorphism in the *KCNE1* gene remains unclear. An experimental study documented that the minK 38G isoform was associated with reduced I_{Ks} current, thus resulting in increased APD,²⁰ whereas data from the KORA study²¹ did not evidence its modifying effect on the QT interval. Indeed, the *KCNE1* S38G SNP might not have a causal effect on the I_{Ks} current, but it could be a marker in linkage disequilibrium with another functional variant. An association between the *KCNE1* S38G SNP and atrial fi-

Table 4 Univariate and multivariate analysis for cardiovascular risk factors and *KCNE1* S38G polymorphism according to dominant, recessive and additive genetic models (Florence HF patients and controls)

Univariate analysis		Odds ratio (95% CI)		P	
Age		0.99 (0.97–1.004)			.2
Sex		0.69 (0.45–1.05)			.08
Hypertension		2.28 (1.59–3.26)			<.0001
Smoking habit		2.12 (1.41–3.18)			<.0001
Diabetes		5.54 (3.23–9.22)			<.0001
Dyslipidemia		1.65 (1.10–2.46)			.01
Genetic polymorphism		Dominant model		Recessive model	
		Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Univariate analysis, <i>KCNE1</i> S38G		2.40 (1.48–3.89)	<.0001	1.45 (1.00–2.12)	.049
Multivariate analysis, ^a <i>KCNE1</i> S38G		2.22 (1.23–3.98)	.008	1.13 (0.72–1.78)	.6
		Additive model			
		Odds ratio (95% CI)	P		
Univariate analysis, <i>KCNE1</i> S38G		2.62 (1.53–4.89)		<.0001	
Multivariate analysis, ^a <i>KCNE1</i> S38G		2.13 (1.09–4.15)		.03	

Note: CI: confidence interval.

^aAdjusted for age, sex, and cardiovascular risk factors.

Table 5 Univariate and multivariate analysis for cardiovascular risk factors and *KCNE1* S38G polymorphism according to dominant, recessive and additive genetic models (Pisa HF patients and controls)

Univariate analysis	Odds ratio (95% CI)		P			
Age	0.99 (0.97–1.01)		.2			
Sex	0.70 (0.46–1.09)		.1			
Hypertension	1.53 (1.01–2.32)		.04			
Smoking habit	2.98 (1.86–4.76)		<.0001			
Diabetes	3.89 (2.12–7.15)		<.0001			
Dyslipidemia	2.03 (1.30–13.7)		.002			
Genetic polymorphism	Dominant model	Recessive model	Additive model			
	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Univariate analysis, <i>KCNE1</i> S38G	1.76 (1.07–2.89)	.03	1.75 (1.12–2.73)	.01	2.32 (1.29–4.15)	.005
Multivariate analysis, ^a <i>KCNE1</i> S38G	1.54 (0.84–2.81)	.2	2.49 (1.45–4.29)	.001	2.63 (1.29–5.35)	.008

Note: CI: confidence interval.

^aAdjusted for age, sex, and cardiovascular risk factors.

brillation has been investigated,^{7,8} but no data are available on HF susceptibility. We previously demonstrated that the *KCNE1* gene was associated with atrial fibrillation, possibly by modulating the ionic properties of the atria and in turn their irregular activation.⁸ The present study, in which patients with atrial fibrillation have been excluded, provides evidence for a role of the same gene in modulating HF predisposition. Therefore, we could hypothesize that the *KCNE1* gene, through the modulation of I_{Ks} current, may affect alterations in ionic properties in both atrial and ventricular myocytes, thus determining two different clinical phenotypes.

A limitation of this study is the lack of information concerning electrocardiogram data, and in particular QTc values, as the *KCNE1* gene has been demonstrated to be associated with QT interval prolongation in HF.⁶

Our results failed to demonstrate the influence of the *KCNE1* locus on the severity of the disease, as no difference in *KCNE1* 38G allele frequency has been observed according to NYHA class. This datum possibly strengthens the observation that the *KCNE1* locus could be involved in the electrical damage in HF but not in its progression.

Findings from the present study, which have been replicated in a different population, are intriguing for the novelty of the data. Nevertheless, owing to the limited sample size, further studies performed in larger populations are required to better define the role of the *KCNE1* S38G SNP in influencing not only HF susceptibility but also the severity of the disease.

Conclusion

This preliminary study shows an association between the *KCNE1* gene, which is involved in the modulation of I_{Ks} delayed-rectifier potassium current, and HF, thus providing new highlights for investigating the contribution of cardiac potassium channels.

References

- Stevenson WG, Stevenson LW, Middlekauff HR, et al. Sudden death prevention in patients with advanced ventricular dysfunction. *Circulation* 1993;88:2953–2961.
- Beuckelmann DJ, Nähbauer M, Erdmann E. Alterations of K⁺ currents in isolated human ventricular myocytes from patients with terminal heart failure. *Circ Res* 1993;73:379–385.
- Akar FG, Spragg DD, Tunin RS, et al. Mechanisms underlying conduction slowing and arrhythmogenesis in nonischemic dilated cardiomyopathy. *Circ Res* 2004;95:717–725.
- Sanguineti MC, Jurkiewicz NK. Two components of cardiac delayed rectifier K⁺ current. Differential sensitivity to block by class III antiarrhythmic agents. *J Gen Physiol* 1990;96:195–215.
- Splawski I, Tristani-Firouzi M, Lehmann MH, et al. Mutations in the hminK gene cause long QT syndrome and suppress IKs function. *Nat Genet* 1997;17:338–340.
- Watanabe E, Yasui K, Kamiya K, et al. Upregulation of KCNE1 induces QT interval prolongation in patients with chronic heart failure. *Circ J* 2007;71:471–478.
- Lai LP, Su MJ, Yeh HM, et al. Association of the human minK gene 38G allele with atrial fibrillation: evidence of possible genetic control on the pathogenesis of atrial fibrillation. *Am Heart J* 2002;144:485–490.
- Fatini C, Sticchi E, Genuardi M, et al. Analysis of minK and eNOS genes as candidate loci for predisposition to non-valvular atrial fibrillation. *Eur Heart J* 2006;27:1712–1718.
- Fatini C, Sticchi E, Marcucci R, et al. ACE insertion/deletion, but not -240A>T polymorphism, modulates the severity in heart failure. *J Investig Med* 2008;56:1004–1010.
- McKee PA, Castelli WP, McNamara PM, et al. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971;285:1441–1446.
- Corrado D, Bassi C, Thiene G. Is it time to include ion channel diseases among cardiomyopathies? *J Electrocardiol* 2005;38:81–87.
- Practice guidelines for primary care physicians: 2003 ESH/ESC Hypertension guidelines. ESH/ESC hypertension guidelines committee. *Hypertension* 2003;21:1779–1786.
- Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adult (Adult Treatment Panel III). Final report. *Circulation* 2002;106:3143–3421.
- Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care* 2003;26:S5–S20.
- Lundquist AL, Manderfield LJ, Vanoye CG, et al. Expression of multiple KCNE genes in human heart may enable variable modulation of I(Ks). *J Mol Cell Cardiol* 2005;38:277–287.
- Rose J, Armoundas AA, Tian Y, et al. Molecular correlates of altered expression of potassium currents in failing rabbit myocardium. *Am J Physiol Heart Circ Physiol* 2005;288:H2077–H2087.
- Furukawa T, Ono Y, Tsuchiya H, et al. Specific interaction of the potassium channel beta-subunit minK with the sarcomeric protein T-cap suggests a T-tubule-myofibril linking system. *J Mol Biol* 2001;313:775–784.
- Vatta M, Faulkner G. Cytoskeletal basis of ion channel function in cardiac muscle. *Future Cardiol* 2006;2:467–476.
- Lai LP, Deng CL, Moss AJ, et al. Polymorphism of the gene encoding a human minimal potassium ion channel (minK). *Gene* 1994;151:339–340.
- Ehrlich JR, Zicha S, Coutu P, et al. Atrial fibrillation-associated minK38G/S polymorphism modulates delayed rectifier current and membrane localization. *Cardiovasc Res* 2005;67:520–528.
- Akyol M, Jalilzadeh S, Sinnet MF, et al. The common non-synonymous variant G38S of the *KCNE1*-(minK)-gene is not associated to QT interval in Central European Caucasians: results from the KORA study. *Eur Heart J* 2007;28:305–309.