



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Aggressive approach and outcome in patients presenting atrial fibrillation and hypertension

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Aggressive approach and outcome in patients presenting atrial fibrillation and hypertension / A.Conti; E.Canuti; Y.Mariannini; M.Zanobetti; F.Innocenti; B.Paladini; G.Pepe; L.Padeletti; G.F.Gensini. - In: INTERNATIONAL JOURNAL OF CARDIOLOGY. - ISSN 0167-5273. - STAMPA. - ...:(2011), pp. 0-0. [10.1016/j.ijcard.2011.09.074]

Availability:

The webpage <https://hdl.handle.net/2158/592657> of the repository was last updated on

Published version:

DOI: 10.1016/j.ijcard.2011.09.074

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:


La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)



Contents lists available at SciVerse ScienceDirect

International Journal of Cardiology

journal homepage: www.elsevier.com/locate/ijcard

Aggressive approach and outcome in patients presenting atrial fibrillation and hypertension

Alberto Conti ^{a,*}, Erica Canuti ^a, Yuri Mariannini ^a, Maurizio Zanobetti ^a, Francesca Innocenti ^a, Barbara Paladini ^b, Giuseppe Pepe ^b, Luigi Padeletti ^c, Gian Franco Gensini ^c

^a Emergency Medicine, Department of Medical and Surgical Critical Care, University of Florence and Careggi University Hospital, Florence, Italy

^b Emergency Department, Careggi University Hospital, Florence, Italy

^c Department of Heart and Vessel Disease, University of Florence and Careggi University Hospital, Florence, Italy

ARTICLE INFO

Article history:

Received 1 August 2011

Accepted 17 September 2011

Available online xxx

Keywords:

Hypertension

Atrial fibrillation

Emergency medicine

Pharmacological treatment

Electrical treatment

Intensive approach

ABSTRACT

Aim: Aggressive approach in patients presenting atrial fibrillation (AF) and hypertension could result in improving rhythm control and reducing admission.

Methods: Out of 3475 patients presenting AF, those with hypertension ($n = 1739$, 52%) underwent standard ($n = 591$, group 1, years 2004–2005) or aggressive pharmacological and electrical approach ($n = 1148$, group 2, years 2006–2009). Overall, in 1071 patients AF duration was less than 48 h. Primary endpoint was rhythm conversion; secondary endpoints were modalities of rhythm conversion and reduction of admissions.

Results: At univariate and multivariate analyses, AF lasting less than 48 h, absence of comorbidities and younger age were independent predictors of sinus rhythm; conversely, lack of sinus rhythm, older age, AF lasting more than 48 h and comorbidities were independent predictors of hospitalization. Overall, 55% of patients achieved sinus rhythm in group 1 versus 62% in group 2 ($p = 0.018$). Interestingly, in patients with AF lasting less than 48 h, 89% achieved sinus rhythm, more likely by class IC than by class III antiarrhythmic drugs ($p < 0.001$). Overall reduction of admission accounts for 60%; 50% of patients need admission in group 1 versus 29% in group 2 ($p < 0.001$).

Conclusions: Aggressive pharmacological and electrical approach in patients presenting AF and hypertension significantly improved rhythm conversion overall up to 62%. Patients with AF lasting less than 48 h eventually achieved sinus rhythm up to 89%, mostly by class IC antiarrhythmic drugs. Admissions eventually reduced up to 60%.

© 2011 Published by Elsevier Ireland Ltd.

1. Introduction

In emergency settings, effective therapeutic management of atrial fibrillation (AF) is not yet clear, because of lack of understanding of several pathophysiological aspects of arrhythmia and because the existing possibility of the underlying presence of structural heart disease to which several pharmacological side effects are related. Moreover, to date there has been no convincing evidence that antiarrhythmic or antihypertensive therapy and 'rhythm control' is better than the rate control approach [1–3]. However, recent data have shown therapeutic success might be associated with rhythm-control and clinical outcomes driven by hospitalizations for arrhythmia and other comorbidities [4]. Atrial fibrillation is established to share strong epidemiological associations with a large panel of risk factors including advanced age, male sex, diabetes mellitus and especially hypertension or with several diseases including heart failure,

coronary artery disease, valvular heart disease and thyroid disease [5,6]. Atrial fibrillation could also be the effect of an underlying 'masked' disorder associated with metabolic syndrome and its components, sleep apnea, inflammation, and aspects of lifestyle as alcohol consumption and physical activity. Eventually, in some patients with AF, no underlying pathology is present and the etiology remains unknown [7]. Detection of the underlying disorder may result in implementation of effective treatment that could improve outcomes, but the presence of structural heart disease as in patients with hypertension should not to be considered as absolute contraindication to aggressive approach and rhythm control. The aim of the present study was to evaluate if aggressive approach in patients with AF and hypertension without moderate to severe structural heart disease could result in improving rhythm control and eventually reducing admission in a large series of patients presenting to the Emergency Department [8].

2. Materials and methods

The study enrolled AF patients presenting to the Emergency Department of the tertiary care teaching hospital, in Florence, Italy. Facilities for triage of patients with suspected dysrhythmia included an Observation Unit with 6 monitor equipped beds

* Corresponding author at: Emergency Medicine, Department of Critical Care Medicine and Surgery, Careggi University Hospital, Florence, Italy. Tel.: +39 338 7472294; fax: +39 055 7947934.

E-mail address: aaaconti@hotmail.com (A. Conti).

and dedicated personnel (resident, faculty, and cardiologist on call). Coded diagnoses are entered into an electronic medical chart, allowing easy selection of all patients with a diagnosis of interest. Data were collected from January 2004 to December 2010.

2.1. Patient selection

Inclusion criteria were the presence of AF documented by ECG. Consecutive patients presenting to the Emergency Department complaining of palpitations or malaise were considered and those with AF documented by ECG were evaluated and enrolled in the study. Exclusion criteria were age <18 years, Killip Class ≥ 2 , New York Heart Association functional class $\geq III$, clinically detectable ongoing complications like acute coronary syndrome, stroke and pulmonary or peripheral embolism, and hemodynamic instability. Patients with moderate to severe structural heart disease were excluded from the study. All patients gave their consent for study participation. The study was conducted according to good clinical practice and the Helsinki Principles.

2.2. Definition of subtype of AF, structural heart disease and comorbidities

Paroxysmal AF was defined by electrocardiographic evidence of AF followed by subsequent ECG showing sinus rhythm or clinical documentation by a physician of paroxysmal AF; persistent AF was defined when lasting longer than 7 days or when terminated by therapeutic treatment; permanent AF (or chronic AF) was defined in the presence of serial electrocardiograms showing only AF and no interim evidence of sinus rhythm; solitary AF was defined by AF in the absence of overt cardiovascular disease or precipitating illness [9]. Systemic hypertension was defined by need for antihypertensive therapy or the presence of systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg. Coronary artery disease was defined by angiographic findings of stenosis $>50\%$ in any of the 3 main arterial distributions, effort angina, or history of myocardial infarction. Moderate to severe valvular heart disease was considered in the presence of substantial murmur on physical heart examination, or by history of documented prior valve repair/replacement or by echocardiography confirmation. Congestive heart failure was defined by a physician's diagnosis or by need for treatment with dedicated drug treatment. Carotid artery disease referred to the presence of stenosis $>50\%$ in the carotid arteries based on neurovascular imaging or previous intervention. Stroke included development of any type of stroke, as defined by clinical documentation of the diagnosis with or without confirmatory findings on imaging studies. Diabetes mellitus was defined by a physician's diagnosis or by need for treatment with insulin or oral hypoglycemic agents. Chronic obstructive pulmonary disease and dysthyroidism were defined by a physician's diagnosis as documented in the medical records. Abnormal kidney function was defined as the presence of equivalent serum creatinine ≥ 200 mmol/L or renal transplant or the presence of chronic dialysis. 'Abnormal liver function' was defined as chronic hepatic disease (e.g. cirrhosis) or biochemical evidence of significant hepatic derangement (e.g. bilirubin $>2\times$ upper limit of normal, in association with aspartate aminotransferase/alanine aminotransferase/alkaline phosphatase $>3\times$ upper limit normal, etc.). Also patients with other chronic diseases (e.g. immunity disorders and cancer) and eventually at short-term life expectancy were considered.

2.3. Management of patients and study protocol

All patients underwent evaluation with medical history, physical examination, blood tests, serial ECGs and plasma levels of Troponin I. Chest radiograph, blood gas analysis and echocardiogram were performed according to the clinical conditions. Patients with AF onset less than 48 h and without moderate to severe structural heart disease were considered eligible for rhythm control and received within the first-line 6-hour period i.v. bolus (in a few minutes) Flecainide (2 mg/kg), Propafenone (2 mg/kg), or Amiodarone (5 mg/kg), [9–13]. Patients who did not recover sinus rhythm could be submitted to a second dose of drugs previously given, eventually followed by oral administration up to 24 h of Flecainide (2 mg/kg plus 100 mg twice a day), Propafenone (2 mg/kg plus 150 mg three times a day), or Amiodarone (5 mg/kg plus 200 mg twice a day). After the first or second-line approach, patients could be submitted to early or elective electrical conversion (previous transesophageal echocardiography or after 3 weeks anticoagulation, respectively). After DC shock, patients were observed for a period of 6 h to promptly detect adverse events. No randomization was used and the therapeutic approach was at the discretion of the physician on duty [9].

2.4. Patients considered for enrollment in the study

Two groups of patients were considered. Group 1 included those patients presenting to the Emergency Department between January 2004 and December 2005 managed with standard approach; conversely group 2 included those patients presenting between January 2006 and December 2009 managed with aggressive treatment strategy in the Intensive Observation Unit and eventually additional Outpatient Clinic. Treatment strategy included intensive pharmacological treatment followed by electrical treatment as required. Hypertension, diabetes mellitus, comorbidities, AF duration (less or more than 48 h), quality of therapeutic approach and rhythm conversion or lack of rhythm control were analyzed.

2.4.1. Endpoint

Primary endpoint included rhythm conversion; secondary endpoints included modalities of rhythm conversion and reduction of admissions.

2.5. Statistical analysis

Summary data are expressed as mean \pm SD. Statistical comparisons of demographic and clinical features among the 2 groups were performed using χ^2 test for non parametric variables, as required. Analysis of variance was used for parametric variables. Univariate and multivariate logistic regression were used to define independent predictors of hospitalization. *P* values are 2 sided. A *p* value <0.05 was considered statistically significant. Calculations were performed with version 17, SPSS statistical package (SPSS Inc, Chicago, IL).

3. Results

From January 2004 to December 2009, 3475 AF patients presenting to the Emergency Department (1.1% of total visits per year) were considered for enrolment in the study. Those with hypertension (1739, 52%) and without moderate to severe structural heart disease were enrolled and stratified into two groups: group 1 (years 2004–2005) included 591 patients and group 2 (years 2006–2009) included 1148 patients. Baseline clinical characteristics of enrolled patients are shown in Table 1. In our series approximately two-third of patients presented AF lasting less than 48 h; one-half showed male gender and one-half comorbidities; mean age was 72 years. Due to the prospective observational design of the study, differences existed among clinical characteristics of the two groups of patients including male gender, comorbidities and duration of AF. Mean age was the same (Table 1). At univariate and multivariate analyses, AF lasting less than 48 h, absence of comorbidities and younger age were independent predictors of sinus rhythm (Table 2). Conversely, lack of sinus rhythm, older age, AF lasting more than 48 h, presence of comorbidities and diabetes were independent predictors of hospitalization (Table 3).

Overall, 1039 (60%) patients achieved sinus rhythm; we found a strong association between lower decades of age and conversion to sinus rhythm in patients with atrial fibrillation and hypertension enrolled in the study (Fig. 1). However, 55% patients achieved sinus rhythm in group 1 versus 62% in group 2 ($p=0.018$, Table 4). Spontaneous, electrical or pharmacological conversion account for 24%, 8% and 68% in group 1 versus 30%, 21% and 49% in group 2 ($p<0.001$, Table 4). Interestingly, out of 1071 patients with AF lasting less than 48 h, 950 (89%) achieved sinus rhythm as follows: 249 (26%) spontaneously, 160 (17%) by electrical conversion and 541 (57%) by intensive pharmacological approach (394 with class 1C and 147 with class III antiarrhythmic drugs, 73% and 27%, respectively, $p<0.001$, Fig. 3). In this subset of patients, spontaneous, electrical or pharmacological conversion account for 23%, 7% and 70% in group 1 versus 28%, 21% and 51% in group 2 ($p<0.001$). In both groups, patients more likely achieved sinus rhythm by class 1C as compared to class III antiarrhythmic drugs ($p<0.001$, Table 5). Overall, only 36% ($n=629$) patients need admission; 50% in group 1 versus 29% in group 2; thus reduction of admission accounts for up to 60% ($p<0.001$, Fig. 2). In our series, we found a strong association between higher decades of age and admission to hospital (Fig. 4).

Table 1

Baseline clinical characteristics of patients with atrial fibrillation and hypertension enrolled in the study ($n=1739$).

Clinical characteristics	Total ($n=1739$)	Group 1 ($n=591$)	<i>p</i>	Group 2 ($n=1148$)
Age (years s.d.)	72 11	72 \pm 11	0.478	73 \pm 11
Gender (male)	793 (46%)	247 (42%)	0.022	546 (48%)
Diabetes mellitus	202 (12%)	57 (10%)	0.069	145 (13%)
Comorbidities	747 (43%)	277 (47%)	0.019	470 (41%)
AF <48 h	1071 (62%)	320 (54%)	<0.001	751 (65%)
Sinus rhythm	1039 (60%)	330 (56%)	0.018	709 (62%)
Discharge	1110 (64%)	296 (50%)	<0.001	814 (71%)

Table 2
Association between clinical variables and sinus rhythm at univariate and multivariate analyses.

	Univariate analysis			Multivariate analysis		
	OR	Confidence interval 95%	p	OR	Confidence interval 95%	p
AF lasting <48 h	51.077	38,132–68,436	<0.001	47.890	35,626–64,377	<0.001
Comorbidities	1.911	1.573–2.322	<0.001	1.687	1.247–2.264	<0.001
Age	0.961	0.951–0.970	<0.001	0.962	0.969–0.996	<0.001
Diabetes mellitus	1.834	0.1366–2.463	<0.001	–	–	–
Male gender	0.960	0.792–1.164	0.680	–	–	–

4. Discussion

Detection of the underlying AF disorder even clinically ‘masked’ could result in implementing effective treatment and improving prognosis [6,8,14]. Suggestion resulted in reducing aggressive approach to AF patients. Indeed in the past decades clinical pathways pursued conversion to sinus rhythm, in patients presenting AF, when no structural heart disease was detected. At the same time this condition was considered to be associated essentially with isolated AF which accounted for almost 30% of all AF cases [1,6,15]. However, recent studies have demonstrated that lone AF is quite a rare disorder. Scientists from the Mayo Clinic of Olmsted County in the American state of Minnesota examined, between 1950 and 1980, 3623 patients with a first episode of AF and followed up them until 2003. Data revealed that only 2% of the total population of patients with AF really presented isolated AF with no concomitant heart disease, diabetes mellitus, hypertension or noncardiac disease that potentially could shorten life expectancy [7]. In the Framingham Heart Study, hypertension and diabetes were the sole cardiovascular risk factors to be predictive of AF after controlling for age and other predisposing conditions [16]. Thus, the role of hypertension as risk factor for AF is established; however, despite its leading importance as a highly prevalent and modifiable risk factor, only a few data are available regarding predictors and outcome of AF in hypertension. Extrapolations of results to uncomplicated hypertensive subjects with possible coexistence of LV hypertrophy could lead to improve aggressive treatment to pursue sinus rhythm [5,6,17].

The present study shows approximately two-third of patients presenting AF and hypertension achieved sinus rhythm when submitted to aggressive approach versus standard approach ($p = 0.018$, Table 2). In addition, four-fifth of patients with AF lasting less than 48 h achieved sinus rhythm without difference between groups; patients more likely achieved sinus rhythm by class 1C than by class III antiarrhythmic drugs ($p < 0.001$, Table 5). Strong independent predictors of sinus rhythm were AF lasting less than 48 h, absence of comorbidities (Table 2) and younger age (Fig. 1). Overall reduction of admission with aggressive versus standard approach accounts for 60%, 50% versus

Table 3
Association between clinical variables and hospital admission at univariate and multivariate analyses.

	Univariate analysis			Multivariate analysis		
	OR	Confidence interval 95%	p	OR	Confidence interval 95%	p
Sinus rhythm	8.993	7.186–11.253	<0.001	6.047	4.382–8.445	<0.001
Age	1.040	1.030–1.051	<0.001	1.022	1.010–1.033	<0.001
AF lasting <48 h	0.170	0.137–0.210	<0.001	0.694	0.502–0.960	<0.001
Comorbidities	0.435	0.356–0.531	<0.001	0.544	0.432–0.684	<0.001
Diabetes mellitus	0.489	0.364–0.657	<0.001	0.616	0.437–0.868	0.006
Male gender	1.093	0.898–1.330	0.376	–	–	–




Fig. 1. Association between decades of age and sinus rhythm in patients with atrial fibrillation and hypertension enrolled in the study ($n = 1739$).

29%, respectively ($p < 0.001$, Fig. 2). Independent predictors of hospitalization were lack of sinus rhythm, AF lasting more than 48 h, presence of comorbidities (Table 3) and older age (Fig. 4). Consistent with our findings, previous studies have reported that hypertension accounts for more than one-half in patients presenting AF, and aggressive approach could result in improving sinus rhythm and reducing admission. In fact, in the Manitoba Follow-up study, prevalence of hypertension was 53%, and the risk of AF in hypertensive subjects was increased by 1.4 times, while increased by 1.9 times in the Framingham Heart Study [18]. Because of its high prevalence in the population, hypertension independently accounts for more AF cases than any other risk factor [18,19]. Several other studies stated Observation Units as a rational choice for improving the utilization of health care resources [20], especially for successful treatment of chest pain [21], asthma [22], syncope [23] and eventually for treatment of AF, in order to reduce in-hospital stay [24]. Nonetheless no strategy to pursue normal sinus rhythm including antiarrhythmic drug therapy, conversion and ablation has definitely being shown to reduce the risk of stroke and hospitalization [25], in our setting physicians were strongly invited to consider pharmacological conversion as first line treatment [9]. Hence, among patients who recovered sinus rhythm, pharmacological approach was the most representative therapeutic strategy (55%). However, in our study we observed greater percentage of successful electrical treatment through the 6-year period (8% in group 1 versus 21% in group 2; $p < 0.001$). This fact related with electrical approach as rescue treatment when pharmacological treatment failed. Recently, an observational survey of management of AF in real life has shown therapeutic success was likely associated with rhythm-control. In this large survey enrolling 5171 patients therapeutic success was 54% overall (rhythm control 60% versus rate control 47%) a result driven by control of AF. Duration of AF, older age and heart failure predicted permanent AF

Table 4
Modalities of rhythm conversion in patients enrolled in the study. Class AAD, Vaughan-Williams Class of Anti Arrhythmic Drugs.

	Total n = 1739	Group 1 n = 591	p	Group 2 n = 1148
Sinus rhythm (n, %)	1039 (60%)	330 (56%)	0.018	709 (62%)
Spontaneous conversion (n, %)	289 (28%)	77 (24%)	0.028	212 (30%)
Electrical conversion (n, %)	175 (17%)	27 (8%)	<0.001	148 (21%)
Pharmacological conversion (n, %)	574 (55%)	225 (68%)	<0.001	349 (49%)
Class IC AAD (n, %)	412 (72%)	135 (60%)		277 (79%)
Class III AAD (n, %)	162 (28%)	90 (40%)		72 (21%)




Fig. 2. Patients enrolled in group 1 and 2 (black bars); patients admitted (gray bars). *p<0.001.

[4]. Prevalence of AF and hospitalizations for AF increased in recent years [26,27]; indeed, a report of the American Heart Association showed hospitalizations, with AF as the first-listed diagnosis, increased by 34% from 1996 to 2001 [25]. This trend appears related to the changing practice pattern with AF management [28] rather than changes over time in the distribution of age, sex, and comorbidities [27,29]. As a consequence, AF is a costly public health problem, with hospitalizations as the primary cost driver [30] especially in western countries [31,32]. To date, avoiding hospital admissions becomes mandatory. In our series, aggressive approach to rhythm conversion was planned to reduce admission to hospital. Primary part of the plan considered adherence to guidelines and appropriateness in administration of antiarrhythmic drugs (Fig. 2). Previous trials demonstrated antiarrhythmic drugs succeeded in reducing admissions [13,33] also in patients with structural heart disease [34]. In addition, electrical treatment was considered as rescue treatment after failure of pharmacological approach. Finally, in present study, a substantial portion of patients of group 2 with comorbidities or lack of rhythm control, in whom admission might be considered a real option, was evaluated in Outpatient Clinic for optimization of rate control or rhythm control in order to reduce frequency and severity of AF symptoms, avoiding current admission.

Table 5
Modalities of rhythm conversion in patients with atrial fibrillation lasting less than 48 h enrolled in the study. Class AAD, Vaughan–Williams Class of Anti-Arrhythmic Drugs.

	Total n = 1071	Group 1 n = 320	p	Group 2 n = 751
Sinus rhythm (n, %)	950 (89%)	280 (88%)	0.460	670 (89%)
Spontaneous conversion (n, %)	249 (26%)	64 (23%)	0.129	185 (28%)
Electrical conversion (n, %)	160 (17%)	20 (7%)	<0.001	140 (21%)
Pharmacological conversion (n, %)	541 (57%)	196 (70%)	<0.001	345 (51%)
Class IC AAD (n, %)	394 (73%)	120 (61%)		274 (79%)
Class III AAD (n, %)	147 (27%)	76 (39%)		71 (21%)

Consistent with this strategy a recent large in-hospital observational registry of AF patients showed that hospitalizations within 2 years after AF diagnosis were associated with increased AF symptom burden and with known limitations of contemporary drug therapy. In this analysis concerning a real world population, the authors suggested that interventions to reduce hospitalization should improve quality of life by reducing symptom frequency and severity rather than just preventing AF recurrences [35]. Although criteria for submission to the Outpatient Clinic were not standardized and at discretion of the emergency physician on duty, we consider this option a remarkable step to avoid current and further hospital admissions.

5. Conclusion

Aggressive pharmacological and electrical approach in patients presenting AF and hypertension significantly improved rhythm conversion overall up to two-third. In those patients presenting AF lasting less than 48 h sinus rhythm accounts for four-fifth. Large part of patients achieved sinus rhythm by pharmacological treatment strategy, mostly by class IC antiarrhythmic drugs. Admissions eventually reduced up to 60%.

Acknowledgement

The authors of this manuscript have certified that they comply with the Principles of Ethical Publishing in the International Journal of Cardiology [36].




Fig. 3. Diagram of management in patients presenting with atrial fibrillation lasting less than 48 h enrolled in the study. Class ADD, Vaughan–Williams Class of Anti-Arrhythmic Drugs.




Fig. 4. Association between decades of age and admission to hospital in patients with atrial fibrillation and hypertension enrolled in the study (n = 1739).

References

- [1] Roy D, Talajic M, Nattel S, et al. Rhythm control versus rate control for atrial fibrillation and heart failure. *N Engl J Med* 2008;358(25):2667–77.
- [2] Hohnloser SH, Kuck KH, Lillenthal J. Rhythm or rate control in atrial fibrillation—Pharmacological Intervention in Atrial Fibrillation (PIAF): a randomised trial. *Lancet* 2000;356(9244):1789–94.
- [3] Schaer BA, Schneider C, Jick SS, Conen D, Osswald S, Meier CR. Risk for incident atrial fibrillation in patients who receive antihypertensive drugs: a nested case-control study. *Ann Intern Med* 2010;152(2):78–84.
- [4] Camm AJ, Breithardt G, Crijns H, et al. Real-life observations of clinical outcomes with rhythm- and rate-control therapies for atrial fibrillation RECORDAF (Registry on Cardiac Rhythm Disorders Assessing the Control of Atrial Fibrillation). *J Am Coll Cardiol* 2011;58(5):493–501.
- [5] Mancia G, Laurent S, Agabiti-Rosei E, et al. Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. *Blood Press* 2009;18(6):308–47.
- [6] Camm AJ, Kirchhof P, Lip GY, et al. Guidelines for the management of atrial fibrillation: the task force for the management of atrial fibrillation of the European Society of Cardiology (ESC). *Eur Heart J* 2010;31(19):2369–429.
- [7] Jahangir A, Lee V, Friedman PA, et al. Long-term progression and outcomes with aging in patients with lone atrial fibrillation: a 30-year follow-up study. *Circulation* 2007;115(24):3050–6.
- [8] Chugh SS, Blackshear JL, Shen WK, Hammill SC, Gersh BJ. Epidemiology and natural history of atrial fibrillation: clinical implications. *J Am Coll Cardiol* 2001;37(2):371–8.
- [9] Wann LS, Curtis AB, January CT, et al. 2011 ACCF/AHA/HRS focused update on the management of patients with atrial fibrillation (Updating the 2006 Guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Heart Rhythm* 2011;8(1):157–76.
- [10] Naccarelli GV, Wolbrette DL, Bhatta L, et al. A review of clinical trials assessing the efficacy and safety of newer antiarrhythmic drugs in atrial fibrillation. *J Interv Card Electrophysiol* 2003;9(2):215–22.
- [11] Naccarelli GV, Wolbrette DL, Khan M, et al. Old and new antiarrhythmic drugs for converting and maintaining sinus rhythm in atrial fibrillation: comparative efficacy and results of trials. *Am J Cardiol* 2003;91(6A):15D–26D.
- [12] del AC, Martin A, Laguna P, Gargantilla P. Analysis of current management of atrial fibrillation in the acute setting: GEFAUR-1 study. *Ann Emerg Med* 2005;46(5):424–30.
- [13] Conti A, Del TB, Mariannini Y, et al. Management of patients with acute atrial fibrillation in the ED. *Am J Emerg Med* 2010;28(8):903–10.
- [14] Estes III NA, Sacco RL, Al-Khatib SM, et al. American heart association atrial fibrillation research summit: a conference report from the american heart association. *Circulation* 2011;124(3):363–72.
- [15] Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991;22(8):983–8.
- [16] Jouven X, Desnos M, Guerot C, Ducimetiere P. Idiopathic atrial fibrillation as a risk factor for mortality. The Paris Prospective Study I. *Eur Heart J* 1999;20(12):896–9.
- [17] Verdecchia P, Reboldi G, Gattobigio R, et al. Atrial fibrillation in hypertension: predictors and outcome. *Hypertension* 2003;41(2):218–23.
- [18] Kannel WB, Abbott RD, Savage DD, McNamara PM. Epidemiologic features of chronic atrial fibrillation: the Framingham study. *N Engl J Med* 1982;306(17):1018–22.
- [19] Kannel WB, Wolf PA, Benjamin EJ, Levy D. Prevalence, incidence, prognosis, and predisposing conditions for atrial fibrillation: population-based estimates. *Am J Cardiol* 1998;82(8A):2N–9N.
- [20] Roberts R, Graff LG. Economic issues in observation unit medicine. *Emerg Med Clin North Am* 2001;19(1):19–33.
- [21] Farkouh ME, Smars PA, Reeder GS, et al. A clinical trial of a chest-pain observation unit for patients with unstable angina. Chest Pain Evaluation in the Emergency Room (CHEER) Investigators. *N Engl J Med* 1998;339(26):1882–8.
- [22] Rydman RJ, Isola ML, Roberts RR, et al. Emergency Department Observation Unit versus hospital inpatient care for a chronic asthmatic population: a randomized trial of health status outcome and cost. *Med Care* 1998;36(4):599–609.
- [23] Smars PA, Decker WW, Shen WK. Syncope evaluation in the emergency department. *Curr Opin Cardiol* 2007;22(1):44–8.
- [24] Raghavan AV, Decker WW, Meloy TD. Management of atrial fibrillation in the emergency department. *Emerg Med Clin North Am* 2005;23(4):1127–39.
- [25] Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics—2011 update: a report from the American Heart Association. *Circulation* 2011;123(4):e18–209.
- [26] Wattigney WA, Mensah GA, Croft JB. Increasing trends in hospitalization for atrial fibrillation in the United States, 1985 through 1999: implications for primary prevention. *Circulation* 2003;108(6):711–6.
- [27] Friberg J, Buch P, Scharling H, Gadsbøhiol N, Jensen GB. Rising rates of hospital admissions for atrial fibrillation. *Epidemiology* 2003;14(6):666–72.
- [28] Miyasaka Y, Barnes ME, Gersh BJ, et al. Changing trends of hospital utilization in patients after their first episode of atrial fibrillation. *Am J Cardiol* 2008;102(5):568–72.
- [29] Humphries KH, Jackevicius C, Gong Y, et al. Population rates of hospitalization for atrial fibrillation/flutter in Canada. *Can J Cardiol* 2004;20(9):869–76.
- [30] Reynolds MR, Essebag V, Zimetbaum P, Cohen DJ. Healthcare resource utilization and costs associated with recurrent episodes of atrial fibrillation: the FRACTAL registry. *J Cardiovasc Electrophysiol* 2007;18(6):628–33.
- [31] Chugh SS, Blackshear JL, Shen WK, Hammill SC, Gersh BJ. Epidemiology and natural history of atrial fibrillation: clinical implications. *J Am Coll Cardiol* 2001;37(2):371–8.
- [32] Tsang TS, Petty GW, Barnes ME, et al. The prevalence of atrial fibrillation in incident stroke cases and matched population controls in Rochester, Minnesota: changes over three decades. *J Am Coll Cardiol* 2003;42(1):93–100.
- [33] Nichol G, McAlister F, Pham B, et al. Meta-analysis of randomised controlled trials of the effectiveness of antiarrhythmic agents at promoting sinus rhythm in patients with atrial fibrillation. *Heart* 2002;87(6):535–43.
- [34] Camm AJ, Capucci A, Hohnloser SH, et al. A randomized active-controlled study comparing the efficacy and safety of vernakalant to amiodarone in recent-onset atrial fibrillation. *J Am Coll Cardiol* 2011;57(3):313–21.
- [35] Reynolds MR, Morais E, Zimetbaum P. Impact of hospitalization on health-related quality of life in atrial fibrillation patients in Canada and the United States: results from an observational registry. *Am Heart J* 2010;160(4):752–8.
- [36] Shewan LG, Coats AJ. Ethics in the authorship and publishing of scientific articles. *Int J Cardiol* 2010;144:1–2.