



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

# FLORE

## Repository istituzionale dell'Università degli Studi di Firenze

### Free graft anoplasty

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Free graft anoplasty / Fucini C;Caminati F;Basiricò F;Bartolini N;Mirasolo VM. - In: COLORECTAL DISEASE. - ISSN 1463-1318. - ELETTRONICO. - 14:(2012), pp. 506-507.

*Availability:*

This version is available at: 2158/777998 since:

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

# Free graft foreskin anoplasty

C. Fucini, F. Caminati, F. Basiricò, N. Bartolini and V. M. Mirasolo

Department of Medical Surgical Critical Care, Careggi Hospital, Florence, Italy

Received 15 December 2011; accepted 28 December 2011; Accepted Article online 23 February 2012

## Abstract

**Aim** Reconstruction of a stenotic anal canal and repair of a stenotic perineal colostomy using a free graft foreskin.

**Method** The use of free graft foreskin anoplasty was described by Freeman for the treatment of mucosal prolapse in paediatric patients. The original surgical technique was modified and employed in two adult patients for the reconstruction of the anal region.

**Results** The graft, in both cases, took well with a satisfactory functional and morphological recovery of the anal canal.

**Conclusion** Free graft foreskin anoplasty, has proved to be an effective solution to stenosis in the anal canal following major local surgery.

**Keywords** Anoplasty, foreskin, graft, anal stenosis


## Indications

The use of a free foreskin graft in the reconstruction of the anal canal was introduced by Freeman in 1984 for the treatment of mucosal prolapse in paediatric patients [1]. Preserving the principle of the free graft, the original surgical technique was modified and employed for the reconstruction of a stenotic anal canal and repair of a stenotic perineal colostomy.


## Method

After a routine circumcision, the foreskin is turned inside, out exposing the mucosal surface on the outside (Fig. 1). After excision of the stenotic part of anal canal, the foreskin graft is positioned with the mucosal sheet coated on the inside of the neo-anal canal. The graft is sutured proximally to the mucosa of the residual anal canal and distally to the perineal skin (Fig. 2). Alternatively, the foreskin graft can be cut longitudinally and positioned with its skin surface on the outside and the mucosal surface on the inside to cover the excised stenotic scar (Fig. 3). The graft is anastomosed to the colonic mucosa and to the perineal skin.

The technique was first employed in a patient aged 58 years, with severe perineal *maladie de Verneuil* (hidradenitis suppurativa), undergoing wide excision includ-



**Figure 1** Routine circumcision and preparation of the foreskin graft.



**Figure 2** First case: positioning of the graft.

Correspondence to: N. Bartolini, Largo Brambilla 3, 50134, Florence, Italy. Cliniche Chirurgiche, AOU Careggi.  
E-mail: n.bartolini@gmail.com



**Figure 3** Second case: positioning of the graft after cutting longitudinally.



**Figure 4** First patient: the appearance of the anal canal.

ing the lower part of the anal canal. The wound was covered with a myocutaneous gracilis flap and a fasciocutaneous gluteal sliding flap. An anastomosis was made between the preserved upper anal canal and the skin of the myocutaneous gracilis flap. After 2 months, the anastomosis showed significant stenosis which was treated by the foreskin anoplasty technique (Fig. 2). The graft took nicely and after 5 years the anal canal showed a complete morphological and functional recovery (Fig. 4).

The second use of the technique was in a patient with anal cancer treated by abdominoperineal excision of the rectum with total anorectal reconstruction with a dynamic graciloplasty. A stenosis of the perineal colostomy occurred after 2 months due to ischaemia of the last 2 cm of the colonid neorectum. The patient underwent a foreskin anoplasty. Currently, after 1 year, the graft has taken well with satisfactory function.

## Discussion

Free graft foreskin anoplasty, after routine circumcision, has proved to be an effective solution to stenosis in the anal canal following major local surgery. The operation is indicated when the loss of anorectal mucosa is extensive and the other reconstruction techniques are not suitable.

## Reference

- 1 Freeman NV. The foreskin anoplasty. *Dis Colon Rectum* 1984; 27: 309–13.